

QUANTITATIVE POLYNOMIAL APPROXIMATION ON CERTAIN PLANAR SETS⁽¹⁾

BY
D. J. NEWMAN AND L. RAYMON

I. Let Q be a compact space in E^k . The *massivity*, $m_n(Q)$, is a sequence defined as follows: Let X_n be a set of $n+1$ elements of Q ; then

$$m_n(Q) = \max_{X_n \subset Q} \min_{x_i, x_j \in X_n; i \neq j} |x_i - x_j|.$$

Note that m_1 is what is normally defined to be the diameter of the space. Also note that m_n decreases to zero monotonely as $n \rightarrow \infty$. E.g., if Q is a set of measure $m > 0$, $m_n(Q)$ is asymptotic to m/n ; if Q is a k -dimensional cube, $m_n(Q)$ is asymptotic to $c/n^{1/k}$ [6, p. 21].

Q_1 will represent the interval $0 \leq x \leq 1$, and Q_2 the square $0 \leq x, y \leq 1$. Given a function $f(x)$ defined on Q and a $\delta > 0$, the *modulus of continuity* of the function $f(x)$, $\omega_f(\delta)$, is defined as follows:

$$\omega_f(\delta) = \sup_{|x-y| \leq \delta} |f(x) - f(y)|.$$

For $f(x)$ in $L^2(Q_1)$, we continue $f(x)$ to have period 1, and we define

$$\omega_f(\delta) = \sup_{|t| \leq \delta} \left(\int_0^1 |f(x+t) - f(x)|^2 dx \right)^{1/2},$$

while for $f(x, y)$ in $L^2(Q_2)$ continued to have period 1 in x and in y

$$\omega_f(\delta) = \sup_{t^2 + s^2 \leq \delta^2} \left(\int_0^1 \int_0^1 |f(x+t, y+s) - f(x, y)|^2 dx dy \right)^{1/2}.$$

$\omega(\delta)$ is a nonnegative, nondecreasing continuous function of δ . $f(x)$ is continuous iff $\omega_f(0^+) = 0$. Also, $\omega(\delta)$ is sublinear, and hence $\omega_f(\delta) \geq c\delta$ for some constant c . If $\omega_f(\delta) \leq \delta$ for all $\delta > 0$, $f(x)$ is said to be a *contraction* on Q .

Let $C(Q)$ be the space of all real valued continuous functions on Q and let P be a finite dimensional subspace. Let K be the class of contractions on Q . We now introduce the *degree of approximation*

$$\rho_P = \sup_{f \in K} \inf_{p \in P} \|f - p\|.$$

Presented to the Society, January 24, 1968; received by the editors March 7, 1967 and, in revised form, August 2, 1967.

⁽¹⁾ This research was partially supported by the National Science Foundation, Grant GP-4391.

We have the following *general lower bound theorem* for the degree of approximation: If the dimension of P is n , then $\rho_P \geq m_n(Q)/2$ [4]. If for some Q there exists c such that $\rho \leq cm_n(Q)$ for all n , where P is the class of n th degree polynomials, we can then say that P *efficiently* approximates all contractions on Q . If for some family of compact sets Q_τ there exists c such that $\rho_P(Q) \leq cm_n(Q)$ for all Q in Q_τ and for all n , then P can be said to approximate contractions *efficiently on the family* Q_τ .

The classic upper bound theorem is that of Dunham Jackson which says, essentially, that polynomials approximate efficiently on Q_1 [3, p. 36]. More recently, further results in this direction have been obtained. Yu. A. Brudnji [2] and D. J. Newman simultaneously discovered that for P_n the class of n th degree polynomials on linear sets Q of positive measure, the corresponding degree of approximation, ρ_n , for n sufficiently large, satisfies the inequality $\rho_n \leq cm/n$, c independent of Q . Newman and H. S. Shapiro have shown that polynomials approximate efficiently on the family of cubes, spheres and balls of all dimensions [6]. On the other hand, polynomials are not efficient on the family of all rectangles in the plane (and, hence, the two-dimensional analog of the result for linear sets of positive measure cannot hold). In fact, there is even a (highly pathological) linear set (of measure zero) on which polynomials are not efficient.

An n th degree polynomial in more than one variable will refer to a polynomial in which the maximum degree in any single variable is less than n . Unless otherwise specified, P and the corresponding ρ will refer to n th degree polynomials. In this case, they will sometimes be denoted, respectively, P_n and ρ_n . (Note that P_n may have dimension greater than n .)

The term *normal curve* will be used for a compact continuous curve of finite length contained in Q_2 . In this article we seek estimates for $\rho_P(Q)$ for Q a normal curve.

II. LEMMA 1. *If Q is a normal curve, there exist $c_1, c_2 > 0$ such that $c_1/n \leq m_n(Q) \leq c_2/n$.*

Proof. The projection of Q on the x -axis (or the y -axis) must be a line segment $[a, b]$. Consider the points $\{a + k(b-a)/n, k=0, 1, \dots, n\}$. Invert back to $n+1$ corresponding points of Q to get the set X_n . The x -coordinate (or the y -coordinate) of the minimum distance between any two points of X_n is $(b-a)/n$, giving $m_n(Q) \geq (b-a)/n$. Also, $m_n(Q) \leq 2L/n$, where L is the length of Q . For, otherwise, there would be $n+1$ nonintersecting discs of radius L/n , with their centers being elements of Q . The portion of Q contained within these discs would have length $\geq 2L$. This contradiction establishes Lemma 1.

LEMMA 2. *On Q_2 , there exist $c_1, c_2 > 0$ such that $c_1/n < \rho_n < c_2/n$.*

This lemma is a special case of the more general theorem of Newman and Shapiro [6, Theorem 4, p. 212].

LEMMA 3. Let $f(x)$ be a function defined on $S \subset Q$, Q a compact set in E^n , where $f(x)$ has modulus of continuity $\omega(\delta)$. Then there is a function $f^*(x)$ defined on Q with the following properties:

(a) $f^*(x) = f(x)$, $x \in S$,

(b) $|f^*(x) - f^*(y)| \leq \omega(d(x, y))$, where $d(x, y)$ is the distance from x to y . (In other words, a function can be extended without changing its modulus of continuity.)

Proof. It is sufficient to show that f can be properly extended to one point \bar{x} not in S , for the result would then follow by transfinite induction.

For all x, y in S ,

$$\begin{aligned} f(x) - f(y) &\leq \omega(d(x, y)) \\ &\leq \omega(d(x, \bar{x}) + d(y, \bar{x})), \text{ by the triangle inequality and by the} \\ &\quad \text{fact that } \omega \text{ is nondecreasing,} \\ &\leq \omega(d(x, \bar{x})) + \omega(d(y, \bar{x})), \text{ by sublinearity of } \omega. \end{aligned}$$

Hence, for any x we can find α so that

$$f(x) - \omega(d(x, \bar{x})) \leq \alpha \leq f(x) + \omega(d(x, \bar{x})).$$

Defining $f^*(\bar{x})$ to be α , we have

$$|f(x) - f^*(\bar{x})| \leq \omega(d(x, \bar{x})). \quad \text{Q.E.D.}$$

COROLLARY. Let f be a contraction on $S \subset Q$, $Q \subset E^n$. Then f can be extended as a contraction on Q .

LEMMA 4. The binomials $x^i y^j$, $0 \leq i, j \leq n$, generate on the curve $\sum_{k=0}^N a_{km} x^k y^m = 0$ a vector space of dimension less than $3Nn$.

Proof. Let S be the space generated by the $x^i y^j$, $0 \leq i, j \leq n$ on the entire plane. Consider the space generated by the $x^i y^j$, $0 \leq i, j \leq n$ on the restriction to the given algebraic curve. A basis of the binomials for this space spans, on the plane, a subspace S_0 of S . Another subspace S_1 of S is generated by those binomials spanning S which are not included in the basis for S_0 . Note that the dimension of S is $(n+1)^2$. We must show that $\dim S_0 < 3Nn$. For $n < N$, the result is trivial. Now,

$$x^i y^j \sum_0^N a_{km} x^k y^m, \quad 0 \leq i, j \leq n - N,$$

are linearly independent elements of S_1 . Hence $\dim S_1 \geq (n - N)^2$. Thus

$$\dim S_0 = \dim S - \dim S_1 \leq (n+1)^2 - (n - N)^2 < 3nN, \quad n \geq N,$$

and the lemma is proved.

LEMMA 5. If Q is a normal curve, then there exist $c_1, c_2 > 0$ such that $c_1/n^2 < \rho_n < c_2/n$ for all n .

Proof. Since the dimension of the space spanned by n th degree polynomials on Q is $\leq n^2$, we have $\rho_n \geq m_{n^2}/2 \geq c_1/n^2$, by the general lower bound theorem and Lemma 1. To prove the right hand inequality, we can consider any contraction on Q to be extended, by the corollary to Lemma 3, as a contraction to Q_2 . Now $\rho_n(Q) \leq \rho_n(Q_2) \leq c_2/n^2$, by Lemma 2, completing the proof.

THEOREM 1. *If Q is the restriction of an algebraic curve*

$$\sum_{i=0; j=0}^N a_{ij}x^i y^j = 0$$

to Q_2 , then there exist $c_1, c_2 > 0$ such that $c_1/n \leq \rho_n(Q) \leq c_2/n$.

Proof. From Lemma 4, the general lower bound theorem and Lemma 1, $\rho_n \geq 1/6Nn$, while $\rho_n \leq c/n$ by Lemma 5.

For Q a normal curve, the trivial estimates of c_1/n and c_2/n^2 as upper and lower bounds for ρ_n have been established. If the dimension of n th degree polynomials on Q is equal to $O(n)$, then $c_1/n < \rho_n < c_2/n$ (as in the case where Q is an algebraic curve). If the dimension of n th degree polynomials on Q is of greater order than n , say of order n^2 , to what extent can the trivial estimates for ρ_n be improved? Theorem 2 and Theorem 5 provide a partial answer to this question. It is toward the establishment of these theorems that we conclude this section with several additional lemmata. In Theorem 1 ρ_n can be taken in the L^2 norm or the sup norm, where what is meant by $\|f\|_{L^2}$ on the normal curve $y = g(x)$ is $(\int_0^1 |f(x, g(x))|^2 dx)^{1/2}$. In all following results ρ_P will be considered in L^2 only.

We denote by C_P the set of all $\varphi(x)$ with $L^2[0, 1]$ norm one which are in the orthogonal complement of P .

LEMMA 6.

$$\rho_P = \sup_{\varphi \in C_P} \left\| \int_0^x \varphi(t) dt \right\|_{L^2}.$$

For proof, cf. [5, Lemma 2, p. 942].

We recall that $F(x)$ is said to be in the Paley-Wiener class for the upper half plane, PW , if

(a) F is analytic in the upper half plane, and

$$\int_{-\infty}^{\infty} |F(x + iy)|^2 dx < M \quad \text{for all } y > 0,$$

or, equivalently,

(b)
$$F(z) = \int_0^{\infty} e^{izx} \varphi(x) dx, \quad \varphi(x) \in L^2[0, \infty].$$

For proof of the equivalence of (a) and (b), and for the general Paley-Wiener theory, the reader is referred to [7, pp. 1-13].

LEMMA 7. Let $P = \{x^\lambda, \lambda \in \Lambda\}$, where $0 \in \Lambda, \lambda \geq 0$. Then

$$\rho_P^2(Q_1) = \sup_{F \in PW} \frac{\int_{-\infty}^{\infty} \frac{|F(x+i)|^2}{x^2+1/4} \prod_{\lambda \in \Lambda} \frac{x^2+(\lambda-1/2)^2}{x^2+(\lambda+3/2)^2} dx}{\int_{-\infty}^{\infty} |F(x)|^2 dx}$$

For proof, cf. [5, Lemma 3, p. 943].

It is convenient to adopt the following notation:

$$M(\Lambda) = \max_x \frac{1}{x^2+1/4} \prod_{\lambda \in \Lambda} \frac{x^2+(\lambda-1/2)^2}{x^2+(\lambda+3/2)^2}$$

LEMMA 8. For P defined as in Lemma 7, $\rho_P^2(Q_1) \leq M(\Lambda)$.

Since, by Parseval's Identity,

$$\int_{-\infty}^{\infty} |F(x+i)|^2 dx \leq \int_{-\infty}^{\infty} |F(x)|^2 dx, \quad \text{for all } F \in PW,$$

Lemma 8 is a corollary to Lemma 7.

LEMMA 9. If $|\sum_{k=0}^n a_k x^k| \leq 1$ whenever $0 \leq x \leq 1$, then

$$|a_k| \leq 2^{2k} \frac{n(n+k-1)!}{(n-k)!(2k)!}$$

Furthermore, these are the best possible bounds for $|a_k|$.

To derive this bound, one demonstrates that $\cos 2n(\arccos x^{1/2})$, which is equal to

$$\sum (-1)^k 2^{2k} \frac{n(n+k-1)!}{(n-k)!(2k)!} x^k,$$

is maximal for each coefficient. For proof, see [1, p. 30]. Note that the above upper bounds yield the estimate $|a_k| \leq 3^{2n}$ for all k .

LEMMA 10. If $|\sum_{k=0}^{n^2} a_k x^{k/n}| \leq 1, 0 < \delta \leq x \leq 1$, then

$$|a_k| \leq \left(\frac{3}{1-\delta}\right)^{2n^2}, \quad k = 0, 1, \dots, n^2.$$

Proof. Assume $|\sum_{k=0}^{n^2} a_k w^k| \leq 1, 0 < \delta \leq w \leq 1$. Let $y = (w-\delta)/(1-\delta)$. Then

$$\left| \sum_0^{n^2} a_k ((1-\delta)y + \delta)^k \right| = \left| \sum b_k y^k \right| \leq 1, \quad 0 \leq y \leq 1.$$

Then, by Lemma 9, $|b_k| \leq 3^{2n^2}$. This, in turn, gives

$$|a_k| < \left(\frac{3}{1-\delta}\right)^{2n^2}.$$

Letting $w = x^{1/n}$, the result follows.

LEMMA 11. *Let p, q, n be integers, $p < q \leq n$, with p, q relatively prime. The inequality $kq < pm_1 + qm_2 < (k+1)q$, m_1, m_2, k positive integers, has at least k solutions in m_1, m_2 whenever $k \leq q-1$. For $q-1 < k \leq n$, the inequality has $q-1$ solutions.*

Proof. The solutions in m_1, m_2 of $kq < pm_1 + qm_2 < (k+1)q$ are the same as the solutions in m_1, m_2 of $k < m_1p/q + m_2 < k+1$. m_1p/q is a nonintegral rational which is less than k for any m_1 in $\{1, 2, \dots, k\}$. Thus, letting $m_2 = 1 + [k - m_1p/q]$, there is exactly one solution for $m_1 = 1, 2, \dots, k$, hence at least k solutions.

Similarly, for $k \geq q$, $m_1p/q + 1 + [k - m_1p/q]$ are solutions for $m = 1, 2, \dots, q-1$, and the lemma is proved.

III. Our main theorems are Theorem 2 and Theorem 5.

THEOREM 2. *If Q is the set (x, e^x) , $0 \leq x \leq 1$, then there exist $c_1, c_2 > 0$ such that*

$$\frac{c_1}{n^{3/2}} \leq \max_{f \in k} \min_{p \in P_n} \|f(x, e^x) - p(x, e^x)\|_{L^2} \leq \frac{c_2}{n^{3/2}},$$

i.e., $c_1/n^{3/2} \leq \rho_n \leq c_2/n^{3/2}$ in the L^2 norm.

Let α be an irrational number. α will be called *of degree $f(n)$* if there exist p, n such that $|\alpha - p/n| < 1/f(n)$ for infinitely many n .

THEOREM 5. *Let α , $0 < \epsilon \leq \alpha \leq 1$ be an irrational of degree $n^4(3/(1-\delta))^{2n^2}$, $0 < \delta < 1$. Let Q be the set of points (x, x^α) , $\delta \leq x \leq 1$. Then there exists a subsequence ρ_{n_i} of ρ_n and constants $c_1, c_2 > 0$ such that, in the L^2 norm, $c_1/n_i^{3/2} \leq \rho_{n_i} \leq c_2/n_i^{3/2}$.*

Proof of Theorem 2. We are approximating with linear combinations of $x^k e^{mx}$, $k, m \leq n$. By Lemma 6,

$$\rho_n = \sup_{\varphi \in C_P} \left\| \int_0^x \varphi(t) dt \right\|, \quad \text{where } \int_0^1 \varphi(x) x^k e^{mx} dx = 0, \quad k, m \leq n,$$

or, letting $te = e^x$, where

$$(1) \quad \int_{1/e}^1 \varphi(t) (\log t)^k t^{m-1} dt = 0, \quad k, m \leq n.$$

Let

$$F(z) = \int_{1/e}^1 \varphi(t) t^{-(iz+1/2)} dt.$$

Letting $t = e^{-u}$, we note that $F \in PW$. Setting $k=0$ in (1), we get $F((m+\frac{1}{2})i) = 0$, $m \leq n-1$. Setting $k=1$ in (1) and integrating by parts, we get $F'((m+\frac{1}{2})i) = 0$, $m \leq n-1$. For general k , integration by parts k times yields

$$(2) \quad F^{(k)}((m+\frac{1}{2})i) = 0, \quad m \leq n-1.$$

(In each case the boundary terms drop out by the orthogonality of φ to $x^k e^{mx}$.)

Since $F(x) = \int_{-\infty}^{\infty} e^{tux} e^{-u/2} \varphi(e^{-u}) du$, Parseval's Identity yields

$$(3) \quad \frac{1}{2\pi} \int_{-\infty}^{\infty} |F(x)|^2 dx = \int_{-\infty}^{\infty} e^{-u} |\varphi(e^{-u})|^2 du = \int_0^1 |\varphi(t)|^2 dt = 1.$$

Integrating by parts,

$$F(z) = t^{-(iz+1/2)} \int_0^t \varphi(u) du + (iz+1/2) \int_{1/e}^1 t^{-(iz+3/2)} \int_0^t \varphi(u) du dt.$$

But φ is orthogonal to 1, hence

$$F(z) = (iz+1/2) \int_{1/e}^1 t^{-(iz+3/2)} \int_0^t \varphi(u) du dt.$$

Thus

$$\begin{aligned} \frac{F(x+i)}{ix+1/2} &= \int_{1/e}^1 t^{-(ix+1/2)} \int_0^t \varphi(u) du dt \\ &= \int_{-\infty}^{\infty} e^{tix-v/2} \int_0^{e^{-v}} \varphi(u) du dv. \end{aligned}$$

Hence, by Parseval's Identity,

$$(4) \quad \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{|F(x+i)|^2}{x^2+1/4} dx = \int_{-\infty}^{\infty} e^{-v} \left| \int_0^{e^{-v}} \varphi(u) du \right|^2 dv = \int_0^1 \left| \int_0^t \varphi(u) du \right|^2 dt.$$

It follows from (2) and the general Paley-Wiener theory that

$$G(z) = F(z) \prod_{m \leq n-1} \left(\frac{z+i(m+1/2)}{z-i(m+1/2)} \right)^n$$

is in *PW*. Further, since $|x+i(m+1/2)| = |x-i(m+1/2)|$, (3) gives

$$(5) \quad \frac{1}{2\pi} \int_{-\infty}^{\infty} |G(x)|^2 dx = 1,$$

and (4) becomes

$$(6) \quad \int_0^1 \left| \int_0^t \varphi(u) du \right|^2 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{|G(x+i)|^2}{x^2+1/4} \prod_{m \leq n-1} \left(\frac{x^2+(m-1/2)^2}{x^2+(m+3/2)^2} \right)^n dx.$$

From (5), (6) and Lemma 6 we get

$$(7) \quad \rho_n^2 = \sup_{G \in PW} \frac{\int_{-\infty}^{\infty} \frac{|G(x+i)|^2}{x^2+1/4} \prod_{m \leq n-1} \left(\frac{x^2+(m-1/2)^2}{x^2+(m+3/2)^2} \right)^n dx}{\int_{-\infty}^{\infty} |G(x)|^2 dx}.$$

But $\|G(x+i)\|_{L^2}^2 \leq \|G(x)\|_{L^2}^2$. Hence

$$\begin{aligned} \rho_n^2 &\leq \max_x \frac{1}{x^2+1/4} \prod_{m \leq n-1} \left(\frac{x^2+(m-1/2)^2}{x^2+(m+3/2)^2} \right)^n \\ &= \max_x \frac{(x^2+1/4)^{2n-1}}{(x^2+(n-1/2)^2)^n (x^2+(n+3/2)^2)^n}. \end{aligned}$$

We note that the function $x = x(n)$ at which the max is taken on must go to infinity with n , for, if not, the upper bound thus obtained for ρ_n would grossly violate the general lower bound theorem. This being the case,

$$\rho_n < c \max_x \frac{x^{2n-1}}{(x^2+n^2)^n}.$$

The max occurs at $x^2 = 2n^3 - n^2$, and the maximum value is $c(2n^3 - n^2)^{-1/2}(1 - 1/2n)^n$, which tends to $c/(2e)^{1/2}n^{3/2}$, giving the desired upper bound.

Choosing $G(z) = 1/(z + in^{3/2})$ we get from (7) that

$$\begin{aligned} \rho_n^2 &\geq \frac{\int_{-\infty}^{\infty} \frac{c}{x^2} \left(\frac{x^2}{x^2+n^2}\right)^{2n} \frac{dx}{x^2+n^3}}{\int_{-\infty}^{\infty} \frac{dx}{x^2+n^3}} \\ &\geq \frac{n^{3/2}}{\pi} \int_{n^{3/2}}^{2n^{3/2}} \frac{c'}{x^2} \left(\frac{x^2}{x^2+n^2}\right)^{2n} \frac{dx}{x^2+n^3} \\ &\geq \frac{c'}{\pi 2e^2 n^3} \int_1^2 \frac{du}{u^2+1} = \frac{c''}{n^3}. \end{aligned} \qquad \text{Q.E.D.}$$

Our interest will now, temporarily, be turned from linear combinations of x^k , $k=0, 1, \dots, n$ as approximating functions to approximation with linear combinations of $x^{k/q}$, $k=0, 1, 2, \dots, nq$, on the interval Q_1 . We refer to the latter space and the corresponding degree of approximation as P_{nq}^* and ρ_{nq}^* .

Since $P_n \subset P_{nq}^*$, it follows from Jackson's theorem that there is a c_1 such that $\rho_{nq}^* \leq c_1/nq$. By the general lower bound theorem, there is a c_2 such that $\rho_{nq}^* \geq c_2/n$. The following result, in addition to being of independent interest, will be used in the proof of Theorem 4.

THEOREM 3. *There exist constants $c_1, c_2 > 0$ such that $c_1/nq^{1/2} \leq \rho_{nq}^* \leq c_2/nq^{1/2}$.*

Proof. By Lemma 7,

$$(8) \quad \rho_{nq}^{*2} = \sup_{F \in PW} \frac{\int_{-\infty}^{\infty} \frac{|F(x+i)|^2}{x^2+1/4} \prod_{k=0}^{nq} \frac{x^2+(k/q-1/2)^2}{x^2+(k/q+3/2)^2} dx}{\int_{-\infty}^{\infty} |F(x)|^2 dx}$$

while, by Lemma 8,

$$(9) \quad \rho_{nq}^{*2} \leq M(\Lambda), \quad \text{where } \Lambda = \{k/q \mid k = 0, 1, \dots, nq\}.$$

But, by cancellation,

$$(10) \quad M(\Lambda) = \max_x (x^2+1/4)^{-1} \prod_{k=0}^{2q-1} (x^2+(k/q-1/2)^2) \prod_{j=qn-2q+1}^{qn} (x^2+(j/q+3/2)^2)^{-1}.$$

By (9) and by the general lower bound theorem, M cannot fall off faster than c/n^4 . The point $x = x(n, q)$ where the max is taken on must go to infinity with n or q ,

for, if not, (10) would yield $M = O(1/n^{2qn})$. We now have, from (10), that there exist positive constants k_1, k_2 such that

$$(11) \quad k_1 M \leq \frac{1}{x^2} \left(\frac{x^2}{x^2 + n^2} \right)^{2q} \leq k_2 M.$$

The max in (11) is determined to be at $x^2 = 2qn^2 - n^2$, yielding $M = O(1/n^2q)$. Thus, from (9) we have

$$(12) \quad \rho_{nq}^* = O(1/nq^{1/2}).$$

Choosing $F(z) = 1/(z + inq^{1/2})$, we have, from (8) and (11),

$$\begin{aligned} \rho_{nq}^{*2} &\geq \frac{\int_{-\infty}^{\infty} \frac{c}{x^2} \left(\frac{x^2}{x^2 + n^2} \right)^{2q} \frac{dx}{x^2 + n^2q}}{\int_{-\infty}^{\infty} \frac{dx}{x^2 + n^2q}} \\ &\geq \frac{nq^{1/2}}{\pi} \int_{nq^{1/2}}^{2nq^{1/2}} \frac{c}{x^2} \left(\frac{x^2}{x^2 + n^2} \right)^{2q} \frac{dx}{x^2 + n^2q} \\ &\geq \frac{c}{\pi nq^{1/2}} (1 + 1/q)^{-2q} \int \frac{dx}{x^2 + n^2q} \geq \frac{c'}{n^2q}, \end{aligned}$$

which, together with (12), proves the theorem.

Let Q be the curve $y = x^{p/q}$, p, q relatively prime, $\delta \leq x \leq 1$. Consider the family of all such Q . Since these curves are algebraic, Theorem 1 implies $c_1/n \leq \rho_n \leq c_2/n$. However, for each Q in the family, the constants, c_1, c_2 , although independent of n , might well depend on which particular curve from the family is under consideration, i.e., c_1 and/or c_2 may be nontrivial functions of p and/or q . We shall now concern ourselves with more precise estimates for ρ_n —estimates that will yield the order of magnitude as a function of which Q from the family is under consideration. The following result in this direction will be crucial to the proof of Theorem 5.

THEOREM 4. *Let Q be the set $(x, x^{p/q})$, $\delta \leq x \leq 1$, p, q relatively prime. Then there exist $c_1, c_2 > 0$ such that, in the L^2 norm, $c_1/nq^{1/2} \leq \rho_n \leq c_2/nq^{1/2}$ for n, p, q such that $n \geq q$.*

Proof. We note that for $y = x^{p/q}$, approximating with n th degree polynomials in x and y is the same as approximating with $x^{i+jp/q}$, $0 \leq i, j \leq n$.

Let $f^*(x) = f(x, x^{p/q})$ be defined for $\delta \leq x \leq 1$. Now

$$\|f(x, x^{p/q}) - p_n(x, x^{p/q})\| = \left\| f^*(x) - \sum a_{ij} x^{i+jp/q} \right\|.$$

Also, the slope of $f^*(x)$,

$$\frac{\|f^*(y) - f^*(x)\|}{d(x, y)} = \frac{\|f(y, y^{p/q}) - f(x, x^{p/q})\|}{d((x, x^{p/q}), (y, y^{p/q}))} \cdot \frac{d((x, x^{p/q}), (y, y^{p/q}))}{d(x, y)} \leq (1 + m^2)^{1/2},$$

where m is the maximum slope for $y = x^{p/q}$ on $[\delta, 1]$. Since $m < 1/\delta$, we have

$\|f^*(y) - f^*(x)\| \leq M|y - x|$, $M = (1 + 1/\delta^2)^{1/2}$. Thus $f^*(x)/M$ is in $K(\delta, 1)$. Hence the assertion of our theorem is equivalent to the existence of c_1, c_2 such that

$$(13) \quad \frac{c_1}{nq^{1/2}} \leq \max_{f \in K(\delta, 1)} \min_{\{a_{ij}\}} \left\| f(x) - \sum_{i,j=0}^n a_{ij}x^{i+jp/q} \right\| \leq \frac{c_2}{nq^{1/2}}.$$

First, the left inequality in (13) will be established. Define

$$\rho_{nq}^*(a, b) = \max_{f \in K(a, b)} \min_{\{a_k\}} \left\| f(x) - \sum_{k=0}^{nq} a_k x^{k/q} \right\|.$$

LEMMA 12. $\rho_{2nq}^*(\delta, 1) \geq (1 - \delta)\rho_{2nq}^*(0, 1)$.

Proof. With respect to $\rho_{2nq}^*(0, 1)$, consider the coefficients $\{b_k\}$ for the function, g , that maximizes. Then

$$\left\| g(x) - \sum_0^{2nq} b_k x^{k/q} \right\| = \rho_{2nq}^*(0, 1).$$

Let $f(x) = (1 - \delta)g((x - \delta)/(1 - \delta))$. $f(x)$ is in $K(\delta, 1)$. Now

$$\sum_0^{2nq} a_k x^{k/q} = \sum (1 - \delta)b_k((x - \delta)/(1 - \delta))^{k/q}$$

is the best approximating polynomial to $f(x)$ on $[\delta, 1]$. For, suppose the contrary, i.e.,

$$\left\| \sum c_k x^{k/q} - f(x) \right\| < \left\| \sum (1 - \delta)b_k((x - \delta)/(1 - \delta))^{k/q} - f(x) \right\|$$

on $[\delta, 1]$. Then, on $[0, 1]$,

$$\begin{aligned} \left\| \sum c_k((1 - \delta)x + \delta)^{k/q} - f((1 - \delta)x + \delta) \right\| &\leq \left\| \sum (1 - \delta)b_k x^{k/q} - f((1 - \delta)x + \delta) \right\| \\ &\leq \left\| \sum (1 - \delta)b_k x^{k/q} - g(x) \right\|, \end{aligned}$$

contrary to the assumption that the b_k minimized for g . Hence,

$$\begin{aligned} \rho_{2nq}^*(\delta, 1) &\geq \left\| \sum (1 - \delta)b_k((x - \delta)/(1 - \delta))^{k/q} - f(x) \right\| \quad \text{on } [\delta, 1] \\ &= (1 - \delta) \left\| \sum b_k((x - \delta)/(1 - \delta))^{k/q} - g((x - \delta)/(1 - \delta)) \right\| \quad \text{on } [\delta, 1] \\ &= (1 - \delta) \left\| \sum b_k x^{k/q} - g(x) \right\| \quad \text{on } [0, 1] \\ &= (1 - \delta)\rho_{2nq}^*(0, 1), \end{aligned}$$

and Lemma 12 is established. Now,

$$\max_{f \in K(\delta, 1)} \min_{\{a_{ij}\}} \left\| f(x) - \sum a_{ij}x^{i+jp/q} \right\| \geq \rho_{2nq}^*(\delta, 1) \geq (1 - \delta)\rho_{2nq}^*(0, 1) \geq \frac{c}{nq^{1/2}},$$

the last inequality following from Theorem 3.

We now prove the upper bound in (13). Let f be extended, as a contraction, to $[0, 1]$. Lemma 8 now reduces the proof of the desired inequality to showing that

$$M(\Lambda) < c/n^2q, \quad \text{where } \Lambda = \{i + jp/q, 0 \leq i, j \leq n\}.$$

Let $x = x(n, q)$ be the point x at which the max in $M(\Lambda)$ is taken on. As in the proof of Theorem 2, $x(n, q)$, for fixed q , must tend to infinity with n . Observe that $(x^2 + (\lambda - 1/2)^2)/(x^2 + (\lambda + 3/2)^2)$ is an increasing function of λ for $\lambda > 4(1 + x^2)^{1/2} - 2$, and decreasing for λ less than this quantity. Let q be fixed. We consider two possible cases:

Case I. $1/x(n, q) = O(1/n)$.

Then there is some c such that $(x^2 + (\lambda - 1/2)^2)/(x^2 + (\lambda + 3/2)^2)$ is decreasing for all $\lambda < cn$. Let Λ_0 be $\{\lambda \in \Lambda \mid \lambda < cn\}$. $M(\Lambda_0) \geq M(\Lambda)$. The facts that

$$(x^2 + (\lambda - 1/2)^2)/(x^2 + (\lambda + 3/2)^2) < 1,$$

that this quotient is a decreasing function of λ , and that, by Lemma 11, there are at least $\min(k, q - 1)$ members of Λ_0 between k and $k + 1$ for $k + 1 < cn$ imply that $M(\Lambda_1) > M(\Lambda_0)$, where

$$\Lambda_1 = \{i + j/q, 0 \leq i \leq [cn], 0 \leq j \leq \min(i, q)\}.$$

That is,

$$\Lambda_1 = \{0, \begin{matrix} 1, & 1 + 1/q, \\ 2, & 2 + 1/q, & 2 + 2/q, \\ 3, & 3 + 1/q, & 3 + 2/q, & 3 + 3/q, \\ 4, & 4 + 1/q, & 4 + 2/q, & 4 + 3/q, \\ \vdots & \vdots & \vdots & \vdots \\ q - 1, & q - 1 + 1/q, & q - 1 + 2/q, & \dots, & q - 1 + (q - 1)/q, \\ q, & q + 1/q, & q + 2/q, & \dots, & q + (q - 1)/q, \\ \vdots & \vdots & \vdots & & \vdots \\ [cn - 1], & [cn - 1] + 1/q, & [cn - 1] + 2/q, & \dots, & [cn - 1] + (q - 1)/q. \end{matrix}$$

By the cancellation of the terms in the product (all but the first two and last two entries in each column cancel) we have

$$M(\Lambda_1) = \max_x \frac{1}{x^2 + \frac{1}{4}} \prod_{k=0}^q \frac{x^2 + (k + k/q - 1/2)^2}{x^2 + ([cn] + k/q + 1/2)^2} \cdot \frac{x^2 + (k + k/q + 1/2)^2}{x^2 + ([cn] + k/q + 3/2)^2}$$

which is of the order of

$$\max_x \frac{1}{x^2} \left(\prod \left(1 - \frac{c^2 n^2 - k^2}{x^2 + c^2 n^2} \right) \right)^2 < \max_x \frac{1}{x^2} \exp \left(-2 \sum \frac{c^2 n^2 - k^2}{x^2 + c^2 n^2} \right),$$

or

$$\max_x \frac{1}{x^2} \exp \left(-\frac{2c^2 n^2 q - q^3/3}{x^2 + n^2} \right).$$

But the maximum (away from the origin) is taken at $x^2 = n^2(2cq^2 - 1) - q^3/3$, where we get $M(\Lambda_1) = O(1/qn^2)$, and the estimate is established for Case I.

Since f is a contraction, it follows from (15) that

$$(18) \quad \|f(x, x^{p/n}) - f(x, x^\alpha)\| \leq 1/\varepsilon n^4(3/(1-\delta))^{2n^2}.$$

Now,

$$\begin{aligned} \|p_n(x, x^\alpha) - p_n(x, x^{p/n})\| &\leq \max_{\delta \leq x \leq 1} |p_n(x, x^\alpha) - p_n(x, x^{p/n})| \\ &\leq \max_{\delta \leq x \leq 1} \sum_{k=0}^n \left| \left(\frac{d^k}{dy^k} p_n(x, y) \right)_{y=x^{p/n}} \right| |x^\alpha - x^{p/n}|^k, \end{aligned}$$

by Taylor's Theorem. But this is

$$< \frac{1}{\varepsilon n^4(3/(1-\delta))^{2n^2}} \sum \left| \frac{d^k}{dy^k} p_n(x, y) \right|_{y=x^{p/n}}$$

by (15). But, by Lemma 10, the coefficients of $p_n(x, x^{p/n})$ are bounded by $(3/(1-\delta))^{2n^2}$. Hence, we have

$$(19) \quad \|p_n(x, x^\alpha) - p_n(x, x^{p/n})\| < 1/\varepsilon n^2.$$

Substitution of (16), (18), and (19) in (17) yields the upper bound asserted in (14).

To establish the lower bound, we note that

$$\begin{aligned} \|p_n(x, x^\alpha) - f(x, x^\alpha)\| + \|p_n(x, x^\alpha) - p_n(x, x^{p/n})\| + \|f(x, x^{p/n}) - f(x, x^\alpha)\| \\ \geq \|p_n(x, x^{p/n}) - f(x, x^{p/n})\|, \end{aligned}$$

and hence that by (18), (19) and Theorem 4,

$$\|p_n(x, x^\alpha) - f(x, x^\alpha)\| \geq \frac{c_1}{n^{3/2}} - \frac{1}{n^4} - \frac{1}{n^2} \geq \frac{c_1}{2n^{3/2}}. \quad \text{Q.E.D.}$$

BIBLIOGRAPHY

1. S. Bernstein, *Leçons sur les propriétés extrémales et la meilleure approximation des fonctions analytiques d'une variable réelle*, Gauthier-Villars, Paris, 1926.
2. Yu. A. Brudnji, "Constructive characteristics of functions given on certain perfect sets on the real axis," pp. 122-126, in *Investigation on contemporary problems of the constructive theory of functions*, edited by V. I. Smirnov, Fizmatgiz, Moscow, 1961.
3. D. Jackson, *The theory of approximation*, Amer. Math. Soc. Colloq. Publ., Vol. 11, Amer. Math. Soc., Providence, R. I., 1930.
4. G. G. Lorentz, *Approximation of smooth functions*, Bull. Amer. Math. Soc. **66** (1960), 124-125.
5. D. J. Newman, *A Müntz-Jackson theorem*, Amer. J. Math. **87** (1965), 940-944.
6. D. J. Newman and H. S. Shapiro, "Jackson's theorem in higher dimensions," pp. 208-219, in *On approximation theory, proceedings of the conference at Oberwolfach*, 1963, Birkhäuser, Basel, 1964.
7. R. E. A. C. Paley and N. Wiener, *Fourier transforms in the complex domain*, pp. 1-13, Amer. Math. Soc. Colloq. Publ., Vol. 19, Amer. Math. Soc., Providence, R. I., 1934; reprint 1960.

YESHIVA UNIVERSITY,

NEW YORK, NEW YORK

TEMPLE UNIVERSITY,

PHILADELPHIA, PENNSYLVANIA