

PIERCING POINTS OF CRUMPLED CUBES

BY
L. D. LOVELAND

McMillan [20] proved that a free 2-sphere in S^3 can be pierced by a tame arc at each of its points. Since each complementary domain of a free 2-sphere is an open 3-cell [22], it seems natural to attempt to prove some theorem analogous to McMillan's with this weaker hypothesis. We show that a crumpled cube C in S^3 has at most one nonpiercing point if $\text{Int } C$ is an open 3-cell. A *crumpled cube* C is the union of a 2-sphere and one of its complementary domains in S^3 , and a point p of $\text{Bd } C$ is a *piercing point* of C if there is a homeomorphism h of C into S^3 such that $h(\text{Bd } C)$ can be pierced by a tame arc at $h(p)$. It follows that a 2-sphere S in S^3 has at most two points where S cannot be pierced by a tame arc if each component of $S^3 - S$ is an open 3-cell (Corollary 3). McMillan [21] obtained the same result independently using an entirely different approach. Our proof follows from Lemmas 1-3 and the main result of [8].

Other results follow from the methods used in the proofs of Lemmas 1-3. For example, if a Cantor set W lies in a 2-sphere S in S^3 such that each component of $S^3 - S$ is an open 3-cell, then W is tame (Theorem 3). Thus each Cantor set on a free 2-sphere is tame. We also show that a continuum F is cellular if F lies in the boundary S of a cellular 3-cell and F does not separate S (Theorem 4).

We also obtain two characterizations of piercing points of crumpled cubes. One that is useful in this paper is that a point p in the boundary S of a crumpled cube C is a piercing point of C if and only if Property $(*, p, \text{Int } C)$ is satisfied (Theorem 2). If F is a closed subset of S , $(*, F, \text{Int } C)$ is defined to mean that Bing's Side Approximation Theorem [6] can be applied relative to S and $\text{Int } C$ in such a way that the intersection with S of the polyhedral approximation to S lies in the union of a finite set of mutually disjoint small disks in $S - F$. A precise definition can be found in [11] or [15]. The other characterization of piercing points (see Corollary 4) follows as a consequence of this one and results from [17].

The references should be consulted for needed definitions.

I. Piercing points of the closure of an open 3-cell.

LEMMA 1. *If $\varepsilon > 0$ and p and q are points in the boundary S of a crumpled cube C in S^3 , then there exist a crumpled cube M , an ε -homeomorphism h taking S onto $\text{Bd } M$, and a Sierpinski curve X such that*

- (1) p and q are inaccessible points of X ,
- (2) $X \subset S \cap \text{Bd } M$,

Received by the editors September 12, 1967.

- (3) $\text{Bd } M$ is locally tame modulo $\{p, q\}$,
- (4) M lies in an ε -neighborhood of C ,
- (5) h is the identity on X , and
- (6) each component of $S - X$ has diameter less than ε .

Proof. We shall use the technique introduced by Martin [18]. Let D be a disk on S such that $p \in \text{Int } D$, $q \in S - D$, and $\text{Bd } D$ is tame [4], and let J_1, J_2, \dots be a sequence of tame simple closed curves in D such that $\text{Bd } J_1 = \text{Bd } D$ and if D_1, D_2, \dots are the disks on D bounded by J_1, J_2, \dots , respectively, then $D_{i+1} \subset D_i$ and $p \in \bigcap D_i$. Using repeatedly the results from [6] and [11] together with the techniques of [4], we obtain a collection of tame annuli A_1, A_2, \dots such that

- $\text{Bd } A_i = \text{Bd } D_i \cup \text{Bd } D_{i+1}$,
- $A_i \cap D_i$ contains a Sierpinski curve X_i in $D_i - \text{Int } D_{i+1}$,
- $A_j \cap \text{Int } A_i = \emptyset$ if $i \neq j$,
- $\text{Cl}(\bigcup A_i)$ is a disk E with boundary J_1 ,
- $\text{Bd } M = (S - D) \cup E$, and each component of $A_i - X_i$ has diameter less than ε/i .

The procedure for obtaining the annuli A_i is given roughly by Martin in [18], so we do not pursue the details.

The same procedure as outlined above relative to q and the disk $S - \text{Int } D$ yields a sequence of Sierpinski curves Y_i such that $\{p, q\} \cup (\bigcup X_i) \cup (\bigcup Y_i)$ is the Sierpinski curve X required in Lemma 1.

LEMMA 2. *If the closure of $S^3 - C$ is a 3-cell, then, in addition to the conditions in Lemma 1, M and X can be selected such that*

- (7) $C \subset M$,
- (8) $\text{Bd } M \cap S = X$, and
- (9) the closure of each component of $M - C$ is a 3-cell.

Proof. We use Lemma 1 to obtain a Sierpinski curve X such that X contains p and q inaccessibly, X is locally tame modulo $\{p, q\}$, and each component of $S - X$ is small. Since S is tame from $S^3 - C$ we can obtain a 2-sphere $\text{Bd } M$ by pushing each component of $S - X$ slightly into $S^3 - C$. Since $\text{Bd } M$ is locally tame modulo X , we see that $\text{Bd } M$ is locally tame modulo $\{p, q\}$ [5]. If we identify M as the crumpled cube containing C and bounded by $\text{Bd } M$, conditions (7) and (8) follow. If Z is the closure of a component of $M - C$, then $\text{Bd } Z$ is a 2-sphere that is locally tame from $\text{Int } Z$ modulo a tame simple closed curve in X . It follows that $\text{Bd } Z$ is tame from $\text{Int } Z$ (see [11, Theorem 2] and [15, Theorem 14]), so Z is a 3-cell.

LEMMA 3. *Suppose K is a crumpled cube in S^3 such that $S^3 - K$ is an open 3-cell. If K_1, K_2, \dots is a sequence of mutually disjoint 3-cells in K such that, for each i , $K_i \cap \text{Bd } K$ is a disk D_i , then $S^3 - \text{Cl}(K - \bigcup_1^\infty K_i)$ is an open 3-cell.*

Proof. For each integer n we let $M_n = \text{Cl}(K - \bigcup_{i=1}^n K_i)$ and $V_n = S^3 - M_n$. The theorem will follow from [7] once we show that each V_n is an open 3-cell or equivalently that each M_n is cellular. In the remainder of the proof we show that

$M = M_1$ is cellular. Since the K_i are disjoint the same procedure can be used inductively to show the cellularity of each M_n .

Let $\alpha > 0$ and let A be a polyhedral arc in $S^3 - K_1$ from a point k in $\text{Int } K - K_1$ to a point m in $S^3 - K$. Since K_1 is a 3-cell, there exist disks D and E such that $D \subset \text{Int } D_1$, $\text{Bd } D$ is tame, $\text{Bd } D = \text{Bd } E$, $E - \text{Bd } E$ lies in $\text{Int } K_1$, and E is homeomorphically within $\alpha/2$ of $\text{Bd } K_1 - \text{Int } D_1$. Using Bing's Side Approximation Theorem [6] relative to the open set $\text{Bd } K - D$ in $\text{Bd } K$, we obtain an annulus F such that $\text{Bd } D \subset \text{Bd } F$, $\text{Bd } F - \text{Bd } D \subset S^3 - K$, $F \cap (A \cup M \cup \text{Int } E \cup \text{Int } D) = \emptyset$, F lies within $\alpha/2$ of M , and F is locally polyhedral modulo $\text{Bd } D$. Without loss in generality we may assume that $G = F \cup E$ is a polyhedral disk near M [10].

Since $S^3 - K$ is an open 3-cell, there is a 2-sphere R within α of $\text{Bd } K$ such that R separates the two boundary components of F , R separates k from m , and R lies in $S^3 - K$. We assume that R is polyhedral [1] and that F and R are in general position. Now we assume that A and R are in general position (i.e., $R \cap A$ consists of a finite number of points where A pierces R), and we choose a component T of $R - F$ such that $T \cap A$ consists of an odd number of points. Then T is a disk with holes lying within α of M . We fill these holes with disks near G to obtain a 2-sphere W that lies within α of M . Since $A \cap W = A \cap T$, it follows that W separates k from m . This means that W bounds a 3-cell X such that $M \subset \text{Int } X$ and X lies within α of M .

THEOREM 1. *If C is a crumpled cube in S^3 such that $\text{Int } C$ is an open 3-cell, then C has at most one nonpiercing point.*

Proof. Since there is a homeomorphism h of C into S^3 such that the closure of $S^3 - h(C)$ is a 3-cell [12], [13] and p is a piercing point of C if and only if p is a piercing point of $h(C)$, we assume without loss in generality that the closure of $S^3 - C$ is a 3-cell K . Let p and q be two points of $\text{Bd } C$. We shall show that one of these two points must be a piercing point of C . It follows from Lemma 2 that there exists a 2-sphere S' , a Sierpinski curve X , and a crumpled cube M such that $M \subset K$, $\text{Bd } M = S'$, p and q are inaccessible points of X , $S' \cap \text{Bd } C = X$, S' is locally tame modulo $\{p, q\}$, and the closure of each component of $K - M$ is a 3-cell. From Lemma 3 we see that $S^3 - M$ is an open 3-cell. Since K is a 3-cell it follows that $\text{Bd } M$ is locally tame from $\text{Int } M$ at both p and q (in fact, M is a 3-cell). For more detail, see the proof of Lemma 5 to follow.

It follows from [8, Theorem 1] that $\text{Bd } M$ is also locally tame from $S^3 - M$ at one of the points p and q , say p . Then p lies in a tame arc in X , so p is a point at which $\text{Bd } C$ can be pierced by a tame arc [11]. This means that p is a piercing point of C .

Property $(*, p, \text{Int } C)$, which is used in the following theorem, was defined roughly in the introduction and can be found in either [11] or [15].

THEOREM 2. *A point p in the boundary of a crumpled cube C is a piercing point of C if and only if $(*, p, \text{Int } C)$ is satisfied.*

Proof. If p is a piercing point of C , there exists a homeomorphism h of C into S^3 such that $h(S)$ can be pierced by a tame arc at $h(p)$. According to Gillman [11] this means that $h(p)$ lies in a tame arc A in $h(S)$. Furthermore Gillman [11] proved that $(*, A, h(\text{Int } C))$ is satisfied since A is tame. Lister [14] showed that $(*, h^{-1}(A), \text{Int } C)$ follows since h is a homeomorphism. Of course this implies $(*, p, \text{Int } C)$.

The other half of the proof of Theorem 2 is merely a rearrangement of the same ideas.

The following result is a consequence of Theorem 2 and a result by Martin [19].

COROLLARY 1. *If p is a point in a 2-sphere S in S^3 and U and V are the components of $S^3 - S$, then either $(*, p, U)$ or $(*, p, V)$ is satisfied.*

COROLLARY 2. *If C and L are the crumpled cubes bounded by a 2-sphere S in S^3 and a point $p \in S$ is a piercing point of both C and L , then S can be pierced by a tame arc at p .*

Proof. It follows from Theorem 2 that both $(*, p, \text{Int } C)$ and $(*, p, \text{Int } L)$ are satisfied. This means that S can be pierced by a tame arc at p [11].

COROLLARY 3. *If each complementary domain of a 2-sphere S in S^3 is an open 3-cell, then S contains two points p and q such that S can be pierced by a tame arc at each point of $S - \{p, q\}$.*

COROLLARY 4. *A point p in the boundary of a crumpled cube C is a piercing point of C if and only if p lies in an arc A in $\text{Bd } C$ such that for each $\varepsilon > 0$ there is a positive number δ such that each unknotted simple closed curve that lies in $\text{Int } C$ and has diameter less than δ can be shrunk to a point in an ε -subset of $S^3 - A$.*

Proof. From Theorem 2 we see that $(*, p, \text{Int } C)$ is satisfied if p is a piercing point of C . Then p lies in an arc A satisfying the conditions of Corollary 4 (see Theorem 1 in [17] and the remark on p. 511 in [15]).

If there exists an arc A as in the statement of Theorem 2, then $(*, A, \text{Int } C)$ follows (see the remark prior to the statement of Theorem 2 in [17]). Thus it follows from Theorem 2 that p is a piercing point of C .

II. Certain Cantor sets are tame. Using the methods of §I, we show that Cantor set W is tame if W lies in a 2-sphere S such that each component of $S^3 - S$ is an open 3-cell.

LEMMA 4. *If $\varepsilon > 0$ and W is a closed 0-dimensional subset of a 2-sphere S in S^3 , then there exists a Sierpinski curve X in S such that*

- (1) W lies inaccessibly in X ,
- (2) X is locally tame modulo W , and
- (3) each component of $S - X$ has diameter less than ε .

Proof. The proof is much the same as that given for Lemma 1. We let B_1 be a finite collection of mutually disjoint disks $D_{11}, D_{12}, \dots, D_{1n_1}$ in S such that each

$\text{Bd } D_{1j}$ is tame, $W \subset \bigcap \text{Int } D_{1j}$, and $\text{diam } D_{1j} < 1$. We inductively define, for each positive integer n , a similar finite collection B_n of disjoint disks each of diameter less than $1/n$. If we denote the union of the disks in B_n by B_n^* , then we insist that B_n^* is in the interior of B_{n-1}^* and that each point of W is a component of $\bigcap_{n=1}^{\infty} B_n^*$.

Now we apply the procedure outlined in the proof of Lemma 1. First we obtain a tame Sierpinski curve X_1 in $S - \bigcup \text{Int } D_{1i}$ such that each component of $S - X_1$ has diameter less than ε . Then X_2 is a finite collection of tame Sierpinski curves each in the closure of a component of $B_1^* - B_2^*$ and having small holes. This process is continued so that, for each n , we obtain a finite collection X_n of tame Sierpinski curves in the closure of $B_n^* - B_{n+1}^*$. If we denote the union of the Sierpinski curves in X_n by X_n^* , then $(\bigcup X_n^*) \cup W$ is the desired Sierpinski curve X .

LEMMA 5. *If W is a closed 0-dimensional subset of the boundary S of a cellular 3-cell C in S^3 and X is a Sierpinski curve in S containing W such that X is locally tame modulo W , then there exists a cellular 3-cell M such that*

- (1) $\text{Bd } M$ is locally tame modulo a point of W ,
- (2) $X = \text{Bd } M \cap S$, and
- (3) $M \subset C$.

Proof. The proof of Lemma 2 shows how to construct a crumpled cube M satisfying conditions (1) through (3), and it follows from Lemma 3 that M is cellular. Since $M \subset C$ and $\text{Int } C$ is 1-ULC it is not difficult to show that for each $\varepsilon > 0$ there exists a $\delta > 0$ such that each simple closed curve lying in $\text{Int } M$ and having diameter less than δ can be shrunk to a point in an ε -subset of $\text{Int } C$. This implies Property $(A, X, \text{Int } M)$, defined in [15], which is equivalent to $(*, X, \text{Int } M)$ [15, Theorems 8–10]. Hence M is a 3-cell [15, Theorem 14]. It follows from [8] that $\text{Bd } M$ is locally tame modulo a point of W .

THEOREM 3. *If W is a closed 0-dimensional subset of a 2-sphere S in S^3 and each component of $S^3 - S$ is an open 3-cell, then W is tame.*

Proof. Let U and V be the components of $S^3 - S$, and let X be a Sierpinski curve satisfying the conditions of Lemma 4 relative to some $\varepsilon > 0$. There is a homeomorphism h of $S \cup V$ into S^3 such that $S^3 - h(V)$ is a cellular 3-cell [12], [13]. Since $h(X)$ is locally tame modulo $h(W)$ it follows from Lemma 5 that $h(X)$ is locally tame modulo a point $h(p) \in h(W)$. Each Sierpinski curve $h(Y)$ in $h(X) - h(p)$ satisfies $(*, h(Y), h(V))$ since $h(Y)$ is tame [11], so it follows from a result by Lister [14] that $(*, Y, V)$ is satisfied.

Applying the same argument, where f is a homeomorphism of $S \cup U$ into S^3 , we obtain a point $q \in W$ such that each Sierpinski curve Y in $X - \{p, q\}$ satisfies both $(*, Y, V)$ and $(*, Y, U)$. Then Y is tame [15]. This means that X is locally tame modulo two points, so W is locally tame modulo two points. A theorem proven by Bing [3] shows W to be tame.

COROLLARY 5. *If a closed 0-dimensional set W lies in the interior of a cellular disk D in S^3 , then W is tame.*

Proof. For each point $p \in W$ there exists a disk D_p in $\text{Int } D$ and a 2-sphere S_p such that $p \in \text{Int } D_p \subset D_p \subset S_p$ and S_p is locally tame modulo D_p [2, Theorem 5]. It follows from the fact that D_p is cellular [21] that each component of $S^3 - S_p$ is an open 3-cell [23]. This means that W is locally tame (Theorem 3), so W is tame [3].

COROLLARY 6. *If W is a closed 0-dimensional subset of a free 2-sphere S in S^3 , then W lies in a tame Sierpinski curve on S .*

Proof. The proof of Theorem 3 shows that W lies in a Sierpinski curve X in S such that X is locally tame modulo two points (each component of a free 2-sphere must be an open 3-cell [22]). Since these two points lie in tame arcs in S (see [20, Theorem 5] and [11, Theorem 6]), they each lie in tame arcs in X [11, Lemma 6.1]. Thus X is tame [10].

III. Other related results.

LEMMA 6. *If $\varepsilon > 0$ and F is a continuum in the boundary S of a crumpled cube C in S^3 such that F does not separate S , then there exists a null sequence of mutually disjoint ε -disks $\{E_i\}$ in $S - F$ and a crumpled cube M such that*

- (1) $F \cup (S - \bigcup \text{Int } E_i) \subset \text{Bd } M$,
- (2) $\text{Bd } M$ is locally tame modulo F , and
- (3) M lies in an ε -neighborhood of C .

Proof. Since F does not separate S there is a sequence $\{D_i\}$ of disks on S such that $F = \bigcap D_i$, $\text{Bd } D_i$ is tame, and $D_{i+1} \subset \text{Int } D_i$. Now we follow the procedure outlined in the proof of Lemma 1.

LEMMA 7. *If $\varepsilon > 0$ and F is a continuum in the boundary S of a cellular 3-cell C in S^3 such that F does not separate S , then there exists a cellular 3-cell M satisfying all the conditions of Lemma 6 and such that $M \subset C$.*

Proof. We use Lemma 6 to obtain the disks E_i in $S - F$. Then the construction of M is indicated in the proofs of Lemmas 2 and 3 where each E_i is replaced with a tame disk in C . Since $M \subset C$ and C is a 3-cell it is easy to see that M is also a 3-cell (see the proof of Lemma 5).

THEOREM 4. *If F is a continuum on the boundary S of a cellular 3-cell in S^3 such that F does not separate S , then F is cellular.*

Proof. It follows from Lemma 7 that there exists a cellular 3-cell M such that $\text{Bd } M$ is locally tame modulo F . Then Theorem 5.2 of [23] insures that F is cellular.

REMARK. The hypothesis that S is the boundary of a 3-cell in the previous theorem cannot be removed. Furthermore, Theorem 4 becomes false if we require

only that each component of $S^3 - S$ be an open 3-cell. For example, one can grow two of the "feelers" described in [9] into opposite complementary domains of a sphere and let F be any arc containing the two wild points of the resulting sphere. It then follows from [16] that F is not cellular. However, F ought to be cellular if each component of $S^3 - S$ is an open 3-cell and F contains at most one (of the two possible) points where S cannot be pierced by a tame arc. Theorem 5 is a special case of this conjecture.

THEOREM 5. *If S is a 2-sphere in S^3 that is locally tame modulo a 0-dimensional set, F is a subcontinuum of S that does not separate S , each component of $S^3 - S$ is an open 3-cell, and F contains at most one of the two possible points where S cannot be pierced by a tame arc, then F is cellular.*

Proof. Under the conditions of the hypothesis S can have at most two wild points [8] and F can contain at most one of these points. Then there exists a disk D on S and a point $p \in F$ such that $F \subset \text{Int } D$ and S is locally tame at each point of $D - p$. It follows from Corollary 1 and Theorem 14 of [15] that S is locally tame from one component V of $S^3 - S$ at p . If $C = S \cup V$, then C is a cellular crumpled cube. Now we are able to use the technique in the proofs of Lemmas 6 and 7 to obtain a cellular 3-cell M such that $M \subset C$, $D \subset \text{Bd } M$, and $\text{Bd } M$ is locally tame modulo D . Then $\text{Bd } M$ is locally tame modulo F [10], and Theorem 5 follows from Theorem 4.

REFERENCES

1. R. H. Bing, *Approximating surfaces with polyhedral ones*, Ann. of Math. **65** (1957), 456-483.
2. ———, *A surface is tame if its complement is 1-ULC*, Trans. Amer. Math. Soc. **101** (1961), 294-305.
3. ———, *Tame Cantor sets in E^3* , Pacific J. Math. **11** (1961), 435-446.
4. ———, *Each disk in E^3 contains a tame arc*, Amer. J. Math. **84** (1962), 583-590.
5. ———, *Pushing a 2-sphere into its complement*, Michigan Math. J. **11** (1964), 33-45.
6. ———, *Improving the side approximation theorem*, Trans. Amer. Math. Soc. **116** (1965), 511-525.
7. M. Brown, *The monotone union of open n -cells is an open n -cell*, Proc. Amer. Math. Soc. **12** (1961), 812-814.
8. C. E. Burgess, *Criteria for a 2-sphere in S^3 to be tame modulo two points*, Michigan Math. J. **14** (1967), 321-330.
9. R. H. Fox and E. Artin, *Some wild cells and spheres in three-dimensional spaces*, Ann. of Math. **49** (1948), 979-990.
10. P. H. Doyle and J. G. Hocking, *Some results on tame disks and spheres in E^3* , Proc. Amer. Math. Soc. **11** (1960), 832-836.
11. D. S. Gillman, *Side approximation, missing an arc*, Amer. J. Math. **85** (1963), 459-476.
12. N. Hosay, *The sum of a real cube and a crumpled cube is S^3* , Notices Amer. Math. Soc. **10** (1963), 668.
13. L. L. Lininger, *Some results on crumpled cubes*, Trans. Amer. Math. Soc. **118** (1965), 534-549.
14. F. M. Lister, *Simplifying the intersection of disks in Bing's Side Approximation Theorem*, Pacific J. Math. **22** (1967), 281-295.

15. L. D. Loveland, *Tame subsets of spheres in E^3* , Pacific J. Math. **19** (1966), 489–517.
16. ———, *Wild points of cellular subsets of 2-spheres in S^3* , Michigan Math. J. **14** (1967), 427–431.
17. ———, *Sufficient conditions for a closed set to lie on the boundary of a 3-cell*, Proc. Amer. Math. Soc. **19** (1968), 649–652.
18. J. Martin, *Tame arcs on disks*, Proc. Amer. Math. Soc. **16** (1965), 131–133.
19. ———, *The sum of two crumpled cubes*, Michigan Math. J. **13** (1966), 147–151.
20. D. R. McMillan, Jr., *Some topological properties of piercing points*, Pacific J. Math. **22** (1967), 313–322.
21. ———, *Piercing a disk along a cellular set*, Proc. Amer. Math. Soc. **19** (1968), 153–157.
22. C. D. Papakyriakopoulos, *On Dehn's lemma and the asphericity of knots*, Ann. of Math. **66** (1957), 1–26.
23. D. G. Stewart, *Cellular subsets of the 3-sphere*, Trans. Amer. Math. Soc. **114** (1965), 10–22.

UTAH STATE UNIVERSITY,
LOGAN, UTAH