

ON A LEMMA OF MILUTIN CONCERNING AVERAGING OPERATORS IN CONTINUOUS FUNCTION SPACES

BY
SEYMOUR Z. DITOR⁽¹⁾

Abstract. We show that any infinite compact Hausdorff space S is the continuous image of a totally disconnected compact Hausdorff space S' , having the same topological weight as S , by a map φ which admits a regular linear operator of averaging, i.e., a projection of norm one of $C(S')$ onto $\varphi^\circ C(S)$, where $\varphi^\circ: C(S) \rightarrow C(S')$ is the isometric embedding which takes $f \in C(S)$ into $f \circ \varphi$. A corollary of this theorem is that if S is an absolute extensor for totally disconnected spaces, the space S' can be taken to be the Cantor space $\{0, 1\}^{\mathfrak{m}}$, where \mathfrak{m} is the topological weight of S . This generalizes a result due to Milutin and Pełczyński. In addition, we show that for compact metric spaces S and T and any continuous surjection $\varphi: S \rightarrow T$, the operator $u: C(S) \rightarrow C(T)$ is a regular averaging operator for φ if and only if u has a representation $uf(t) = \int_0^1 f(\theta(t, x)) dx$ for a suitable function $\theta: T \times [0, 1] \rightarrow S$.

1. Introduction. In his proof that for uncountable compact metric spaces S , all Banach spaces $C(S)$ of continuous (scalar-valued) functions on S with the sup norm are linearly homeomorphic, Milutin [5], [6] establishes the following result:

MILUTIN'S LEMMA. *There is a continuous map φ of the Cantor set K onto the unit interval I for which there exists a projection of norm one of $C(K)$ onto $\varphi^\circ C(I)$, where $\varphi^\circ: C(I) \rightarrow C(K)$ is the isometric embedding which takes $f \in C(I)$ into its composition with φ , $\varphi^\circ(f) = f\varphi$.*

An equivalent formulation of the above is that there is a linear operator u of norm one from $C(K)$ to $C(I)$ with the property that $u(f\varphi) = f$ for all $f \in C(I)$, i.e. $u\varphi^\circ = \text{id}_{C(I)}$. Following Pełczyński [7], we call such an operator u a *regular linear operator of averaging for φ* . Using this crucial lemma, Pełczyński [7, Theorem 5.6] has shown that the same result holds if the unit interval is replaced by any compact

Received by the editors May 2, 1969, and, in revised form, November 19, 1969.

AMS Subject Classifications. Primary 4725, 4625.

Key Words and Phrases. Averaging operator, extension operator, linear exave, projection operator, regular operator, continuous function space, Milutin space, totally disconnected space, inverse and direct limits, integral representation of regular averaging and extension operators.

⁽¹⁾ This research was supported by NSF Grant GP-5138.

It is part of the author's Ph.D. Dissertation, *Linear operators of averaging and extension*, written at the University of California at Berkeley in 1968 under the direction of Professor W. G. Bade.

Copyright © 1970, American Mathematical Society

metric space. In fact, Pełczyński shows that even if S is the product of $m > \aleph_0$ compact metric spaces, there is a continuous map of the Cantor space D^m onto S which admits a regular linear operator of averaging, where D is the two-point space $\{0, 1\}$. Such a space S which is the continuous image of some Cantor space D^m by a map which admits a regular averaging operator is called a *Milutin space* (S , as for all topological spaces mentioned here, is assumed to be compact Hausdorff).

Here we shall prove the following generalization of Milutin's Lemma:

THEOREM 1. *Any infinite compact Hausdorff space is the continuous image of a totally disconnected perfect compact Hausdorff space of the same topological weight by a map which admits a regular linear operator of averaging.*

By the *topological weight* (or simply, *weight*) of a space S we mean, as usual, the smallest cardinal number which is the cardinality of a base for the topology of S . That Theorem 1 generalizes Milutin's Lemma (and Pełczyński's extension to compact metric spaces) follows from the well-known characterization (cf. [2, p. 99]) of the Cantor set as a totally disconnected perfect compact metric space and the fact that a compact Hausdorff space is metrizable if and only if its weight is $\leq \aleph_0$.

If S is an *absolute extensor for totally disconnected spaces* (i.e., if for any totally disconnected space X and closed subspace A , any continuous map from A to S can be extended to a continuous map from X to S), then an easy corollary of Theorem 1 is that S is a Milutin space. Since Michael [4] has shown that any compact metric space, and therefore any product of compact metric spaces, is an absolute extensor for totally disconnected spaces, this corollary includes the above mentioned result of Pełczyński.

The proof of Theorem 1 is in a sense almost "natural" and in the metric case provides a new proof of Milutin's Lemma. The proof of the lemma which is due to Milutin (cf. [7, Lemma 5.5]) depends on an ingenious construction of a map $\varphi: K \rightarrow I$ for which the regular averaging operator $u: C(K) \rightarrow C(I)$ can be defined by a formula, $uf(t) = \int_0^1 f(\theta(t, x)) dx$, for $f \in C(K)$, $t \in I$, where $\theta: I \times I \rightarrow K$ is a Lebesgue measurable function whose definition is tied to that for φ . In Theorem 2 we show that every regular averaging operator $u: C(S) \rightarrow C(T)$, for compact metric spaces S and T , has a representation $uf(t) = \int_0^1 f(\theta(t, x)) dx$ for a suitable function $\theta: T \times I \rightarrow S$, and conversely, every such representation defines a regular averaging operator.

2. Preliminaries. We need two technical lemmas. One of these (Lemma 1) appears in [7] but not quite in the form we require. However, as the proof given in [7] (a partition of unity argument) carries over verbatim to the more general formulation, we shall state this lemma without proof. As for the other result (Lemma 2), although half of it appears in [7] (cf. [8] also) we shall for the sake of completeness give the entire proof. Since both lemmas have application in a more

general situation and no further effort is required, we shall introduce Pelczyński's idea of a linear exave and state the lemmas in this more general setting.

DEFINITION. Let S, T be compact Hausdorff spaces and $\varphi: S \rightarrow T$ a continuous map. A bounded linear operator $u: C(S) \rightarrow C(T)$ is called a *linear exave* for φ if $\varphi^\circ u \varphi^\circ = \varphi^\circ$. If φ is onto (resp. one-to-one), then u is called a *linear operator of averaging* (resp. *extension*) for φ . u is called *regular* if $\|u\| = 1$ and $u(1_S) = 1_T$ (where 1_X is the constant function on X everywhere equal to one).

Note that if φ is onto (resp. one-to-one), then φ° is one-to-one (resp. onto) and $\varphi^\circ u \varphi^\circ = \varphi^\circ$ reduces to $u \varphi^\circ = \text{id}_{C(T)}$ (resp. $\varphi^\circ u = \text{id}_{C(S)}$). If $S \subset T$ and φ is the inclusion map, then $\varphi^\circ: C(T) \rightarrow C(S)$ is the restriction operator, $\varphi^\circ(g) = g|_S$, and for each $f \in C(S)$, $u(f)$ is indeed an extension of f to all of T , i.e. $u(f)|_S = \varphi^\circ u(f) = f$.

The connection with projections is immediate (cf. [7, §2]): averaging operators correspond to bounded projections onto closed selfadjoint subalgebras (φ admits a linear operator of averaging if and only if there is a bounded projection of $C(S)$ onto $\varphi^\circ C(T)$); extension operators correspond to bounded projections onto closed ideals (φ admits a linear operator of extension if and only if there is a bounded projection of $C(T)$ onto the null space of φ° , i.e., the ideal of functions which vanish on $\varphi(S)$).

LEMMA 1. *Given a continuous map $\varphi: S \rightarrow T$, where S, T are compact Hausdorff spaces, suppose that $\{T_\alpha\}_{\alpha \in A}$ is a family of closed subsets of T , that $\{U_\alpha\}_{\alpha \in A}$ is a family of open subsets of T which cover $\varphi(S)$, and that $U_\alpha \subset T_\alpha$ for all $\alpha \in A$. Suppose further that $\{S_\alpha\}_{\alpha \in A}$ is a family of closed subsets of S such that for all $\alpha \in A$, $U_\alpha \cap \varphi(S) \subset \varphi(S_\alpha) \subset T_\alpha$, and that each $\varphi|_{S_\alpha}: S_\alpha \rightarrow T_\alpha$ admits a (regular) linear exave. Then φ admits a (regular) linear exave.*

In what follows we require the idea of an inverse limit system. The basic facts we use can be found in [1, pp. 212–220].

LEMMA 2. *Let $(\{S_\beta\}, \{\pi_\alpha^\beta\})$ and $(\{T_\beta\}, \{\theta_\alpha^\beta\})$ be inverse limit systems of compact Hausdorff spaces and continuous maps over the directed set A with each π_α^β onto. Let $(S, \{\pi_\alpha\}) = \text{inv lim } (\{S_\beta\}, \{\pi_\alpha^\beta\})$ and $(T, \{\theta_\alpha\}) = \text{inv lim } (\{T_\beta\}, \{\theta_\alpha^\beta\})$.*

(i) *If for every $\alpha \in A$, $u_\alpha: C(S_\alpha) \rightarrow C(T_\alpha)$ is a bounded linear operator such that $u_\beta \pi_\alpha^\beta = \theta_\alpha^\beta u_\alpha$ for $\alpha \leq \beta$ and $\sup \{\|u_\alpha\|\}_{\alpha \in A} < \infty$, then there is a unique linear operator $u = \text{dir lim } u_\alpha: C(S) \rightarrow C(T)$ such that $u \pi_\alpha^\circ = \theta_\alpha^\circ u_\alpha$ for all $\alpha \in A$ and $\|u\| \leq \sup \{\|u_\alpha\|\}_{\alpha \in A}$; if each u_α is regular, then so is u .*

(ii) *If for every $\alpha \in A$, $\varphi_\alpha: S_\alpha \rightarrow T_\alpha$ is continuous with $\varphi_\alpha \pi_\alpha^\beta = \theta_\alpha^\beta \varphi_\beta$ for $\alpha \leq \beta$ and the u_α 's in (i) are linear exaves for the φ_α 's, then $u = \text{dir lim } u_\alpha$ is a linear exave for $\varphi = \text{inv lim } \varphi_\alpha: S \rightarrow T$. If each u_α is an operator of averaging (resp. extension), then so is u .*

Proof. (i) Let $M = \bigcup \{\pi_\alpha^\circ C(S_\alpha)\}_{\alpha \in A}$. If $\alpha \leq \gamma$, then $\pi_\alpha = \pi_\alpha^\gamma \pi_\gamma$ and $\pi_\alpha^\circ C(S_\alpha) = \pi_\gamma^\circ \pi_\alpha^\gamma C(S_\alpha) \subset \pi_\gamma^\circ C(S_\gamma)$. Since A is directed it follows easily that M is a selfadjoint subalgebra of $C(S)$ which contains the identity. Moreover, if $x \neq y$ in S , then

$\pi_\alpha(x) \neq \pi_\alpha(y)$ for some $\alpha \in A$ and hence, for some $h \in C(S_\alpha)$, $h\pi_\alpha(x) \neq h\pi_\alpha(y)$, i.e., $\pi_\alpha^\circ(h)(x) \neq \pi_\alpha^\circ(h)(y)$. Therefore M separates the points of S so that, by the Stone-Weierstrass theorem, M is dense in $C(S)$ and it suffices to define u on M . Since each π_α^β is onto it follows that each π_α is onto so that the π_α° 's are isometric embeddings. Our requirement that $u\pi_\alpha^\circ = \theta_\alpha^\circ u_\alpha$ for all $\alpha \in A$ then dictates that we define u on M by setting $u(f) = \theta_\alpha^\circ u_\alpha (\pi_\alpha^\circ)^{-1}(f)$ for $f \in \pi_\alpha^\circ C(S_\alpha)$. It is clear that, providing our definition makes sense, the unique extension of this u to all of $C(S)$ is the one and only operator which satisfies all our requirements. Suppose then that $f = \pi_\alpha^\circ(g) = \pi_\beta^\circ(h)$, where $g \in C(S_\alpha)$, $h \in C(S_\beta)$. Then for $\gamma \geq$ both α and β ,

$$\pi_\gamma^\circ \pi_\alpha^\gamma(g) = \pi_\alpha^\circ(g) = \pi_\beta^\circ(h) = \pi_\gamma^\circ \pi_\beta^\gamma(h)$$

so that $\pi_\alpha^\gamma(g) = \pi_\beta^\gamma(h)$. Hence

$$\begin{aligned} \theta_\alpha^\circ u_\alpha (\pi_\alpha^\circ)^{-1}(f) &= \theta_\gamma^\circ \theta_\alpha^\gamma u_\alpha(g) = \theta_\gamma^\circ u_\gamma \pi_\alpha^\gamma(g) \\ &= \theta_\gamma^\circ u_\gamma \pi_\beta^\gamma(h) = \theta_\gamma^\circ \theta_\beta^\gamma u_\beta(h) = \theta_\beta^\circ u_\beta (\pi_\beta^\circ)^{-1}(f), \end{aligned}$$

which shows that u is well-defined.

(ii) We must show that $\varphi^\circ u \varphi^\circ = \varphi^\circ$ and since $\bigcup \{\theta_\alpha^\circ C(T_\alpha)\}_{\alpha \in A}$ is dense in $C(T)$, it suffices to verify that $\varphi^\circ u \varphi^\circ \theta_\alpha^\circ = \varphi^\circ \theta_\alpha^\circ$ for all $\alpha \in A$. But

$$\varphi^\circ \theta_\alpha^\circ = (\theta_\alpha \varphi)^\circ = (\varphi_\alpha \pi_\alpha)^\circ = \pi_\alpha^\circ \varphi_\alpha^\circ$$

so that

$$\varphi^\circ u \varphi^\circ \theta_\alpha^\circ = \varphi^\circ u \pi_\alpha^\circ \varphi_\alpha^\circ = \varphi^\circ \theta_\alpha^\circ u_\alpha \varphi_\alpha^\circ = \pi_\alpha^\circ \varphi_\alpha^\circ u_\alpha \varphi_\alpha^\circ = \pi_\alpha^\circ \varphi_\alpha^\circ = \varphi^\circ \theta_\alpha^\circ.$$

The last assertion in (ii) merely says that $\text{inv lim } \varphi_\alpha$ is onto (resp. one-to-one) whenever each φ_α is onto (resp. one-to-one).

COROLLARY 1. *Let $A = \{\alpha : 0 \leq \alpha < \Omega\}$ be an interval of ordinals and let $(\{S_\beta\}, \{\pi_\alpha^\beta\})$ be an inverse limit system of compact Hausdorff spaces and continuous maps over A , where each $\pi_\alpha^{\alpha+1}$ is onto and admits a regular linear operator of averaging. Suppose further that for every limit ordinal $\gamma < \Omega$, $(S_\gamma, \{\pi_\alpha^\gamma\}) = \text{inv lim}_{\beta < \gamma} (\{S_\beta\}, \{\pi_\alpha^\beta\})$. Then, if $(S_\Omega, \{\pi_\alpha\}) = \text{inv lim}_{\beta < \Omega} (\{S_\beta\}, \{\pi_\alpha^\beta\})$, the map $\pi_0: S_\Omega \rightarrow S_0$ is onto and admits a regular averaging operator.*

Proof. Let $T_\alpha = S_0$, $\theta_\alpha^\beta = \text{id}_{S_0}$, $\varphi_\alpha = \pi_0^\alpha$ for $0 < \alpha \leq \beta$ and let $u_\alpha^{\alpha+1}$ be a regular averaging operator for $\pi_\alpha^{\alpha+1}$. Inductively define $u_\alpha: C(S_\alpha) \rightarrow C(T_\alpha)$ for $0 < \alpha < \Omega$ by: $u_1 = u_0^1$, $u_\alpha = u_{\alpha-1} u_{\alpha-1}^\alpha$ if α is not a limit ordinal, and $u_\alpha = \text{dir lim}_{\beta < \alpha} (u_\beta)$ if α is a limit ordinal. We note that

(a) if α is a limit ordinal we may identify $(S_0, \{\text{id}_{S_0}\}) = (T_\gamma, \{\theta_\alpha^\gamma\})$ with $\text{inv lim}_{\beta < \gamma} (\{T_\beta\}, \{\theta_\alpha^\beta\})$ and then $\pi_0^\gamma = \text{inv lim}_{\beta < \gamma} (\varphi_\beta)$,

(b) the composition of regular averaging operators of a finite sequence of maps is a regular averaging operator of the composition of the maps,

(c) if

$$(*) \quad u_\beta \text{ is well-defined and } u_\beta \pi_\alpha^{\beta \circ} = \theta_\alpha^{\beta \circ} u_\alpha = u_\alpha \text{ for } \alpha \leq \beta < \gamma,$$

then if γ is a limit ordinal, Lemma 2 implies that (*) holds for $\beta \leq \gamma$, whereas if γ is not a limit ordinal, then for $\alpha < \gamma$,

$$u_\gamma \pi_\alpha^{\gamma^\circ} = u_{\gamma-1} u_{\gamma-1}^{\gamma^\circ} \pi_{\gamma-1}^{\gamma^\circ} \pi_\alpha^{\gamma-1^\circ} = u_{\gamma-1} \pi_\alpha^{\gamma-1^\circ} = u_\alpha$$

so that again (*) holds for $\beta \leq \gamma$.

It is clear then that the resulting system is well-defined and satisfies the hypotheses of Lemma 2 so that $u_\Omega = \text{dir lim } u_\alpha$ is a regular averaging operator for $\text{inv lim } \varphi_\alpha = \pi_\Omega$.

3. Proof of Theorem 1. Let S be a compact Hausdorff space of weight $m \geq \aleph_0$. Choose a neighborhood base of the diagonal in $S \times S$ having cardinality m and consisting of open symmetric sets (see [3, Chapter 6] for the terminology and notation of this paragraph). Suppose B_0, B_1, B_2, \dots is a well-ordering of this base where we may assume that the index set A consists of all ordinals $< \Omega$, the first ordinal of cardinality m . Each member B_α of the base gives rise to an open cover $\{B_\alpha(s)\}_{s \in S}$ of S , where $B_\alpha(s) = \{s' \in S : (s, s') \in B_\alpha\}$. For each α choose a finite subcover \mathcal{U}_α of $\{B_\alpha(s)\}_{s \in S}$. The collection $\{\mathcal{U}_\alpha\}$ has the following property: if $s_1 \neq s_2$ in S , then for some α the closure of no member of \mathcal{U}_α contains both s_1 and s_2 ; for we can find B_{α_1} with $(s_1, s_2) \notin B_{\alpha_1}$ and then choose B_α so that $B_\alpha \circ B_{\alpha_1} \circ B_\alpha \circ B_{\alpha_1} \subset B_{\alpha_1}$, from which the assumption that for some $s \in S$ both s_1 and s_2 are in $(B_\alpha(s))^- \subset (B_\alpha \circ B_{\alpha_1})(s)$ gives that (s_1, s) and (s, s_2) are in $B_\alpha \circ B_{\alpha_1}$, hence that (s_1, s_2) is in B_{α_1} , a contradiction.

We proceed by induction to define an inverse limit system $(\{S_\beta\}, \{\pi_\alpha^\beta\})$ over A satisfying the hypotheses of Corollary 1.

Define \mathcal{V}_n for $0 \leq n < \omega$ as follows:

$$\mathcal{V}_0 = \{U^- : U \in \mathcal{U}_0\}, \quad \mathcal{V}_n = \{(U \cap V)^- : U \in \mathcal{U}_n, V \in \mathcal{V}_{n-1}\}, \quad n \geq 1.$$

We note that

- (i) each \mathcal{V}_n is a finite cover of S by closed sets whose interiors cover S ,
- (ii) \mathcal{V}_{n+1} refines \mathcal{V}_n , and
- (iii) each member V of \mathcal{V}_n is a union of members of \mathcal{V}_{n+1} whose interiors in the relative topology of V cover V .

Now set $S_0 = S$ and for $0 \leq n < \omega$ let C_n be the collection of all chains $c: V_n \subset V_{n-1} \subset \dots \subset V_0$, where $V_i \in \mathcal{V}_i, i=0, 1, \dots, n$. Given such a chain $c \in C_n$, let $V(c) = V_n$ and for $n \geq 1$ let $c^* \in C_{n-1}$ be the chain $V_{n-1} \subset \dots \subset V_0$. For each chain $c \in C_n$, let $S_{n+1,c} = V(c) \times \{c\}$, the topological product of $V(c)$ (as a subspace of S) and the one-point space $\{c\}$. Let S_{n+1} be the topological sum (or union) of all the spaces $S_{n+1,c}$ for $c \in C_n$. Then each S_{n+1} is a compact Hausdorff space of weight $\leq m$ and the sets $S_{n+1,c}$ are both closed and open (clopen) in S_{n+1} .

Define $\pi_0^1: S_1 \rightarrow S_0$ by $\pi_0^1(x, c) = x$, and for $n > 0, \pi_n^{n+1}: S_{n+1} \rightarrow S_n$ by $\pi_n^{n+1}(x, c) = (x, c^*)$. (Note that if $(x, c) \in S_{n+1,c}$ then $x \in V(c) \subset V(c^*)$ so that $(x, c^*) \in V(c^*) \times \{c^*\} = S_{n,c^*}$.) It is easily seen that each $\pi_n^{n+1}, n \geq 0$, is a continuous surjection

which by Lemma 1 admits a regular averaging operator (using (i), (ii), (iii) above and the fact that each $\pi_n^{n+1}|_{S_{n+1,c}}$ is a homeomorphism onto its image).

Now let $(S_\omega, \{\pi_n^\omega\}) = \text{inv lim}_{n < \omega} (\{S_n\}, \{\pi_k^n\})$, where $\pi_k^n = \pi_k^{k+1} \cdot \dots \cdot \pi_{n-2}^{n-1}$. Continue by applying the same process to S_ω as was applied to S , using now the sequence of open covers $(\pi_0^\omega)^{-1}(\mathcal{U}_{\omega+n})$, $n=0, 1, 2, \dots$, and get $S_{\omega+1} = (S_\omega)_1$, $S_{\omega+2} = (S_\omega)_2, \dots$ with maps $\pi_\omega^{\omega+1}, \pi_{\omega+1}^{\omega+2}, \dots$. Then define $(S_{\omega 2}, \{\pi_\alpha^{\omega 2}\}) = \text{inv lim}_{\beta < \omega 2} (\{S_\beta\}, \{\pi_\alpha^\beta\})$ and continue in the obvious manner.

It is clear that this process gives us a system satisfying the hypotheses of Corollary 1. Hence if we let $(S_\Omega, \{\pi_\alpha\}) = \text{inv lim}_{\beta < \Omega} (\{S_\beta\}, \{\pi_\alpha^\beta\})$, then $\pi_0: S_\Omega \rightarrow S_0 = S$ is a continuous surjection which admits a regular averaging operator. Since the continuous image of a compact Hausdorff space has weight \leq the weight of the space and since the product of m spaces of weight $\leq m$ has weight $\leq m$ for $m \geq \aleph_0$, it is clear that S_Ω has weight m . We show S_Ω is totally disconnected by showing that any two distinct points x, y in S_Ω can be separated by clopen sets:

Case 1. $\pi_0(x) = \pi_0(y)$. Since $x \neq y$, there is an $\alpha < \Omega$ such that $x_\alpha \neq y_\alpha$ (where we write x_α for $\pi_\alpha(x)$) and we may suppose that α is the first ordinal for which this is the case. Then α is not a limit ordinal (for a limit ordinal α , S_α is by definition $\text{inv lim}_{\beta < \alpha} (S_\beta)$, so that $x_\beta = y_\beta$ for all $\beta < \alpha$ implies that $x_\alpha = y_\alpha$). Hence $\alpha = \beta + n$, where β is either a limit ordinal or 0 and n is an integer ≥ 1 . It follows then that x_α and y_α are in different $(S_\beta)_{n,c}$'s in $S_\alpha = (S_\beta)_n$, for $\pi_{\alpha-1}^\alpha$ is one-to-one on each $(S_\beta)_{n,c}$ so that, if x_α and y_α were in the same $(S_\beta)_{n,c}$, we would have $\pi_{\alpha-1}^\alpha(x_\alpha) \neq \pi_{\alpha-1}^\alpha(y_\alpha)$, i.e. $x_{\alpha-1} \neq y_{\alpha-1}$, which contradicts our choice of α if $\alpha > 1$ and contradicts $\pi_0(x) = \pi_0(y)$ if $\alpha = 1$. Hence for some c , x and y are separated by the clopen set $\pi_\alpha^{-1}(S_\beta)_{n,c}$.

Case 2. $\pi_0(x) \neq \pi_0(y)$. Choose \mathcal{U}_α so that the closure of no member of \mathcal{U}_α contains both $\pi_0(x)$ and $\pi_0(y)$. Then if $\alpha + 1 = \beta + n$, where β is either a limit ordinal or 0 and n is an integer ≥ 1 , we must have $x_{\alpha+1}$ and $y_{\alpha+1}$ in different $(S_\beta)_{n,c}$'s in $S_{\alpha+1}$, for otherwise $x_\beta = \pi_\beta^{\alpha+1}(x_{\alpha+1})$ and $y_\beta = \pi_\beta^{\alpha+1}(y_{\alpha+1})$ would both belong to the closure of some member of $(\pi_0^\beta)^{-1}(\mathcal{U}_\alpha)$ and hence $\pi_0(x) = \pi_0^\beta(x_\beta)$ and $\pi_0(y) = \pi_0^\beta(y_\beta)$ would both belong to the closure of some member of \mathcal{U}_α . Hence for some c , x and y are separated by the clopen set $\pi_{\alpha+1}^{-1}(S_\beta)_{n,c}$.

Finally, if K is the Cantor set, $S' = S_\Omega \times K$ is a totally disconnected perfect compact Hausdorff space of weight m , and if $\theta: S_\Omega \rightarrow S'$ is any continuous section of the projection map $\pi: S' \rightarrow S_\Omega$, so that $\pi\theta$ is the identity on S_Ω , then $\theta^\circ: C(S') \rightarrow C(S_\Omega)$ is a regular averaging operator for π . Hence for any regular averaging operator u for π_0 , $u\theta^\circ$ is a regular averaging operator for the continuous surjection $\pi_0\pi: S' \rightarrow S$. This completes the proof.

REMARK. In the metric case the above proof can be simplified by taking the index set A to be the natural numbers and for each n in A taking \mathcal{U}_n to be any finite cover of S by open sets of diameter $\leq 1/n$.

Let us note before proceeding that if X is totally disconnected and of weight m , then X can be regarded as a closed subset of D^m . For we can find a base B for the

topology of X which has cardinality m and consists entirely of clopen sets and, letting k_U be the characteristic function of U , the map from X to $D^B = D^m$ which takes $x \in X$ into $(k_U(x))_{U \in B}$ is an embedding.

COROLLARY 2. *If S is an absolute extensor for totally disconnected spaces, then S is a Milutin space.*

Proof. Let S have weight m . Then by Theorem 1 and our remarks above, there is a closed subset S' of D^m and a continuous surjection $\varphi_1: S' \rightarrow S$ which admits a regular averaging operator u_1 . Let $\varphi: D^m \rightarrow S$ be a continuous extension of φ_1 and let $\theta: S' \rightarrow D^m$ be the inclusion map. Then $\varphi_1 = \varphi\theta$ and $\text{id}_{C(S)} = u_1\varphi_1^\circ = u_1\theta^\circ\varphi^\circ$ so that $u = u_1\theta^\circ$ is a regular averaging operator for φ .

4. A representation theorem for regular linear exaves. The theorem of this section is motivated by the explicit construction given by Milutin in the proof of his lemma. To give a general idea of what is involved we sketch this construction here (see [7, Lemma 5.5] for details). First, since the Cantor set K is homeomorphic to $K \times K$ (both are homeomorphic to D^{\aleph_0}), $K \times K$ can be used instead of K . Now let h be the Cantor map of K onto the unit interval I , i.e. $h(\sum_{n=1}^{\infty} 2\xi_n 3^{-n}) = \sum_{n=1}^{\infty} \xi_n 2^{-n}$, where $(\xi_n) \in D^{\aleph_0}$, and define $\eta: I \rightarrow K$ by $\eta(t) = \min h^{-1}(t)$. The map $\varphi: K \times K \rightarrow I$ is then defined by letting $\varphi(\xi, \zeta)$ be the nonnegative root of the equation $h(\zeta) = \lambda h(\xi) + \lambda^2(1 - h(\xi))$ and the operator $u: C(K \times K) \rightarrow C(I)$ is obtained by setting $uf(t) = \int_0^1 f(\theta(t, x)) dx$, where $\theta: I \times I \rightarrow K \times K$ is given by $\theta(t, x) = (\eta(x), \eta(tx + t^2(1 - x)))$. φ and u are the required map and operator. The following theorem shows that this kind of integral representation of u is a general phenomenon.

THEOREM 2. *If S and T are compact metric spaces, a necessary and sufficient condition that $u: C(S) \rightarrow C(T)$ be a regular linear exave for a continuous map $\varphi: S \rightarrow T$ is that for $f \in C(S)$, $t \in T$,*

$$uf(t) = \int_0^1 f(\theta(t, x)) dx,$$

where $\theta: T \times I \rightarrow S$ is a function having the following properties:

- (1) $\theta(t, x) \in \varphi^{-1}(t)$ for all $t \in \varphi(S)$, $x \in I$.
- (2) $\theta(t, \cdot): I \rightarrow S$ is continuous off a countable set N_t and is left continuous everywhere, for all $t \in T$.
- (3) $\theta(\cdot, x): T \rightarrow S$ is continuous at t for $x \notin N_t$, for all $t \in T$,

Proof. (a) If such a function θ is given, then for any $t \in T$, $\theta(t, \cdot)$ is Borel measurable and $\theta(\cdot, x)$ is continuous at t for almost all x (with respect to Lebesgue measure). Hence for any $f \in C(S)$ the function $f\theta(t, \cdot)$ is integrable and the definition $uf(t) = \int_0^1 f\theta(t, x) dx$ makes sense. Moreover, if the sequence $t_n \rightarrow t$ in T , then $\theta(t_n, x) \rightarrow \theta(t, x)$ for almost all x so that $f\theta(t_n, \cdot) \rightarrow f\theta(t, \cdot)$ almost everywhere

and, by Lebesgue's dominated convergence theorem, it follows that $uf(t_n) \rightarrow uf(t)$. Therefore $uf \in C(T)$. In addition, it is clear that u is linear, $\|u\| \leq 1$, and $u(1_S) = 1_T$, so that u is a regular linear exave. Finally, if $g \in C(T)$ and $s \in S$, then, by (1), $\varphi\theta(\varphi(s), x) = \varphi(s)$ for all x so that

$$\begin{aligned} \varphi^\circ u\varphi^\circ(g)(s) &= u(g\varphi)(\varphi(s)) = \int_0^1 g\varphi\theta(\varphi(s), x) dx \\ &= \int_0^1 g\varphi(s) dx = \varphi^\circ(g)(s), \end{aligned}$$

i.e. u is a linear exave for φ .

(b) To prove the converse we may assume that $S=K$, the Cantor set. For let us suppose that the theorem has already been proved when the domain space is the Cantor set and that we are given the map $\varphi: S \rightarrow T$ with regular linear exave $u: C(S) \rightarrow C(T)$. By Theorem 5.6 of [7] (or Theorem 1 above) there is a continuous surjection $\varphi_1: K \rightarrow S$ which admits a regular averaging operator $u_1: C(K) \rightarrow C(S)$. Since u_1 is an averaging operator, it follows easily (cf. Proposition 4.4 of [7]) that uu_1 is a regular linear exave for $\varphi\varphi_1: K \rightarrow T$. Hence there exists a function $\theta_1: T \times I \rightarrow K$ satisfying the conditions

$$(1)' \theta_1(t, x) \in (\varphi\varphi_1)^{-1}(t) \text{ for all } t \in \varphi\varphi_1(K) = \varphi(S), x \in I,$$

(2)' $\theta_1(t, \cdot): I \rightarrow K$ is continuous off a countable set N_t and is left continuous everywhere, for all $t \in T$,

(3)' $\theta_1(\cdot, x): T \rightarrow K$ is continuous at t for $x \notin N_t$, for all $t \in T$, and $uu_1(h)(t) = \int_0^1 h\theta_1(t, x) dx$ for all $h \in C(K)$, $t \in T$. If we now let

$$\theta = \varphi_1\theta_1: T \times I \rightarrow S,$$

then the conditions (1)', (2)', (3)' imply that θ satisfies (1), (2), and (3), and if $f \in C(S)$, $t \in T$,

$$\int_0^1 f\theta(t, x) dx = \int_0^1 f\varphi_1\theta_1(t, x) dx = uu_1(f\varphi_1)(t) = u(u_1\varphi_1^\circ f)(t) = uf(t).$$

Therefore, we may assume that $S=K$ and the problem is reduced to constructing the function θ for a continuous map $\varphi: K \rightarrow T$ with regular linear exave $u: C(K) \rightarrow C(T)$.

For $t \in T$, let μ_t be the continuous linear functional on $C(K)$ defined by $\mu_t(f) = uf(t)$. By the Riesz representation theorem, μ_t may be identified with a finite regular Borel measure on K . Since u is a regular linear exave for φ , it follows that, under this identification, μ_t is a nonnegative measure of total variation one which is concentrated on $\varphi^{-1}(t)$ if $t \in \varphi(K)$ (see [7, Proposition 4.1]). If $t_n \rightarrow t_0$ in T , then for any $f \in C(K)$, $\mu_{t_n}(f) \rightarrow \mu_{t_0}(f)$ and, in particular, $\mu_{t_n}(E) \rightarrow \mu_{t_0}(E)$ if E is a clopen subset of K . In what follows we shall use the standard notation $[\]$, $(\)$, etc. both for intervals in I and for intervals in K .

Now, for $t \in T$, let K_t denote the support of μ_t and let us define the functions

$$\begin{aligned} \delta_t: K_t &\rightarrow I & \text{by} & \quad \delta_t(\xi) = \mu_t[0, \xi], \\ \theta_t: I &\rightarrow K_t & \text{by} & \quad \theta_t(x) = \inf \delta_t^{-1}[x, 1], \text{ and} \\ \theta: T \times I &\rightarrow K & \text{by} & \quad \theta(t, x) = \theta_t(x). \end{aligned}$$

Since $K_t \subset \varphi^{-1}(t)$ for $t \in \varphi(K)$, θ satisfies condition (1).

Since μ_t is a nonnegative regular Borel measure, it is clear that

(i) δ_t is increasing and right continuous, for all $t \in T$,

and, by the right continuity of δ_t , it follows that

(ii) $\delta_t \theta_t(x) \geq x$ for all $t \in T, x \in I$.

Using (i) and (ii) we proceed to establish

(iii) θ_t is increasing and left continuous, for all $t \in T$.

If $x_1 \leq x_2$, then $\delta_t^{-1}[x_1, 1] \supset \delta_t^{-1}[x_2, 1]$ so that $\theta_t(x_1) = \inf \delta_t^{-1}[x_1, 1] \leq \inf \delta_t^{-1}[x_2, 1] = \theta_t(x_2)$. Hence θ_t is increasing. To show θ_t is left continuous we must show that if $x_n \uparrow x_0$, so that $\theta_t(x_n) \uparrow$ some $\xi_0 \in K_t$, then $\xi_0 = \theta_t(x_0) = \inf \delta_t^{-1}[x_0, 1]$. First, $\xi_0 \leq \inf \delta_t^{-1}[x_0, 1]$, for if $\xi \in \delta_t^{-1}[x_0, 1]$, then for all $n, x_n \leq x_0 \leq \delta_t(\xi)$, i.e. $\xi \in \delta_t^{-1}[x_n, 1]$, so that $\theta_t(x_n) \leq \xi$ for all n and hence $\xi_0 \leq \xi$. Therefore it suffices to show that $\xi_0 \in \delta_t^{-1}[x_0, 1]$ and this follows since, by (ii), $x_n \leq \delta_t \theta_t(x_n)$ for all n and hence $x_0 = \lim_n x_n \leq \lim_n \delta_t \theta_t(x_n) \leq \delta_t(\xi_0)$ (since $\theta_t(x_n) \uparrow \xi_0$ and δ_t is increasing), i.e. $\xi_0 \in \delta_t^{-1}[x_0, 1]$. This establishes (iii).

From (iii) it follows that θ satisfies condition (2).

Before verifying that condition (3) is satisfied we prove

(iv) if $x > 0$, then $\theta_t(x) \leq \xi$ if and only if $x \leq \mu_t[0, \xi]$.

For, if $\mu_t[0, \xi] \geq x > 0$, then $[0, \xi] \cap K_t \neq \emptyset$. Letting $\xi' = \sup ([0, \xi] \cap K_t)$, we have $\xi \geq \xi' \in K_t, \delta_t(\xi') = \mu_t[0, \xi'] = \mu_t[0, \xi] \geq x$, and hence, $\theta_t(x) = \inf \delta_t^{-1}[x, 1] \leq \xi' \leq \xi$. Conversely, if $\theta_t(x) \leq \xi$, then $x \leq \delta_t \theta_t(x) = \mu_t[0, \theta_t(x)] \leq \mu_t[0, \xi]$.

To prove that θ satisfies condition (3), we must show that if $t_0 \in T$ and x_0 is a point of continuity of $\theta(t_0, \cdot) = \theta_{t_0}$, then $\theta(\cdot, x_0)$ is continuous at t_0 . By compactness, it suffices to show that if $t_n \rightarrow t_0$ and $\theta_{t_n}(x_0) \rightarrow \xi_0$, then $\xi_0 = \theta_{t_0}(x_0)$. By including 0 and 1 in N_t , we can assume that $x_0 \neq 0$ or 1.

Case 1. $\theta_{t_0}(x_0) > \xi_0$. Then there is a ξ in K such that $\xi_0 \leq \xi < \theta_{t_0}(x_0)$ and $[0, \xi]$ is clopen. Hence, for large $n, \theta_{t_n}(x_0) \leq \xi$ so that

$$x_0 \leq \delta_{t_n} \theta_{t_n}(x_0) = \mu_{t_n}[0, \theta_{t_n}(x_0)] \leq \mu_{t_n}[0, \xi] \rightarrow \mu_{t_0}[0, \xi]$$

and therefore $x_0 \leq \mu_{t_0}[0, \xi]$. But, by (iv), this implies that $\theta_{t_0}(x_0) \leq \xi$, a contradiction.

Case 2. $\theta_{t_0}(x_0) < \xi_0$. Then there is a ξ in K such that $\theta_{t_0}(x_0) \leq \xi < \xi_0$ and $[0, \xi]$ is clopen. Since x_0 is a point of continuity of θ_{t_0} and $x_0 < 1$, there is an $x_1 > x_0$ such that $\theta_{t_0}(x_1) \leq \xi$, and since $\theta_{t_n}(x_0) \rightarrow \xi_0$, we eventually have $\xi < \theta_{t_n}(x_0)$ which, by (iv) gives $\mu_{t_n}[0, \xi] < x_0$. But this is impossible since we would then have

$$x_1 \leq \delta_{t_0} \theta_{t_0}(x_1) \leq \mu_{t_0}[0, \xi] = \lim_n \mu_{t_n}[0, \xi] \leq x_0.$$

Therefore $\theta_{t_0}(x_0) = \xi_0$ and θ satisfies condition (3).

We must now verify that for all $f \in C(K)$, $t \in T$,

$$uf(t) = \mu_t(f) = \int_0^1 f\theta(t, x) dx.$$

However, the topology of K has a base consisting of clopen sets of the form $(\xi_1, \xi_2]$ and $[0, \xi]$, and the subspace of $C(K)$ generated by the characteristic functions k_V of such sets V is dense in $C(K)$. Since $k_{(\xi_1, \xi_2]} = k_{[0, \xi_2]} - k_{[0, \xi_1]}$, this subspace is also generated by the characteristic functions of the clopen sets $[0, \xi]$. Moreover, both u and the function $f \rightarrow \int_0^1 f\theta(\cdot, x) dx$ are bounded linear transformations from $C(K)$ to $C(T)$ (see part (a) of this proof). Hence it suffices to verify that for $t \in T$, $\xi \in K$,

$$\mu_t[0, \xi] = \int_0^1 k_{[0, \xi]}(\theta(t, x)) dx.$$

But, if $x > 0$ and we let $A = [0, \xi] \subset K$ and $B = [0, \mu_t A] \subset I$, then, by (iv) above, we have $k_A \theta(t, x) = k_B(x)$, so that

$$\int_0^1 k_A \theta(t, x) dx = \int_0^1 k_B(x) dx = \mu_t[0, \xi].$$

This completes the proof.

REFERENCES

1. S. Eilenberg and N. E. Steenrod, *Foundations of algebraic topology*, Princeton Univ. Press, Princeton, N. J., 1952. MR 14, 398.
2. J. G. Hocking and G. S. Young, *Topology*, Addison-Wesley, Reading, Mass., 1961. MR 23 #A2857.
3. J. L. Kelley, *General topology*, Van Nostrand, Princeton, N. J., 1955. MR 16, 1136.
4. E. Michael, *Selected selection theorems*, Amer. Math. Monthly 63 (1956), 233–238.
5. A. A. Milutin, *On spaces of continuous functions*, Dissertation, Moscow State University, Moscow, 1952. (Russian)
6. ———, *Isomorphism of spaces of continuous functions on compacta having the power of the continuum*, Teor. Funkcii Funkcional. Anal. i Priložen 2 (1966), 150–156. (Russian)
7. A. Pelczyński, *Linear extensions, linear averagings, and their applications to linear topological classification of spaces of continuous functions*, Dissertationes Math. Rozprawy Mat. 58 (1968). MR 37 #3335.
8. Z. Semadeni, *Inverse limits of compact spaces and direct limits of spaces of continuous functions*, Studia Math. 31 (1968), 373–382.

LOUISIANA STATE UNIVERSITY,
BATON ROUGE, LOUISIANA 70803