

STIEFEL-WHITNEY NUMBERS OF QUATERNIONIC AND RELATED MANIFOLDS

BY
 E. E. FLOYD

Abstract. There is considered the image of the symplectic cobordism ring Ω_*^{Sp} in the unoriented cobordism ring N_* . A polynomial subalgebra of N_* is exhibited, with all generators in dimensions divisible by 16, such that the image is contained in the polynomial subalgebra. The methods combine the K -theory characteristic numbers as used by Stong with the use of the Landweber-Novikov ring.

Let M be a closed smooth manifold of dimension k . For n large, M can be embedded in S^{4n+k} in essentially only one way. Let ν denote the normal bundle of the embedding. We say that M is quaternionic if ν has the structure of a quaternionic vector bundle. Stong [11] proved that in dimensions $k < 24$ all Stiefel-Whitney numbers of a quaternionic manifold M^k vanish. In his thesis, David Segal [8] extended this to dimensions $k < 32$. He also showed the existence of a quaternionic M^{32} unorientedly cobordant to $[RP(2)]^{16}$.

To put the situation more generally, there is the unoriented cobordism ring $N_* = \sum N_k$, and there is a quaternionic cobordism group $\Omega_*^{Sp} = \sum \Omega_k^{Sp}$ [10]. There is a natural homomorphism $\Omega_*^{Sp} \rightarrow N_*$, and $\text{Im}(\Omega_*^{Sp} \rightarrow N_*)$ is the subalgebra of N_* consisting of all cobordism classes represented by quaternionic manifolds M . I am not able to compute this image, but I do give an upper bound to the image and now turn to an outline of the method. In their more widely ranging studies of symplectic cobordism, both Porter [14] and Segal have recently obtained very closely related results.

Let A' denote a graded vector space over Z_2 with basis consisting of all s^R , where R ranges over all sequences $R = (r_1, r_2, \dots)$ of nonnegative integers with $\sum r_k < \infty$ and where $\deg s^R = \sum kr_k$. Landweber [5] and Novikov [7] put a Hopf algebra structure on A' . The product can be defined as composition of operators. Namely each s^R acts on any polynomial algebra $Z_2[x_1, \dots, x_m]$ by

$$\begin{aligned} s^0(x_i) &= x_i, \\ s^{\Delta_k}(x_i) &= x_i^{k+1} \quad \text{where } \Delta_k = (0, \dots, 0, 1, 0, \dots), \\ s^R(x_i) &= 0 \quad \text{if } R \neq 0 \text{ and } R \neq \Delta_k, \\ s^R(yz) &= \sum_{R_1 + R_2 = R} s^{R_1}(y)s^{R_2}(z). \end{aligned}$$

Received by the editors March 26, 1970.

AMS 1968 subject classifications. Primary 5710, 5732; Secondary 5730.

Key words and phrases. Quaternionic manifolds, symplectic cobordism, unoriented cobordism, Stiefel-Whitney numbers, K -theory characteristic numbers.

Copyright © 1971, American Mathematical Society

The mod 2 Steenrod algebra \mathcal{S} can be considered as a subalgebra of A' by identifying Sq^k with $s^{k\Delta_1}$. There is also the subalgebra T of A' generated by s^{Δ_2} and all the Sq^k . As always, $A'/T^+ \cdot A'$ has a natural coproduct where T^+ consists of all sums of elements of T of positive degree. Its dual algebra is a polynomial algebra with generators in each dimension 2^k ($k > 1$), each $2(2k-1)$ where $k \neq 2^j$, and each $2k$ with $k \neq 2^j$. More specifically, we show in §4 that A' is a free left T -module with basis elements s^R for all R with $r_{2^j-1} = 0$, $r_{2k-1} = 0 \pmod 2$ for $k \neq 2^j$, $r_{2^k} = 0 \pmod 2$.

Now let MO denote the Thom spectrum of the orthogonal group, so that $\pi_k(MO)$ is the unoriented cobordism group N_k . There is the pairing

$$\tilde{H}^*(MO; \mathbb{Z}_2) \otimes N_* \rightarrow \mathbb{Z}_2$$

sending $s^R \otimes [M]$ into the normal characteristic number mod 2 $s^R[M]$. We may identify $\tilde{H}^*(MO; \mathbb{Z}_2)$ with A' and obtain $A'_k \otimes N_k \rightarrow \mathbb{Z}_2$. According to Thom [12],

$$(A'/\mathcal{S}^+ \cdot A')_k \otimes N_k \rightarrow \mathbb{Z}_2$$

is a dual pairing.

Define $P_k \subset N_k$ to be all cobordism classes $[M]$ of N_k such that $(s^{\Delta_2 s^R})[M] = 0$, for all R of degree $k-2$. Then $P = \sum P_k$ is a subalgebra of N_* and P_k is the dual group of $(A'/T^+ A')_k$. In §5 we show that we can select a basis $[M^2], [M^4], [M^5], \dots$ for the polynomial algebra N_* such that P is the subalgebra generated by all $[M^{2^k-1}]^2$ for $k \neq 2^j$, $[M^{2^k}]^2$, $[M^{2^k}]$ for $k \neq 2^j$.

Similarly define U to be the subalgebra of A' generated by $s^{\Delta_2}, s^{2\Delta_2}$ and all Sq^k . In §4 the structure of $A'/U^+ A'$ is computed. In §5 we let $Q_k \subset N_k$ be all $[M]$ with

$$(s^{\Delta_2 s^R})[M] = 0, \quad \text{degree } R = k-2,$$

$$(s^{2\Delta_2 s^R})[M] = 0, \quad \text{degree } R = k-4.$$

We show that $Q = \sum Q_k$ is given by $Q = P^2$.

Consider now the spectrum MSP coming from the symplectic group. Then $\Omega_*^{Sp} = \pi_*(MSP)$. There is the pairing

$$\tilde{H}^*(MSP; \mathbb{Z}_2) \otimes \Omega_*^{Sp} \rightarrow \mathbb{Z}_2.$$

We can regard $\tilde{H}^*(MSP; \mathbb{Z}_2)$ as another copy of the Hopf algebra A' , with generators $'S^R$. The two key remarks of §3 which make possible the use of A' are the following: if $[M] \in \Omega_{8k}^{Sp}$ then

$$('S^{\Delta_2} \cdot 'S^R)[M] = 0 \pmod 2, \quad R \text{ of degree } k-2;$$

if $[M] \in \Omega_{8k}^{Sp}$ then

$$('S^{2\Delta_2} \cdot 'S^R)[M] = 0 \pmod 2, \quad \text{degree } R = k-4.$$

These assertions are proved by a very standard use of K -theory, in particular the Adams operations. §§2 and 3 are largely devoted to a proof of these remarks.

The homomorphism $\Omega_*^{Sp} \rightarrow N_*$ comes from a map of spectra $g: MSp \rightarrow MO$. We have

$$g^*(s^{4R}) = 'S^R, \quad g^*(s^{R'}) = 0 \quad \text{if } R' \neq 4R.$$

Hence if $[M] \in \text{Im}(\Omega_*^{Sp} \rightarrow N_*)$ then

$$\begin{aligned} (s^{4\Delta_2} \cdot s^R)[M] &= 0, & (s^{8\Delta_2} \cdot s^R)[M] &= 0, \\ s^{R'}[M] &= 0, & R' &\neq 4R. \end{aligned}$$

It follows without difficulty that

$$\text{Im}(\Omega_*^{Sp} \rightarrow N_*) \subset Q^4 = P^8.$$

That is, there is a basis $[M^2], [M^4], [M^5], \dots$ for N_* such that $\text{Im}(\Omega_*^{Sp} \rightarrow N_*)$ is contained in the polynomial subalgebra generated by

$$[M^2]^{16}, [M^4]^{16}, [M^5]^{16}, [M^6]^8, \dots$$

Here M^{2k} and M^{2l-1} for $l \neq 2^j$ occur to the power 16, M^{2k} for $k \neq 2^j$ to the power 8. This is one of the two main theorems of the paper.

While about it, it is easy to prove a similar assertion for another natural subalgebra E of N_* . Let E_{2k} consist of all cobordism classes N_{2k} which are represented by weakly almost complex manifolds M for which all Chern numbers

$$c_{2r+1}c_{i_1} \cdots c_{i_p}[M]$$

vanish. We prove that $E \subset P^4$ so that E is contained in the subalgebra generated by

$$[M^2]^8, [M^4]^8, [M^5]^8, [M^6]^4, \dots$$

This is the other principal theorem of the paper. We conjecture that $E = P^4$; at least as in §1 one can construct a few nonzero elements of E . In fact, in a later paper we hope to prove $E = P^4$ using methods similar to those of Stong [11].

1. **A few examples.** The purpose of this section is to prove the following.

THEOREM. *There is a closed smooth manifold M^{16n} , unorientedly cobordant to $[RP(2n)]^8$, whose stable tangent bundle*

$$\tau(M) - 16n \in \tilde{K}O(M)$$

is of the form $2y$ where $y \in \tilde{K}O(M)$.

Proof. Consider the complex Grassmann manifold

$$M^{16n} = CM(4n, 2)$$

consisting of all 2-dimensional vector subspaces V of C^{4n+2} . There is the bundle ξ associating with each $V \in M$ all $x \in V$, the trivial bundle $4n+2$ associating with V

all $x \in C^{4n+2}$, and ξ^\perp associating with V all $x \in V^\perp$ where V^\perp is the orthogonal complement of V in C^{4n+2} . The tangent bundle to M is [4]

$$\begin{aligned}\tau(M) &= \text{Hom}_C(\xi, \xi^\perp) \\ &= \text{Hom}_C(\xi, 4n+2) - \text{Hom}_C(\xi, \xi) \\ &= (4n+2)\xi - \text{Hom}_C(\xi, \xi).\end{aligned}$$

There is the operator $\alpha: \text{Hom}_C(\xi, \xi) \rightarrow \text{Hom}_C(\xi, \xi)$ where $\alpha(\varphi)$ is the adjoint φ^* of φ . Clearly α is a conjugate linear involution. Let

$$A_+ = \{\varphi: \alpha(\varphi) = \varphi\}, \quad A_- = \{\varphi: \alpha(\varphi) = -\varphi\}.$$

Then

$$\text{Hom}_C(\xi, \xi) = A_+ + A_-$$

and multiplication by i is a real isomorphism between A_+ and A_- . Hence $\text{Hom}_C(\xi, \xi)$ is of the form $2x$ in $\tilde{K}O(M)$, and $\tau(M) - 16n = 2y$ for some $y \in \tilde{K}O(M)$.

The equation $[CM(4n, 2)] = [RP(2n)]^8$ follows readily from the equations $[CM(n, k)] = [RM(n, k)]^2$, $[RM(2n, 2k)] = [CM(n, k)]^2$. Here $CM(n, k)$ consists of all k -dimensional complex vector subspaces of C^{n-k} and $RM(n, k)$ of all real k -dimensional real subspaces of R^{n-k} . Each of these follows from the theorem [1]: Let $T: M \rightarrow M$ be a smooth involution on the closed smooth manifold M^{2n} , with fixed point set F of dimension n ; if the normal bundle to F has the same Stiefel-Whitney classes as the tangent bundle of F , then $[M] = [F]^2$.

Consider $R^{2n+2k} = C^{n+k}$, and let $M = RM(2n, 2k)$. There is $T: M \rightarrow M$ defined by $T(V) = I(V)$ where I is multiplication by i . The fixed point set F is clearly $CM(n, k)$. The inclusion $i: F \subset M$ induces $i^*\tau(M)$ and

$$i^*\tau(M) = \text{Hom}_R(i^*\xi, i^*\xi^\perp).$$

For $V \in F$, we have V is a complex vector space and

$$\text{Hom}_R(V, V^\perp) = \text{Hom}_C(V, V^\perp) + \text{Hom}'_C(V, V^\perp)$$

where the second term denotes all conjugate linear homomorphisms. Now $\text{Hom}_C(i^*\xi, i^*\xi^\perp)$ is the tangent bundle $\tau(F)$, hence the normal bundle η to F in M is

$$\eta = \text{Hom}'_C(i^*\xi, i^*\xi^\perp).$$

Hence

$$\xi(F) = \overline{i^*(\xi)} \otimes_C i^*\xi^\perp, \quad \eta = \overline{i^*\xi} \otimes_C \overline{i^*\xi^\perp}.$$

These two bundles have the same Chern classes reduced mod 2, hence they have the same Stiefel-Whitney classes. The second equation follows. The first follows readily from [1, pp. 64–65].

It is possible to put the above construction a little more generally, but I do not know how to take advantage of the added generality. Namely suppose M is a closed smooth manifold whose stable tangent bundle is of the form $2y$ for some $y \in \tilde{K}O(M)$. Let ξ be a smooth complex bundle over M and let $CM(2\xi, 2)$ denote the space of all two-dimensional vector subspaces of fibers of 2ξ . Then $M' = CM(2\xi, 2)$ has stable tangent bundle of the form $2z$ for some $z \in \tilde{K}O(M')$.

Note that we can make M^{16n} weakly almost complex by assigning to its stable tangent bundle the complex structure of the complexification of y . In this complex structure we have $c_{2r+1}(M^{16n}) = 0$ since on the one hand $c_{2r+1}(M^{16n})$ is torsion and on the other hand $H^*(M^{16n})$ has no torsion. Hence

$$c_{2r+1}c_{i_1} \cdots c_{i_p}[M^{16n}] = 0$$

whether the Chern numbers are those of the tangent bundle or the normal bundle. That is, $[RP(2n)]^8 \in E_{16n}$.

2. The spectra of Thom spaces. Let ξ be the canonical complex line bundle over $BU(1)$, and let η be the canonical quaternionic line bundle over $BSp(1)$. There is a map

$$g: BU(1) \rightarrow BSp(1)$$

with $g^*(\eta) = \xi + \bar{\xi}$. Let $v = \eta - 2$ in $K(BSp(1))$. Then $K(BSp(1))$ is the ring of formal power series $Z[[v]]$. We may take $H^*(BSp(1))$ as $Z[u]$ for $u \in H^4(BSp(1))$, and suppose

$$\text{ch } v = u + \text{higher order terms.}$$

Also if ψ^2 is the Adams operation,

$$g^*\psi^2(\eta - 2) = \psi^2(\xi + \bar{\xi} - 2) = \xi^2 + \bar{\xi}^2 - 2 = (\xi + \bar{\xi} - 2)^2 + 4(\xi + \bar{\xi} - 2)$$

so that $\psi^2(v) = 4v + v^2$. More generally we can identify $K^*(BSp(1))$ with $K^*(pt)[[v]]$.

Similarly we can consider $v' = \eta - 1$ as an element of $\tilde{K}O^4(\) \approx \tilde{K}Sp(\)$ and

$$KO^*(BSp(1)) = KO^*(pt)[[v']].$$

The natural homomorphism $KO^*(\) \rightarrow K^*(\)$ maps v' into v and v'^n into v^n .

One can next check that

$$H^*(BSp(1) \times \cdots \times BSp(1)) \approx Z[u_1, \dots, u_n],$$

$$KO^*(BSp(1) \times \cdots \times BSp(1)) \approx KO^*(pt)[[v'_1, \dots, v'_n]],$$

$$K^*(BSp(1) \times \cdots \times BSp(1)) \approx K^*(pt)[[v_1, \dots, v_n]],$$

where $\text{ch } v_i = u_i + \text{higher order terms}$, $\psi^2(v_i) = 4v_i + v_i^2$, $KO^*(\) \rightarrow K^*(\)$ maps v'_i into v_i .

One can use Dold's theorem [2] on the natural map

$$BSp(1) \times \cdots \times BSp(1) \rightarrow BSp(n)$$

to check that $H^*(BSp(n))$, $KO^*(BSp(n))$, $K^*(BSp(n))$ can be identified with the symmetric terms of

$$Z[u_1, \dots, u_n], \quad KO^*(pt)[[v'_1, \dots, v'_n]], \quad K^*(pt)[[v_1, \dots, v_n]]$$

respectively.

Finally we identify $MSp(n)$ with $BSp(n)/BSp(n-1)$ to obtain an exact sequence

$$\dots \longrightarrow h^*(MSp(n)) \longrightarrow h^*(BSp(n)) \xrightarrow{f^*} h^*(BSp(n-1)) \longrightarrow \dots$$

for $h=H, KO$ or K . In each case f^* is an epimorphism with kernel the ideal generated by $u_1 \cdots u_n, v'_1 \cdots v'_n, v_1 \cdots v_n$, respectively. Hence we identify $\tilde{H}^*(MSp)$ with the ideal of symmetric polynomials of $Z[u_1, \dots, u_n]$ generated by $u_1 \cdots u_n$, $\tilde{K}(MSp(n))$ with the ideal of symmetric formal series in $Z[[v_1, \dots, v_n]]$ generated by $v_1 \cdots v_n$, and $\tilde{KO}^*(MSp(n))$ with the ideal of symmetric elements of $KO^*[[v'_1, \dots, v'_n]]$ generated by $v'_1 \cdots v'_n$.

There is the spectrum MSp generated by the $MSp(n)$, and the groups $\tilde{H}^*(MSp)$, $\tilde{KO}^*(MSp)$, $\tilde{K}^*(MSp)$ defined as inverse limits.

For each sequence $R=(r_1, r_2, \dots, r_k, \dots)$ of nonnegative integers with $|R|=\sum r_k < \infty$, there is the element $'S^R$ of $\tilde{H}^*(MSp)$ represented in $\tilde{H}^*(MSp(n))$ by

$$'S^R = \sum u_1^2 \cdots u_{r_1}^2 u_{r_1+1}^3 \cdots u_{r_1+r_2}^3 \cdots u_{r_1+r_2+1}^4 \cdots u_{|R|+1} \cdots u_n$$

for n large.

Define $\|R\| = \sum kr_k$, so that $'S^R \in H^{4\|R\|}(MSp)$.

Hence we identify $\tilde{H}^*(MSp)$ with the free abelian group A generated by all $'S^R$ for all R . Similarly $\tilde{K}(MSp)$ contains a free abelian group A generated by all S^R , where S^R is represented in $\tilde{K}(MSp(n))$ by

$$S^R = \sum v_1^2 \cdots v_{r_1}^2 v_{r_1+1}^3 \cdots v_{r_1+r_2}^3 \cdots v_{|R|+1} \cdots v_n.$$

Moreover $\text{ch } S^R = 'S^R + \text{higher order terms}$.

There is the quaternionic cobordism group Ω_k^{Sp} [10], which may either be regarded as all bordism classes of closed smooth k -manifolds with given quaternionic structure on the stable normal bundle, or as all homotopy classes of maps

$$f: S^{4n+4k} \rightarrow MSp(n), \quad n \text{ large.}$$

Given $[M] \in \Omega_{4k}^{Sp}$ represented by $f: S^{4n+4k} \rightarrow MSp(n)$ and given $'S^R$, we get the integer

$$'S^R[M] = \langle f^* 'S^R, \sigma_{4n+4k} \rangle$$

where σ_{4n+4k} is the orientation class of S^{4n+4k} . Note that $'S^R[M]=0$ unless $\|R\|=k$. Similarly for each $S^R \in \tilde{K}(MSp)$, we get the integer

$$S^R[M] = \langle \text{ch } f^* S^R, \sigma_{4n+4k} \rangle,$$

and $S^R[M]=0$ unless $\|R\| \leq k$. If $\|R\|=k$, then $S^R[M] = 'S^R[M]$.

The above completes the notation concerning the spectrum MSp . Note that an equivalent discussion holds for MU except for the remarks about KO^* . Namely $\tilde{H}^*(MU)$ can be identified with the free abelian group A generated by all $'S^R$, where $'S^R$ is represented in $H^*(MU(n))$ by

$$'S^R = \sum x_1^2 \cdots x_{r_1}^2 x_{r_1+1}^3 \cdots x_{r_1+r_2}^3 \cdots x_{|R|+1} \cdots x_n.$$

Similarly if $[M] \in \Omega_{2k}^U$ is represented by $f: S^{2n+2k} \rightarrow MU(n)$, there are the integers $'S^R[M], S^R[M]$ defined as before.

Finally there is the spectrum MO . The cohomology group $\tilde{H}^*(MO; Z_2)$ can be identified with the vector space over Z_2 with basis all s^R where s^R is represented in $\tilde{H}^*(MO(n); Z_2)$ by

$$s^R = \sum z_1^2 \cdots z_{r_1}^2 z_{r_1+1}^3 \cdots z_{r_1+r_2}^3 \cdots z_{|R|+1} \cdots z_n.$$

For $[M] \in N_k$, there is the integer mod 2

$$s^R[M] = \langle f^* s^R, \sigma'_{n+k} \rangle.$$

There are natural maps of spectra

$$MSp \xrightarrow{g} MU \xrightarrow{h} MO.$$

We assume the following facts:

- (a) $g^*: \tilde{K}(MU) \rightarrow \tilde{K}(MSp)$ is an epimorphism,
- (b) we have $h^*: \tilde{H}^*(MO; Z_2) \rightarrow \tilde{H}^*(MU; Z_2)$ given by $h^*(s^{2R}) = 'S^R \text{ mod } 2, h^*(s^R) = 0$ if $R \neq 2R'$,
- (c) $g^*: \tilde{H}^*(MU; Z_2) \rightarrow \tilde{H}^*(MSp; Z_2)$ is given by $g^*('S^{2R}) = s^{2R} \text{ mod } 2, g^*('S^R) = 0$ if $R \neq 2R'$.

3. Divisibility relations among characteristic numbers. Denote by A a free abelian group with basis elements S^R , one for each sequence $R=(r_1, r_2, \dots)$ of nonnegative integers with $|R| = \sum r_k < \infty$. Following Landweber [5] and Novikov [7], there is a multiplication defined on A . Each S^R operates as a group homomorphism on any polynomial algebra $Z[x_1, \dots, x_n]$ with given generators by

$$\begin{aligned} S^0(x_i) &= x_i, \\ S^{\Delta_k}(x_i) &= x_i^{k+1} \quad \text{where } \Delta_k = (0, \dots, 0, 1, 0, \dots), \\ S^R(x_i) &= 0 \quad \text{if } R \neq 0 \text{ and } R \neq \Delta_k, \\ S^R(yz) &= \sum_{R_1 + R_2 = R} S^{R_1}(y) \cdot S^{R_2}(z), \end{aligned}$$

for all $y, z \in Z[x_1, \dots, x_n]$.

Given R, R' there exists a unique $T \in A$ with $S^R \circ S^{R'} = T$ on all $Z[x_1, \dots, x_n]$. In this fashion A becomes an associative ring with unit.

For each nonnegative integer k there is also an operator

$$SQ^k: Z[x_1, \dots, x_n] \rightarrow Z[x_1, \dots, x_n]$$

defined by

$$\begin{aligned} SQ^0(x_i) &= x_i, & SQ^1(x_i) &= x_i^2, \\ SQ^k(x_i) &= 0 \quad \text{for } k > 1, \\ SQ^k(yz) &= \sum_{i+j=k} SQ^i(y) \cdot SQ^j(z). \end{aligned}$$

As operators, we have $SQ^k = S^{(k,0,0,\dots)}$ and thus we identify SQ^k with $S^{(k,0,\dots)} \in A$. There are the formulas

$$S^R(x_i^m) = \frac{m!}{r_1!r_2! \dots (m-|R|)!} x_i^{m+\|R\|}, \quad SQ^k(x_i^m) = \binom{m}{k} x_i^{m+k}.$$

An element $T \in A$ is uniquely determined by its operator value $T(x_1x_2 \dots x_n)$ in $Z[x_1, \dots, x_n]$ for n large. Thus

$$\begin{aligned} S^{\Delta_2}(x_1 \dots x_n) &= \sum x_1^3 x_2 \dots x_n, \\ (SQ^1 \cdot SQ^1 - 2SQ^2)(x_1 \dots x_n) &= 2 \sum x_1^3 x_2 \dots x_n, \end{aligned}$$

and hence

$$(3.1) \quad S^{\Delta_2} = \frac{1}{2}SQ^1 \cdot SQ^1 - SQ^2.$$

Similarly

$$\begin{aligned} S^{2\Delta_2}(x_1 \dots x_n) &= \sum x_1^3 x_2^3 x_3 \dots x_n, \\ (SQ^2 \cdot SQ^2 - SQ^3 \cdot SQ^1 - SQ^1 \cdot SQ^3 + 2SQ^4)(x_1 \dots x_n) &= 4 \sum x_1^3 x_2^3 x_3 \dots x_n, \end{aligned}$$

and hence

$$(3.2) \quad S^{2\Delta_2} = \frac{1}{4}(SQ^2 \cdot SQ^2 - SQ^3 \cdot SQ^1 - SQ^1 \cdot SQ^3 + 2SQ^4).$$

We now turn to the pairing

$$A \otimes \Omega_*^{Sp} \rightarrow Z, \quad S^R \otimes [M] \rightarrow S^R[M]$$

of §2 where $S^R \in \tilde{K}(MSp)$. Let $f: S^{4n+4k} \rightarrow MSp(n)$ and let $S^R \in \tilde{K}(MSp(n))$ where $r = \|R\| \leq k$. Then

$$\begin{aligned} S^R &= \sum v_1^2 \dots v_{r_1+1}^3 \dots v_{|R|+1} \dots v_n, \\ \psi^2 S^R &= \sum (4v_1 + v_1^2)^2 \dots (4v_{r_1+1} + v_{r_1+1}^2)^3 \dots \\ &= 4^{n+r} S^R + 4^{n+r-1} (SQ^1 \cdot S^R) + \dots + 4^{n-k+2r} (SQ^{k-r} \cdot S^R) + \dots \end{aligned}$$

Also

$$\begin{aligned}
 (\psi^2 S^R)[M] &= \langle \text{ch } f^* \psi^2 S^R, \sigma_{4n+4k} \rangle \\
 &= 4^{n+k} \langle \text{ch } f^* S^R, \sigma_{4n+4k} \rangle = 4^{n+k} S^R[M].
 \end{aligned}$$

Combining the two equations and dividing by 4^{n-k+2r} , we get

$$(3.3) \quad 4^{2k-2r} S^R[M] = 4^{k-r} S^R[M] + \dots + 4(SQ^{k-r-1} \cdot S^R)[M] + (SQ^{k-r} \cdot S^R)[M].$$

Recall here that $[M] \in \Omega_{4k}^{Sp}$ and that $\|R\| = r \leq k$.

(3.4) COROLLARY. For each $[M] \in \Omega_{4k}^{Sp}$ and each S^R of degree $r < k$, we have $(SQ^{k-r} \cdot S^R)[M] = 0 \pmod 4$.

(3.5) If $[M] \in \Omega_{8k}^{Sp}$ and if $S^R \in \tilde{K}(MSp)$ is of degree $r < 2k$, then $(SQ^{2k-r} \cdot S^R)[M] = 0 \pmod 8$.

Proof. Consider $MSp(2n)$, and consider first R with $\|R\|$ odd. Since $\|R\|$ is odd, in $\tilde{K}O^4(MSp(2n))$ there is the element

$$z = \sum v_1'^2 \cdots v_{r_1+1}'^3 \cdots v_{|R|+1}' \cdots v_{2n}'$$

and $\tilde{K}O^*() \rightarrow \tilde{K}()$ maps z into S^R . Since $\tilde{K}O^4() \approx \tilde{K}Sp()$, we may as well suppose that S^R is a quaternionic bundle. Then

$$S^R[M^{8k}] = \langle \text{ch } f^*(S^R), \sigma_{8n+8k} \rangle$$

and since we can consider $f^*(S^R) \in \tilde{K}Sp(S^{8n+8k})$, we get $S^R[M] = 0 \pmod 2$. In particular if $\|R\| = 2k - 1$, (3.3) then gives $12S^R[M^{8k}] = (SQ^1 \cdot S^R)[M^{8k}]$ and hence

$$(SQ^1 \cdot S^R)[M] = 0 \pmod 8, \quad \|R\| = 2k - 1.$$

The general case now follows from (3.3), for $(SQ^{2k-r-1} \cdot S^R)[M] = 0 \pmod 2$, by the case already treated.

We can now convert (3.4) and (3.5) into remarks about cohomology characteristic numbers.

(3.6) For all $[M^{4k}] \in \Omega_{4k}^{Sp}$ and all $'S^R \in H^{4k-8}(MSp)$ we have

$$('S^{\Delta_2} \cdot 'S^R)[M^{4k}] = 0 \pmod 2.$$

Here the product is that of $A = \tilde{H}^*(MSp)$.

Proof. We have

$$\begin{aligned}
 ('S^{\Delta_2} \cdot 'S^R)[M] &= (S^{\Delta_2} \cdot S^R)[M] \\
 &= ((\frac{1}{2}SQ^1 \cdot SQ^1 - SQ^2) \cdot S^R)[M] = 0 \pmod 2
 \end{aligned}$$

by (3.4).

(3.7) For all $[M^{8k}] \in \Omega_{8k}^{Sp}$ and all $'S^R \in \tilde{H}^{8k-16}(MSp)$ we have

$$('S^{2\Delta_2} \cdot 'S^R)[M^{8k}] = 0 \pmod 2.$$

Proof. We have

$$\begin{aligned} ('S^{2\Delta_2} \cdot 'S^R)[M] &= (S^{2\Delta_2} \cdot S^R)[M] \\ &= (\frac{1}{4}(SQ^2 \cdot SQ^2 - SQ^3 \cdot SQ^1 - SQ^1 \cdot SQ^3 + 2SQ^4) \cdot S^R)[M] \\ &= 0 \pmod 2 \end{aligned}$$

by (3.5).

We shall next see that (3.6) applies to a considerably more general class of manifolds M .

Consider $\tilde{H}^*(MU(2n))$ as the ideal of $H^*(BU(2n))$ generated by c_{2n} . In particular, consider $\tilde{H}^*(MU(2n))$ as a module over $H^*(BU(2n))$. The natural map $g: MSp(n) \rightarrow MU(2n)$ then has

$$g^*: H^*(MU(2n)) \rightarrow H^*(MSp(n))$$

an epimorphism with kernel the submodule generated by $c_1, c_3, \dots, c_{2k+1}, \dots$. Hence if

$$g^*: \tilde{K}(MU(2n)) \rightarrow \tilde{K}(MSp(n))$$

has $g^*(T)=0$, then $\text{ch } T$ has every term in the submodule generated by $c_1, c_3, \dots, c_{2k+1}, \dots$.

Consider now the pairing

$$\tilde{K}(MU) \otimes \Omega_{4k}^U \rightarrow Z, \quad T \otimes [M] \rightarrow T[M].$$

It follows readily from the above that if $g^*(T)=0$ and if every Chern number of M^{4k} involving any c_{2r+1} vanishes, then $T[M^{4k}]=0$. Recall that

$$g^*: \tilde{K}(MU) \rightarrow \tilde{K}(MSp)$$

is also an epimorphism. Given $S^R \in \tilde{K}(MSp)$, select $T \in \tilde{K}(MU)$ with $g^*(T)=S^R$. Given $[M^{4k}] \in \Omega_{4k}^U$ such that every Chern number of M involving any c_{2r+1} vanishes, then define $S^R[M^{4k}]=T[M^{4k}]$. Note that $S^R[M^{4k}]$ is an integer.

We next see that (3.3) holds for this wider class of manifolds. Recall that $\Omega_*^{Sp} \rightarrow \Omega_*^U$ maps Ω_*^{Sp} into the subalgebra W of Ω_*^U consisting of all $[M^{4k}]$ such that every Chern number of $[M^{4k}]$ involving any c_{2r+1} vanishes. Moreover $\Omega_*^{Sp} \rightarrow W$ is an isomorphism modulo 2-torsion groups. Hence given $[M^{4k}] \in W$ there is 2^s with $2^s[M^{4k}] \in \text{Im } \Omega_*^{Sp}$. Then (3.3) holds for $2^s M^{4k}$ and hence also for M^{4k} . We can now easily obtain the following.

(3.8) *Suppose that $[M^{4k}] \in \Omega_{4k}^U$ has all Chern numbers involving any c_{2r+1} vanishing. Then for every $'S^R$ in $\tilde{H}^*(MSp)$ with $\|R\|=k$, the number $'S^R[M^{4k}]$ is well defined and $('S^{\Delta_2} \cdot 'S^R)[M^{4k}]=0 \pmod 2$, $\|R\|=k-2$.*

4. The mod 2 algebra $A \otimes Z_2$. We now set up the machinery to properly utilize (3.7) and (3.8). The starting point is the work of Landweber on the Landweber-Novikov algebra.

Let A' be a vector space over Z_2 with basis consisting of all s^R where R ranges over all sequences $R=(r_1, r_2, \dots)$ of nonnegative integers with $\sum r_k < \infty$. Every element of A' acts on any polynomial algebra $Z_2[x_1, \dots, x_n]$ exactly as in §3. This determines an associative ring structure on A' , and $A' = A \otimes Z_2$. There are also operators

$$Sq^k: Z_2[x_1, \dots, x_n] \rightarrow Z_2[x_1, \dots, x_n]$$

exactly as before, and we consider $Sq^k = s^{(k, 0, \dots)} \in A'$. Moreover A' is a Hopf algebra with coproduct

$$\psi(s^R) = \sum_{R_1 + R_2 = R} s^{R_1} \otimes s^{R_2}.$$

Denote by $\mathcal{S} \subset A'$ the subalgebra generated by all the $Sq^k, k \geq 0$. Reasoning of the type of Milnor (see Landweber again) shows that \mathcal{S} has as additive basis all s^R for all $R=(r_1, r_2, \dots)$ such that $r_k = 0$ whenever k is not of the form $2^j - 1$.

We can now set the problem of this section. Denote by $T \subset A'$ the subalgebra generated by s^{Δ_2} and by all Sq^k ; denote by $U \subset A'$ the subalgebra generated by s^{Δ_2} , by $s^{2\Delta_2}$, and by all Sq^k . Then A' is a left T -module and there is $A'/T^+ \cdot A'$. Moreover A' is a left U -module and there is $A'/U^+ \cdot A'$. In this section we determine the structure of $A'/T^+ \cdot A'$ and $A'/U^+ \cdot A'$.

Linearly order the sequences R by

- (i) $R < R'$ if $\|R\| < \|R'\|$,
- (ii) $R < R'$ if $\|R\| = \|R'\|$ and if $r_1 = r'_1, \dots, r_k = r'_k, r_{k+1} > r'_{k+1}$.

Note that this is a well-ordering, since for any k there are only a finite number of R with $\|R\| = k$.

For any R, R' we have

$$\begin{aligned} s^R s^{R'}(x_1 \cdots x_m) &= s^R \left(\sum x_1^2 \cdots x_{r'_1}^2 x_{r'_1+1}^3 \cdots \right) \\ &= a(R, R') s^{R+R'} + \sum_{T > R+R'} a_T s^T, \end{aligned}$$

where

$$a(R, R') = \prod \binom{r_k + r'_k}{r_k} \pmod 2.$$

Hence if $a(R, R') = 1 \pmod 2$ then

$$s^R s^{R'} = s^{R+R'} + \sum_{T > R+R'} a_T s^T.$$

(4.1) Suppose that x_i and y_j are sequences of elements of A' , where

$$x_i y_j = s^{R_{i,j}} + \sum_{T > R_{i,j}} a_T s^T.$$

Suppose for $i \neq k$ or for $j \neq l$ that $R_{i,j} \neq R_{k,l}$. The elements x_i, y_j are then linearly independent in A' .

The proof is quite clear. For in any set of elements $R_{i,j}$ there is at least one.

We need now a few computations (see Landweber). Let s_k denote s^{Δ_k} and let $s_{k,k}$ denote $s^{2\Delta_k}$.

$$\begin{aligned} (s_{2p}s_{2q+1})(x_1x_2 \cdots x_n) &= s_{2p}(\sum x_1^{2q+2}x_2 \cdots x_n) \\ &= \sum x_1^{2q+2}x_2^{2p+1} \cdots x_n, \\ (s_{2q+1}s_{2p})(x_1 \cdots x_n) &= s_{2q+1}(\sum x_1^{2p+1}x_2 \cdots x_n) \\ &= \sum x_1^{2p+2q+2}x_2 \cdots x_n + \sum x_1^{2p+1}x_2^{2q+2} \cdots x_n, \end{aligned}$$

and

$$(1) \quad s_{2p}s_{2q+1} + s_{2q+1}s_{2p} = s_{2p+2q+1}.$$

Similarly

$$(2) \quad s_{2p}s_{2p} = s_{4p}$$

so that $s_2^{k+1} = (s_2)^{2^k}$. Also

$$(3) \quad s_{2p+1}s_{2q+1} = s_{2q+1}s_{2p+1}, \quad (s_{2p+1})^2 = 0.$$

Moreover for $k > 1$,

$$\begin{aligned} (Sq^n \cdot s_k)(x_1 \cdots x_m) &= Sq^n(\sum x_1^{k+1}x_2 \cdots x_n) \\ &= \sum_r \left(\binom{k+1}{r} \sum x_1^{k+r+1}x_2^2 \cdots x_{n-r+1}^2x_{n-r+2} \cdots \right) \\ &= \sum \binom{k+1}{r} s_{k+r} \cdot Sq^{n-r}(x_1 \cdots x_m) \end{aligned}$$

and

$$(4) \quad Sq^n s_k = \sum \binom{k+1}{r} s_{k+r} \cdot Sq^{n-r}, \quad k > 1.$$

Recall that T is generated by s_2 and the Sq^k . Since $s_1 \in \mathcal{S} \subset T$ and $s_2 \in T$ we obtain inductively that $s_{2k+1} \in T$ from $s_{2k+1} = s_2s_{2k} + s_{2k-1}s_2$. Since $s_{2^j-1} \in \mathcal{S}$, the new information here is that $s_{2^k-1} \in T$ for $k \neq 2^j$. Also $s_{2^k+1} \in T$.

It follows from (4) that

$$(5) \quad Sq^n \cdot s_{2^k} = s_{2^k} \cdot Sq^n + s_{2^k+1} \cdot Sq^{n-1} + s_{2^k+1} \cdot Sq^{n-2^k} + s_{2^k+1+1} \cdot Sq^{n-2^k-1},$$

$$\begin{aligned} (6) \quad Sq^n \cdot s_{2k+1} &= \sum \binom{2k+2}{r} s_{2k+r+1} \cdot Sq^{n-r} \\ &= \sum n_{2r} s_{2k+2r+1} \cdot Sq^{n-2r}. \end{aligned}$$

(4.2) T has as additive basis all

$$X = (s_2^{\varepsilon_2} s_4^{\varepsilon_4} \cdots s_{2^k}^{\varepsilon_{2^k}} \cdots) (s_5^{\varepsilon_5} s_9^{\varepsilon_9} s_{11}^{\varepsilon_{11}} \cdots) s^{(r_1, 0, r_3, 0, 0, 0, r_7, 0, \dots)}$$

where $\varepsilon_k = 0$ or 1 and where all but a finite number of the parameters are zero.

Proof. Letting

$$\begin{aligned} x_i &= (s_2^{\epsilon_2} s_4^{\epsilon_4} \dots)(s_5^{\epsilon_5} s_9^{\epsilon_9} \dots), \\ y_j &= s^{(r_1, 0, r_3, 0, 0, 0, r_7, \dots)}, \\ R_{i,j} &= (r_1, \epsilon_2, r_3, \epsilon_4, \epsilon_5, 0, r_7, \epsilon_8, \dots), \end{aligned}$$

we see that $x_i y_j = s^{R_{i,j}}$ + higher terms. Linear independence then follows from (4.1). That every element of T can be written as a linear combination of elements in the form X follows from (1), (2), (3), (5), (6).

(4.3) THEOREM. Consider A' as a T -module under left multiplication by T . Then A' is a free T -module with basis all elements s^R for all R with $r_{2^j-1} = 0, r_{2^k} = 0 \pmod 2, r_{2s-1} = 0 \pmod 2$ for $s \neq 2^j$.

Proof. That A' is a free T -module follows from Milnor-Moore [6], but this is also self-contained. Let

$$\begin{aligned} x_i &= (s_2^{\epsilon_2} s_4^{\epsilon_4} \dots)(s_5^{\epsilon_5} s_9^{\epsilon_9} \dots) s^{(r_1, 0, r_3, 0, \dots)}, \\ y_j &= s^{(0, 2r_2, 0, 2r_4, 2r_5, r_6, \dots)}, \\ R_{i,j} &= (r_1, 2r_2 + \epsilon_2, r_3, 2r_4 + \epsilon_4, 2r_5 + \epsilon_5, r_6, \dots). \end{aligned}$$

Then $x_i y_j = s^{R_{i,j}}$ + larger terms. Hence the elements y_j are linearly independent in A' considered as a T -module, using (4.1). Since every R is of the form $R_{i,j}$ it is clear from induction that the elements $x_i y_j$ generate A' as a vector space, and hence the y_j generate A' as a T -module. The theorem follows.

There is a ring homomorphism $\alpha: A' \rightarrow A'$ defined by

$$\alpha(s^{2R}) = s^R, \quad \alpha(s^{R'}) = 0 \quad \text{if } R' \neq 2R.$$

For consider a polynomial algebra

$$M = \mathbb{Z}_2[x_1, \dots, x_k, \dots]$$

and in it consider the subalgebra

$$N = \{y^2 \mid y \in M\}.$$

Now each s^R acts on M and $s^{2R}(y^2) = (s^R(y))^2, s^{R'}(y^2) = 0$ for $R' \neq 2R$. Thus if $R' \neq 2R$ then $s^{R'}$ acts trivially on N , and there is the commutative diagram

$$\begin{array}{ccc} M & \xrightarrow{s^R} & M \\ \approx \downarrow & & \downarrow \approx \\ N & \xrightarrow{s^{2R}} & N \end{array}$$

where $M \approx N$ is the squaring map. Thus we let $\alpha(s^R)$ be the unique s^R with commutativity holding in

$$\begin{array}{ccc} M & \xrightarrow{s^R} & M \\ \approx \downarrow & & \downarrow \approx \\ N & \xrightarrow{s^{R'}} & N \end{array}$$

In particular, $\alpha(s_{2,2}) = s_2$, $\alpha(Sq^{2k}) = Sq^k$, $\alpha(s_2) = 0$, $\alpha(Sq^{2k+1}) = 0$, and $\alpha(U) = T$ where U is the subalgebra generated by $s_2, s_{2,2}$ and the Sq^k . It follows that α maps $A'/U^+ \cdot A'$ onto $A'/T^+ \cdot A'$.

(4.4) THEOREM. *The homomorphism $\alpha: A' \rightarrow A'$ induces an isomorphism*

$$A'/U^+ \cdot A' \approx A'/T^+ \cdot A'$$

of vector spaces.

Proof. We need the following multiplication formulas.

- (1) $s_{2,2} s_{4k-2} + s_{4k-2} s_{2,2} = s_{4k+2}$,
- (2) $s_{2,2} s_{2,2} + s_{2,2} s_4 + s_2 s_{2,2} = s_{4,4}$,
- (3) $(s_{2^k, 2^k})^2 = s_{2^{k+1}, 2^{k+1}}$, $k \geq 2$,
- (4) $s_{2k-3, 2k-3} s_{4,4} + s_4 s_{2k+1, 2k-3} + s_{4,4} s_{2k-3, 2k-3} + \binom{2k-2}{2} s_{2k+5, 2k-3} = s_{2k+1, 2k+1}$.

It follows from (1) that $s_{4k+2} \in U$. Hence also $s_{8k+4} = (s_{4k+2})^2 \in U, \dots$ and all $s_k \in U$. It follows from (2) that $s_{4,4} \in U$ and from (3) that $s_{2^k, 2^k} \in U$ for all k . Finally (4) implies that $s_{2k+1, 2k+1} \in U$ for all k .

To prove the theorem, we have only to check that the elements s^R for which $r_{2^j-1} = 0, r_{2^j-1} = 0 \pmod 4$ for $s \neq 2^j, r_{2^k} = 0 \pmod 4, r_{2^t} = 0 \pmod 2$ for $t \neq 2^j$ generate A' as a U -module. For clearly the epimorphism $A'/U^+ \cdot A' \rightarrow A'/T^+ \cdot A'$ maps these elements into a basis.

Fix m and consider all R with $\|R\| = m$. The largest such s^R is s^{Δ_m} and $s^{\Delta_m} \in U^+$. Fix R and assume that for every $R' > R$ with $\|R'\| = m$ that $s^{R'}$ is expressible in terms of the proposed basis elements. Also suppose that every $s^{R''}$ is expressed in terms of the proposed base elements if $\|R''\| < m$. Let $R = (r_1, r_2, \dots)$.

Case 1. If some $r_{2^k-1} = t \neq 0$ then

$$s^{t \Delta_{2^k-1}} (r_1, \dots, r_{2^k-2}, 0, r_{2^k}, \dots) = s^R + \text{larger terms.}$$

Hence in this case s^R can be suitably expressed.

Case 2. $r_{2^k} = t = 4r + \varepsilon$ where $\varepsilon = 1, 2$ or 3 . Taking the case $\varepsilon = 3$ we have

$$s_{2^k} \cdot s^{(r_1, \dots, r_{2^k-1}, 4r+2, r_{2^k+1}, \dots)} = s^R + \text{larger terms}$$

and s^R is expressed in terms of the proposed base elements. Similarly if $\varepsilon = 1$.

Taking the case $\epsilon=2$ we have

$$s_{2^k, 2^k} s^{(r_1, \dots, r_{2^k-1}, 4r, \dots)} = s^R + \text{larger terms}$$

and this case follows.

Case 3. $r_{2^k+1} = 4r + \epsilon$ where $\epsilon = 1, 2$ or 3 . This goes exactly as Case 2.

Case 4. $r_{2^k} = 2r + 1, k \neq 2^j$.

$$s_{2^k} s^{(r_1, \dots, r_{2^k-1}, 2r, \dots)} = s^R + \text{larger terms.}$$

Since $s_{2^k} \in U^+$, this case follows.

If none of the above hold, then s^R is one of the proposed base elements. The theorem follows.

5. **The main theorems.** Recall that in §2 we have identified $\tilde{H}^*(MO; Z_2)$ with the algebra $A' = A \otimes Z_2$ of §4. If N_* is the unoriented cobordism group, there is also the pairing $A' \otimes N_* \rightarrow Z_2$ mapping $s^R \otimes [M]$ into $s^R[M]$. Letting $A' = \sum A'_k$ where A'_k is generated by all s^R with $\|R\| = k$, we have

$$A'_k \otimes N_k \rightarrow Z_2.$$

Recalling that $\mathcal{S} \subset A'$ is the Steenrod algebra, we know from Thom that $(\mathcal{S}^+ \cdot A')_k \otimes N_k \rightarrow 0$ and that the induced map

$$(5.1) \quad (A'/\mathcal{S}^+ \cdot A')_k \otimes N_k \rightarrow Z_2$$

is a dual pairing.

Consider now the ring $T \supset \mathcal{S}$ of §2. Define P_k as the subgroup of N_k consisting of all $[M] \in N_k$ such that $(s_2 \cdot s^R)[M] = 0$ for all s^R of degree $k-2$. Alternatively, P_k is the annihilator of $(T^+ \cdot A')_k$ in $A' \otimes N_k \rightarrow Z_2$ and thus there is the dual pairing

$$(A'/T^+ \cdot A')_k \otimes P_k \rightarrow Z_2.$$

$$(5.2) \quad P = \sum P_k \text{ is a subalgebra of } N_*.$$

Proof. If $[M], [M'] \in P$, then

$$\begin{aligned} (s_2 \cdot s^R)[M \times M'] &= \sum_{R_1 + R_2 = R} (s_2 s^{R_1})[M] \cdot s^{R_2}[M'] \\ &+ \sum_{R_1 + R_2 = R} s^{R_1}[M] \cdot (s_2 s^{R_2})[M'] \end{aligned}$$

and $[M \times M'] \in P$.

(5.3) **THEOREM.** *There exist generators $[M^2], [M^4], [M^5], \dots$ for N_* as a polynomial algebra over Z_2 such that P is the polynomial subalgebra with generators $[M^2]^2, [M^4]^2, [M^6], [M^8]^2, [M^9]^2, [M^{10}], \dots$. Here every $[M^{2^k+1}]$ and every $[M^{2^k}]$ occurs to the power two and every $[M^{2^k}]$ with $k \neq 2^j$ to the power one.*

Proof. We know that $(A'/T^+ \cdot A')_k$ and P_k are dually paired. In (4.3) the dimension of $(A'/T^+ \cdot A')_k$ has been obtained and it agrees with that of the above polynomial subalgebra. Hence we have only to show the existence of a polynomial

basis for N_* such that $[M^{2k+1}]^2, [M^{2k}]^2 \in P$ and $[M^{2k}] \in P$ for $k \neq 2^j$. In the first cases the choice is arbitrary since

$$(s_2 s^R)[M \times M] = \sum s_2 s^{R_1}[M] \cdot s^{R_2}[M] + \sum s^{R_1}[M] \cdot s_2 s^{R_2}[M] = 0.$$

Thus we have only to show the existence for $k \neq 2^j$ of an $[M^{2k}] \in P_{2k}$ with $s_{2k}[M^{2k}] = 1$. From (4.3) we know that $s_{2k} \notin T^+ \cdot A'$ for $k \neq 2^j$. Since

$$(A'/T^+ \cdot A')_k \otimes P_k \rightarrow Z_2$$

is a dual pairing, there exists $[M^{2k}] \in P_{2k}$ with $s_{2k}[M^{2k}] = 1$. The theorem follows.

(5.4) THEOREM. Define Q_k to be the set of all cobordism classes $[M] \in N_k$ such that

- (i) $(s_2 \cdot s^R)[M] = 0$, all R with $\|R\| = k - 2$, and
- (ii) $(s_{2,2} \cdot s^R)[M] = 0$, all R with $\|R\| = k - 4$.

Let $Q = \sum Q_k$. Then Q is a subalgebra of N_* and $Q = P^2 = \{x^2 \mid x \in P\}$.

Proof. It is readily checked that Q is a subalgebra, and this is left to the reader. We have a dual pairing

$$(A'/U^+ \cdot A')_k \otimes Q_k \rightarrow Z_2$$

and an isomorphism

$$\alpha: A'/U^+ \cdot A' \approx A'/T^+ \cdot A'.$$

It is also readily checked that if $M \in P_k$ then $[M]^2 \in Q_{2k}$. The diagram

$$\begin{array}{ccc} P_k & \longrightarrow & \text{Hom} [(A'/T^+ A')_k, Z_2] \\ \downarrow f & & \approx \downarrow \alpha' \\ Q_{2k} & \longrightarrow & \text{Hom} [(A'/U^+ A')_{2k}, Z_2] \end{array}$$

is commutative, where $f(x) = x^2$. Hence f is an isomorphism and $Q_{2k} = (P_k)^2$. The theorem follows.

It follows immediately from (5.3) that $P_{2k+1} = 0$. Hence in (5.4) we may replace (ii) by

- (ii)' $(s_{2,2} \cdot s^R)[M] = 0$ when $\dim M = 2k, \|R\| = 2k - 4$.

Let E_{2k} denote the subgroup of N_{2k} consisting of all cobordism classes represented by a weakly almost complex manifold M^{2k} such that its Chern numbers satisfy

$$c_{2r+1} c_{1_1} \cdots c_{1_r} [M] = 0, \quad r = 0, 1, 2, \dots$$

Note that this condition is equivalent whether the stable tangent bundle or the normal bundle is used to define Chern classes. Let $E = \sum E_{2k}$.

(5.5) THEOREM. The subalgebras E, P and $\text{Im} (\Omega_*^{Sp} \rightarrow N_*)$ of N_* satisfy $E \subset P^4$, $\text{Im} (\Omega_*^{Sp} \rightarrow N_*) \subset P^8$.

Proof. The map $g: MSP \rightarrow MO$ has $g^*(s^{4R}) = s^{R'} \pmod 2$, $g^*(s^{R'}) = 0$ for $R' \neq 4R$. Hence for each $[M]$ in $\text{Im}(\Omega_*^{Sp} \rightarrow N_*)$ we have $s^{R'}[M] = 0$ for $R' \neq 4R$. From (3.6) and (3.7) we have

$$(s^{4\Delta_2} \cdot s^R)[M] = 0, \quad (s^{8\Delta_2} \cdot s^R)[M] = 0,$$

the second holding in case $\dim M = 8k$ and $\|R\| = 8k - 16$.

From $s^{R'}[M] = 0$ unless $R' = 4R$ it follows that $[M] = [M']^4$ for some $[M']$. For define $A' \rightarrow Z_2$ by $s^R \rightarrow s^{4R}[M]$. Alternatively there is $\alpha^2: A' \rightarrow A'$ and $\alpha^2(x) \rightarrow x[M]$, which is well defined. Then $\alpha^2(Sq^{4n} s^{4R}) = Sq^n \cdot s^R$, hence $Sq^n \cdot s^R \rightarrow 0$ and the homomorphism $A' \rightarrow Z_2$ kills $\mathcal{S}^+ \cdot A'$. Hence there is an $[M']$ with $s^R[M'] = s^{4R}[M]$, and $[M] = [M']^4$. Then

$$(s^{\Delta_2} \cdot s^R)[M'] = 0, \quad (s^{2\Delta_2} \cdot s^R)[M'] = 0,$$

the latter holding in case $\dim M = 8k$. Then by (5.4) and the remark following it, we have $[M'] \in P^2$ and $\text{Im}(\Omega_*^{Sp} \rightarrow N_*) \subset P^8$. In exactly the same way using (3.8) we see that $E \subset P^4$.

6. Remarks and questions. (a) It can be shown that every $[M] \in P_k$ is a Wall manifold, i.e.

$$w_1^2 w_{i_1} \cdots w_p [M] = 0, \quad \text{all } i_1, \dots, i_p.$$

Thus P is contained in the Wall subalgebra W of N_* . Clearly $P \neq W$ since $P_5 = 0$ and $W_5 \neq 0$. It would be interesting to have a characterization of P as a subalgebra of W . A generator for W_6 can also serve as generator for P_6 . Hence $[M^6]$ can be taken to be the $CP(2)$ bundle over the Klein bottle used by Wall. I have not tried to obtain models for all the generators M^{2k} .

(b) From §3 one gets the impression that one might try to study divisibility relations between Chern numbers in terms of the algebra A . The advantage is that the Adams operations on $A \subset \tilde{K}(MU)$, when suitably stabilized, can be put in terms of the multiplication of A . Can one understand the Stong results [9] on

$$\tilde{K}(MU) \otimes \Omega_*^U \rightarrow Z$$

purely arithmetically? Can one use A to make a general arithmetic of divisibility relations?

(c) In (5.5) it has been shown only that $E \subset P^4$, but it seems safe to conjecture $E = P^4$. Since P^4 is a polynomial algebra with generators of dimensions 16, 32, 40, 24, 64, 72, 40, . . . , it would be enough to exhibit generators in E in these dimensions. Generators in dimension 16, 32, 64, . . . were constructed in §1. By messy calculations I have shown the existence of an M^{24} . Hence in dimensions $k < 40$ we have $E_k = (P^4)_k$. Are there polynomial generators of P_*^4 whose stable tangent bundles are complexifications of real bundles? Here the first unanswered dimension is 24.

(d) Is $\text{Im}(\Omega_*^{Sp} \rightarrow N_*) = P^8$?

REFERENCES

1. P. E. Conner and E. E. Floyd, *Differentiable periodic maps*, Ergebnisse der Math. und ihrer Grenzgebiete, Band 33, Academic Press, New York; Springer-Verlag, Berlin, 1964. MR 31 #750.
2. ———, *The relation of cobordism to K-theories*, Lecture Notes in Math., no. 28, Springer-Verlag, Berlin and New York, 1966. MR 35 #7344.
3. ———, *Torsion in SU-bordism*, Mem. Amer. Math. Soc. No. 60 (1966). MR 32 #6471.
4. W. C. Hsiang and R. H. Szczarba, *On the tangent bundle of a Grassmann manifold*, Amer. J. Math. 86 (1964), 698–704. MR 30 #2523.
5. P. S. Landweber, *Cobordism operations and Hopf algebras*, Trans. Amer. Math. Soc. 129 (1967), 94–110. MR 36 #2145.
6. J. W. Milnor and J. C. Moore, *On the structure of Hopf algebras*, Ann. of Math. (2) 81 (1965), 211–264. MR 30 #4259.
7. S. P. Novikov, *The method of algebraic topology from the viewpoint of cobordism theories*, Izv. Akad. Nauk SSSR Ser. Mat. 31 (1967), 855–951. = Math. USSR Izv. 1 (1967), 827–913. MR 36 #4561.
8. David M. Segal, *Symplectic cobordism of manifolds*, Thesis, Massachusetts Institute of Technology, Cambridge, Mass., 1968; Comment. Math. Helv. 45 (1970), 159–169.
9. R. E. Strong, *Relations among characteristic numbers*. I, II, Topology 4 (1965), 267–281; Topology 5 (1966), 133–148. MR 33 #740; MR 33 #741.
10. ———, *Notes on cobordism theory*, Princeton Univ. Press, Princeton, N. J.; Univ. of Tokyo Press, Tokyo, 1968.
11. ———, *Some remarks on symplectic cobordism*, Ann. of Math. (2) 86 (1967), 425–433. MR 36 #2162.
12. R. Thom, *Quelques propriétés globales des variétés différentiables*, Comment. Math. Helv. 28 (1954), 17–86. MR 15, 890.
13. C. T. C. Wall, *Determination of the cobordism ring*, Ann. of Math (2) 72 (1960), 292–311. MR 22 #11403.
14. Don D. Porter, *Novikov resolutions for symplectic cobordism*, Thesis, Northwestern University, Evanston, Ill., 1969.

UNIVERSITY OF VIRGINIA,
CHARLOTTESVILLE, VIRGINIA 22903