

ON THE EXISTENCE OF STRONGLY SERIES SUMMABLE MARKUSCHEVICH BASES IN BANACH SPACES

BY
WILLIAM B. JOHNSON

Abstract. The main result is: *Let X be a complex separable Banach space. If the identity operator on X^* is the limit in the strong operator topology of a uniformly bounded net of linear operators of finite rank, then X admits a strongly series summable Markushevich basis.*

I. Introduction. Let X be a separable Banach space. A biorthogonal sequence $\{x_i, f_i\}_{i=1}^\infty$ in (X, X^*) is called a Markushevich basis (M -basis) for X provided $\{x_i\}_{i=1}^\infty$ is fundamental in X and $\{f_i\}_{i=1}^\infty$ is total over X . Following Ruckle [8], we say that an M -basis $\{x_i, f_i\}_{i=1}^\infty$ for X is strongly series summable (s.s.s.) provided there exists a set $\{\lambda_{i,n} : i=1, 2, \dots, n; n=1, 2, \dots\}$ of scalars (called a summation matrix for $\{x_i, f_i\}_{i=1}^\infty$) such that, for each x in X , $x = \lim_{n \rightarrow \infty} \sum_{i=1}^n \lambda_{i,n} f_i(x) x_i$. Note that a Schauder basis is a s.s.s. M -basis for which each $\lambda_{i,n}$ can be chosen to be 1.

The results of [8], [9], and [4] indicate that the duality theory of a space which has a s.s.s. M -basis is essentially the same as that of a space which admits a Schauder basis. The reason for this appears to be that if $\{x_i, f_i\}_{i=1}^\infty$ is a s.s.s. M -basis for X with summation matrix $(\lambda_{i,n})$ then, for each f in the coefficient space⁽¹⁾ of the basis, f is the norm limit of $\{\sum_{i=1}^n \lambda_{i,n} f(x_i) f_i\}_{n=1}^\infty$. Thus the adjoints of the "partial sum" operators defined by $T_n(x) = \sum_{i=1}^n \lambda_{i,n} f_i(x) x_i$ also act like partial sum operators. In this respect s.s.s. M -bases behave more like Schauder bases than do such weaker structures as generalized summation bases (see [3]).

In this paper we prove the following rather strong existence theorem for s.s.s. M -bases:

THEOREM 1. *Let X be a separable complex Banach space such that X^* has the λ -metric approximation property for some $\lambda \geq 1$. If Y is a separable subspace of X^* , then there exists a strongly series summable Markushevich basis for X whose coefficient space contains Y .*

If $\lambda \geq 1$, we say that the Banach space X has the λ -metric approximation property (λ -m.a.p.) if there is a net $\{S_d : d \in D\}$ of linear operators of finite rank on X

Received by the editors April 20, 1970.

AMS 1969 subject classifications. Primary 4610.

Key words and phrases. Strongly series summable Markushevich bases, complete biorthogonal sequences, Schauder bases, metric approximation property.

(¹) The coefficient space of an M -basis $\{x_i, f_i\}_{i=1}^\infty$ for X is the norm closure in X^* of the linear span of $\{f_i\}_{i=1}^\infty$.

Copyright © 1971, American Mathematical Society

uniformly bounded by λ which converges pointwise (i.e., in the strong operator topology) to the identity operator on X . Equivalently, X has the λ -m.a.p. provided that, for each finite-dimensional subspace F of X and positive number ε , there is an operator S of finite rank on X such that $\|S\| \leq \lambda$ and $\|S(x) - x\| \leq \varepsilon\|x\|$ for each $x \in F$.

The 1-m.a.p. was introduced by Grothendieck [2] under the name metric approximation property. Grothendieck showed that if X is reflexive and has the (topological) approximation property, then in fact both X and X^* have the 1-m.a.p. This result together with Theorem 1 implies that every separable, reflexive complex Banach space which has the approximation property also admits a s.s.s. M -basis. Of course, it may be that Theorem 1 is always applicable, for it is not even known that there exists a Banach space which does not have the 1-m.a.p.

We use the following notation: X represents a complex Banach space and X^* is the dual to X . The complex assumption is used in an essential way in Lemma 4, and we do not know whether the real version of Theorem 1 is true. I denotes the identity operator on either X or X^* . "Operator" means "bounded linear operator". The range space and null space of an operator, L , are denoted by, respectively, $\mathcal{R}(L)$ and $\ker L$. If L is an operator on X and S is a subspace of X , $L|_S$ denotes the restriction of L to S . The linear span of a subset, A , of a linear space is denoted by $\text{sp } A$. The canonical embedding of X into X^{**} is denoted by " \wedge ".

II. The existence theorem. Our first lemma is both a generalization and a special case of Helly's theorem [11, p. 103].

LEMMA 1. *Let F be a finite-dimensional Banach space, S a finite-dimensional subspace of X^* , L an operator from X^* into F , and $\varepsilon > 0$. There exists a weak*-continuous operator T from X^* into F such that $T|_S = L|_S$ and $\|T\| \leq \|L\| + \varepsilon$.*

Proof. We use the notation of [10] in this proof. We identify the weak*-continuous operators from X^* to F with $X \otimes_\lambda F$ and the operators from X^* to F with $X^{**} \otimes_\lambda F$ [10, p. 30]. Since F is finite dimensional, $X^{**} \otimes_\lambda F$ is thereby identified with $(X \otimes_\lambda F)^{**}$. Now $S \otimes F^*$ is identified with a (finite-dimensional) subspace of $(X \otimes_\lambda F)^*$, so by Helly's theorem [11, p. 103], there is T in $X \otimes_\lambda F$ such that $\|T\| \leq \|L\| + \varepsilon$ and $f(T(s)) = f(L(s))$ for each $s \in S$ and $f \in F^*$. Since F^* is total over F , $T(s) = L(s)$ for each $s \in S$ and hence $T|_S = L|_S$. Q.E.D.

A Banach space X is said to have the λ duality metric approximation property ($\lambda \geq 1$) provided there is a net $\{S_d : d \in D\}$ of operators of finite rank on X uniformly bounded by λ such that $\{S_d : d \in D\}$ converges pointwise to I and $\{S_d^* : d \in D\}$ converges pointwise to I . Equivalently, X has the λ duality m.a.p. provided that, for each $\varepsilon > 0$ and each pair of finite-dimensional subspaces E of X and F of X^* , there is an operator L of finite rank on X such that $\|L\| \leq \lambda$, $\|L(x) - x\| \leq \varepsilon\|x\|$ for each $x \in E$, and $\|L^*(f) - f\| \leq \varepsilon\|f\|$ for each $f \in F$.

LEMMA 2. *Suppose that X^* has the λ -m.a.p. Then X has the λ duality m.a.p.*

Proof. Using the hypothesis and Lemma 1, we can construct a net $\{S_d : d \in D\}$ of operators of finite rank on X uniformly bounded by λ such that $\{S_d^* : d \in D\}$ is pointwise convergent on X^* to I . For each $x \in X$, the net $\{S_d(x) : d \in D\}$ weakly converges to x , hence (cf., e.g., [1, p. 477]) there is a net $\{T_e : e \in E\}$ of operators on X such that $\{T_e : e \in E\}$ is pointwise convergent on X to I ; each T_e is a convex combination of a subset $\{S_{e(i)}\}_{i=1}^{n_e}$ of $\{S_d : d \in D\}$; and for each $d \in D$ there is $e' \in E$ such that if $e \geq e'$ then $e(i) \geq d$ for $i=1, 2, \dots, n_e$. Thus $\{T_e : e \in E\}$ is uniformly bounded by λ and $\{T_e^* : e \in E\}$ is pointwise convergent on X^* to I . Q.E.D.

The proof of the next lemma is suggested by the proof of Lemma 3.1 of [5].

LEMMA 3. *Suppose that X has the λ duality m.a.p., E is a finite-dimensional subspace of X , F is a finite-dimensional subspace of X^* , and $\varepsilon > 0$. Then there exists an operator L of finite rank on X such that $\|L\| \leq \lambda + \varepsilon$, $L_{|E} = I_{|E}$, and $L_{|F}^* = I_{|F}$.*

Proof. Let $n = \dim E$ and $m = \dim F$. Choose $1 > \beta > 0$ small enough so that $\beta + \beta m(\lambda + \beta)/(1 - \beta) \leq \varepsilon$ and choose $1 > \alpha > 0$ small enough so that $(n\alpha/(1 - \alpha))\lambda \leq \beta/2$.

Let M be an operator of finite rank on X such that $\|M\| \leq \lambda$ and, for each $x \in E$ and $f \in F$,

- (1) $\|x - M(x)\| \leq \alpha\|x\|$ and
- (2) $\|f - M^*(f)\| \leq \beta/2\|f\|$.

By (1), for each $x \in E$, $(1 - \alpha)\|x\| \leq \|M(x)\|$, hence $M_{|E}$ has an inverse, Q , satisfying $\|Q\| \leq 1/(1 - \alpha)$. Also, for each $y \in M[E]$, $\|Q(y) - y\| \leq (\alpha/(1 - \alpha))\|y\|$.

Let P be a projection of X onto $M[E]$ such that $\|P\| \leq n$. Let $N = QPM + (I - P)M$. Clearly N has finite rank and $N_{|E} = I_{|E}$. Now if $x \in X$,

$$\|N(x) - M(x)\| = \|QPM(x) - PM(x)\| \leq \frac{\alpha}{1 - \alpha} \|P\| \|M\| \|x\| \leq \frac{n\alpha\lambda}{1 - \alpha} \|x\|.$$

Thus $\|N - M\| \leq n\alpha\lambda/(1 - \alpha)$, from which it follows that $\|N^*\| \leq \lambda + \beta/2$ and $\|N^* - M^*\| \leq \beta/2$. This last inequality and (2) imply that, for each $f \in F$, $\|N^*(f) - f\| \leq \beta\|f\|$.

As in the first part of the proof, we have that $N_{|F}^*$ is an isomorphism with inverse, Q' , satisfying $\|Q'(f) - f\| \leq (\beta/(1 - \beta))\|f\|$ for each $f \in N^*[F]$. Let P' be a projection of X^* onto $N^*[F]$ such that $\|P'\| \leq m$ and let $L^* = Q'P'N^* + (I - P')N^*$. (Note that L^* is indeed weak*-continuous because N^* is weak*-continuous and has finite rank.) Then $L_{|F}^* = I_{|F}$ and, for each $f \in X^*$,

$$\|L^*(f) - N^*(f)\| \leq \frac{\beta}{1 - \beta} \|P'\| \|N^*\| \|f\| \leq \frac{\beta m(\lambda + \beta)}{1 - \beta} \|f\|.$$

Thus $\|L^*\| \leq \lambda + \beta + \beta m(\lambda + \beta)/(1 - \beta) \leq \lambda + \varepsilon$.

Since $\|L\| = \|L^*\|$, it remains to be seen only that $L_{|E} = I_{|E}$. Let $x \in E$ and suppose that $f \in X^*$. Then using the fact that $x = N(x)$, we have

$$\begin{aligned} f(L(x)) &= L^*(f)(x) = L^*(f)(N(x)) \\ &= N^*Q'P'N^*(f)(x) + N^*(I - P')N^*(f)(x) \\ &= P'N^*(f)(x) + f(N(N(x))) - P'N^*(f)(N(x)) = f(x). \end{aligned}$$

Since X^* is total over E , $L(x) = x$. Q.E.D.

LEMMA 4. Let $\{x_i, f_i\}_{i=1}^n$ be a finite biorthogonal set in (X, X^*) , let T be an operator of finite rank on X such that $T(x_i) = x_i$ and $T^*(f_i) = f_i$ for $i = 1, 2, \dots, n$, and let $\varepsilon > 0$. Then there exists a finite biorthogonal set $\{x_i, f_i\}_{i=n+1}^{n+m}$ in (X, X^*) and a set $\{\lambda_i\}_{i=1}^m$ of complex numbers such that $\{x_i, f_i\}_{i=1}^{n+m}$ is biorthogonal and $\|L - T\| \leq \varepsilon$, where L is the operator on X defined by

$$L(x) = \sum_{i=1}^n f_i(x)x_i + \sum_{i=n+1}^{n+m} \lambda_{i-n} f_i(x)x_i.$$

Proof. Define a projection U on X by $U(x) = \sum_{i=1}^n f_i(x)x_i$ and let $X_0 = \mathcal{R}(I - U)$. Note that $TU = UT = U$, so $T[X_0] \subset X_0$ and $\ker T \subset X_0$. Let P be a projection of finite rank on X_0 such that $PT(I - U) = TP(I - U) = T(I - U)$. (For example, choose $\ker P$ to be a closed complement in $\ker T$ to $\mathcal{R}(T) \cap \ker T$ and choose $\mathcal{R}(P)$ to be a complement in X_0 to $\ker P$ which contains $\mathcal{R}(T) \cap X_0$.)

Let $m = \dim \mathcal{R}(P)$ and choose a basis $\{z_i\}_{i=1}^m$ for $\mathcal{R}(P)$ such that the matrix representation $(\alpha_{ij})_{i,j=1}^m$ of $T|_{\mathcal{R}(P)}$ with respect to $\{z_i\}_{i=1}^m$ is lower triangular—i.e., $\alpha_{ij} = 0$ if $j > i$. Now pick a sequence $\{\lambda_i\}_{i=1}^m$ of pairwise distinct complex numbers sufficiently close to $\{\alpha_{ii}\}_{i=1}^m$ so that

$$\|Q - T|_{\mathcal{R}(P)}\| \leq \varepsilon / \|P\| \|I - U\|,$$

where Q is the operator on $\mathcal{R}(P)$ whose matrix representation, (β_{ij}) , with respect to $\{z_i\}_{i=1}^m$ is given by

$$\begin{aligned} \beta_{ij} &= \lambda_i & \text{if } i = j, \\ &= \alpha_{ij} & \text{if } i \neq j. \end{aligned}$$

Since (β_{ij}) is lower triangular, $\{\lambda_i\}_{i=1}^m$ is the set of eigenvalues for Q . The λ_i 's are distinct, so there is a basis $\{x_i\}_{i=n+1}^{n+m}$ for $\mathcal{R}(P)$ such that $Q(x_i) = \lambda_{i-n} x_i$ for $i = n + 1, \dots, n + m$. Picking $\{f_i\}_{i=n+1}^{n+m}$ in $\mathcal{R}([P(I - U)]^*)$ biorthogonal to $\{x_i\}_{i=n+1}^{n+m}$, we have that, for each $x \in \mathcal{R}(P)$, $Q(x) = \sum_{i=n+1}^{n+m} \lambda_{i-n} f_i(x)x_i$.

Now $\{x_i, f_i\}_{i=1}^{n+m}$ is biorthogonal and if L is defined by

$$L(x) = \sum_{i=1}^n f_i(x)x_i + \sum_{i=n+1}^{n+m} \lambda_{i-n} f_i(x)x_i,$$

then clearly $L = TU + QP(I - U)$. Thus

$$\begin{aligned} \|L - T\| &= \|TU + QP(I - U) - TU - TP(I - U)\| \\ &\leq \|Q - T|_{\mathcal{R}(P)}\| \|P\| \|I - U\| \leq \varepsilon. \end{aligned} \qquad \text{Q.E.D.}$$

Proof of Theorem 1. Let λ be such that X has the λ duality m.a.p. (Lemma 2). Let $\{z_i\}_{i=1}^\infty$ be fundamental in X and let $\{g_i\}_{i=1}^\infty$ be a subset of X^* such that Y is contained in the closure of the linear span of $\{g_i\}_{i=1}^\infty$. Assume, without loss of generality, that $\|z_1\| = \|g_1\| = g_1(z_1) = 1$. We define the desired s.s.s. M -basis $\{x_i, f_i\}_{i=1}^\infty$ for X and a summability matrix $(\lambda_{i,n})$ for $\{x_i, f_i\}_{i=1}^\infty$ by induction. Set $k(1) = 1$, $x_1 = z_1$, $f_1 = g_1$, $\lambda_{1,1} = 1$. Now suppose $k(m)$, $\{x_i, f_i\}_{i=1}^{k(m)}$, and

$\{\lambda_{i,n} : i \leq n; n = 1, 2, \dots, k(m)\}$ have been defined. Extend $\{x_i, f_i\}_{i=1}^{k(m)}$ to a biorthogonal set $\{x_i, f_i\}_{i=1}^j$ ($j = k(m), k(m) + 1$, or $k(m) + 2$) so that $z_{m+1} \in \text{sp } \{x_i\}_{i=1}^j$ and $g_{m+1} \in \text{sp } \{f_i\}_{i=1}^j$ (cf., e.g., the proof of Theorem III.1 in [3]). Now by Lemma 3 and Lemma 4 there are a positive integer $k(m+1) \geq j$, a biorthogonal set $\{x_i, f_i\}_{i=j+1}^{k(m+1)}$ in (X, X^*) , and complex numbers $\{\alpha_i\}_{i=j+1}^{k(m+1)}$ such that $\{x_i, f_i\}_{i=1}^{k(m+1)}$ is biorthogonal and if T is defined on X by $T(x) = \sum_{i=1}^j f_i(x)x_i + \sum_{i=j+1}^{k(m+1)} \alpha_i f_i(x)x_i$, then $\|T\| \leq \lambda + 1/m$. We complete the induction by defining

$$\begin{aligned} \lambda_{i,n} &= \lambda_{i,k(m)} && \text{if } i \leq k(m) < n < k(m+1), \\ &= 0 && \text{if } k(m) < i \leq n < k(m+1), \\ &= 1 && \text{if } i \leq k(m) \text{ and } n = k(m+1), \\ &= \alpha_i && \text{if } k(m) < i \leq k(m+1) = n. \end{aligned}$$

It is easy to check that $\{x_i, f_i\}_{i=1}^\infty$ has the desired properties. Q.E.D.

REMARK 1. Suppose that X is separable and X^* has the λ -m.a.p. for some λ . Theorem 1 shows that there are s.s.s. M -bases for X whose coefficient spaces are "arbitrarily large". One might guess that if Y is a separable subspace of X^* and Y contains a subspace which is the coefficient space of some s.s.s. M -basis for X , then Y is itself the coefficient space for some s.s.s. M -basis for X , because the corresponding statement for generalized summation bases is true (cf. [3, proof of Theorem IV.1]). This is not the case: Let $X = l_1$. It is a rather easy consequence of Theorem 4.3 of [6] that the coefficient space of any s.s.s. M -basis for l_1 is an \mathcal{L}_∞ space in the sense of [6]. Simply pick Y to be a separable subspace of $l_\infty (= l_1^*)$ which contains c_0 but is not an \mathcal{L}_∞ space. (For example, Y can be the closed span of $c_0 \cup K$, where K is a subspace of l_∞ isomorphic to l_2 . It follows from Theorem 1 of [7] that Y is isomorphic to $c_0 \oplus l_2$ and is thus not an \mathcal{L}_∞ space.)

Recall that an M -basis $\{x_i, f_i\}_{i=1}^\infty$ for X whose coefficient space is X^* is called shrinking (see [3]). Now if $\{x_i, f_i\}_{i=1}^\infty$ is a s.s.s. M -basis then the remarks in the introduction show that $\{f_i, \hat{x}_i\}_{i=1}^\infty$ is a s.s.s. M -basis for the coefficient space of the basis. Thus a shrinking M -basis $\{x_i, f_i\}_{i=1}^\infty$ which is s.s.s. is also shrinking as a s.s.s. M -basis in the sense that $\{f_i, \hat{x}_i\}_{i=1}^\infty$ is a s.s.s. M -basis for X^* . In view of Theorem 1, we thus have

COROLLARY 1. *If X^* is separable and has the λ -m.a.p. for some $\lambda \geq 1$, then X admits a shrinking s.s.s. M -basis.*

Let us say that a s.s.s. M -basis $\{x_i, f_i\}_{i=1}^\infty$ is *boundedly complete* provided there is a summability matrix $(\lambda_{i,n})$ for $\{x_i, f_i\}_{i=1}^\infty$ such that for every sequence $\{t_i\}_{i=1}^\infty$ of scalars, if $\{\sum_{i=1}^n t_i \lambda_{i,n} x_i\}_{n=1}^\infty$ is bounded then it is convergent. A simple modification of Theorem II.3 of [3] shows that a s.s.s. M -basis is *boundedly complete* if and only if it is boundedly complete as an M -basis in the sense of [3] (and thus in the above definition of *boundedly complete* "there is a summability matrix" can be replaced by "for each summability matrix"). Thus using Corollary 1 and the results of [3] we have

THEOREM 2. *X admits a boundedly complete s.s.s. M-basis if and only if X has the λ -m.a.p. for some $\lambda \geq 1$ and X is isomorphic to a separable conjugate Banach space.*

We conclude with a conjecture which, by Theorem 2, has an affirmative answer if X is a conjugate space:

CONJECTURE 1. *If X is separable and has the λ -m.a.p. for some $\lambda \geq 1$, then X admits a s.s.s. M-basis.*

REFERENCES

1. N. Dunford and J. T. Schwartz, *Linear operators. I: General theory*, Pure and Appl. Math., vol. 7, Interscience, New York, 1958. MR 22 #8302.
2. A. Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc. No. 16 (1955). MR 17, 763.
3. W. B. Johnson, *Markushevich bases and duality theory*, Trans. Amer. Math. Soc. **149** (1970), 171–177.
4. ———, *Markushevich bases and duality theory. II* (in preparation).
5. J. Lindenstrauss, *Extension of compact operators*, Mem. Amer. Math. Soc. No. 48 (1964). MR 31 #3828.
6. J. Lindenstrauss and H. P. Rosenthal, *The \mathcal{L}_p spaces*, Israel J. Math. **8** (1970).
7. H. P. Rosenthal, *On totally incomparable Banach spaces*, J. Functional Analysis **4** (1969), 167–175. MR 40 #1758.
8. W. H. Ruckle, *Representation and series summability of complete biorthogonal sequences*, Pacific J. Math. **34** (1970), 511–528.
9. ———, *The tensor product of complete biorthogonal sequences*, Clemson University Technical Report #25, Clemson, S.C., 1969.
10. R. Schatten, *A theory of cross-spaces*, Ann. of Math. Studies, no. 26, Princeton Univ. Press, Princeton, N. J., 1950. MR 12, 186.
11. A. Wilanski, *Functional analysis*, Blaisdell, Waltham, Mass., 1964. MR 30 #425.

UNIVERSITY OF HOUSTON,
HOUSTON, TEXAS 77004