

ADDENDUM TO
"BEHNKE-STEIN THEOREM FOR ANALYTIC SPACES"

BY

ALESSANDRO SILVA

ABSTRACT. A very simple argument shows that Theorem 3.1 in my paper *Behnke-Stein theorem for analytic spaces*, (these Transactions, 199 (1974), pp. 317-326) is enough, via a Narasimhan result, to obtain information about the torsion of the homology groups of a Runge pair of Stein spaces.

Let (X, Y) , with $Y \subset X$, Y open, be a pair of reduced complex analytic spaces of (complex) dimension n . Andreotti and Narasimhan proved in [1], among many others, the following result:

(1.1) If (X, Y) is a Runge pair of Stein spaces (a 1-Runge pair in the terminology of [3]) and every singularity of X outside Y is isolated, then

$$H_r(X \text{ mod } Y, \mathbb{Z}) = 0$$

for $r \geq n + 1$.

We wish to show the following statements (1.2) and (1.3),

(1.2) (Narasimhan [2, Theorem 3]): if X is a Stein space, then:

$$H_r(X, \mathbb{Z}) = 0 \quad \text{for } r \geq n + 1$$

and

$H_n(X, \mathbb{Z})$ is without torsion;

(1.3) (Silva [3, Theorem 3.1]): if (X, Y) is a Runge pair of Stein spaces (or, equivalently, in the terminology of [3], a 1-Runge pair of cohomologically 1-complete spaces) then

$$H_{n+1}(X \text{ mod } Y, \mathbb{C}) = 0;$$

make us able to remove from (1.1) the assumption that the singularities of X outside Y are isolated.

Indeed, if we write the exact homology sequence for the pair (X, Y) :

Received by the editors December 9, 1974.

AMS (MOS) subject classifications (1970). Primary 32E15.

Key words and phrases. Runge pairs.

Copyright © 1975, American Mathematical Society

$$\cdots \rightarrow H_r(X, \mathbf{Z}) \rightarrow H_r(X \bmod Y, \mathbf{Z}) \xrightarrow{\varphi_r} H_{r-1}(Y, \mathbf{Z}) \rightarrow \cdots$$

from (1.2) we obtain that $H_r(X \bmod Y, \mathbf{Z}) = 0$ for $r > n + 1$. Suppose now $r = n + 1$. (1.3) implies that $H_{n+1}(X \bmod Y, \mathbf{Z})$ is a torsion group. If we look again at the exact homology sequence for (X, Y) we see that the morphism:

$$\varphi_{n+1}: H_{n+1}(X \bmod Y, \mathbf{Z}) \rightarrow H_n(Y, \mathbf{Z})$$

is injective, so that, $H_n(Y, \mathbf{Z})$ being without torsion, we must have $H_{n+1}(X \bmod Y, \mathbf{Z}) = 0$.

In conclusion we have obtained the following

THEOREM. *If (X, Y) is a Runge pair of Stein spaces, then*

$$H_r(X \bmod Y, \mathbf{Z}) = 0,$$

for $r \geq n + 1$.

BIBLIOGRAPHY

1. A. Andreotti and R. Narasimhan, *A topological property of Runge pairs*, Ann. of Math. (2) **76** (1962), 499–509.
2. R. Narasimhan, *On the homology groups of Stein spaces*, Invent. Math. **2** (1967), 377–385. MR 35 #7356.
3. A. Silva, *Behnke-Stein theorem for analytic spaces*, Trans. Amer. Math. Soc. **199** (1974), 317–326.

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON,
NEW JERSEY 08540