

MAXIMAL CHAINS OF PRIME IDEALS
IN INTEGRAL EXTENSION DOMAINS. I

BY

L. J. RATLIFF, JR. AND S. McADAM⁽¹⁾

ABSTRACT. Let (R, M) be a local domain, let k be a positive integer, and let Q be a prime ideal in $R_k = R[X_1, \dots, X_k]$ such that $MR_k \subset Q$. Then the following statements are equivalent: (1) There exists an integral extension domain of R which has a maximal chain of prime ideals of length n . (2) There exists a minimal prime ideal z in the completion of R such that $\text{depth } z = n$. (3) There exists a minimal prime ideal w in the completion of $(R_k)_Q$ such that $\text{depth } w = n + k - \text{depth } Q$. (4) There exists an integral extension domain of $(R_k)_Q$ which has a maximal chain of prime ideals of length $n + k - \text{depth } Q$. (5) There exists a maximal chain of prime ideals of length $n + k - \text{depth } Q$ in $(R_k)_Q$. (6) There exists a maximal chain of prime ideals of length $n + 1$ in $R[X_1]_{(M, X_1)}$.

1. Introduction. In [6, Theorem 3.6], it is shown that if (R, M) is a local domain, then $R_1 = R[X_1]$ is catenary if and only if $R_k = R[X_1, \dots, X_k]$ is catenary, for all $k \geq 0$. This result is closely related to the equivalence of the following statements (see (1.1) for the definitions): R satisfies the second chain condition for prime ideals (s.c.c.); R satisfies the altitude formula; and, R is quasi-unmixed [6, Theorem 3.6]. Thus it is of some interest.

The original goal of the authors in starting this paper was to see how the lengths of maximal chains of prime ideals in R_1 compare to the lengths of maximal chains of prime ideals in R_k (without assuming that R_1 is catenary). The answer we obtained is included in the result mentioned at the start of this paper, and, as is seen by that result, it is closely related to the lengths of maximal chains of prime ideals in integral extension domains of R and to the depths of the minimal prime ideals in the completion of R . (This result is given in (2.3), (2.5), (2.9), and, more specifically (2.14).)

Received by the editors May 28, 1975.

AMS (MOS) subject classifications (1970). Primary 13A15, 13B20, 13C15; Secondary 13B25.

Key words and phrases. Altitude formula, catenary ring, catenary chain conjecture, chain condition for prime ideals, chain conjecture, completion of a local ring, depth conjecture, first chain condition for prime ideals, integral extension, local ring, maximal chain of prime ideals, Noetherian ring, polynomial extension ring, quasi-unmixed local ring, second chain condition for prime ideals, semilocal ring, unmixed local ring, upper conjecture.

⁽¹⁾ Research supported in part by NSF-MCS 76-06009.

Copyright © 1976, American Mathematical Society

After the main theorem (the above six equivalent statements) is proved, the rest of the paper is devoted to showing that a large number of important known results follow as easy corollaries. Among these are: (a) R satisfies the s.c.c. if and only if some $(R_k)_Q$ is catenary if and only if all $(R_k)_Q$ satisfy the s.c.c. (2.16) and (2.17); (b) R satisfies the s.c.c. if and only if R is quasi-unmixed (2.19); (c) if R satisfies the s.c.c., then, for each prime ideal p in R , every local domain which is a locality over R/p satisfies the s.c.c. (2.20) (in particular, this holds for R/p and R_p (2.26)); (d) the above mentioned [6, Theorem 3.6] (2.18); (e) two equivalences to the upper conjecture (2.22) (see (2.21)); and, (f) the relationship between lengths of maximal chains of prime ideals in $(R_k)_Q$ and in $R[X_1]_{(M, X_1)}$ (2.9.2) (our original goal). (A number of quite deep theorems are used to prove the equivalence of the above six statements, so the authors do not claim that this method affords easier proofs than those originally given for the above-mentioned corollaries. However, once it is proved that the above six statements are equivalent, the corollaries are nearly obvious.)

All rings in this paper are assumed to be commutative rings with an identity element. The undefined terminology is the same as that in [5]. We mention, in particular, that $A \subset B$ means that the set A is a proper subset of the set B .

We close this introduction with the following less well-known definitions, since they are frequently used in the remainder of this paper.

(1.1) **DEFINITIONS.** Let A be an integral domain, and let $a = \text{altitude } A < \infty$.

(1.1.1) A satisfies the *first chain condition for prime ideals* (f.c.c.) in case every maximal chain of prime ideals in A has length $= a$.

(1.1.2) A is *catenary* in case, for each pair of prime ideals $P \subset Q$ in A , $(A/P)_{Q/P}$ satisfies the f.c.c.

(1.1.3) A satisfies the *second chain condition for prime ideals* (s.c.c.) in case every integral extension domain of A satisfies the f.c.c.

(1.1.4) A satisfies the *chain condition for prime ideals* (c.c.) in case, for each pair of prime ideals $P \subset Q$ in A , $(A/P)_{Q/P}$ satisfies the s.c.c.

(1.1.5) A is *quasi-unmixed* (resp., *unmixed*) in case A is semilocal and every minimal (resp., every) prime divisor of zero in the completion of A has depth $= a$.

(1.1.6) A satisfies the *altitude formula* in case, for each finitely-generated integral domain B over A and for each prime ideal P in B , altitude $B_P + \text{trd}(B/P)/(A/(P \cap A)) = \text{altitude } A_{P \cap A} + \text{trd } B/A$, where $\text{trd } C/D$ denotes the transcendence degree of the quotient field of the integral domain C over the quotient field of its subdomain D .

A number of facts concerning these definitions are given in [8, Remarks

2.22–2.25 and 3.7–3.8]. We mention at this point only that, for a local domain R , R is catenary if and only if R satisfies the f.c.c.

2. Main results. We will be greatly concerned with the lengths of maximal chains of prime ideals in a ring. Therefore, since we use the expression so often, we adopt the following abbreviation.

(2.1) **DEFINITION.** It will be said that a ring A has a *mcpil* n in case there exists a maximal chain of prime ideals of length n in A (that is, a chain of prime ideals $p_0 \subset p_1 \subset \cdots \subset p_n$ such that p_0 is minimal, p_n is maximal, and height $p_i/p_{i-1} = 1$ ($i = 1, \dots, n$)).

The following lemma combines, for future reference, two known and quite useful results [1, Corollary 4] and [2, Theorem 1.10]. Only the last sentence of the lemma is new.

(2.2) **LEMMA (CF. [1, COROLLARY 4] AND [2, THEOREM 1.10]).** *Let $A \subseteq B$ be integral domains such that A is Noetherian and B is integral over A . Assume that B has a mcpil n . Then there exists a principal integral extension domain $C = A[c]$ of A which has a mcpil n , say $(0) \subset p_1 \subset \cdots \subset p_n$, such that height $p_i = i = \text{height } p_i \cap A$ ($i < n$). Further, c may be chosen so as to satisfy either of the conditions: $c \in p_n$; or, $c \notin p_n$.*

PROOF. Let N be a maximal ideal in B such that there exists a mcpil n in B_N , and let $M = N \cap A$. Then it follows from [2, Theorem 1.10] (applied to A_M) that there exists a finite integral extension domain D of A which has a mcpil n , say $(0) \subset Q_1 \subset \cdots \subset Q_n$. Therefore, there exists a maximal chain of prime ideals $(0) \subset q_1 \subset \cdots \subset q_{n-1} \subset q_n = Q_n$ in D such that height $q_i = i = \text{height } q_i \cap A$ ($i < n$) [1, Corollary 4]. Let $c \in q_n$ such that c is not in any other maximal ideal in D which lies over $q_n \cap A$. Then it is readily seen that $C = A[c]$ has a mcpil n (namely, $(0) \subset p_1 \subset \cdots \subset p_{n-1} \subset p_n$, where $p_i = q_i \cap C$) such that height $p_i = i = \text{height } p_i \cap A$ ($i < n$) and $c \in p_n$. Then also, $C = A[c - 1]$ has a mcpil n and $c - 1 \notin p_n$. Q.E.D.

The following result is the first part of the main theorem in this paper.

(2.3) **THEOREM.** *Let R be a semilocal domain, and let n be a positive integer. Then there exists an integral extension domain D of R which has a mcpil n if and only if there exists a minimal prime ideal z in the completion R^* of R such that depth $z = n$.*

PROOF. If R is local, then it is known [6, Proposition 3.5] that there exists a height one maximal ideal in the integral closure R' of R if and only if there exists a depth one minimal prime ideal in R^* . Therefore, for the semilocal case, there exists a depth one minimal prime ideal in R^* if and only if there exists a depth one minimal prime ideal in R_M^* , for some maximal ideal M^* in

R^* , if and only if, by [6, Proposition 3.5], there exists a height one maximal ideal in the integral closure of $R_{M^* \cap R}$ (since the completion of $R_{M^* \cap R}$ is $R_{M^*}^*$) if and only if there exists a height one maximal ideal in R' . Therefore, assume that $n > 1$, and assume that the lemma holds for $n - 1$.

Assume first that such a D exists. Then, by (2.2), it may be assumed that $D = R[c]$ and $(0) \subset Q_1 \subset \cdots \subset Q_n$ is a mcpil n in D such that height $Q_i = i = \text{height } Q_i \cap R$ ($i < n$). Let $q = Q_1 \cap R$, so $D/Q_1 = (R/q)[c']$ (with $c' = c + Q_1$) has a mcpil $n - 1$. Therefore, by the case $n - 1$, there exists a depth $n - 1$ minimal prime ideal in the completion $(R/q)^* \cong R^*/qR^*$ of R/q . Therefore qR^* has a minimal prime divisor of depth $n - 1$, say q^* , and height $q^* = \text{height } q = 1$. Then there exists a minimal prime ideal z in R^* such that $z \subset q^*$, so depth $z = n$ (since R^*/z is a complete local domain, so is catenary (1.1.2)), hence $n = 1 + \text{depth } q^* = \text{height } q^*/z + \text{depth } q^*/z = \text{altitude } R^*/z = \text{depth } z$.

Conversely, assume that z is a depth n minimal prime ideal in R^* ($n > 1$). Then there exists a height one prime ideal p^* in R^* such that $z \subset p^*$ and $p = p^* \cap R \neq (0)$. (For, there exist only a finite number of prime ideals p^* in R^* such that $z \subset p^*$ and $\text{height } p^*/z = 1 < \text{height } p^*$ [1, Theorem 1], so let p_1^*, \dots, p_k^* be these. (No p_i^* is maximal, since $n > 1$ and R^*/z is *local*.) Then, with J the Jacobson radical of R , $J \not\subseteq \bigcup(p_i^* \cap R)$, so let $b \in J, \notin \bigcup(p_i^* \cap R)$. Then $(z, b)R^*$ has a height one prime divisor p^* (by the choice of b and the principal ideal theorem in R^*/z), so $p^* \supset z$ and $p^* \cap R \neq (0)$.) Then p^* is a minimal prime divisor of pR^* ; hence height $p = 1$ [5, (22.9)]. Now, as above, depth $p^* = \text{depth } z - 1 = n - 1$. Now, if depth $p = n - 1$, then R has a mcpil n . Therefore assume that depth $p > n - 1$ ($\text{depth } p \geq \text{depth } p^*$). Then in $(R/p)^* \cong R^*/pR^*$ there exists a depth $n - 1$ minimal prime ideal (namely, p^*/pR^*), so, by the case $n - 1$ and (2.2), there exists an integral extension domain $D' = (R/p)[c']$ of R/p which has a mcpil $n - 1$. Thus there exists an integral extension domain $D = R[c]$ of R which has a prime ideal q such that $q \cap R = p$ and such that $D/q = D'$. Then height $q = 1$, since height $p = 1$, so D has a mcpil n (since D' has a mcpil $n - 1$). Q.E.D.

(2.4) REMARK. The case $n = 1$ of (2.3) was essentially proved in [6, Proposition 3.5], as was mentioned at the start of the proof of (2.3). A hoped for generalization of the case $n = 1$ is:

There exists a depth n minimal prime ideal in the completion R^* of R
 $(*)$ if and only if there exists a height n maximal ideal in the integral closure R' of R .

This generalization is, in fact, equivalent to the *chain conjecture* (that is, the integral closure of a semilocal domain satisfies the c.c. (1.1.4)).

PROOF. Assume first that the chain conjecture holds. If there exists a depth n minimal prime ideal in R^* , then, by (2.3), there exists an integral extension

domain D of R which has a mcpil n . Then $D[R']$ is an integral extension domain of R' which has a mcpil n (by the going-up theorem), so, since R' satisfies the c.c., there exists a height n maximal ideal in R' . (And, if there exists a height n maximal ideal in R' , then there exists a depth n minimal prime ideal in R^* , by (2.3).) Hence $(*)$ holds.

For the converse, to prove that the chain conjecture holds, it suffices to prove that if H is a Henselian local domain, then H satisfies the s.c.c. (1.1.3) [9, (2.4)]. For this, if $(*)$ holds, then, for each minimal prime ideal z in the completion of H , $\text{depth } z = \text{altitude } H$ (since there is only one maximal ideal in the integral closure of H). Therefore, H is quasi-unmixed (1.1.5); hence H satisfies the s.c.c. [6, Theorem 3.1]. Q.E.D.

To prove the second part of our main theorem we need two facts. The first of these is that a transcendental extension domain of a Noetherian domain A satisfies the altitude formula relative to A [11, Proposition 2, p. 326].

Also, to prove (2.5), we need the following result: If C is a class of local domains R such that each $R \in C$ is a locality over some complete local domain, then each $R \in C$ is analytically unramified and unmixed (1.1.5). This follows from [3, Proposition 4]. (It is straightforward to check that the hypotheses of [3, Proposition 4] are satisfied by C .)

(2.5) THEOREM. *Let (R, M) be a local ring, let n and k be positive integers, and let Q be a prime ideal in $R_k = R[X_1, \dots, X_k]$ such that $Q \cap R = M$. Then there exists a depth n minimal prime ideal in the completion (R^*, M^*) of R if and only if there exists a minimal prime ideal w in the completion L^* of $L = (R_k)_Q$ such that $\text{depth } w = n + k - \text{depth } Q$.*

PROOF. Let $I = R^*[X_1, \dots, X_k]$. Then, by [7, Lemma 3.2], $Q^* = QI$ is a prime ideal and L is a dense subspace of $S = I_{Q^*}$, so $S^* = L^*$, where S^* is the completion of S . Further, $\text{depth } Q^* = \text{depth } Q$, since $I/Q^* = R_k/Q$.

Assume first that z is a depth n minimal prime ideal in R^* . Then zS is a minimal prime ideal in S and $\text{depth } zS = n + k - \text{depth } Q^*$, as will now be shown. Namely, $zI \subseteq M^*I \subseteq Q^*$, so Q^*/zI is a prime ideal in $I/zI \cong (R^*/z)[X_1, \dots, X_k]$ which lies over M^*/z , so $\text{height } Q^*/zI + \text{trd}(I/Q^*)/(R^*/M^*) = \text{height } M^*/z + k$, by the altitude formula (1.1.6) [11, Proposition 2, p. 326]; that is, $\text{depth } zS = \text{depth } z + k - \text{depth } Q^*$, since $\text{depth } Q^* = \text{altitude } I/Q^* = \text{trd}(I/Q^*)/(R^*/M^*)$ [5, (14.6)] (since R^*/M^* is a field). Therefore, there exists a minimal prime ideal w in $L^* = S^*$ such that $\text{depth } w = n + k - \text{depth } Q$, since $\text{depth } Q = \text{depth } Q^*$, by the preceding paragraph. (Any minimal prime divisor w of $(zS)L^*$ will do, by [3, Proposition 4].)

Conversely, let w be a minimal prime ideal in $L^* = S^*$, and let $\text{depth } w = d$. Let $q = w \cap S$, so q is a minimal prime ideal in S and w is a minimal prime

divisor of qS^* . Hence, since S/q is unmixed, by the comment preceding this theorem, $\text{depth } q = \text{depth } w = d$. Also, $z = q \cap R^*$ is a minimal prime ideal and $q = zS$. Therefore, by the preceding paragraph, $d = \text{depth } q = \text{depth } zS = \text{depth } z + k - \text{depth } Q$. Q.E.D.

(2.6) COROLLARY. *Let (R, M) , k , Q , and L be as in (2.5), and assume that R is an integral domain. Then there exists an integral extension domain of R which has a mcpil n if and only if there exists an integral extension domain of L which has a mcpil $n + k - \text{depth } Q$.*

PROOF. This follows immediately from (2.3) and (2.5). Q.E.D.

It follows from (2.6) that there exists an integral extension domain of R which has a mcpil n if and only if there exists an integral extension domain of $R[X_1, \dots, X_k]_{(M, X_1, \dots, X_k)}$ which has a mcpil $n + k$. This fact will be considerably sharpened in (2.14) below. For the present, it suffices to emphasize that (2.6) depends only on $\text{depth } Q$, and not on the ideal Q itself. It should also be noted, in particular, that, by (2.6), if there exists a mcpil m in $(R_k)_Q$ ($MR_k \subset Q$), then there exists an integral extension domain of R which has a mcpil $m - k + \text{depth } Q$.

We need one further result to complete the proof of our main theorem. The needed result is a weak version of a result given in [10, (3.1)]. However, since a proof of the weak version of the result is not long, we include it at this point.

(2.7) PROPOSITION. *Let (R, M) and (S, N) be local domains such that S is a finite integral extension of R . Then there exists a mcpil n in R if and only if there exists a mcpil n in S .*

PROOF. Assume that there exists a mcpil n in S . Then, by [1, Corollary 4], there exists a mcpil n in S , say $(0) \subset p_1 \subset \dots \subset p_n = N$ such that $\text{height } p_i = i = \text{height } p_i \cap R$ ($i < n$). Then $(0) \subset p_1 \cap R \subset \dots \subset p_{n-1} \cap R \subset N \cap R = M$ is a mcpil n in R , since $\text{depth } p_{n-1} \cap R = \text{depth } p_{n-1} = 1$.

The converse follows from the going-up theorem [5, (10.9)]. Q.E.D.

It is now expedient to introduce notation for the set of lengths of maximal chains of prime ideals in localizations of transcendental extensions of a ring. We do so in the following definition.

(2.8) DEFINITION. Let A be a ring, k a nonnegative integer and Q a prime ideal in $A_k = A[X_1, \dots, X_k]$ ($A_0 = A$). Then $W(k, Q) = \{n; \text{there exists a mcpil } n \text{ in } (A_k)_Q\}$.

The following theorem completes the proof of the main result in this paper. (The main theorem is summarized in (2.14) below.) A number of important known results follow easily from it, as will be seen in the subsequent corollaries.

(2.9.1) was proved in [1, Theorem 6]. We include it here for completeness.

(2.9) THEOREM. *Let (R, M) be a local domain, let k be a positive integer, and let Q be a prime ideal in R_k such that $Q \cap R = M$. Then the following statements hold:*

- (2.9.1) *If $Q = MR_k$, then $W(k, Q) = W(0, M)$.*
- (2.9.2) *If $Q \supset MR_k$, then $W(k, Q) = W'(k, Q) = W''(k, Q)$, where $W'(k, Q) = \{n; \text{there exists a mcplil } n \text{ in some integral extension domain of } (R_k)_Q\}$ and $W''(k, Q) = \{m + k - \text{depth } Q - 1; m \in W(1, (M, X_1))\}$.*

PROOF. (2.9.1) was proved in [1, Theorem 6].

(2.9.2) For $k = 1$ and $Q = (M, X_1)R[X_1]$, $W(k, Q) = W''(k, Q) = W(1, (M, X_1))$, so it suffices to prove that $W(k, Q) = W'(k, Q)$ in this case. Assume temporarily that this is known. Then $W'(k, Q) = W''(k, Q)$ in general, since $n \in W'(k, Q)$ if and only if there exists an integral extension domain of $(R_k)_Q$ which has a mcplil n if and only if (by (2.6)) there exists an integral extension domain of R which has a mcplil $n - k + \text{depth } Q$ if and only if (by (2.6) and assumption) $n - k + \text{depth } Q + 1 \in W(1, (M, X_1))$ if and only if $n \in W''(k, Q)$. Therefore it suffices to prove that $W(1, (M, X_1)) = W'(1, (M, X_1))$ and $W(k, Q) = W'(k, Q)$ ($k > 0$ and $MR_k \subset Q$). However, the proofs of both of these are essentially the same, so it will be shown that $W(k, Q) = W'(k, Q)$.

For this, it is clear that $W(k, Q) \subseteq W'(k, Q)$. For the opposite inclusion, let $d = \text{depth } Q$. Then $d = \text{trd}(R_k/Q)/(R/M)$ [5, (14.6)], so renumber the X_i such that X_1, \dots, X_d modulo Q are a transcendence basis for R_k/Q over R/M . Then $Q \cap R_d = MR_d$ [7, Lemma 4.2] (where $R_d = R[X_1, \dots, X_d]$). Also, with $A = (R_d)_{MR_d}$ and $B = A[X_{d+1}, \dots, X_k]$, $(R_k)_Q = B_P$, where $P = Q(R_k)_Q \cap B$, and P is a maximal ideal in B (since $P \cap A = MA$ and B/P is algebraic over the field A/MA). Let $D = A[X_{d+1}, \dots, X_{k-1}]$ ($= A$, if $d = k - 1$; $d \leq k - 1$, since $Q \supset MR_k$), and let $N = P \cap D$, so N is a maximal ideal in D .

Now assume that $n \in W'(k, Q)$; that is, assume that there exists an integral extension domain of $(R_k)_Q$ which has a mcplil n . Then, since $B \cong D[X]$ and P and $N = P \cap D$ are maximal ideals, and since $(R_k)_Q = B_P$, there exists an integral extension domain of $S = D_N$ which has a mcplil $n - 1$ (2.6). Therefore, by (2.2), there exists a principal integral extension domain $S[c]$ of S which has a mcplil $n - 1$ such that $c \in p_{n-1}$ (where $(0) \subset p_1 \subset \dots \subset p_{n-1}$ is the chain). Hence, considering the preimage of this chain in $S[X]$, there exists a mcplil n in $S[X]_{(NS, X)} \cong D[X]_{(N, X)}$.

Since $B \cong D[X]$, let P' be the maximal ideal in $D[X]$ which corresponds to P , so $P' \cap D = P \cap D = N$. Also, $P' = (N, f)D[X]$, for some monic polynomial $f \in D[X]$, so, since $T = D[f]_{(N, f)} \cong D[X]_{(N, X)}$, there exists a mcplil n

in T . Then $D[X]_{P'}$ is integral over T (since $P' = P''D[X]$, where $P'' = P' \cap D[f] = (N, f)D[f]$), so there exists a mcpil n in $D[X]_{P'}$ (2.7); hence there exists a mcpil n in $(R_k)_Q = B_P \cong D[X]_{P'}$. Therefore $n \in W(k, Q)$. Q.E.D.

(2.10) COROLLARY. *Let (R, M) be a local domain. Then $m \in W(1, (M, X_1))$ if and only if $m + k - 1 \in W(k, (M, X_1, \dots, X_k))$.*

PROOF. Let $Q = (M, X_1, \dots, X_k)R_k$. Then, by (2.9.2),

$$\begin{aligned} W(k, Q) &= W'' = \{m + k - \text{depth } Q - 1; m \in W(1, (M, X_1))\} \\ &= \{m + k - 1; m \in W(1, (M, X_1))\}, \end{aligned}$$

since $\text{depth } Q = 0$. Q.E.D.

(2.11) COROLLARY. *Let (R, M) and k be as in (2.9), and let P and Q be prime ideals in R_k such that $\text{depth } P = \text{depth } Q$ and $P \cap R = M = Q \cap R$. Then $W(k, P) = W(k, Q)$.*

PROOF. This follows immediately from (2.9). Q.E.D.

The following corollary contains (2.10) and (2.11) as special cases.

(2.12) COROLLARY. *Let (R, M) be a local domain, let $k \leq n$ be positive integers, let P be a prime ideal in R_k such that $MR_k \subset P$, and let Q be a prime ideal in R_n such that $MR_n \subset Q$. Then $W(n, Q) = \{i + n - \text{depth } Q - k + \text{depth } P; i \in W(k, P)\}$. In particular, if $\text{depth } P = \text{depth } Q$, then $W(n, Q) = \{i + n - k; i \in W(k, P)\}$.*

PROOF. By (2.9.2), $j \in W(n, Q)$ if and only if $j = m + n - \text{depth } Q - 1$ and $i \in W(k, P)$ if and only if $i = m + k - \text{depth } P - 1$, where $m \in W(1, (M, X_1))$. The conclusion follows from this. Q.E.D.

The following corollary will be sharpened in (2.16)–(2.18) below.

(2.13) COROLLARY. *The following statements are equivalent for a local domain (R, M) :*

(2.13.1) $R[X_1]_{(M, X_1)}$ is catenary.

(2.13.2) *There exists a positive integer k and a prime ideal Q in R_k such that $MR_k \subset Q$ and $(R_k)_Q$ is catenary.*

(2.13.3) *For all integers $k \geq 0$, and for all prime ideals Q in R_k such that $Q \cap R = M$, $(R_k)_Q$ is catenary.*

PROOF. It is clear that (2.13.3) \Rightarrow (2.13.1) \Rightarrow (2.13.2). Finally, if (2.13.2) holds, then $W(k, Q)$ is a one-element set; hence $W(1, (M, X_1))$ is a one-element set (2.9.2), and so, for all $n > 0$ and for all prime ideals P in R_n such that $MR_n \subset P$, $W(n, P)$ is a one-element set (2.9.2); hence $(R_n)_P$ is catenary. Further, R is catenary if and only if $(R_n)_{MR_n}$ is catenary (2.9.1). Hence, since the catenary

property is inherited by quotient rings and $(R_n)_{MR_n}$ is a quotient ring of $(R_n)_P$, (2.13.2) implies (2.13.3). Q.E.D.

Before giving further corollaries of (2.9), it will be helpful at this point to summarize the main theorem.

(2.14) **SUMMARY OF MAIN THEOREM.** *Let (R, M) be a local domain, let k be a positive integer, let Q be a prime ideal in R_k such that $MR_k \subset Q$, and let $S = (R_k)_Q$. Then the following statements are equivalent:*

- (2.14.1) *There exists an integral extension domain of R which has a mcpl n .*
- (2.14.2) *There exists a minimal prime ideal z in the completion of R such that $\text{depth } z = n$.*
- (2.14.3) *There exists a minimal prime ideal w in the completion of S such that $\text{depth } w = n + k - \text{depth } Q$.*
- (2.14.4) *There exists an integral extension domain of S which has a mcpl $n + k - \text{depth } Q$.*
- (2.14.5) *There exists a mcpl $n + k - \text{depth } Q$ in S .*
- (2.14.6) *There exists a mcpl $n + 1$ in $R[X_1]_{(M, X_1)}$.*

PROOF. (2.14.1) \Leftrightarrow (2.14.2), by (2.3). (2.14.2) \Leftrightarrow (2.14.3), by (2.5). (2.14.3) \Leftrightarrow (2.14.4), by (2.3). (2.14.4) \Leftrightarrow (2.14.5), by (2.9.2). (2.14.5) \Leftrightarrow (2.14.6), by (2.12). Q.E.D.

(2.14.1) \Leftrightarrow (2.14.6) was given in [2, Theorem 1.8].

The following summary is included for the sake of completeness.

(2.15) **SUMMARY.** *With (R, M) and k as in (2.14), let $S = (R_k)_{MR_k}$. Then the following statements are equivalent:*

- (2.15.1) *There exists an integral extension domain of R which has a mcpl n .*
- (2.15.2) *There exists an integral extension domain of S which has a mcpl n .*
- (2.15.3) *There exists a minimal prime ideal z in the completion of R such that $\text{depth } z = n$.*
- (2.15.4) *There exists a minimal prime ideal w in the completion of S such that $\text{depth } w = n$.*

PROOF. (2.15.1) \Leftrightarrow (2.15.3) and (2.15.2) \Leftrightarrow (2.15.4), by (2.3); and (2.15.3) \Leftrightarrow (2.15.4), by (2.5). Q.E.D.

Before giving further corollaries, it should be noted that the corollaries given below are *really* corollaries of (2.14); that is, we do not have a built-in circle. (The proofs of the results used to prove (2.14) are not based on the corollaries.) The only place where this is not exactly true is in the reference to [6, Proposition 3.5] in the proof of (2.3). However, this reference can be replaced by [6, Lemma 2.17], and then [6, Proposition 3.5] follows from a short computation.

(2.16) COROLLARY (CF. [7, THEOREM 2.21(5) \iff (8)]). *Let (R, M) , k , Q , and S be as in (2.14). Then S is catenary if and only if S satisfies the s.c.c.*

PROOF. S is catenary if and only if every integral extension domain of S satisfies the f.c.c. (by (2.14.4) \iff (2.14.5)) if and only if S satisfies the s.c.c. Q.E.D.

(2.17) COROLLARY (CF. [7, THEOREM 2.21 (1) \iff (8)]). *Let (R, M) , k , Q , and S be as in (2.14). Then R satisfies the s.c.c. if and only if S is catenary.*

PROOF. S is catenary if and only if every integral extension domain of R satisfies the f.c.c. (by (2.14.1) \iff (2.14.5)) if and only if R satisfies the s.c.c. Q.E.D.

In the following corollary we use the fact that if p is a prime ideal in a local domain R which satisfies the s.c.c., then R_p satisfies the s.c.c. This is proved in (2.26) below.

(2.18) COROLLARY (CF. [6, THEOREM 3.6]). *The following statements are equivalent for a local domain (R, M) :*

- (2.18.1) R satisfies the s.c.c.
- (2.18.2) R_k is catenary, for some $k > 0$.
- (2.18.3) R_k is catenary, for all $k \geq 0$.

PROOF. It is clear that (2.18.3) implies (2.18.2). Also, (2.18.2) implies (2.18.1), by (2.17), since if R_k is catenary, for some $k > 0$, then $S = (R_k)_{(M, X_1, \dots, X_k)}$ is catenary.

Finally, assume that R satisfies the s.c.c. and let $k \geq 0$. Then to prove that R_k is catenary, it suffices to prove that if N is a maximal ideal in R_k , then $(R_k)_N$ is catenary. For this, it may clearly be assumed that $k > 0$. Also, if $N \cap R = M$, then $(R_k)_N$ is catenary (by (2.17), since N is maximal (hence $MR_k \subset N$)), so assume that $N \cap R = P \neq M$. Then R_P satisfies the s.c.c. and $N^* = NR_P[X_1, \dots, X_k]$ is a maximal ideal which lies over PR_P . Hence, by (2.17) applied to R_P , $(R_k)_N = ((R_P)_k)_N$ is catenary. Q.E.D.

(2.19) COROLLARY (CF. [6, THEOREM 3.1]). *A local domain R satisfies the s.c.c. if and only if R is quasi-unmixed.*

PROOF. R satisfies the s.c.c. if and only if, by (2.17), $R[X_1]_{(M, X_1)}$ is catenary if and only if $W(1, (M, X_1))$ is a one-point set if and only if, by (2.14.2) \iff (2.14.6), all minimal prime ideals in the completion of R have the same depth if and only if R is quasi-unmixed. Q.E.D.

In (2.26) below it will be shown that if a local domain R satisfies the s.c.c., then, for each prime ideal p in R , R_p and R/p satisfy the s.c.c. Using this, a

fairly easy proof of the following known and important result can be given using (2.16) and (2.17).

(2.20) COROLLARY (CF. [6, COROLLARY 3.7]). *If a local domain R satisfies the s.c.c., then, for each prime ideal p in R , and for each local domain L which is a locality over R/p , L satisfies the s.c.c.*

PROOF. Assume that R satisfies the s.c.c., let p be a prime ideal in R , and let L be a local domain which is a locality over R/p , say $L = ((R/p)[a_1, \dots, a_k])_q$. Then there exists a prime ideal Q in R_k such that L is a homomorphic image of $(R_k)_Q$. Let $P = Q \cap R$, so $(R_k)_Q$ is a quotient ring of $A = R_P[X_1, \dots, X_k]$ and R_P satisfies the s.c.c. Therefore, if N is a maximal ideal in A such that $QA \subseteq N$, then $D = A_N$ satisfies the s.c.c. (by (2.16) and (2.17)); hence $(R_k)_Q = D_{QD}$ satisfies the s.c.c., and so L satisfies the s.c.c. (see (2.26)). Q.E.D.

(2.9.2) says that the lengths of maximal chains of prime ideals in $(R_k)_Q$ ($k > 0$ and $MR_k \subset Q$) are known, once the lengths of maximal chains of prime ideals in $D = R[X_1]_{(M, X_1)}$ are known. It is, of course, of much interest to know how the lengths of maximal chains of prime ideals in D compare to the lengths of maximal chains of prime ideals in R . The only case in which we know how these lengths compare is when R satisfies the s.c.c.; that is, by (2.17), when $W(1, (M, X_1)) = \{\text{altitude } R + 1\}$. In all other cases this is a very difficult problem, but we make the following conjecture.

(2.21) CONJECTURE. $\{n + 1; n \in W(0, M)\} \subseteq W(1, (M, X_1)) \subseteq \{n + 1; n \in W(0, M)\} \cup \{2\}$.

The first containment is obvious.

This conjecture was called the upper conjecture in [2], and it was shown there that the depth conjecture implies the upper conjecture which, in turn, implies the catenary chain conjecture [2, Propositions 3.5 and 3.7]. (*Depth conjecture*: If P is a prime ideal in a local domain R such that height $P > 1$, then there exists a prime ideal p in R such that $p \subset P$ and $\text{depth } p = \text{depth } P + 1$. *Catenary chain conjecture*: The integral closure of a catenary local domain satisfies the c.c.)

Nagata's examples [5, Example 2, pp. 203–205] support the conjecture. That is, in these examples, the first inclusion is an equality for $m > 0$ (m as in [5]) and the second inclusion is an equality for $m = 0$.

The following result shows that only the last step in a maximal chain of prime ideals keeps us from verifying the upper conjecture for a catenary local domain:

If (R, M) is a catenary local domain and $(0) \subset Q_1 \subset \cdots \subset Q_n$ is a maxi-

mal chain of prime ideals in an integral extension domain of R , then $\text{height } Q_i = i = \text{height } Q_i \cap R$ ($i < n$). For, $R_{Q_i \cap R}$ satisfies the s.c.c. [8, Theorem 3.9]. (Also, $\text{height } Q_n = \text{height } M$, if $n > 1$. Now $n = 1$ is possible; that is, possibly $Q_1 = Q_n$ is a height one maximal ideal and altitude $R > 1$, but this does not cause any real problem. The real problem lies in showing that if $n > 1$, then $\text{depth } Q_{n-1} = 1$.)

We close the discussion of the upper conjecture with the following result.

(2.22) COROLLARY. *The following statements are equivalent:*

(2.22.1) *The upper conjecture holds.*

(2.22.2) *If R' is an integral extension domain of a local domain R such that R' has a mcpil n , then either there exists a mcpil n in R or $n = 1$.*

(2.22.3) *If there exists a depth n minimal prime ideal in the completion of a local domain R , then there either exists a mcpil n in R or $n = 1$.*

PROOF. Assume that the upper conjecture holds, let R be a local domain, and let R' be an integral extension domain of R . If there exists a mcpil n in R' , then, by (2.14.1) \Leftrightarrow (2.14.6), $n + 1 \in W(1, (M, X_1)) \subseteq \{i + 1; i \in W(0, M)\} \cup \{2\}$, so (2.22.1) implies (2.22.2). Conversely, let (R, M) be a local domain and let $m \in W(1, (M, X_1))$. Then there exists an integral extension domain of R which has a mcpil $m - 1$, by (2.14.1) \Leftrightarrow (2.14.6). Therefore, by hypothesis, either $m - 1 \in W(0, M)$ or $m - 1 = 1$; hence (2.22.2) implies (2.22.1).

Again assume that the upper conjecture holds, let R be a local domain, and let z be a depth n minimal prime ideal in the completion of R . Then, by (2.14.2) \Leftrightarrow (2.14.6), $n + 1 \in W(1, (M, X_1)) \subseteq \{i + 1; i \in W(0, M)\} \cup \{2\}$. Therefore (2.22.1) implies (2.22.3). Conversely, let (R, M) be a local domain and let $m \in W(1, (M, X_1))$. Then there exists a depth $m - 1$ minimal prime ideal in the completion of R , by (2.14.2) \Leftrightarrow (2.14.6). Therefore, by hypothesis, either $m - 1 \in W(0, M)$ or $m - 1 = 1$; hence (2.22.3) implies (2.22.1). Q.E.D.

If there exists a mcpil m in $D = R[X_1]_{(M, X_1)}$, then there exists a mcpil m in D , say $(0) \subset p_1 \subset \cdots \subset p_m = (M, X_1)D$ such that $R[X_1]/(p_1 \cap R[X_1])$ is integral over R . This follows by taking $S = D$ in the following corollary.

(2.23) COROLLARY. *Let (R, M) , k , Q , and S be as in (2.14). Assume that there exists a mcpil n in S . Then there exists a mcpil $m = n - (k - 1) + \text{depth } Q$ in $R[X_1]_{(M, X_1)}$, say $(0) \subset p_1 \subset \cdots \subset p_m = (M, X_1)$, such that $R[X_1]/(p_1 \cap R[X_1])$ is integral over R .*

PROOF. By (2.14.1) \Leftrightarrow (2.14.5), there exists an integral extension domain of R which has a mcpil $m - 1$. Therefore, by (2.2), there exists a principal

integral extension domain $R[c]$ of R which has a mcpil $m - 1$, say $(0) \subset \cdots \subset q_{m-1} = N$ such that $c \in N$. Then $N = (M, c)R[c]$, so the conclusion follows by considering the preimage of this chain in $R[X_1]$. Q.E.D.

(2.14) allows us to determine possible lengths of maximal chains of prime ideals in integral extension domains of R/p and of R_p (where p is a prime ideal in a local domain R). This is explicitly stated in the next two remarks.

(2.24) REMARK. Let p be a prime ideal in a local domain (R, M) , and let $D = R[X_1]_{(M, X_1)}$.

(2.24.1) There exists an integral extension domain of R/p which has a mcpil n if and only if in D , there exists a saturated chain of prime ideals of length $n + 1$ with pD the smallest term and $(M, X_1)D$ the largest term.

(2.24.2) There exists an integral extension domain of R_p which has a mcpil n if and only if in D there exists a saturated chain of prime ideals of length $n + 1$ with $(p, X_1)D$ the largest term and (0) the smallest term.

PROOF. (2.24.1) There exists an integral extension domain of R/p which has a mcpil n if and only if in $D/pD = (R/p)[X_1]_{(M/p, X_1)}$ there exists a mcpil $n + 1$, by (2.14.1) \Leftrightarrow (2.14.6), and (2.24.1) clearly follows from this.

(2.24.2) There exists an integral extension domain of R_p which has a mcpil n if and only if in $D_{(p, X_1)D} = R_p[X_1]_{(pR_p, X_1)}$ there exists a mcpil $n + 1$, by (2.14.1) \Leftrightarrow (2.14.6), and (2.24.2) clearly follows from this. Q.E.D.

(2.25) REMARK. Let p be a prime ideal in a local domain (R, M) .

(2.25.1) If there exists an integral extension domain of R/p which has a mcpil n , then there exists an integral extension domain of R which has a mcpil $n + \text{height } p$.

(2.25.2) If there exists an integral extension domain of R_p which has a mcpil n , then there exists an integral extension domain of R which has a mcpil $n + \text{depth } p$.

PROOF. (2.25.1) If there exists an integral extension domain of R/p which has a mcpil n , then, by (2.24.1), there exists a mcpil $n + 1 + \text{height } p$ in D ; hence, by (2.14.1) \Leftrightarrow (2.14.6), there exists an integral extension domain of R which has a mcpil $n + \text{height } p$.

The proof of (2.25.2) is similar. Q.E.D.

(2.26) COROLLARY (CF. [4, LEMMA 1]). Let p be a prime ideal in a local domain R . If R satisfies the s.c.c., then R_p and R/p satisfy the s.c.c.

PROOF. Assume that there exists an integral extension domain of R/p which has a mcpil n . Then, by (2.25.1), there exists an integral extension domain of R which has a mcpil $n + \text{height } p$, so $n + \text{height } p = \text{altitude } R$, if R satisfies the s.c.c., and $\text{height } p + \text{depth } p = \text{altitude } R$. Therefore $\text{depth } p = n$; hence R/p satisfies the s.c.c.

The proof that R_p satisfies the s.c.c. is similar. Q.E.D.

(2.9) was stated for polynomial rings over a local domain. However, it is straightforward to see that it holds for polynomial rings over a Noetherian domain. Specifically, we close this paper with the following remark.

(2.27) REMARK. Let P be a prime ideal in a Noetherian domain A , and let Q be a prime ideal in $A_k = A[X_1, \dots, X_k]$ such that $Q \cap A = P$. Let $B = (A_k)_{(A-P)}$, and let $d = \text{depth } QB$, so $\text{trd}(B/QB)/(A_P/PA_P) = d$. Renumber the X_i such that X_1, \dots, X_d modulo QB are a transcendence basis for B/QB over A_P/PA_P , and let $W_i = \{n_i\}$; there exists an integral extension domain of $(A_i)_{Q \cap A_i}$ which has a mcpil $n_i\}$ ($i = 0, 1, \dots, d$ with $A_0 = A$). Then $W_0 = \dots = W_d$ and $n \in W_d = W_0$ if and only if $n + j \in W(d + j, Q \cap A_{d+j})$ ($j = 1, \dots, k - d$).

PROOF. By [7, Lemma 4.2], $Q \cap A_i = PA_i$ ($i = 0, 1, \dots, d$); hence the W_i are equal (2.9.1). Let $C = (A_d)_{PA_d}$, and let $C_j = C[X_{d+1}, \dots, X_{d+j}]$ ($j = 0, 1, \dots, k - d$ and $C_0 = C$). Then $(A_k)_Q = (C_{k-d})_{Q'}$, where $Q' = Q(A_k)_Q \cap C_{k-d}$, so, since $(Q \cap A_{d+j})A_{d+j+1} \subset Q \cap A_{d+j+1}$ ($j = 0, 1, \dots, k - d - 1$), the ideals $Q' \cap C_j$ are maximal ($j = 0, 1, \dots, k - d$). Therefore $n \in W_d$ if and only if $n + j \in W(d + j, Q \cap A_{d+j})$ ($j = 1, \dots, k - d$), by (2.14.1) \Leftrightarrow (2.14.5) applied to C ($(A_{d+j})_{Q \cap A_{d+j}} = (C_j)_{Q' \cap C_j}$). Q.E.D.

BIBLIOGRAPHY

1. S. McAdam, *Saturated chains in Noetherian rings*, Indiana Univ. Math. J. 23 (1973/74), 719–728. MR 48 #11094.
2. S. McAdam and E. G. Houston, *Chains of primes in Noetherian rings*, Indiana Univ. Math. J. 24 (1975), 741–753.
3. M. Nagata, *On the chain problem of prime ideals*, Nagoya Math. J. 10 (1956), 51–64. MR 18, 8.
4. ———, *Note on a chain condition for prime ideals*, Mem. Coll. Sci. Univ. Kyoto Ser. A. Math. 32 (1959), 85–90. MR 21 #7227.
5. ———, *Local rings*, Interscience Tracts in Pure and Appl. Math., no. 13, Interscience, New York, 1962. MR 27 #5790.
6. L. J. Ratliff, Jr., *On quasi-unmixed local domains, the altitude formula and the chain condition for prime ideals*. I, Amer. J. Math. 91 (1969), 508–528. MR 40 #136.
7. ———, *On quasi-unmixed local domains, the altitude formula, and the chain condition for prime ideals*. II, Amer. J. Math. 92 (1970), 99–144. MR 42 #249.
8. ———, *Characterizations of catenary rings*, Amer. J. Math. 93 (1971), 1070–1108. MR 45 #6804.
9. ———, *Chain conjectures and H-domains*, Conf. on Commutative Algebra (Univ. Kansas, Lawrence, Kan., 1972), Lecture Notes in Math., vol. 311, Springer-Verlag, Berlin, 1973, pp. 222–238. MR 49 #2714.
10. ———, *Four notes on saturated chains of prime ideals*, J. Algebra 39 (1976), 75–93.
11. O. Zariski and P. Samuel, *Commutative algebra*, Vol. II, University Ser. in Higher Math., Van Nostrand, Princeton, N. J., 1960. MR 22 #11006.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE, CALIFORNIA 92502

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS, AUSTIN, TEXAS