

## THE $p$ -ADIC LOG GAMMA FUNCTION AND $p$ -ADIC EULER CONSTANTS<sup>1</sup>

BY

JACK DIAMOND

**ABSTRACT.** We define  $G_p$ , a  $p$ -adic analog of the classical log gamma function and show it satisfies relations similar to the standard formulas for log gamma. We also define  $p$ -adic Euler constants and use them to obtain results on  $G'_p$  and on the logarithmic derivative of Morita's  $\Gamma_p$ .

**1. Introduction.** Leopoldt and Kubota defined  $p$ -adic  $L$ -functions by summing a function of two variables with respect to one of the variables. We present a general theorem on this technique and then use it to define  $G_p$ , a  $p$ -adic analog of the classical log  $\Gamma$  function. We work with  $\log \Gamma$  rather than  $\Gamma$  because the only continuous  $p$ -adic function defined on a subset of  $\Omega_p$  and satisfying  $f(x + 1) = xf(x)$  is the zero function. It is possible to construct an analog of  $\Gamma$  by modifying the functional equation (see Morita [7]), but then we do not have close analogs of the standard formulas for  $\Gamma$  or  $\log \Gamma$ . For  $G_p$ , which is not the log of Morita's gamma function, we have the functional equation, an extension theorem, the Stirling series, the Gauss multiplication theorem, a power series, certain "Laurent" series and a formula due to Gauss which is valid for  $G'_p$  at rational points.

This last formula was discussed by Lehmer in [6], where he defined Euler constants for arithmetic progressions. We define  $p$ -adic Euler constants and present a proof of Gauss' theorem which is valid in both the  $p$ -adic and complex systems. We also apply the results on Euler constants to obtain a finite expression for the logarithmic derivative of Morita's  $p$ -adic gamma function at certain rational values in its domain.

**2. Notation and definitions.** We will use  $Q$ ,  $Q_p$ ,  $Z$ ,  $Z_p$ ,  $C$  and  $\Omega_p$  for, respectively, the field of rational numbers, the  $p$ -adic completion of  $Q$ , the ring of rational integers, the  $p$ -adic completion of  $Z$  in  $Q_p$ , the field of complex numbers and the completion of the algebraic closure of  $Q_p$ .  $B_n$  will be the  $n$ th Bernoulli number defined by  $te^t/(e^t - 1)$ .  $\nu$  will be the  $p$ -adic valuation on

---

Received by the editors April 12, 1976.

*AMS (MOS) subject classifications* (1970). Primary 12B40; Secondary 33A15.

*Key words and phrases.* Log gamma function, Euler constants,  $p$ -adic functions.

<sup>1</sup>This paper is part of the author's Ph.D. dissertation, submitted to Princeton University in July, 1975.

$\Omega_p$  with  $\nu(p) = 1$  and  $|\cdot|_p$  will be the absolute value on  $\Omega_p$  with  $|p|_p = p^{-1}$ . We will use boldface letters to indicate  $r$ -tuples.

A polydisc about  $\mathbf{c} \in \Omega_p^r$  is a set of the form

$$\{(x_1, \dots, x_r): |x_i - c_i|_p \leq \rho_i, i = 1, 2, \dots, r\}$$

where  $\mathbf{c} = (c_1, \dots, c_r)$  and all  $\rho_i > 0$ .  $(\rho_1, \dots, \rho_r)$  is called the radius of the polydisc.  $\mathbf{a}$  and  $\mathbf{M}$  will denote  $(a_1, \dots, a_r)$  and  $(M_1, \dots, M_r)$ , respectively.

We call a function defined on a subset of  $\Omega_p^r$  holomorphic if it can be represented by a single power series and locally holomorphic if at each point in the domain we can represent the function by a power series on some polydisc containing the point.

The author wishes to express his gratitude to Professor Bernard Dwork for many helpful discussions on this material.

**3.  $p$ -adic sums.** We begin by considering sums of the type used by Leopoldt and Kubota [5]. The first theorem is a generalization of a result in [2, p. 309].

**THEOREM 1.** *Suppose we have rational integers  $a_i, b_i, M_i$  with  $a_i > 0, b_i > 1, M_i > 1$  for  $i = 1, 2, \dots, r$ . Let  $R$  be an open set in  $\Omega_p^r$  with  $\mathbf{a} + \mathbf{MZ}_p \subset R$ .  $B$  is a Banach space over  $\Omega_p$  and  $f: R \rightarrow B$  is locally holomorphic.*

We define

$$S(k_1, \dots, k_r, b_1, \dots, b_r) = \frac{1}{b_1 \dots b_r p_1^{k_1} \dots p_r^{k_r}} \sum_{i=1}^r \sum_{n_i=0; n_i \equiv a_i \pmod{M_i}}^{M_i b_i p^{k_i-1}} f(n_1, \dots, n_r).$$

Then

- (i)  $L = \lim_{(k_i) \rightarrow \infty} S(k_1, \dots, b_r)$  exists;
- (ii)  $L$  is independent of the  $\{b_i\}$  used;
- (iii)  $L$  may be calculated by iteration of the limit in any order.

**PROOF.** We begin with  $a_1, \dots, a_r = 0, M_1, \dots, M_r = 1$  and  $f$  holomorphic on  $R$  with  $Z_p^r \subset R$ . We can write

$$f(\mathbf{x}) = \sum_J a_J \mathbf{x}^J,$$

where the right side represents a power series in  $r$  variables with  $J$  running through the  $r$ -tuples of nonnegative integers.

After we substitute the series for  $f$  in the formula for  $S$  and use the fact (see [2]) that

$$\left| (-1)^j B_j - \frac{1}{bp^k} \sum_{n=0}^{bp^k-1} n^j \right|_p < p^{2-k}$$

for  $k = 0, 1, \dots$  and  $j = 0, 1, \dots$ , it is easy to verify that

$$\lim_{k_r \rightarrow \infty} \dots \lim_{k_1 \rightarrow \infty} S(k_1, \dots, b_r) = \sum_J a_J (-1)^J B_J,$$

and each limit is uniform with respect to the remaining variables. We can now conclude that

$$L = \lim_{\{k_i\} \rightarrow \infty} S(k_1, \dots, b_r) \text{ exists.}$$

The next two parts of the theorem are obvious.

Now suppose *f* is locally holomorphic on *R* and  $R \supset Z'_p$ . Then there is a finite covering of  $Z'_p$  by polydiscs whereby *f* is holomorphic on each polydisc, each polydisc has the same radius and the radius has the form  $(p^{-N}, \dots, p^{-N})$ . We let  $A = \{0, \dots, p^N - 1\}$  and for each  $W \in A'$  we define

$$f_W(\mathbf{x}) = p^{-Nr} f(W + p^N \mathbf{x}).$$

Each  $f_W$  is holomorphic on the disc with center  $(0, \dots, 0)$  and radius  $(1, \dots, 1)$ .

It is convenient now to introduce an integral type notation.

We define

$$\int_{\mathbf{a}, \mathbf{M}} f(\mathbf{x}) \, d\mathbf{x} = L,$$

where *L* is defined in Theorem 1.

We have

$$\int_{\mathbf{0}, \mathbf{1}} f(\mathbf{x}) \, d\mathbf{x} = \sum_{W \in A'} \int_{\mathbf{0}, \mathbf{1}} f_W(\mathbf{x}) \, d\mathbf{x},$$

and the conclusion of the theorem follows directly.

Finally, if  $f: R \rightarrow B$  is locally holomorphic and  $\mathbf{a} + \mathbf{M}Z_p \subset R$ , we define  $g(\mathbf{x}) = f(\mathbf{a}' + \mathbf{xM})$  where  $\mathbf{xM} = (x_1 M_1, \dots, x_r M_r)$  and  $\mathbf{a}'$  is the least nonnegative residue of  $\mathbf{a}$  mod  $\mathbf{M}$ . Since *g* satisfies the conditions needed earlier in this proof, and  $\int_{\mathbf{a}, \mathbf{M}} f(\mathbf{x}) \, d\mathbf{x} = \int_{\mathbf{0}, \mathbf{1}} g(\mathbf{x}) \, d\mathbf{x}$ , we have established Theorem 1.

The next result is our basic device for constructing *p*-adic functions. We will use it to define a *p*-adic analog of  $\log \Gamma$  and to define *p*-adic Euler constants. It can be used to show the existence and holomorphy of the *p*-adic *L*-functions and similarly constructed functions occurring in the works cited as references.

**THEOREM 2.** *Suppose  $a_i, M_i$  are rational integers with  $a_i \geq 0, M_i \geq 1$  for  $i = 1, 2, \dots, r, \{C_1, \dots, C_t\}$  is a set of polydiscs in  $\Omega_p^r, R = \cup_{i=1}^t C_i$  and  $\mathbf{a} + \mathbf{M}Z_p \subset R$ . Let *D* be a polydisc in  $\Omega_p^s$  and suppose  $f: R \times D \rightarrow \Omega_p$  is holomorphic on each  $C_i \times D, i = 1, 2, \dots, t$ . Then*

$$F(x) = \int_{a, M} f(u, x) du$$

exists and is holomorphic on  $D$ .

PROOF. We let  $\Lambda(D)$  = Banach space of holomorphic functions from  $D \rightarrow \Omega_p$ . For  $u \in R$  we define

$$\phi(u) = \text{the mapping } x \rightarrow f(u, x).$$

For a fixed  $u_i \in C_i$  we have

$$f(u, x) = \sum_j a_{i,j}(x)(u - u_i)^j$$

for all  $u \in C_i$  and  $x \in D$ .

If  $a_{i,j}$  denotes the map on  $D$ ,  $x \rightarrow a_{i,j}(x)$ , then

$$\phi(u) = \sum_j a_{i,j}(u - u_i)^j$$

for  $u \in C_i$ .

Each  $a_{i,j} \in \Lambda(D)$ , so  $\phi: R \rightarrow \Lambda(D)$  is locally holomorphic and we may apply Theorem 1. Since

$$\int_{a, M} f(u, x) du = \left( \int_{a, M} \phi(u) du \right)(x),$$

we conclude that  $F \in \Lambda(D)$ .

The following corollary is a useful form of Theorem 2.

COROLLARY. Suppose  $a, b, M$  are rational integers with  $a > 0, b > 1, M > 1$ . Let  $f$  be locally holomorphic on a set  $A \subset \Omega_p$ . Let  $x \in \Omega_p^s$  and  $T(u, x)$  be locally holomorphic on some subset of  $\Omega_p^{s+1}$ . Define  $A^* = \{x | T(a + MZ_p, x) \subset A\}$ .

Then  $A^*$  is open, and if  $A^* \neq \emptyset$ ,

$$F(x) = \lim_{k \rightarrow \infty} \frac{1}{bp^k} \sum_{\substack{n=0 \\ n \equiv a \pmod{M}}}^{Mbp^k-1} f(T(n, x))$$

is independent of  $b$  and locally holomorphic on  $A^*$ .

PROOF. Given  $c \in A^*$  and  $u \in a + MZ_p$  there is a polydisc  $D(u, c)$  containing  $(u, c)$  on which  $f \circ T$  is holomorphic. Holding  $c$  fixed, a finite number of  $D(u_i, c)$  cover  $(a + MZ_p, c)$ . Each  $D(u_i, c)$  has the form  $C_i \times D_i$  where  $C_i$  is a disc in  $\Omega_p$  containing  $u_i$  and  $D_i$  is a polydisc about  $c$ . Let  $D = \cap D_i$ . We know the following:

- (i)  $T(C_i \times D) \subset A$  for each  $i$ , so  $D \subset A^*$ ;
- (ii)  $\cup C_i$  covers  $a + MZ_p$ ;
- (iii)  $f \circ T$  is holomorphic on each  $C_i \times D$ .

From Theorem 2 we see  $F(x)$  is holomorphic on  $D$  and therefore locally holomorphic on  $A^*$ .

We will occasionally wish to differentiate  $F(x)$ . We have

**THEOREM 3.** *Using the definitions and conditions of Theorem 2,*

$$\frac{\partial F(x)}{\partial x_i} = \int_{a,M} \frac{\partial f(u, x)}{\partial x_i} du.$$

**PROOF.** We fix  $x \in D$  and for  $t \in \Omega_p$  we let  $t^* = (0, \dots, t, \dots, 0)$ ,  $t$  being in the  $i$ th position,  $t^* \in \Omega_p^*$ . We define

$$h(u, t) = \frac{f(u, x + t^*) - f(u, x)}{t} \quad \text{for } t \neq 0$$

and  $h(u, 0) = \lim_{t \rightarrow 0} h(u, t)$ .

We observe that  $h(u, 0) = \partial f(u, x) / \partial x_i$  and that there is a neighborhood  $D_0$ , of zero, so  $h$  is holomorphic on each  $C_i \times D_0$ .

From the definition of derivative we have

$$\frac{\partial F(x)}{\partial x_i} = \lim_{t \rightarrow 0} \int_{a,M} h(u, t) du.$$

Now, if we let  $H(t) = \int_{a,M} h(u, t) du$  we can use Theorem 2 to see that  $H$  is holomorphic on  $D_0$  and, in particular, continuous at 0.

Thus we have

$$\frac{\partial F(x)}{\partial x_i} = \lim_{t \rightarrow 0} H(t) = H(0) = \int_{a,M} \frac{\partial f(u, x)}{\partial x_i} du.$$

The next result shows how certain sums can be used to solve difference equations.

**THEOREM 4.** *If  $a, M$  are rational integers where  $M > a \geq 0$ ,  $f'(x + a)$  exists and  $F(x) = \int_{a,M} f(x + u) du$ , then  $F(x + M)$  exists and  $F(x + M) = F(x) + Mf'(x + a)$ .*

**PROOF.** This follows directly from the definition of the right side.

**4. The *p*-adic log  $\Gamma$  function.** We now consider the problem of constructing a *p*-adic analog of  $\log \Gamma(x)$ .

In looking for a *p*-adic analog of  $\log \Gamma(x)$  we want a function  $G_p$  which sends a subset of  $\Omega_p$  into  $\Omega_p$  and satisfies the functional equation  $G_p(x + 1) = G_p(x) + \log(x)$ .  $\log(x)$  is defined by the usual power series when  $|x - 1|_p < 1$ , and by setting  $\log(p) = 0$  and using the functional equations for  $\log(x)$  when  $|x - 1|_p > 1$  and  $x \neq 0$ . There is a complete discussion of this idea in [3]. Just as in the complex case, this functional equation forces  $G_p$  to be discontinuous on either the positive integers or the negative integers. This is somewhat unfortunate in the *p*-adic case because if we want a locally

holomorphic function we must exclude  $Z_p$  from the domain of  $G_p$ . However, this is all we need exclude because on the domain  $\Omega_p - Z_p$  we have a locally holomorphic function which satisfies  $G_p(x + 1) = G_p(x) + \log(x)$  and several other relations similar to those of the complex  $\log \Gamma(x)$ . We will use the construction given in §3 to define  $G_p(x)$  and demonstrate its properties.

An alternative approach is to slightly modify the functional equation to obtain a functional locally holomorphic on all of  $\Omega_p$ . After considering  $G_p(x)$  we will exhibit a sequence of such functions. We have the relation that the sequence of functions locally holomorphic on  $\Omega_p$  converges pointwise to  $G_p$ .

The technique of changing the functional equation has been used by Morita [7] to define  $\Gamma_p$ , a function on  $Z_p$ , which is an analog of  $\Gamma$ . Our  $G_p$  is clearly not  $\log \Gamma_p$ .

DEFINITION OF  $G_p$ . We use the corollary of Theorem 2 with  $T(u, x) = u + x$  and  $f(x) = x \log(x) - x$ .  $f$  is locally holomorphic on  $\Omega_p - \{0\}$ . We then have

$$G_p(x) = \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{n=0}^{p^k-1} (x+n)\log(x+n) - (x+n).$$

$G_p$  is locally holomorphic on  $\Omega_p - Z_p$ , and at each  $c \in \Omega_p - Z_p$  the disc of holomorphy is the largest (open) disc  $D(c)$  such that  $D(c) \cap Z_p = \emptyset$ .

An immediate consequence of Theorem 4 is the functional equation:

THEOREM 5.  $G_p(x + 1) = G_p(x) + \log x$ .

Stirling's Theorem, which is an asymptotic formula for  $\log \Gamma(x)$ , is simpler in  $\Omega_p$ . We have

THEOREM 6. *When  $|x|_p > 1$ ,*

$$G_p(x) = \left(x - \frac{1}{2}\right) \log(x) - x + \sum_{r=1}^{\infty} \frac{B_{r+1}}{r(r+1)x^r}.$$

PROOF.

$$G_p(x) = \frac{1}{2} - x + \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{n=0}^{p^k-1} (x+n)(\log(x) + \log(1 + n/x)).$$

Using the power series for  $\log(1 + n/x)$  will lead to the result.

If we match Theorem 6 and the next result with the corresponding classical formulas we see that it is more accurate to speak of  $G_p(x)$  as the analog of  $-\frac{1}{2} \log(2\pi) + \log \Gamma(x)$ . However, for simplicity we will continue to refer to  $G_p$  as the analog of  $\log \Gamma$ .

The following relation is the  $p$ -adic version of Gauss' Multiplication Theorem.

THEOREM 7. *Given any  $m \in Z^+$  we have*

$$G_p(x) = (x - \frac{1}{2})\log(m) + \sum_{a=0}^{m-1} G_p\left(\frac{x+a}{m}\right)$$

provided the right side is defined.

PROOF. We can write

$$\begin{aligned} G_p(x) &= \lim_{k \rightarrow \infty} \frac{1}{mp^k} \sum_{n=0}^{mp^k-1} (x+n) \log(x+n) - (x+n) \\ &= \lim_{k \rightarrow \infty} \frac{1}{mp^k} \sum_{n=0}^{p^k-1} \sum_{a=1}^{m-1} (x+a+mn) \log(x+a+mn) - (x+a+mn). \end{aligned}$$

With a little rearranging, Theorem 7 is easily obtained.

COROLLARY.

$$G_p(x) = \sum_{a=0}^{p^r-1} G_p\left(\frac{x+a}{p^r}\right) \text{ for } r = 0, 1, 2, \dots$$

This last corollary provides us with a means for transferring results about  $G_p(x)$  when  $|x|_p > 1$  to  $G_p(x)$  with  $|x|_p < 1$ .

For the extension theorem we have

**THEOREM 8.**  $G_p(x) + G_p(1-x) = 0$ .

PROOF. We can see immediately from Theorem 6 that  $G_p(x) + G_p(-x) = -\log(x)$  when  $|x|_p > 1$ . Combining this with  $G_p(x+1) = G_p(x) + \log(x)$  we have  $G_p(x) + G_p(1-x) = 0$  when  $|x|_p > 1$ .

Given any  $x \in \Omega_p - Z_p$  with  $|x|_p < 1$  we can choose an  $r \in \mathbb{Z}^+$  so  $|(x+a)/p^r|_p > 1$  for all  $a \in \mathbb{Z}$ . Then

$$\begin{aligned} G_p(x) + G_p(1-x) &= \sum_{a=0}^{p^r-1} G_p\left(\frac{x+a}{p^r}\right) + G_p\left(\frac{1-x+a}{p^r}\right) \\ &= \sum_{a=0}^{p^r-1} G_p\left(\frac{x+a}{p^r}\right) - G_p\left(1 - \frac{1-x+a}{p^r}\right) \\ &= \sum_{a=0}^{p^r-1} G_p\left(\frac{x+a}{p^r}\right) - G_p\left(\frac{x+p^r-a-1}{p^r}\right), \end{aligned}$$

and, since as  $a$  goes from 0 to  $p^r - 1$ ,  $p^r - a - 1$  goes from  $p^r - 1$  to 0, Theorem 8 is proven.

The complex log  $\Gamma(x)$  has a simple power series about 1, with values of the Riemann  $\zeta$ -function appearing in the coefficients. We will now find the power series for  $G_p(x)$  about  $1/p$ .

We use Theorem 4 to obtain

$$D^{(1)}G_p(x) = \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{n=0}^{p^k-1} \log(x+n)$$

and, in particular,

$$D^{(1)}G_p(1/p) = p \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{\substack{m=0 \\ m \equiv 1 \pmod{p}}}^{p^k-1} \log(m).$$

We write this as

$$D^{(1)}G_p(1/p) = -p\gamma_p(1, p),$$

$$D^{(r)}G_p(x) = (-1)^r (r-2)! \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{n=0}^{p^k-1} \frac{1}{(x+n)^{r-1}} \quad \text{for } r > 2,$$

$$\begin{aligned} \frac{D^{(r)}G_p(1/p)}{r!} &= \frac{(-1)^r}{r(r-1)} \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{n=0}^{p^k-1} \frac{1}{(n+1/p)^{r-1}} \\ &= \frac{(-1)^r p^r}{r} \cdot \frac{1}{r-1} \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{\substack{m=0 \\ m \equiv 1 \pmod{p}}}^{p^k-1} \frac{1}{m^{r-1}}. \end{aligned}$$

We will write this last expression as  $(-1)^r p^r \zeta_p(r)/r$ . Using the notation introduced above we have

**THEOREM 9.**

$$G_p(x) = G_p\left(\frac{1}{p}\right) - p\gamma_p(1, p)\left(x - \frac{1}{p}\right) + \sum_{r=2}^{\infty} \frac{(-1)^r \zeta_p(r) p^r}{r} \left(x - \frac{1}{p}\right)^r.$$

*This series converges for  $|x - 1/p|_p < p$ .*

It is interesting to compare this with the classical formula for  $\log \Gamma(x)$ :

$$\log \Gamma(x) = -\gamma(x-1) + \sum_{r=2}^{\infty} \frac{(-1)^r \zeta(r)}{r} (x-1)^r \quad \text{for } |x-1| < 1.$$

The idea of having  $p = 1$  give us classical results from a  $p$ -adic formula, while only formal here, is valid in certain formulas for  $\zeta(n)$  and  $\zeta_p(n)$  discussed in [1].

Our next result for  $G_p(x)$  is a set of formulas for  $G_p(x)$  valid on the annular regions  $A_n = \{x: n-1 < \nu(x) < n\}$  for  $n = 1, 2, 3, \dots$ . Since these regions have no points of  $Z_p$  we are able to find series which are almost Laurent series. To simplify the discussion we introduce a function  $G^*$  defined by

$$G^*(x) = \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{\substack{n=0 \\ p \nmid n}}^{p^k-1} f(x+n)$$

where  $f(x) = x \log(x) - x$ .

We can write

$$G^*(x) = \sum_{a=1}^{p-1} \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{\substack{n=0 \\ n \equiv a \pmod{p}}}^{p^k-1} f(x+n).$$

For each value of  $a$  the inner lim is locally analytic for  $x$  with  $x+a \notin pZ_p$ . Therefore  $G^*$  is locally analytic for  $x \in \Omega_p - V_p$ , where  $V_p$  is the set of units in  $Z_p$ .

For  $|x|_p < 1$ ,  $G^*$  coincides with a function defined by Morita [7] in the study of the function he calls  $\Gamma_p(x)$ .

To obtain our formulas for  $G_p$  we need the power series for  $G^*(x)$  at  $x = 0$ .

**THEOREM 10.** *If  $|x|_p < 1$ , then*

$$G^*(x) = M_{\epsilon_p}(\log)(x) + \sum_{r=3}^{\infty} \frac{L_p(r, \bar{\omega}^{r-1})(-1)^r}{r} x^r.$$

$L_p(r, \chi)$  is the Leopoldt  $L$ -function for the character  $\chi$ ,  $\epsilon_p$  is the principal character mod  $p$ ,  $M_{\chi}(f)$  is the Leopoldt  $\chi$ -mean [5], and  $\bar{\omega}$  is the character mod  $p$  defined by

$$\bar{\omega}(n) = \begin{cases} \lim_{k \rightarrow \infty} n^{-p^k} & \text{for } (n, p) = 1, \text{ if } p > 2, \\ 1 & \text{if } n \equiv 1 \pmod{4}, \\ -1 & \text{if } n \equiv 3 \pmod{4} \text{ for } p = 2. \end{cases}$$

**PROOF.** Apply Theorem 3.

This result has also been found by Morita [7].

We are now prepared to find series for  $G_p(x)$  on the annular domains  $A_n = \{x: n-1 < \nu(x) < n\}$ ,  $n \in Z^+$ .

With  $x \in A_n$  we write the equations

$$G_p(x/p^i) - G_p(x/p^{i+1}) = G^*(x/p^i) \quad \text{for } i = 0, 1, \dots, n-1;$$

adding these equations we obtain

$$G_p(x) = G_p\left(\frac{x}{p^n}\right) + \sum_{i=0}^{n-1} G^*\left(\frac{x}{p^i}\right).$$

We now use Theorems 6 and 10 to obtain

**THEOREM 11.** *On the annulus  $A_n$  we have the formula*

$$G_p(x) = \left(\frac{x}{p^n} - \frac{1}{2}\right)\log(x) - \frac{x}{p^n} + M_{\epsilon_p}(\log)\left(\frac{p^n - 1}{p^n - p^{n-1}}\right)x \\ + \sum_{r=3}^{\infty} \frac{L_p(r, \bar{\omega}^{r-1})(-1)^r}{r} \left(\frac{p^n - 1}{p^n - p^{r(n-1)}}\right)x^r + \sum_{r=1}^{\infty} \frac{B_{r+1}p^{nr}x^{-r}}{r(r+1)}.$$

If we define  $A_0$  as  $\{x: |x|_p > 1\}$ , then the above formula is valid for  $n = 0, 1, 2, \dots$ .

5. **Analyticity.** The function  $G_p$  is not an analytic function in the sense of Krasner [4], but its second derivative  $G_p''$  is an analytic function on  $\Omega_p - Z_p$ .

**THEOREM 12.**  $G_p''$  is an analytic function on  $\Omega_p - Z_p$ .

**PROOF.** First for  $a, m \in Z$  we define  $D(a, m) = \{x: x \in \Omega_p, \nu(x - a) > m\}$ . For each  $m \in Z^+$  the set  $A_m = \Omega_p - \cup_{a=0}^{p^m-1} D(a, m)$  is a quasi-connected set.  $\{A_m: m \in Z^+\}$  is nested and  $\cup_{m=1}^\infty A_m = \Omega_p - Z_p$ . Therefore if we can prove  $G_p''$  is an analytic element on each  $A_m$ , i.e. the uniform limit of a sequence of rational functions having no poles in  $A_m$ , then we will know  $G_p''$  is an analytic function on  $\Omega_p - Z_p$ .

If we apply Theorem 7 we can write

$$G_p''(x) = \frac{1}{p^{2m+2}} \sum_{a=0}^{p^{m+1}-1} G_p''\left(\frac{x+a}{p^{m+1}}\right)$$

for each  $m \in Z^+$  and  $x \in \Omega_p - Z_p$ .

If we consider just  $x \in A_m$ , then  $|(x+a)/p^{m+1}|_p > p > 1$  for all  $a \in Z$ . Therefore we may use Theorem 6 and obtain

$$G_p''\left(\frac{x+a}{p^{m+1}}\right) = \sum_{r=0}^\infty \frac{B_r}{[(x+a)/p^{m+1}]^{r+1}}.$$

Since  $|(x+a)/p^{m+1}|_p > p$  for all  $x \in A_m$ , this last series converges uniformly on  $A_m$ .

Thus  $G_p''$  is an analytic element on  $A_m$  and  $G_p''$  is an analytic function on  $\Omega_p - Z_p$ .

6. **An alternative approach.** Earlier we mentioned another approach to the idea of  $p$ -adic  $\log \Gamma(x)$ : to change the functional equation. Of course, it must only be a slight change so we can associate it with  $\log \Gamma(x)$ . We will construct a sequence of such functions, which will be locally holomorphic on  $\Omega_p$  and have  $G_p(x)$  as their pointwise limit.

**DEFINITION.** Let

$$H_N(x) = \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{n=0}^{p^k-1} f_N(x+n) \quad \text{for } N = 1, 2, \dots,$$

where

$$f_N(x) = \begin{cases} x \log(x) - x & \text{if } \nu(x) < N, \\ 0 & \text{if } \nu(x) \geq N. \end{cases}$$

Each  $f_N$  is locally analytic on  $\Omega_p$ , so each  $H_N$  is also locally analytic on  $\Omega_p$ .

We have the following equation

$$H_N(x + 1) = \begin{cases} H_N(x) + \log(x) & \text{if } \nu(x) < N, \\ H_N(x) & \text{if } \nu(x) \geq N. \end{cases}$$

It can be shown that  $H_N(0) = 0$  for each  $N$ , so we have for  $n \in \mathbb{Z}^+$ ,

$$H_N(n + 1) = \log \prod_{t=1}^n t, \\ p^N \nmid t$$

in particular,  $H_N(n + 1) = \log(n!)$  if  $n < p^N$ . The following theorem shows the relation between  $G_p(x)$  and  $H_N(x)$ .

**THEOREM 13.** *If  $x$  is such that  $|x - a|_p > p^{-N}$  for all  $a \in \mathbb{Z}_p$  then  $H_N(x) = G_p(x)$ .*

**PROOF.** Inspection of the definitions of  $G_p(x)$  and  $H_N(x)$ .

Theorem 13 shows us that the sequence  $H_N(x)$ ,  $N = 1, 2, \dots$ , and  $x$  fixed with  $x \notin \mathbb{Z}_p$ , eventually becomes constant with the value  $G_p(x)$ .

For  $x$  with  $\nu(x) \geq 1$  the functions  $H_1$  and  $G^*$  coincide. However, for other  $x$  they are not the same and it is  $H_1$  which is the log of the function on  $\mathbb{Z}_p$  which Morita has called  $\Gamma_p(x)$  [7].

**7. *p*-adic Euler constants.** In a recent paper [6], D. H. Lehmer proves a theorem of Gauss by defining a generalization of Euler's constant. Gauss' theorem is a formula for the logarithmic derivative of the Gamma function at rational points  $r/k$  with  $0 < r < k$ . The formula is notable because it is a constant plus a linear combination of logarithms of integers in  $Q(\sqrt[k]{1})$ .

We shall define *p*-adic Euler constants, give the basic results for them and then prove the *p*-adic version of Gauss' theorem: a formula for  $G'_p(r/f)$  with  $0 < r < f$  and  $\nu(r/f) < 0$ .

We show how Gauss' theorem follows from the classical formula for  $L(1, \chi)$ , and since we have the same expression for  $L_p(1, \chi)$  [3], we have a proof valid in both  $C$  and  $\Omega_p$ .

Lehmer defines the (generalized) Euler constants by

$$\gamma(r, k) = \lim_{\chi \rightarrow \infty} \left[ \sum_{\substack{0 < n < \chi \\ n \equiv r \pmod{k}}} \frac{1}{n} - \frac{1}{k} \log \chi \right]$$

and then, using the relation

$$\psi(r/k) = D^{(1)} \log \Gamma(r/k) = \log k - k\gamma(r, k) \quad \text{for } r, k \in \mathbb{Z}^+, r < k,$$

he proves: If  $r, k \in \mathbb{Z}^+$ ,  $r \leq k$ , then

$$\psi\left(\frac{r}{k}\right) = -\gamma - \log\left(\frac{k}{2}\right) - \frac{\pi}{2} \cot\left(\frac{\pi r}{k}\right) + 2 \sum_{0 < j < k/2} \cos\left(\frac{2\pi rj}{k}\right) \log \sin\left(\frac{\pi j}{k}\right).$$

Gauss' theorem, combined with the functional equation, enables us to calculate  $\psi(x)$  in closed form at every rational value of  $x$  for which the function is defined.

Working in  $\Omega_p$  we can define  $\gamma_p(r, f)$  when  $r, f \in Z, f > 1$ , and find a similar formula for  $G'_p(r/f)$ .

When  $\nu(r/f) < 0$  we define

$$\gamma_p(r, f) = - \lim_{k \rightarrow \infty} \frac{1}{fp^k} \sum_{\substack{m=0 \\ m \equiv r \pmod{f}}}^{fp^k-1} \log(m).$$

When  $\nu(r/f) > 0$  we write  $f = p^k f^*$  with  $(p, f^*) = 1$  and let  $\phi = \phi(f^*)$  (the Euler function). We then define

$$\gamma_p(r, f) = \frac{p^\phi}{p^\phi - 1} \sum_{n \in N(r, f)} \gamma_p(r + nf, p^\phi f)$$

where

$$N(r, f) = \{n: 0 \leq n < p^\phi, nf + r \not\equiv 0 \pmod{p^{\phi+k}}\}.$$

Theorem 1 applies to show  $\gamma_p(r, f)$  exists.

To obtain Gauss' theorem in  $\Omega_p$  we need several results which are mostly the same as Lehmer has given for  $C$ . The proofs follow from the definition of  $\gamma_p(r, f)$ , previous results for  $G'_p(x)$  and Theorem 18. We will write  $\psi_p(x) = G'_p(x)$ .

- THEOREM 14.** (i) If  $d|(r, f)$ , then  $f\gamma_p(r, f) = (f/d)\gamma_p(r/d, f/d) - \log d$ .  
 (ii) If  $\nu(r/f) < 0$  and  $0 < r < f$ , then  $\psi_p(r/f) = -\log f - f\gamma_p(r, f)$ .  
 (iii)  $\gamma_p(r, f) = \gamma_p(f - r, f)$ .  
 (iv) If  $b \in Z^+$ , then

$$\gamma_p(r, f) = \sum_{n=0}^{b-1} \gamma_p(r + nf, bf).$$

- (v) If  $p^\mu \equiv 1 \pmod{f^*}$  and  $\nu(r/f) > 0$ , then

$$\gamma_p(r, f) = \frac{p^\mu}{p^\mu - 1} \sum_{\substack{n=0 \\ nf+r \not\equiv 0 \pmod{p^{\mu+k}}} }^{p^\mu-1} \gamma_p(r + nf, p^\mu f).$$

We are going to need a  $p$ -adic analog of Euler's constant. The value

$$\gamma_p = \gamma_p(0, 1) = -\frac{p}{p-1} \lim_{k \rightarrow \infty} \frac{1}{p^k} \sum_{\substack{m=1 \\ (m,p)=1}}^{p^k-1} \log(m)$$

fits our formulas precisely as we need. Morita has also realized the connection between Euler's constant and  $M_p(\log)$  and he gives a slightly different value [7].

Lehmer has defined the formula  $\Phi(f)$  by

$$\Phi(f) = \sum_{\substack{r=1 \\ (r,f)=1}}^f \gamma(r, f).$$

He then proves

$$\Phi(f) = \frac{\phi(f)}{f} \gamma + \frac{\phi(f)}{f} \sum_{q|f} \frac{\log q}{q-1}.$$

In this formula,  $q$  is prime,  $\phi$  is the Euler  $\phi$ -function and  $\gamma$  is Euler's constant.

We define

$$\Phi_p(f) = \sum_{\substack{r=1 \\ (r,f)=1}}^f \gamma_p(r, f) \quad \text{when } \nu(f) > 0.$$

We then have

**THEOREM 15.**

$$\Phi_p(f) = \frac{\phi(f)}{f} \gamma_p + \frac{\phi(f)}{f} \sum_{q|f} \frac{\log q}{q-1}.$$

**PROOF.** We can use Theorem 14(iv) with  $b = f_1/f$  to show that if  $f$  has the same distinct prime factors as  $f_1$  and  $f|f_1$  then  $\Phi_p(f) = \Phi_p(f_1)$ .

It is then sufficient to consider square free  $f$ . This is accomplished by induction on the number of prime factors of  $f$ .

We will need the following algebraic identity.

**THEOREM 16.** *If  $\zeta$  is a primitive  $f$ th root of unity,  $f > 1$ ,  $\epsilon_f$  the principal character mod  $f$  and  $\tau_a(\epsilon_f)$  the Gauss sum,*

$$\tau_a(\epsilon_f) = \sum_{n=1}^f \epsilon_f(n) \zeta^{an},$$

then

$$\prod_{a=1}^{f-1} (1 - \zeta^{-a})^{\tau_a(\epsilon_f)} = \prod_{q|f} q^{-\phi(f)/(q-1)}.$$

*The product on the right side is over the distinct prime divisors of  $f$ .*

PROOF. Let  $\omega_r$  be a primitive  $r$ th root of unity and  $Q_r = Q(\omega_r)$  for  $r = 2, 3, \dots$

We observe that

$$\tau_a(\varepsilon_f) = \sum_{\substack{n=1 \\ (n,f)=1}}^{f-1} \zeta^{an} = \text{tr}_{Q_f/Q}(\zeta^a).$$

When we group together conjugate elements we have

$$(1) \quad \prod_{a=1}^{f-1} (1 - \zeta^{-a})^{\tau_a(\varepsilon_f)} = \prod_{r|f} (N_{Q_r/Q}(1 - \omega_r))^{\text{tr}_{Q_r/Q}(\omega_r)}.$$

Examination of the minimal polynomial of  $\omega_r$  shows that:

- (i) if  $r$  is not square free, then  $\text{tr}_{Q_r/Q}(\omega_r) = 0$ ;
- (ii) if  $r$  is square free, but not prime, then  $N_{Q_r/Q}(1 - \omega_r) = 1$ ;
- (iii) if  $r$  is prime,  $N_{Q_r/Q}(1 - \omega_r) = r$ ;
- (iv) if  $r$  is prime

$$\text{tr}_{Q_r/Q}(\omega_r) = \frac{\phi(f)}{r-1} \text{tr}_{Q_r/Q}(\omega_r) = \frac{-\phi(f)}{r-1}.$$

Placing these four values into (1) establishes the theorem.

Now we state Gauss' theorem in  $\Omega_p$ .

THEOREM 17. *If  $r, f \in \mathbb{Z}^+$ ,  $r < f$  and  $v(r/f) < 0$ , then*

$$\psi_p(r/f) = -\log f - \gamma_p + \sum_{a=1}^{f-1} \zeta^{-ar} \log(1 - \zeta^a).$$

If  $\psi_p$  is replaced by  $\psi$  and  $\gamma_p$  by  $\gamma$  we have Gauss' theorem in  $C$ . Of course  $\log$  is either  $p$ -adic or complex as required.

PROOF. Since we have shown (Theorem 14(ii)) that  $\psi_p(r/f) = -\log f - f\gamma_p(r, f)$ , it will be sufficient to prove

THEOREM 18. *If  $f > 1$  and  $\zeta$  is a primitive  $f$ th root of unity, then*

$$f\gamma_p(r, f) = \gamma_p - \sum_{a=1}^{f-1} \zeta^{-ar} \log(1 - \zeta^a).$$

*Notice that we do not need the restriction  $v(r/f) < 0$  for this result.*

PROOF. We begin by assuming  $(r, f) = 1$  and  $v(r/f) < 0$  and proceed to evaluate  $\sum_{\chi \neq \varepsilon_f} \bar{\chi}(r) L_p(1, \chi)$  in two different ways. The sum is over all non-principal characters mod  $f$ .

For  $\chi$  not principal and if  $p|f$ , we have the formula [3]

$$L_p(1, \chi) = -\frac{1}{f} \sum_{a=1}^{f-1} \tau_a(\chi) \log(1 - \zeta^{-a}).$$

(NOTE: Iwasawa gives this formula in a form valid only for primitive characters, but if  $\bar{\chi}(a)\tau(\chi)$  is replaced by  $\tau_a(\chi)$ , then his proof can be modified to be valid for all nonprincipal characters.)

This is the same as the formula for  $L(1, \chi)$  in  $C$ . Using this result and Theorems 15 and 16 we have

$$(*) \quad \sum_{\chi \neq \epsilon_f} \bar{\chi}(r)L_p(1, \chi) = \frac{\phi(f)\gamma_p}{f} - \Phi_p(f) - \frac{\phi(f)}{f} \sum_{a=1}^{f-1} \zeta^{ar} \log(1 - \zeta^{-a}).$$

This result is also valid, correctly interpreted, in  $C$ .

If we use the expression [5] that

$$L_p(1, \chi) = - \lim_{k \rightarrow \infty} \frac{1}{fp^k} \sum_{n=0}^{fp^k-1} \chi(n) \log n \quad \text{where } (n, p) = 1,$$

we obtain

$$(**) \quad \sum_{\chi \neq \epsilon_f} \bar{\chi}(r)L_p(1, \chi) = \phi(f)\gamma_p(r, f) - \Phi_p(f).$$

(\*\*) is obtained in  $C$  by using

$$L(1, \chi) = \lim_{x \rightarrow \infty} \sum_{0 < n < x} \frac{\chi(n)}{n}.$$

Equating (\*) and (\*\*) yields Theorem 18 in the case where  $(r, f) = 1$  and  $\nu(r/f) < 0$ .

Now suppose  $(r, f) = d > 1$  and  $\nu(r/f) < 0$ . Then we can use 14(i) to obtain

$$f\gamma_p(r, f) = -\log d + \gamma_p - \sum_{a=1}^{f/d-1} \zeta^{-ar} \log(1 - \zeta^{ad}).$$

We can factor  $1 - \zeta^{ad}$  and obtain

$$f\gamma_p(r, f) = -\log d + \gamma_p - \sum_{a=1}^{f/d-1} \sum_{b=0}^{d-1} (\zeta^a \lambda^b)^{-r} \log(1 - \zeta^a \lambda^b)$$

where  $\lambda$  is a primitive  $d$ th root of unity.

Since

$$\{\zeta^a \lambda^b : 0 \leq a < f/d, 0 \leq b < d-1\} = \{\zeta^a : 0 \leq a < f\},$$

$$\begin{aligned} f\gamma_p(r, f) &= -\log d + \gamma_p - \sum_{a=1}^{f-1} \zeta^{-ar} \log(1 - \zeta^a) + \sum_{b=1}^{d-1} \log(1 - \lambda^b) \\ &= \gamma_p - \sum_{a=1}^{f-1} \zeta^{-ar} \log(1 - \zeta^a). \end{aligned}$$

This completes the proof of Gauss' theorem, but we have not yet completed the proof of Theorem 18.

If  $\nu(r, f) \geq 0$  we can use the definition of  $\gamma_p(r, f)$  and the case of Theorem 18 already proven to show

$$f\gamma_p(r, f) = \gamma_p - \frac{1}{p^\phi - 1} \sum_{a=1}^{p^\phi f - 1} \eta^{-ar} \log(1 - \eta^a) \sum_{n \in N(r, f)} \eta^{-anf},$$

where  $\eta$  is a primitive  $p^\phi f$  root of unity.

The last sum on the right is  $p^\phi - 1$  if  $p^\phi | a$  and  $-\eta^{ar-ap^\phi}$  if  $p^\phi \nmid a$ . Thus

$$f\gamma_p(r, f) = \gamma_p - \sum_{a=1}^{f-1} \zeta^{-ar} \log(1 - \zeta^a) + \frac{1}{p^\phi - 1} \sum_{\substack{a=1 \\ p^\phi \nmid a}}^{p^\phi f - 1} \zeta^{-ar} \log(1 - \eta^a).$$

The last sum on the right is 0.  $\square$

We have seen that  $\psi_p$  is locally holomorphic on  $\Omega_p - Z_p$  and  $\psi'_p$  is Krasner-analytic on this domain. We have also shown that the formula

$$(*) \quad -\log f - f\gamma_p(r, f)$$

depends only on the ratio  $r/f$  and that for  $\nu(r/f) < 0$ , and  $0 < r < f$ ,

$$\psi_p(r/f) = (*) = -\log f - \gamma_p + \sum_{a=1}^{f-1} \zeta^{-ar} \log(1 - \zeta^a).$$

Since  $(*)$  is defined for  $r, f$  with  $\nu(r/f) \geq 0$ , it is tempting to use  $(*)$  to extend the definition of  $\psi_p$  onto the rational numbers in  $Z_p$ . However, this "continuation" would not retain the other properties of  $\psi_p$ . The values of  $(*)$ , though, are related to functions similar to  $\psi_p$  when  $\nu(r, f) \geq 0$  and we have

**THEOREM 19.** *Given  $\nu(r/f) \geq 0$ , then for any  $\mu$  such that  $p^\mu \equiv 1 \pmod{f^*}$  we have*

$$(*) = \frac{p^\mu}{p^\mu - 1} H'_\mu\left(\frac{r}{f}\right) = -\log f - \gamma_p + \sum_{a=1}^{f-1} \zeta^{-ar} \log(1 - \zeta^a).$$

$H_N$  is discussed at the end of §4.

**PROOF.** This follows directly from previous results.

Since  $H_1$  (on  $Z_p$ ) is the logarithm of Morita's  $\Gamma_p$  [7], we have a

**COROLLARY.** *If  $0 < r < f$ ,  $\nu(r/f) \geq 0$  and  $f^* | (p - 1)$ , then*

$$\frac{\Gamma'_p}{\Gamma_p}\left(\frac{r}{f}\right) = (1 - 1/p) \left( -\log f - \gamma_p + \sum_{a=1}^{f-1} \zeta^{-ar} \log(1 - \zeta^a) \right).$$

### REFERENCES

1. J. Diamond, *On the values of p-adic L-functions at positive integers* (to appear).
2. J. Fresnel, *Nombres de Bernoulli et fonctions L p-adiques*, Ann. Inst. Fourier (Grenoble) 17 (1967), fasc. 2, 281-333. (1968). MR 37 #169.

3. K. Iwasawa, *Lectures on  $p$ -adic  $L$ -functions*, Ann. of Math. Studies, no. 74, Princeton Univ. Press, Princeton, N. J., 1972. MR 50 #12974.
4. M. Krasner, *Rapport sur le prolongement analytique dans les corps valués complets par la méthode des éléments analytiques quasi-connexes*, Table Ronde d'Analyse non archimédienne (Paris, 1972), Bull. Soc. Math. France, Mém. No. 39-40, Soc. Math. France, Paris, 1974, pp. 131-254. MR 52 #6033.
5. T. Kubota and H. W. Leopoldt, *Eine  $p$ -adische Theorie der Zetawerte*. I. J. Reine Angew. Math. 214/215 (1964), 328-339. MR 29 #1199.
6. D. H. Lehmer, *Euler constants for arithmetical progressions*, Acta Arith. 27 (1975), 125-142. MR 51 #5468.
7. Y. Morita, *A  $p$ -adic analog of the  $\Gamma$  function*, (to appear).
8. N. Nielsen, *Die Gammafunktion*, Bände I, II, Chelsea, New York, 1965. MR 32 #2622.

DEPARTMENT OF MATHEMATICS, QUEENS COLLEGE, FLUSHING, NEW YORK 11367