

## THE $\alpha$ -UNION THEOREM AND GENERALIZED PRIMITIVE RECURSION<sup>(1)</sup>

BY

BARRY E. JACOBS<sup>(2)</sup>

**ABSTRACT.** A generalization to  $\alpha$ -recursion theory of the McCreight-Meyer Union Theorem is proved. **THEOREM.** Let  $\Phi$  be an  $\alpha$ -computational complexity measure and  $\{f_e \mid e < \alpha\}$  an  $\alpha$ -r.e. strictly increasing sequence of  $\alpha$ -recursive functions. Then there exists an  $\alpha$ -recursive function  $k$  such that  $C_k^\Phi = \bigcup_{e < \alpha} C_e^\Phi$ . The proof entails a no-injury cancellation atop a finite-injury priority construction and necessitates a blocking strategy to insure proper convergence.

Two infinite analogues to  $(\omega)$ -primitive recursive functions are studied. Although these generalizations coincide at  $\omega$ , they diverge on all admissible  $\alpha > \omega$ . Several well-known complexity properties of primitive recursive functions hold for one class but fail for the other. It is seen that the Jensen-Karp ordinally primitive recursive functions restricted to admissible  $\alpha > \omega$  cannot possess natural analogues to Grzegorzczuk's hierarchy.

**0. Introduction.** The motivation for the study of computation on infinite ordinals stems from several areas of mathematical logic. Takeuti [26], [27] was concerned with the problem of the reduction of the consistency of set theory to that of a theory of ordinal numbers. Machover [18], seeking to generalize model and recursion theoretic concepts to infinitary languages, developed a recursion theory on regular infinite cardinals. Questions of definability and their relation to higher logics and languages moved Kreisel [11], and later Kreisel and Sacks [12], to develop a recursion theory on Church and Kleene's  $\omega^{ck}$ , called metarecursion theory.

From a set theoretic point of view, Jensen and Karp [9] developed the notion of an ordinally primitive recursive function ( $\text{Prim}_0$ ). Karp's motivation came from an investigation into the classification of infinitary languages, Jensen's from the study of levels of Gödel's constructible hierarchy. A key

---

Received by the editors April 16, 1976.

*AMS (MOS) subject classifications* (1970). Primary 02F27, 68A20; Secondary 02F35, 02F20.

*Key words and phrases.*  $\alpha$ -recursion theory, admissible ordinal,  $\alpha$ -computational complexity measure,  $\alpha$ -recursively enumerable, priority argument,  $\alpha$ -complexity class, primitive recursive ordinal functions, Grzegorzczuk hierarchy.

<sup>(1)</sup>The results of this paper (with the exception of §5) are contained in the author's Ph.D. Thesis, Courant Institute of Mathematical Sciences, New York University, 1975 which was written under the supervision of Professor Martin D. Davis.

<sup>(2)</sup>This research was supported by the National Science Foundation, Grants NSF-DCR71-02039-AOB and MCS 76-07129.

result was a general bounding phenomenon (Stability Theorem) of  $\text{Prim}_0$  functions on the primitive recursive set functions.

Kripke [13] (and independently Platek [20]) arrived at a unifying concept for the aforementioned cases; namely, the notion of admissible ordinal. The theory of computation on admissible ordinals  $\alpha$  became known as  $\alpha$ -recursion theory. By setting up an equation calculus (similar to Kleene (cf. [10]) for  $\alpha = \omega$ ), Kripke was able to develop enough  $\alpha$ -recursion theory to establish an infinite analogue to Kleene's  $T$  predicate and subsequently a Normal Form Theorem. From this he was able to assert that all of the results of unrelativized ordinary recursion theory (as found in Kleene [10]) hold in  $\alpha$ -recursion theory.

There is strong interaction between the Jensen-Karp primitive recursive ordinal functions and  $\alpha$ -recursion theory. Specifically, for admissible  $\alpha$ , the closure of  $\text{Prim}_0$  under a regular (cf. [3, p. 38])  $\alpha$ -bounded min operator yields the  $\alpha$ -recursive functions. Further, the construction of analogues to Kleene's  $T$  predicate for  $\alpha$ -recursion theory may be developed from only  $\text{Prim}_0$  relations and functions, independent of  $\alpha$ .

Deep results of ordinary recursion theory concern the notion of relativization and often require use of the powerful technique called the priority argument. Sacks and Simpson [21] introduced the priority method into  $\alpha$ -recursion theory in their  $\alpha$ -analogue to the Friedberg-Muchnik solution to Post's problem. Since then Sacks' students and coworkers have successfully demonstrated that major priority argument results generalize to  $\alpha$ . The proofs of these results generally require vast modification from the  $\omega$  situation in both the construction and their verifications. (See Lerman [16], Shore [23], Leggett and Shore [15].) An excellent survey can be found in Shore [25].

Another subarea of ordinary recursion theory, abstract complexity theory, has its major theorems obtained in a manner similar to that of relativized recursion theory. Founded upon several axioms of measure of complexity of computation (cf. [1]), deep results are established through constructions also based upon priority mechanisms. (See [1], [2], [19] and [29].) It was shown in [7] and [8] that the major results of this area generalize to  $\alpha$ -recursion theory.

Many recursion theorists, however, are quick to point out differences between the two types of constructions. Consequently, they contend that the name "priority argument" be reserved only for the former type. Specifically, the former generally involves multiple (priority based) cancellations or injuries, the latter single (sometimes double) no-injury cancellations. However, an interesting phenomenon occurs upon generalizing the McCreight-Meyer Union Theorem [19] to  $\alpha$ -recursion theory. While the  $\omega$ -proof employs a typical no-injury cancellation construction, its lift to  $\alpha$  requires expansion to finite injury.

An outline of the paper is as follows.

In §1 we introduce the basic notions of  $\alpha$ -recursion theory and  $\alpha$ -complexity theory that underlie this paper.

In §2 we present a statement of the  $\alpha$ -Union Theorem together with a discussion of the differences between our proof and the McCreight-Meyer one for the case  $\alpha = \omega$ . While the latter employs a no-injury cancellation construction the former requires a no-injury atop a finite injury cancellation construction. Furthermore, the introduction of finite injury to the proof necessitates a blocking strategy in order to ensure proper convergence.

In §3 we present the actual priority construction that yields an  $\alpha$ -recursive  $k$  required by the  $\alpha$ -Union Theorem. This is followed, in §4, by the verification that the construction is correct.

In §5 two generalizations of the class of (ordinary) primitive recursive functions ( $\omega$ -Prim) are studied. Though both coincide on  $\omega$ , they are seen to diverge on all admissible  $\alpha > \omega$ . The basis of the investigation is a sequence of three propositions (A, B and C) characterizing an arbitrary class, Prim, of functions. Any class which is a model of all three is seen, via an application of the  $\alpha$ -Union Theorem, to decompose into an  $\alpha$ -hierarchy based on computational complexity. Further, there also exists a single  $\alpha$ -recursive bound on the complexity of any function in the class.

The class  $\omega$ -Prim serves as a model for the three propositions implying the above two consequences. Of the two generalizations, it is seen that one satisfies A but neither B nor C; the second satisfies A and B while that of C is left open. A consequence of the failure of a class to satisfy Proposition C is the nonexistence of an  $\alpha$ -analogue ( $\alpha > \omega$ ) to Grzegorzczk's hierarchy for that class. In particular, it is shown that this deficiency holds for the well-known class Prim<sub>0</sub> of Jensen and Karp.

Throughout the paper several open problems are proposed.

**1. Preliminaries.** Let  $L_\alpha$  be the collection of sets obtained from Gödel's [4] transfinite hierarchy of constructible sets before  $\alpha$ .  $\alpha$  is  $\Sigma_1$  *admissible* if  $L_\alpha$  satisfies the replacement axiom schema of ZF for  $\Sigma_1$  formulae. From now on  $\alpha$  is taken as a fixed  $\Sigma_1$  admissible ordinal.

We employ usual set theoretic notation:  $\cup A$  for the union of  $A$ ;  $\cup_{\varepsilon < \tau} G_\varepsilon$  for the indexed union of  $G_\varepsilon$ ,  $\varepsilon < \tau$ ;  $A - B$  for the set difference between  $A$  and  $B$ ;  $f|B$  for the mapping  $f$  restricted to  $B$ ;  $f[B]$  for the range of  $f|B$ ;  $\delta \in B$  for  $\delta$  an element of  $B$ ;  $A \subseteq B$  for  $A$  a subset of  $B$ ,  $A \subset B$  for  $A$  a proper subset of  $B$ ;  $\text{dom}(f)$ ,  $\text{rng}(f)$  for the domain and range of  $f$ ; and  $f: A \rightarrow B$  for  $f$  a map from  $A$  to  $B$ .

A partial map  $f: \alpha \rightarrow \alpha$  is  $\alpha$ -partial recursive if its graph has a  $\Sigma_1$  definition over  $L_\alpha$  (with parameters in  $L_\alpha$ ) and is  $\alpha$ -recursive if it is also total on  $\alpha$ . A nonempty subset  $A$  of  $\alpha$  is  $\alpha$ -recursively enumerable ( $\alpha$ -r.e.) if it is the range of an  $\alpha$ -recursive function or, equivalently, the domain of an  $\alpha$ -partial

recursive function.  $A$  is  $\alpha$ -recursive if it and its complement (with respect to  $\alpha$ ) are  $\alpha$ -recursively enumerable. Since there exists a one-one  $\alpha$ -recursive map from  $\alpha$  to  $L_\alpha$ , we need only concern ourselves with functions on  $\alpha$  and subsets of  $\alpha$ .

The main point about any  $\Sigma_1$  admissible  $\alpha$  is that one can perform  $\Delta_1$  ( $\alpha$ -recursive) recursions in  $L_\alpha$ . In particular, one can Gödel number the  $\alpha$ -recursively enumerable subsets of  $\alpha$  (employing  $\alpha$ -recursive pairing  $\langle \cdot \cdot \cdot \rangle$  and projection  $\pi_i$  functions) and, consequently, all the  $\alpha$ -partial recursive functions.

We call a subset  $A \subset \alpha$   $\alpha$ -bounded (or simply bounded) if there exists a  $\beta < \alpha$  so that  $\sigma \in A \rightarrow \sigma < \beta$ .  $A$  is  $\alpha$ -finite if it is both  $\alpha$ -recursive and  $\alpha$ -bounded, alternatively, if  $A$  is a member of  $L_\alpha$ . A consequence of the definitions of  $\Sigma_1$  admissibility and  $\alpha$ -partial recursiveness is

1.1 FACT. If  $f: \alpha \rightarrow \alpha$  is  $\alpha$ -partial recursive and  $A$  an  $\alpha$ -finite subset of  $\text{dom}(f)$ , then  $f[A]$  is  $\alpha$ -finite.

A key notion in  $\alpha$ -recursion theory is that of projecta. The  $\Sigma_1$ -projectum (or simply projectum) of  $\alpha, \alpha^*$ , is the least ordinal  $\beta < \alpha$  such that there exists a one-one  $\alpha$ -recursive  $t$  (referred to as the projection) mapping from  $\alpha$  into  $\beta$ . An  $\alpha$ -recursive projection map  $t: \alpha \rightarrow \alpha^*$  often serves as a vehicle in  $\alpha$ -recursive constructions which hinge upon the notion of "priority".

For many  $\Sigma_1$  admissibles there exist subsets of  $\alpha$  that are  $\alpha$ -recursively enumerable and bounded (below  $\alpha$ ), but not  $\alpha$ -finite. However, if the bound is small enough,  $\alpha$ -finiteness must occur.

1.2 FACT. If  $\eta < \alpha^*$  and  $A$  is a  $\Sigma_1$  subset of  $\eta$ , then  $A$  is  $\alpha$ -finite.

As is often the case  $\alpha$ -priority constructions (this paper, in particular) require shorter listings than  $\alpha^*$ . One such ordering is supplied by  $\Sigma_2$  cofinality. For  $\gamma < \alpha$ ,  $h: \gamma \rightarrow \alpha$  is a  $\Sigma_2$  cofinality function if  $h$  is  $\Sigma_2$  and its range is unbounded in  $\alpha$ . The  $\Sigma_2$  cofinality of  $\alpha$ , called  $\sigma_2\text{cf}(\alpha)$ , is the least  $\gamma < \alpha$  for which there is a  $\Sigma_2$  cofinality function  $h: \gamma \rightarrow \alpha$ .

In [1] Blum axiomatizes several properties common to most interesting measures of the complexity of computation on partial recursive functions. A generalization to  $\alpha$ -recursion theory of his notion of abstract complexity measure forms a basis of this paper.

1.3 DEFINITION. An  $\alpha$ -computational complexity measure  $\Phi$  is an enumeration (in  $\alpha$ ) of the  $\alpha$ -partial recursive functions  $\{\phi_\varepsilon \mid \varepsilon < \alpha\}$  to which are associated the  $\alpha$ -partial recursive  $\alpha$ -step counting functions  $\{\Phi_\varepsilon \mid \varepsilon < \alpha\}$  for which the following axioms hold.

- (1) for all  $\beta, \varepsilon < \alpha$ ,  $\phi_\varepsilon(\beta)$  is defined if and only if  $\Phi_\varepsilon(\beta)$  is defined;
- (2) the predicate  $M(\varepsilon, \beta, \gamma) \Leftrightarrow \Phi_\varepsilon(\beta) = \gamma$  is  $\alpha$ -recursive; and
- (3) the  $\alpha$ -recursive analogues to the  $S_n^m$ -, Universal Function, and Recursion Theorems hold for the enumerations  $\{\phi_\varepsilon\}$  and  $\{\Phi_\varepsilon\}$  (cf. [10]).

Implicit in the above definition is the capability to retrieve, given any index  $\varepsilon < \alpha$ , both the function  $\phi_\varepsilon$ , in the form of an algorithm, and its  $\alpha$ -step counter,  $\Phi_\varepsilon$ . Clearly when  $\alpha = \omega$ , the definition reduces to that of Blum's. Several illustrations of  $\alpha$ -computational complexity measures appear in [7].

Another generalization of a complexity oriented notion is that of  $\alpha$ -complexity class.

**DEFINITION.** For an  $\alpha$ -complexity measure  $\Phi$  and  $\alpha$ -recursive function  $s$  the  $\alpha$ -complexity class,  $C_s^\Phi$ , is the set  $\{\phi_\sigma | \phi_\sigma \text{ is total and } \Phi_\sigma(\beta) \leq s(\beta) \text{ for all but an } \alpha\text{-finite set of } \beta\}$ .

Hence,  $C_s^\Phi$  ( $C_s$  when  $\Phi$  is understood) is the set of all  $\alpha$ -recursive functions whose  $\alpha$ -complexity is bounded by  $s$  on all but an  $\alpha$ -finite subset of  $\alpha$ . Since  $s$  is  $\alpha$ -recursive, the substitution of bounded for  $\alpha$ -finite yields an equivalent definition for  $C_s^\Phi$ .

Finally, we call a sequence of  $\alpha$ -partial recursive functions  $\{f_\varepsilon | \varepsilon < \alpha\}$   $\alpha$ -recursively enumerable if there is an  $\alpha$ -recursively enumerable set of algorithms so that each function is named at least once.

**2. Discussion.** One objective of this paper is to generalize or "lift" to  $\alpha$  the well-known McCreight-Meyer Union Theorem [19]. Namely, that the  $\alpha$ -complexity classes of an increasing  $\alpha$ -recursively enumerable sequence of  $\alpha$ -recursive functions constitute a single  $\alpha$ -complexity class.

**2.1  $\alpha$ -UNION THEOREM.** *Let  $\Phi$  be an  $\alpha$ -computational complexity measure and  $\{f_\varepsilon | \varepsilon < \alpha\}$  an  $\alpha$ -recursively enumerable sequence of  $\alpha$ -recursive functions. Suppose for  $\varepsilon, \tau$  and  $\beta < \alpha$ ,  $f_\varepsilon(\beta) < f_\tau(\beta)$ , whenever  $\varepsilon < \tau$ . Then there exists an  $\alpha$ -recursive function  $k(\beta)$  such that  $C_k^\Phi = \bigcup_{\varepsilon < \alpha} C_{f_\varepsilon}^\Phi$ .*

First consider  $k(\beta) = f_\beta(\beta)$  as a possible candidate for the bounding function. Observe that for  $\varepsilon < \alpha$  the set  $\{\beta | k(\beta) < f_\varepsilon(\beta)\}$  is  $\alpha$ -recursive and bounded (by  $\varepsilon$ ), hence  $\alpha$ -finite. Therefore,  $C_{f_\varepsilon} \subseteq C_k$  and  $\bigcup_{\varepsilon < \alpha} C_{f_\varepsilon} \subseteq C_k$  for all  $\varepsilon < \alpha$ .

However, the opposite inclusion may not necessarily hold. For there might exist some  $\alpha$ -recursive  $\phi_v$  for which  $\Phi_v(\beta) \leq k(\beta)$  on all but an  $\alpha$ -finite set but for each  $\varepsilon < \alpha$ ,  $\Phi_v(\beta)$  exceeds  $f_\varepsilon(\beta)$  for an unbounded set of  $\beta$ . Therefore  $\phi_v$  would be in  $C_k$  but *not* in the union of all the  $C_{f_\varepsilon}$ .

One possible remedy is seen in the McCreight-Meyer proof of the ( $\omega$ -) Union Theorem. There a recursive function  $k$  is developed via a construction founded upon a priority mechanism. Namely, at stage  $s$  ( $< \omega$ ) of the construction a guess is established ( $\langle s, s \rangle$ ) that  $f_s(n) \geq \Phi_s(n)$  almost everywhere (a.e.). Next a search is made through previous guesses ( $\langle v, t \rangle$ ) for those which prove to be incorrect on input  $s$ ; that is,  $f_t(s) < \Phi_v(s)$ . If none are found the value of  $k$  on  $s$  is  $f_s(s)$ . Otherwise, the incorrect guess with highest priority,  $\langle v', t' \rangle$  (lowest value of  $v'$ ), is replaced by a new guess,  $f_s(n) \geq \Phi_{v'}(n)$

a.e.  $(\langle v', s \rangle)$ , and the value of  $k(s)$  becomes  $f_{i'}(s)$ .

Now if  $\phi_v$  were not in the  $\omega$ -union, then it cannot be in  $C_k$ . Since  $\phi_v \notin \bigcup_{i < \omega} C_{f_i}$ , then for all  $i < \omega$ ,  $\phi_v \notin C_{f_i}$ , hence  $f_i(n) < \Phi_v(n)$  infinitely often. Consequently, there will be infinite sequences  $\{s_i | i < \omega\}$  and  $\{\langle v, t_i \rangle | i < \omega\}$  such that  $\langle v, t_i \rangle$  is the incorrect guess of highest priority at stage  $s_i$ , forcing  $k$  below  $\Phi_v$  at least as often.

On the other hand,  $\phi_v$  in  $\bigcup_{i < \omega} C_{f_i}$  implies  $\phi_v$  is in  $C_k$ . Since  $\phi_v$  is in the union, there exist  $n_0$  and  $t_0 < \omega$  such that  $\Phi_v(n) < f_{i_0}(n)$  for all  $n > n_0$ . Ultimately some stage of the construction must be reached so that all assignments to  $k$  will be made either through  $f_s(s)$ ,  $s > t_0$  or  $f_i(s)$ ,  $t > t_0$ . The increasingness of the  $\{f_i\}$  therefore implies  $k$  exceeds  $\Phi_v$  almost everywhere.

A generalization of the Union Theorem to  $\alpha$  necessitates a complete overhaul of the  $\omega$  proof. Although we have enumerations in  $\alpha$  of the functions  $\{\phi_\beta\}$ ,  $\{\Phi_\beta\}$  and  $\{f_\beta\}$ , we cannot use  $\alpha$  (as we did  $\omega$  above) as a basis of our priorities. A difficulty is manifested in the fact that segments bounded below  $\alpha$  can be mapped one-to-one onto unbounded (hence,  $\alpha$ -infinite) sequences of  $\alpha$ . Thus, we are *not* able to claim that if  $\{\beta | \Phi_v(\beta) > f_\tau(\beta)\}$  is unbounded then ultimately a stage will have to be reached at which the guess  $\langle v, \tau \rangle$  will have highest priority and therefore be cancelled.

The usual solution to this is to make the priority listing shorter than  $\alpha$  with one such vehicle being the  $\Sigma_1$  projectum  $\alpha^*$  of  $\alpha$ . This, in fact, forms the basis of the lifts to  $\alpha$  of the Blum-Rabin Complexity and the Borodin Gap Theorems in [8]. The key point is that below  $\alpha^*$ ,  $\alpha$ -recursive enumerability is tantamount to  $\alpha$ -finiteness. Thus we cancel indices  $\varepsilon < \alpha$  only when their images under the  $\Sigma_1$  projection map are the smallest cancellable. Standard arguments then show that for any  $\varepsilon < \alpha$  the collection of stages at which images of indices having higher priority than  $\varepsilon$  (images below that of  $\varepsilon$  in  $\alpha^*$ ) are cancelled is  $\alpha$ -finite.

However, this approach will not totally suffice in our situation. For although we know  $\Phi_\varepsilon(\beta) > f_\tau(\beta)$  unboundedly often implies that guess  $\langle v', \tau \rangle$ ,  $v' < \alpha^*$ , will ultimately be cancelled, another problem arises. Namely, that for any  $\tau_0 < \alpha$ , there must be a stage  $\sigma_0$  such that subsequent assignments to  $k$  are made *through cancelled guesses*  $\langle v, \tau \rangle$  where  $\tau > \tau_0$ . In other words, there is no reason that for some  $\tau_0$ , an  $\alpha$ -infinite number of guesses  $\langle v, \tau \rangle$  may exist for which  $\tau$  is smaller than  $\tau_0$ .

We wrestle with this latter problem by formulating  $k$  through a *finite-injury-atop-a-no-injury-cancellation-construction*. The latter feature ensures that unboundedly often wrong guesses will ultimately be cancelled (or "popped"). The former ensures that " $\tau$ " components of cancelled incorrect guesses never  $\alpha$ -infinitely often regress below any  $\tau_0 < \alpha$ .

However (as demonstrated in [17], [23] and [24]) when injury arguments are

lifted to  $\alpha$ , one often requires even shorter listings than the projectum. For although one may have  $\alpha$ -finitely many injury sets (here, sets of stages at which guesses with fixed priority values  $< \alpha^*$  are injured) and that each set may be proven  $\alpha$ -finite, the union of these sets may grow without bound. (See Shore [25] for the usual illustration of  $\aleph_\omega$ .)

Throughout this paper  $\mu = \sigma 2cf(\alpha)$ ,  $h$  the corresponding  $\Sigma_2$  cofinality function ( $h: \mu \rightarrow \alpha$ ) and  $t$  the one-one  $\alpha$ -recursive projection map ( $t: \alpha \rightarrow \alpha^*$ ). The approach we follow is to formulate a construction that implements a strategy first introduced by Sacks and Simpson [21] and later developed and strongly expedited by Shore [22], [23], [24]. The idea is to segment  $\alpha^*$  into a chain of blocks having type equal to the  $\Sigma_2$  cofinality of  $\alpha$ . The block associated with  $\rho < \mu$ , called  $B_\rho$  (the  $\rho$ th block), is simply the initial segment of  $\alpha^*$  bounded above by  $t \circ h(\rho)$ .

2.2 LEMMA.  $t \circ h$  projects all of  $\mu$  into an unbounded sequence in  $\alpha^*$ .

PROOF. Let  $\delta < \alpha^*$  to find some  $\rho < \mu$  where  $t \circ h(\rho) > \delta$ . Since  $\delta < \alpha^*$ , by  $\Sigma_1$  admissibility and the  $\Sigma_1$ -ness of  $t^{-1}$ , there is a  $\beta_0$  in  $\alpha$  where  $t(\beta') > \delta$  for  $\beta' > \beta_0$ . Since  $h$  is a  $\Sigma_2$  cofinality map, there is a  $\rho < \mu$  such that  $h(\rho) > \beta_0$ .  $\square$

Since our construction is to be  $\alpha$ -effective, we require an  $\alpha$ -recursive approximation to  $h$ . Namely, let

$$h(\beta) = \delta \Leftrightarrow (\exists \sigma_1)(\sigma_2)R(\sigma_1, \sigma_2, \beta, \delta)$$

for  $\alpha$ -recursive  $R$ . Then the  $\sigma$ th approximation to  $h$  ( $\sigma < \alpha$ ) is defined as

$$h^\sigma(\beta) = \delta \Leftrightarrow (\exists \sigma_1)_{< \sigma} (\sigma_2)_{< \sigma} R(\sigma_1, \sigma_2, \beta, \delta) \\ \& (\forall \gamma)_{< \delta} (\forall \sigma_1)_{< \sigma} (\exists \sigma_2)_{< \sigma} \neg R(\sigma_1, \sigma_2, \beta, \gamma);$$

$h^\sigma$  is  $\alpha$ -recursive (by admissibility) and  $h$  is the limit (as  $\sigma \rightarrow \alpha$ ) of  $h^\sigma$ .

2.3 LEMMA. For all  $\rho < \mu$  there exists a  $\sigma_\rho < \alpha$  such that  $(\forall \sigma)[\sigma > \sigma_\rho \rightarrow h(\rho) = h^\sigma(\rho)]$ .  $\square$

For  $\sigma < \alpha$  and  $\rho < \mu$ ,  $B_\rho(\sigma)$  denotes the  $\sigma$ th approximation to block  $B_\rho$ . Namely, the initial segment of  $\alpha^*$  bounded above by  $t \circ h^\sigma(\rho)$ . An immediate consequence is the eventual stability of each  $B_\rho(\sigma)$ .

2.4 COROLLARY. For all  $\rho < \mu$  there is a  $\sigma_\rho < \alpha$  such that  $(\forall \sigma)[\sigma > \sigma_\rho \rightarrow B_\rho = B_\rho(\sigma)]$ .  $\square$

3. Construction. An  $\alpha$ -recursive function  $k$  will be defined in terms of a construction given below. Throughout the execution of the construction several sets will be accumulating.

The set  $K^\sigma$ , at stage  $\sigma$ , represents the function  $k$  being built. A pair  $\langle \beta, \theta \rangle$

is placed into  $K^\beta$  at some stage  $\beta$  if and only if  $k(\beta) = \theta$ . The set  $I^\sigma$ , at stage  $\sigma$ , is a collection of encodings of triples  $\langle \nu, \kappa, \gamma \rangle$ ,  $\nu, \kappa, \gamma < \alpha$ , such that we have made a guess  $f_\kappa(\beta) \geq \Phi_\nu(\beta)$  on all but an  $\alpha$ -finite set of  $\beta$ .

A triple  $\langle \nu, \kappa, \gamma \rangle$  is said to have *priority value*  $t(\gamma) < \alpha^*$ . If for some  $\rho < \mu$  ( $= \sigma 2cf(\alpha)$ ),  $t(\gamma) \in B_\rho$ , we call the triple a  $\rho$ -triple, and if  $t(\gamma) \in B_\rho(\sigma)$  we refer to it as a  $\rho$ -triple at stage  $\sigma$ . When the first component of triple  $\langle \nu, \kappa, \gamma \rangle$  is to be emphasized, we say it is a  $\Phi_\nu$ -triple.

The set  $TO^\sigma$ , at stage  $\sigma$ , consists of triples of guesses which have been rejected. The reason for this is to have sets which increase as  $\sigma \rightarrow \alpha$  instead of pulsating. The set  $TO^{<\sigma}$  is  $TO^\sigma$  just prior to stage  $\sigma$ , that is,  $TO^{<\sigma} = \bigcup_{\tau < \sigma} TO^\tau$ . When  $\sigma = \alpha$  we denote  $TO^{<\sigma}$  by  $TO$ . Similarly, for  $K^{<\sigma}$ ,  $I^{<\sigma}$ ,  $K$  and  $I$ . Finally, if a triple is in  $I^{<\sigma} - TO^{<\sigma}$ , we say that it is *active at stage*  $\sigma$  or simply *active* when the context is clear.

The construction which computes  $k(\eta)$  for  $\eta < \alpha$  is defined by transfinite recursion on stages  $\sigma < \alpha$ . The key ideas are expressed in the following.

At every stage  $\sigma < \alpha$ , attempts are made to eliminate bad guesses. If no bad ones are discovered then  $k(\sigma)$  takes on the value  $f_\sigma(\sigma)$ . Otherwise, for each  $\rho < \mu$  ( $= \sigma 2cf(\alpha)$ ) the incorrect guess  $\langle \nu_\rho, \kappa_\rho, \delta_\rho \rangle$  of highest priority in  $B_\rho(\sigma)$  is snuffed out. If  $f_{\kappa_\rho}(\sigma)$  exceeds the complexity of all correct guesses in blocks  $B_{\rho'}$ ,  $\rho' < \rho$ , then the triple is cancelled or "popped". Otherwise, the triple is "injured" in the hope that at another time, the  $\Phi_{\nu_\rho}$ -triple will be popped. If some triple  $\langle \nu_\rho, \kappa_\rho, \gamma_\rho \rangle$ ,  $\rho < \mu$ , does get popped, then  $k(\sigma)$  becomes no larger than  $f_{\kappa_\rho}(\sigma)$ ; otherwise,  $k(\sigma)$  is  $f_\sigma(\sigma)$ .

A more formal exposition is as follows:

*Stage 0.* Set  $I^0 = TO^0 = K^0 = \emptyset$ .

*Stage  $\sigma$ .* Compute the set

$$V = \{ \xi | \xi \in I^{<\sigma} - TO^{<\sigma} \ \& \ f_{\pi_2(\xi)}(\sigma) < \Phi_{\pi_1(\xi)}(\sigma) \}.$$

These are the guesses active at stage  $\sigma$  which we currently discover to be incorrect.

If  $V = \emptyset$  then set  $\theta = f_\sigma(\sigma)$ ;  $K^\sigma = K^{<\sigma} \cup \{ \langle \sigma, \theta \rangle \}$ ;  $I^\sigma = I^{<\sigma} \cup \{ \langle \sigma, \sigma, \sigma \rangle \}$ ;  $TO^\sigma = TO^{<\sigma}$  and go to stage  $\sigma + 1$ . So far we have guessed correctly. We therefore set  $k(\sigma) = f_\sigma(\sigma)$  and establish a new guess that  $f_\sigma(\beta) \geq \Phi_\sigma(\beta)$  on all but an  $\alpha$ -finite set giving it a priority value  $t(\sigma)$ .

Otherwise, for each  $\rho < \mu$  compute

$$\begin{aligned} \gamma'_\rho &= \min \{ t \circ \pi_3(\xi) | \xi \in V \ \& \ t \circ \pi_3(\xi) \in B_\rho(\sigma) \}; \\ \nu_\rho &= \min \{ \pi_1(\xi) | \xi \in V \ \& \ t \circ \pi_3(\xi) = \gamma'_\rho \} \quad \text{and} \\ \kappa_\rho &= \min \{ \kappa | \langle \nu_\rho, \kappa, t^{-1}(\gamma'_\rho) \rangle \in V \}. \end{aligned}$$

The triple  $\xi_\rho$  just chosen from  $V$  is the one in  $B_\rho(\sigma)$  with highest priority (i.e. lowest priority value) according to its third component. If more than one

member of  $V$  is such, we choose the triple appearing earlier in this list.

Set

$$J_\rho = \{ \langle \beta_1, \beta_2, \beta_3 \rangle \mid \langle \beta_1, \beta_2, \beta_3 \rangle \in I^{<\sigma} - TO^{<\sigma} \\ \&t(\beta_3) \in B_{\rho'}(\sigma) \text{ for } \rho' < \rho \& f_{\beta_2}(\sigma) \geq \Phi_{\beta_1}(\sigma) \}$$

and

$$m_\rho = \sup \{ \Phi_{\pi_1(\xi)}(\sigma) \mid \xi \in J_\rho \}.$$

The set  $J_\rho$  is comprised of all active triples which (1) have priority values in earlier blocks than  $B_\rho(\sigma)$ , and (2) for input  $\sigma$ ,  $f_{\beta_2}(\sigma) \geq \Phi_{\beta_1}(\sigma)$  (i.e. represents a correct guess).  $m_\rho$  exceeds the amount of work performed by these correct guesses.

If  $f_{\kappa_\rho}(\sigma) \geq m_\rho$  then set  $I^\sigma = I^{<\sigma} \cup \{ \langle \nu_\rho, \sigma, \sigma \rangle \}$  and  $TO^\sigma = TO^{<\sigma} \cup \{ \langle \nu_\rho, \kappa_\rho, \gamma_\rho \rangle \}$ . Here we form a new guess that  $f_\sigma(\beta) \geq \Phi_{\nu_\rho}(\beta)$  and append a priority value of  $t(\sigma)$ . In such a situation a  $\Phi_{\nu_\rho}$ -triple is said to be *popped at stage*  $\sigma$ .

If  $f_{\kappa_\rho}(\sigma) < m_\rho$  we set  $\tau_\rho = \sup \{ \pi_2(\xi) \mid \xi \in J_\rho \} \cup \{ \kappa_\rho \}$ ,  $TO^\sigma = TO^{<\sigma} \cup \{ \langle \nu_\rho, \kappa_\rho, \gamma_\rho \rangle \}$  and  $I^\sigma = I^{<\sigma} \cup \{ \langle \nu_\rho, \tau_\rho, \gamma_\rho \rangle \}$ . Here  $\tau_\rho$  is larger than any middle component of a triple in  $J_\rho$ . We eliminate our guess that  $f_{\kappa_\rho}(\beta) \geq \Phi_{\nu_\rho}(\beta)$  and replace it with the guess that  $f_{\tau_\rho}(\beta) \geq \Phi_{\nu_\rho}(\beta)$  on all but an  $\alpha$ -finite set. In such a case we say a  $\Phi_{\nu_\rho}$ -triple is *injured at stage*  $\sigma$ . (Observe that the injured  $\Phi_{\nu_\rho}$ -triple retains its priority but only changes its middle component.)

If for some  $\rho < \mu$ ,  $f_{\kappa_\rho}(\sigma) \geq m_\rho$ , we set  $\theta$  to the least such  $f_{\kappa_\rho}(\sigma)$ ; otherwise  $\theta = f_\sigma(\sigma)$ . In either case, set  $K^\sigma = K^{<\sigma} \cup \{ \langle \sigma, \theta \rangle \}$  and  $I^\sigma = I^{<\sigma} \cup \{ \langle \sigma, \sigma, \sigma \rangle \}$ . In the former the value of  $k(\sigma)$  is the smallest  $f_{\kappa_\rho}(\sigma)$  representing a popped  $\Phi_{\nu_\rho}$ -triple. Otherwise,  $k(\sigma) = f_\sigma(\sigma)$ . An important point is that at any stage  $\sigma$ , if a  $\Phi_{\nu_\rho}$ -triple is *popped*, then  $k(\sigma) < \Phi_{\nu_\rho}(\sigma)$ .

This concludes the construction.  $\square$

**4. Verification.** Clearly  $k$  is a well-defined  $\alpha$ -recursive function; for to compute  $k(\beta)$ ,  $\beta < \alpha$ , we simply run the  $\alpha$ -effective construction (which at any stage assigns exactly one value to  $k$ ), up until stage  $\beta$ . The remainder of the proof is the demonstration that  $C_k = \bigcup_{\varepsilon < \alpha} C_{f_\varepsilon}$ .

The central convergence result is

**4.1 LEMMA.** *For any  $\rho < \sigma 2cf(\alpha)$  there exists a stage  $\sigma_\rho < \alpha$  such that for all stages  $\sigma > \sigma_\rho$  and  $\rho' < \rho$  blocks  $B_{\rho'}$  are stable and no  $\rho'$ -triples are injured or popped.*

**PROOF.** By induction on  $\rho'$ . Assume that for all  $\rho' < \rho$  the lemma holds and consider the map  $m: \sigma 2cf(\alpha) \rightarrow \alpha$  defined as  $m(\delta) \equiv$  the least stage  $\sigma_\delta$  at which the block  $B_\delta$  stabilizes and no  $\delta$ -triple is either injured or popped at any stage  $\sigma > \sigma_\delta$ .

By the induction hypothesis  $m$  is total on  $\rho$ . Furthermore,  $m$  is  $\Sigma_2$  from

$$m(\delta) = \sigma_1 \Leftrightarrow (\sigma_2) [\sigma_2 > \sigma_1 \rightarrow B_\delta(\sigma_2) = B_\delta(\sigma_1) \text{ and at stage } \sigma_2 \\ \text{no } \delta\text{-triple is injured or popped}] \\ \text{and } \sigma_1 \text{ is the least such;}$$

and since  $B_\delta = t \circ h(\delta)$  is  $\Sigma_2$ ,  $B_\delta(\sigma)$  is  $\Sigma_1$ . From  $\rho < \sigma 2cf(\alpha)$  and the definition of  $\Sigma_2$  cofinality,  $m[\rho]$  is bounded below  $\alpha$ . Since  $t$  is  $\Sigma_1$ , by admissibility, we can find a bound on the set of all  $\sigma < \alpha$  whose image under  $t$  falls below  $t \circ h(\rho) + 1$ . By Corollary 2.4, there exists a stage after which  $B_\rho(\sigma)$  stabilizes to  $B_\rho$ . Pick  $\sigma_1$  to exceed the three aforementioned bounds.

Let  $\sigma > \sigma_1$  and  $\rho' < \rho$  to see that, if  $\langle \nu', \kappa', \gamma' \rangle$  is an active  $\rho'$ -triple at stage  $\sigma$ , then  $\Phi_{\nu'}(\sigma) \leq f_{\kappa'}(\sigma)$ . For otherwise, by the details of the construction, some  $\rho'$ -triple will either be popped or injured at stage  $\sigma$ , contradicting the role of  $\sigma_1$  as a bound on such stages.

For any  $\rho$ -triple,  $\langle \nu, \kappa, \gamma \rangle$  involved in any activity following stage  $\sigma_1$ , there are only two possible situations: One is that it is popped. In this case, the  $\Phi_{\nu}$ -triple has as its second and third components, the stage  $\sigma$  of the popping. However, since  $t(\sigma) > t \circ h(\rho)$  the  $\Phi_{\nu}$ -triple is popped out of block  $B_\rho$ . In the other case, it is injured and therefore just the middle component is altered. However, this new component is chosen so that it *exceeds all middle components of  $\rho'$ -triples ( $\rho' < \rho$ ) currently active* (and hence forever). Consequently, the next time this  $\Phi_{\nu}$ -triple sees action, it will have to be popped, and as in the previous case, it will be popped out of block  $B_\rho$ .

These two cases tell us that after stage  $\sigma_1$  any  $\Phi_{\nu}$ -triple which is a  $\rho$ -triple will at most be popped from  $B_\rho$ , injured once, or injured once and then popped from  $B_\rho$ . Since *at most* two active  $\rho$ -triples can have the same priority value (i.e. third components equal), we argue that  $\rho$ -triples can only contribute an  $\alpha$ -finite amount.

Specifically, for  $i = 1, 2$  define  $P_\rho^i = \{\delta \mid \delta \in B_\rho \text{ and } i \text{ } \rho\text{-triple(s) with priority value } \delta \text{ is (are) popped after } \sigma_1\}$ . Let  $IN_\rho^i = \{\delta \mid \delta \in B_\rho \text{ and } i \text{ } \rho\text{-triple(s) with priority value } \delta \text{ is (are) injured after } \sigma_1\}$ . Since these sets are  $\alpha$ -r.e. and bounded below  $\alpha^*$  ( $t \circ h(\rho) < \alpha^*$ ) by Fact 1.2 they are  $\alpha$ -finite. For  $i = 1, 2$ , define  $p_\rho^i(\delta)$  [ $in_\rho^i(\delta)$ ] = the stage  $\sigma$  at which a  $\rho$ -triple with priority value  $\delta$  is popped [injured] for the  $i$ th time. By the  $\alpha$ -effectiveness of the construction,  $in_\rho^i$  and  $p_\rho^i$ ,  $i = 1, 2$ , are  $\alpha$ -partial recursive. For  $i = 1, 2$ , since  $P_\rho^i \subseteq \text{dom}(p_\rho^i)$  and  $IN_\rho^i \subseteq \text{dom}(in_\rho^i)$  by Fact 1.1 both  $p_\rho^i$  [ $P_\rho^i$ ] and  $in_\rho^i$  [ $IN_\rho^i$ ],  $i = 1, 2$ , are  $\alpha$ -finite. Hence they are all bounded. Thus letting  $\sigma_\rho$  be the maximum of their bounds, it follows by the above definitions that no  $\rho'$ -triple,  $\rho' < \rho$ , is either injured or popped past stage  $\sigma_\rho$ . Thus the induction step is complete.  $\square$

A consequence of the above proof is

**4.2 COROLLARY.** *For any  $\rho < \sigma 2cf(\alpha)$  and any  $\delta$  in  $B_\rho$ , there exists a stage*

$\sigma_0 < \alpha$  such that for all stages  $\sigma > \sigma_0$ ,

- (i) all blocks  $B_{\rho'}$ ,  $\rho' \leq \rho$ , have stabilized.
- (ii) no  $\rho'$ -triples are injured or popped for  $\rho' < \rho$ , and
- (iii) no  $\rho$ -triples having priority value less than  $\delta$  are injured or popped.

PROOF. Let  $\sigma_1$  be as in the induction step of the proof of Lemma 4.1. The only modifications needed are in the definitions of  $P_{\rho}^i$ ,  $IN_{\rho}^i$ ,  $p_{\rho}^i$  and  $in_{\rho}^i$ , where we only need concern ourselves with those  $\rho$ -triples with priority values less than  $\delta$ . The remainder of the argument carries over.  $\square$

In the next lemma we argue that if  $k$  dominates the complexity of an  $\alpha$ -recursive function, then there must exist some  $f_{\kappa_0}$  also dominating it.

4.3 LEMMA.  $C_k \subseteq \bigcup_{\epsilon < \alpha} C_{f_{\epsilon}}$ .

PROOF. Suppose  $\phi_{\nu} \in C_k$  to show the existence of an  $f_{\kappa_0}$  where  $\Phi_{\nu}(\beta) < f_{\kappa_0}(\beta)$  on all but an  $\alpha$ -finite set of  $\beta$ . This will imply  $\phi_{\nu} \in C_{\kappa_0}$  and hence  $\phi_{\nu} \in \bigcup_{\epsilon < \alpha} C_{f_{\epsilon}}$ . Assume to the contrary that this is not the case. That is, for each  $\kappa < \alpha$  the sets  $D_{\nu}^{\kappa} = \{\beta \mid f_{\kappa}(\beta) < \Phi_{\nu}(\beta)\}$  are not  $\alpha$ -finite. The  $\alpha$ -recursiveness of the  $f_{\kappa}$  implies that these must be unbounded.

We prove that the set  $A_{\nu} = \{\sigma \mid \text{at stage } \sigma \text{ a } \Phi_{\nu}\text{-triple is popped}\}$  is unbounded. By the details of the construction this would imply  $\{\sigma \mid k(\sigma) < \Phi_{\nu}(\sigma)\}$  is also unbounded. Since we assumed  $\phi_{\nu} \in C_k$  this leads to a contradiction completing the proof of the lemma.

Given a stage  $\sigma_1 < \alpha$ , we show the existence of a stage  $\sigma_2 > \sigma_1$  such that at stage  $\sigma_2$  a  $\Phi_{\nu}$ -triple is popped, verifying the unboundedness of  $A_{\nu}$ . Without loss of generality, we can assume that at the conclusion of stage  $\sigma_1$  a  $\Phi_{\nu}$ -triple  $\langle \nu, \kappa, \gamma \rangle$  exists. Let  $\rho < \sigma_2 \text{cf}(\alpha)$  be such that  $t(\gamma) \in B_{\rho}$ . Let  $\sigma_0$  be the stage obtained from Corollary 4.2 and assume  $\sigma_0$  is at least as large as  $\sigma_1$ . Since there can be at most a second triple having the same priority value  $t(\gamma)$ , we can assume that  $\sigma_0$  is large enough so that if this triple is popped, it has done so by stage  $\sigma_0$ . If in between  $\sigma_1$  and  $\sigma_0$  a  $\Phi_{\nu}$ -triple is popped, we are done. Otherwise, assume  $\langle \nu, \kappa_1, \gamma \rangle \in I^{\sigma_0} - TO^{\sigma_0}$  for some  $\kappa_1 < \alpha$ .

Observe that at stages  $\sigma > \sigma_0$  for any  $\rho'$ -triple  $\langle \nu', \kappa', \gamma' \rangle$ ,  $\rho' < \rho$  active at stage  $\sigma$ , that  $\Phi_{\nu'}(\sigma) < f_{\kappa'}(\sigma)$ . For otherwise, a  $\rho'$ -triple would be either popped or injured contradicting the choice of  $\sigma_0$ . Similarly, for any  $\rho$ -triple with priority value less than  $t(\gamma)$ . By our original assumption, the set  $D_{\nu}^{\kappa_1} = \{\beta \mid f_{\kappa_1}(\beta) < \Phi_{\nu}(\beta)\}$  is unbounded. Thus there must be a smallest  $\eta \in D_{\nu}^{\kappa_1}$  such that  $\eta > \sigma_0$ . By the details of the construction, and the above remarks, at stage  $\eta$ , the  $\Phi_{\nu}$ -triple  $\langle \nu, \kappa_1, \gamma \rangle$  will be the  $\rho$ -triple with lowest priority value active at stage  $\sigma$  such that  $f_{\kappa_1}(\eta) < \Phi_{\nu}(\eta)$ . As in the proof of Lemma 4.1, there are two possibilities.

First, if the value  $f_{\kappa_1}(\eta)$  is greater than or equal to all values  $\Phi_{\nu}(\eta)$  where at

stage  $\eta$  the  $\Phi_{\nu}$ -triple is an active  $\rho'$ -triple  $\rho' < \rho$ . In this case the  $\Phi_{\nu}$ -triple would certainly be popped.

Second, if the value of  $f_{\kappa_1}(\eta)$  is less than some  $\Phi_{\nu}(\eta)$  where the  $\Phi_{\nu}$ -triple is an active  $\rho'$ -triple,  $\rho' < \rho$ . By the details of the construction the triple  $\langle \nu, \kappa_1, \gamma \rangle$  is ejected (i.e., put into  $TO^\eta$ ) and replaced by the  $\Phi_{\nu}$ -triple  $\langle \nu, \kappa_2, \gamma \rangle$  where  $\kappa_2 > \sup\{\kappa' \mid \kappa' \text{ is the second component of an active } \rho'\text{-triple } \rho' < \rho \text{ at stage } \eta\}$ . Since  $D_{\nu}^{\kappa_2} = \{\beta \mid f_{\kappa_2}(\beta) < \Phi_{\nu}(\beta)\}$  is also unbounded, there must be a least  $\beta > \eta$  such that  $\beta \in D_{\nu}^{\kappa_2}$ . Therefore, at stage  $\beta$ , the  $\Phi_{\nu}$ -triple  $\langle \nu, \kappa_2, \gamma \rangle$  would surely be popped.

Since both possibilities ultimately lead to popping, we are done.  $\square$

Our next result shows that if the  $\alpha$ -complexity of an  $\alpha$ -recursive function is dominated by at least one  $f_{\varepsilon_0}$  (and consequently all  $f_{\varepsilon}$ ,  $\varepsilon > \varepsilon_0$ ), it must also be dominated by  $k$ .

4.4 LEMMA.  $\bigcup_{\varepsilon < \alpha} C_f \subseteq C_k$ .

PROOF. Let  $\kappa$  be an index such that  $\Phi_{\nu}(\beta) < f_{\kappa}(\beta)$  on all but an  $\alpha$ -finite set of  $\beta$ . We show that  $\{\beta \mid \Phi_{\nu}(\beta) > k(\beta)\}$  is an  $\alpha$ -finite set. Without loss of generality, we will assume that  $\kappa$  is the least such index.

*Claim.* There exist a stage  $\sigma_1$  and ordinals  $\kappa_1, \gamma < \alpha$  such that (1)  $\langle \nu, \kappa_1, \gamma \rangle \in I^{\sigma_1} - TO^{\sigma_1}$  and (2) for all stages  $\sigma > \sigma_1$ ,  $\langle \nu, \kappa_1, \gamma \rangle \in I^{\sigma} - TO^{\sigma}$  (i.e., after stage  $\sigma_1$  the  $\Phi_{\nu}$ -triple  $\langle \nu, \kappa_1, \gamma \rangle$  remains active).

To prove the claim we first see that the set of stages at which a  $\Phi_{\nu}$ -triple is popped is bounded. By the hypothesis, the set  $\{\beta \mid \Phi_{\nu}(\beta) > f_{\kappa}(\beta)\}$  is  $\alpha$ -finite, hence bounded by some  $\beta' < \alpha$ . By this and the increasingness of the  $\{f_{\varepsilon}\}$ , for all  $\lambda > \kappa$ , the sets  $\{\beta \mid \Phi_{\nu}(\beta) > f_{\lambda}(\beta)\}$  are bounded above by  $\beta'$ . Suppose an unbounded sequence of stages  $\sigma_1 < \sigma_2 < \sigma_3 < \dots$  exists so that at each stage  $\sigma_i$  a  $\Phi_{\nu}$ -triple is popped. Consequently, there will be a sequence of  $\Phi_{\nu}$ -triples for which the second components take on values  $\sigma_i$ . However, once  $\sigma_i > \kappa$  and  $\sigma > \beta'$ ,  $\Phi_{\nu}(\sigma) < f_{\sigma_i}(\sigma)$ , the popping ceases.

Suppose after stage  $\sigma'$  the third component of the  $\Phi_{\nu}$ -triple remains fixed at value  $\gamma$ . Let  $t(\gamma) \in B_{\rho}$ ,  $\rho < \sigma 2cf(\alpha)$ , and let  $\sigma_{\rho}$  be the stage obtained by Lemma 4.1. Since after stage  $\sigma'$ , the  $\Phi_{\nu}$ -triple is always a  $\rho$ -triple it follows that it can no longer be injured following stage  $\sigma_{\rho}$ . Hence, the claim is proven.

Observe that  $\kappa_1$  of the claim is  $\geq \kappa$ . For otherwise, the set  $\{\beta \mid \Phi_{\nu}(\beta) > f_{\kappa_1}(\beta)\}$  would be unbounded by the minimality of  $\kappa$ . As in the proof of Lemma 4.3, there would then be a stage  $\beta > \sigma_1$  at which the  $\Phi_{\nu}$ -triple  $\langle \nu, \kappa_1, \gamma \rangle$  is popped or injured contradicting the claim.

Let  $t(\gamma) \in B_{\rho}$  and let  $\sigma_{\rho}$  be the stage obtained from Lemma 4.1 which bounds injuries and pops resulting from  $\rho'$ -triples  $\rho' < \rho$ . Since  $\{\beta \mid \Phi_{\nu}(\beta) > f_{\kappa}(\beta)\}$  is  $\alpha$ -finite, let  $\sigma_2$  be a bound. Next set  $\sigma' = \max\{\sigma_{\rho}, \sigma_1, \sigma_2\}$  to see that for all  $\sigma > \sigma'$ ,  $k(\sigma) > \Phi_{\nu}(\sigma)$ . By the details of the construction a value  $k(\sigma)$  is

assigned in one of two ways. First, if  $k(\sigma)$  is  $f_\sigma(\sigma)$ . Then since  $\sigma \succ \sigma' \succ \sigma_1 \succ \kappa_1 \succ \kappa$ ,

$$k(\sigma) = f_\sigma(\sigma) \succ f_{\kappa_1}(\sigma) \succ f_\kappa(\sigma) \succ \Phi_\nu(\sigma) \quad (\text{since } \sigma \succ \sigma_2).$$

Second, if  $k(\sigma) = f_{\kappa'}(\sigma)$  for some  $\kappa' < \alpha$ . At stages  $\sigma \succ \sigma'$ , the  $\Phi_\nu$ -triple  $\langle \nu, \kappa_1, \gamma \rangle$ , as well as all  $\rho'$  triples,  $\rho' < \rho$ , have already stabilized. Thus, for all  $\sigma \succ \sigma'$ , if  $\langle \nu', \kappa', \gamma' \rangle$  is one of these triples,  $f_{\kappa'}(\sigma) \succ \Phi_\nu(\sigma)$ . By the construction, if a value is assigned to  $k(\sigma)$  by popping a  $\tilde{\rho}$ -triple  $\langle \tilde{\nu}, \tilde{\kappa}, \tilde{\gamma} \rangle$ , then it must be that  $\tilde{\rho} > \rho$ . Thus again by the construction,  $\Phi_\nu(\sigma) < f_{\kappa'}(\sigma) = k(\sigma)$ .

Therefore, in both situations  $k(\sigma) \succ \Phi_\nu(\sigma)$  for  $\sigma \succ \sigma'$ . Hence, the  $\alpha$ -recursive set  $\{\beta \mid \Phi_\nu(\beta) > k(\beta)\}$  is bounded and thus  $\alpha$ -finite.  $\square$

Lemmas 4.3 and 4.4 combine to yield  $C_k = \bigcup_{\epsilon < \alpha} C_{f_\epsilon}$ .

**5. Complexity of generalized primitive recursive functions.** In this section we examine two infinite analogues to the ordinary primitive recursive functions. Although these generalizations coincide on  $\omega$ , we show them to diverge on all admissible  $\alpha > \omega$ . This investigation is complexity oriented and centers around a well-known application of the Union Theorem.

Our first generalization is that of Jensen and Karp [9] with present formulation due to Gandy.

**DEFINITION.** A function  $f: ON^n \rightarrow ON$  is *ordinally primitive recursive* ( $\text{Prim}_0$ ) if it can be obtained from the initial functions

- (1)  $U_i^n(\bar{\beta}) = \beta_i, \bar{\beta} = (\beta_1, \dots, \beta_n), 1 \leq n < \omega, 1 \leq i \leq n;$
- (2)  $N(\beta) = 0;$
- (3)  $S(\beta) = \beta + 1 = \beta \cup \{\beta\};$
- (4)  $C(\beta, \gamma, \mu, \eta) = \beta$  if  $\mu \in \eta, \gamma$  otherwise; under the operations of
- (5) substitution:

$$(5a) \quad F(\bar{\beta}, \bar{\gamma}) = G(\bar{\beta}, H(\bar{\beta}), \bar{\gamma}), \quad \bar{\beta} = (\beta_1, \dots, \beta_n), \quad \bar{\gamma} = (\gamma_1, \dots, \gamma_n),$$

$m, n < \omega;$

$$(5b) \quad F(\bar{\beta}, \bar{\gamma}) = G(H(\bar{\beta}), \bar{\gamma});$$

(6) recursion:

$$F(\beta, \bar{\gamma}) = G\left(\bigcup \{F(\mu, \bar{\gamma}) \mid \mu \in \beta\}, \beta, \bar{\gamma}\right), \quad n < \omega, \quad \bar{\gamma} = (\gamma_1, \dots, \gamma_n).$$

A relation on ordinals is *ordinally primitive recursive* just in case its characteristic function is  $\text{Prim}_0$ .

It is clear from the above definition that there are at most  $\aleph_0$   $\text{Prim}_0$  functions. Nevertheless, this small class has enough power to provide analogues to Kleene's  $T$  predicates for formalisms of  $\alpha$ -recursion theory.

One such development may be obtained through the Kripke equation

calculus (EC) (cf. [14]). Specifically,  $\phi_\varepsilon$  is the  $\alpha$ -partial recursive function computed in the EC from the finite system of equations  $E$  with Gödel number  $\varepsilon$  ( $= \text{GN}(E)$ )  $< \alpha$ . The ordinal  $\varepsilon < \alpha$  is called an  $\alpha$ -index of  $\phi_\varepsilon$ . If the equations  $E$  contain no constants then  $\varepsilon < \omega$ .

Let  $S_{\sigma,\gamma}^E$  be the  $\alpha$ -finite collection of equations derived from set  $E$  (according to rules of EC) by stage  $\sigma < \alpha$  substituting no constants greater than  $\gamma < \sigma$ . Let  $\pi_i$ ,  $i \leq n$ , and  $\langle \cdot, \cdot \rangle$  be the usual  $\text{Prim}_0$  projection and pairing functions. Following the usual development (cf. Jensen and Karp [9], Kripke [14], Tugué [28]) one shows existence of a  $\text{Prim}_0$   $T$ :

$T(\varepsilon, \beta, \gamma) \leftrightarrow$  the equation with Gödel number  $\pi_3(\gamma)$  giving value of

$\alpha$ -partial recursive  $f$  for  $\beta$  lies in  $S_{\pi_1(\gamma), \pi_2(\gamma)}^E$  where  $\text{GN}(E) = \varepsilon$ ;

and a  $\text{Prim}_0$  function,  $U(\langle \beta_1, \beta_2, \text{GN}("f(\beta) = \delta") \rangle) = \delta$ , both of which combine with an  $\alpha$ -min operator to yield a Normal Form Theorem.

There is an inherent uniformity about the  $\text{Prim}_0$  functions. Namely, if  $f: ON \rightarrow ON$  is ordinally primitive recursive, then there is an  $\varepsilon < \omega$  such that  $f|_\alpha$  is  $\alpha$ -recursive with index  $\varepsilon$  for all admissibles  $\alpha$ . As a consequence,  $f$  will always map any admissible  $\alpha$  (since  $\alpha$  is  $\text{Prim}_0$  closed [9]) into itself.

As powerful as they appear, the class  $\text{Prim}_0$ , when regarded as maps from  $\alpha$  to  $\alpha$ , lack one property of the ordinary primitive recursives. Namely, they are void of the constant functions  $\lambda\beta.\gamma$ , for  $\omega \leq \gamma < \alpha$ . In the  $\omega$ -case constant  $\lambda x.n$  is derived via  $n$  compositions to successor from the null function. For  $\alpha > \omega$ , the finiteness of the definitions of the  $\text{Prim}_0$  functions precludes the derivations of such functions.

**DEFINITION.** A function  $f: \alpha \rightarrow \alpha$  is  $\alpha$ -primitive recursive ( $\alpha$ -Prim) if it can be obtained from the initial functions  $U_i^n, N, S, C$  and the constant functions  $\{\lambda\beta.\gamma | \gamma < \alpha\}$  from the operations of composition and recursion. A relation on  $\alpha^n$  ( $n < \omega$ ) is  $\alpha$ -primitive recursive just in case its characteristic function is  $\alpha$ -Prim.

We define an  $\alpha$ -complexity measure  $\Phi$  based upon the Kripke formalism. Specifically,  $\phi_\varepsilon$  is the  $\alpha$ -partial recursive function computed within the EC from equations having Gödel number  $\varepsilon$ ; the corresponding step counter is

$$\Phi_\varepsilon(\beta) = \min_\gamma T(\varepsilon, \beta, \gamma), \quad \beta, \varepsilon < \alpha.$$

It is easily seen that  $\Phi = \langle \phi_\varepsilon, \Phi_\varepsilon \rangle$  constitutes an  $\alpha$ -complexity measure.

In the following  $\alpha$  is some  $\Sigma_1$  admissible ordinal,  $\Phi$  the Kripke complexity measure defined above and  $\text{Prim}$  some generalization to  $\alpha$  of the primitive recursive functions.

**PROPOSITION A.** Let  $\phi$  be a unary  $\text{Prim}$  function. Then there is a unary  $\text{Prim} g$  such that  $\phi \in C_g^\Phi$ .

PROPOSITION B. For  $g$  a unary Prim function,  $f \in C_g^\Phi$  implies  $f \in \text{Prim}$ .

PROPOSITION C. There exists an  $\alpha$ -recursively enumerable strictly increasing sequence of Prim functions  $\{f_\epsilon | \epsilon < \alpha\}$  such that every Prim function is majorized by some  $f_\epsilon$ .

Let  $\text{Prim} \models \mathfrak{S}$  ( $\text{Prim} \not\models \mathfrak{S}$ ) denote that Prim is (is not) a model of statement  $\mathfrak{S}$ . The key fact is

5.1 THEOREM. Let Prim be a class of functions defined on  $\alpha$  such that  $\text{Prim} \models$  Proposition A & Proposition B & Proposition C. Then

- (i)  $\text{Prim} = \bigcup_{\epsilon < \alpha} C_{f_\epsilon}^\Phi$ , where  $\{f_\epsilon\}$  is the sequence of Proposition C, and
- (ii)  $\text{Prim} = C_t^\Phi$ , for some  $\alpha$ -recursive  $t$ .

PROOF. (i) Let  $\phi \in \text{Prim}$  to see  $\text{Prim} \subseteq \bigcup_{\epsilon < \alpha} C_{f_\epsilon}^\Phi$ . Since  $\text{Prim} \models$  Proposition A, there exists  $g \in \text{Prim}$  such that  $\phi \in C_g^\Phi$ . By  $\text{Prim} \models$  Proposition C,  $g < f_\epsilon$  for some  $\epsilon < \alpha$ ; thus,  $\phi \in C_{f_\epsilon}^\Phi$  and  $\phi \in \bigcup_{\epsilon < \alpha} C_{f_\epsilon}^\Phi$ . For the opposite inclusion, since  $\text{Prim} \models$  Proposition C,  $f_\epsilon \in \text{Prim}$  for all  $\epsilon < \alpha$ . Thus by  $\text{Prim} \models$  Proposition B,  $C_{f_\epsilon}^\Phi$  contains only Prim functions; hence,  $\bigcup_{\epsilon < \alpha} C_{f_\epsilon}^\Phi \subseteq \text{Prim}$ .

(ii) Let  $\{f_\epsilon\}$  be the  $\alpha$ -recursively enumerable strictly increasing sequence of Proposition C. Then, by the  $\alpha$ -Union Theorem there exists an  $\alpha$ -recursive  $t$  such that  $C_t^\Phi = \bigcup_{\epsilon < \alpha} C_{f_\epsilon}^\Phi \models \text{Prim}$ .  $\square$

We next examine which of the various instances of Prim are models of the three propositions.

- 5.2 LEMMA. (i)  $\alpha\text{-Prim} \models$  Proposition A for all admissible  $\alpha$ , and
- (ii)  $\text{Prim}_0 \models$  Proposition A.

PROOF. The demonstration that  $\alpha\text{-Prim} \models$  Proposition A is by induction.

(1) For  $U(x) = x$  let  $\epsilon = \text{GN}("U(x) = x")$  and  $g_U(\beta) = \langle 1, \beta + 1, \text{GN}("U(\beta) = \beta") \rangle$ . By results of Jensen and Karp  $g_U$  is  $\text{Prim}_0$ . Further, in EC,  $U = \phi_\epsilon$  and  $\Phi_\epsilon = g_U$ ; hence,  $U \in C_g^\Phi$ .

(2) For  $N(x) = 0$ , let  $\epsilon = \text{GN}("N(x) = 0")$  and  $g_N(\beta) = \langle 1, \beta + 1, \text{GN}("N(\beta) = 0") \rangle$ .

(3) For  $\lambda x.\gamma$ ,  $\gamma < \alpha$ , let  $\epsilon = \text{GN}("f(x) = \gamma")$  and  $g_f(\beta) = \langle 1, \beta + 1, \text{GN}("f(\beta) = \gamma") \rangle$ .

(4) The case  $S(x) = x + 1$  is a bit more complicated and we omit details here. Essentially, one defines successor in EC (cf. Kripke [14]) via a set of 13 equations and then shows  $g_S(\beta) = \langle m(\beta), \beta + 2, \text{GN}("S(\beta) = \beta + 1") \rangle$ , where  $m$  is  $\text{Prim}_0$ , is the accompanying  $\alpha$ -Prim bound.

Implicit in (5) and (6) is a property of pairing functions that for all  $\beta < \alpha$ ,  $\beta > \pi_1(\beta), \pi_2(\beta)$ .

(5) Assume  $k = \phi_{\varepsilon_k}$ ,  $h = \phi_{\varepsilon_h}$ ,  $\varepsilon_k = \text{GN}(E_k)$ ,  $\varepsilon_h = \text{GN}(E_h)$ ,  $k \in C_{g_k}^\Phi$  and  $h \in C_{g_h}^\Phi$  where  $k, h, g_k$  and  $g_h$  are  $\alpha$ -Prim. Let

$$\varepsilon_{k \circ h} = \text{GN}(E_k \cup E_h \cup \{“f(x) = k(h(x))”\})$$

(assuming no name-conflicts). Let

$$g_f(\beta) = \langle g_k(h(\beta)) + g_h(\beta), \max\{g_k(h(\beta)), g_h(\beta)\}, \text{GN}(“f(\beta) = k \circ h(\beta)”)\rangle.$$

(6) Suppose  $f$  is defined by recursion equations,

$$(*) \quad “f(0) = \gamma” \quad (\gamma < \alpha), \quad “f(\beta) = h(\cup\{f(\delta)|\delta \in \beta\})”$$

where  $\varepsilon_h = \text{GN}(E_h)$  and  $h \in C_{g_h}^\Phi$ . Let  $\varepsilon_f = \text{GN}((*) \cup E_h)$  and  $g_f(\beta)$  be defined by

$$\begin{aligned} g_f(0) &= \langle 1, \gamma, \text{GN}(“f(0) = \gamma”) \rangle, \\ g_f(\beta) &= \langle \max\{g_h(\cup\{f(\delta)|\delta \in \beta\}), \sum\{g_f(\delta)|\delta \in \beta\}\}, \\ &\quad \max\{\cup\{g_f(\delta)|\delta \in \beta\}, g_h(\cup\{f(\delta)|\delta \in \beta\})\}, \\ &\quad \text{GN}(“f(\beta) = h(\cup\{f(\delta)|\delta \in \beta\})”) \rangle. \end{aligned}$$

Since all functions mentioned (with the exception of those in (3)) involve no infinite constants, the above carries over for  $\text{Prim}_0$ .  $\square$

We next see that complexity classes bounded by  $\alpha$ -Prim functions contain only  $\alpha$ -Prim functions. Although this implies Proposition B holds for the ordinary primitive recursive functions, we find that this is not so for  $\text{Prim}_0$ .

5.3 LEMMA. (i)  $\alpha$ -Prim  $\models$  Proposition B.

(ii)  $\text{Prim}_0 \not\models$  Proposition B.

PROOF. (i) Since Kripke's  $T(\varepsilon, \beta, \gamma)$  predicate above is  $\alpha$ -Prim, it is  $\alpha$ -Prim for any  $\alpha$ . Let  $\phi_\varepsilon \in C_g^\Phi$  for some  $g \in \alpha$ -Prim. Since  $\Phi_\varepsilon^\alpha < g$  on all but an  $\alpha$ -finite subset of  $\alpha$ , let  $\delta_0$  be a bound. Then clearly,

$$\phi_\varepsilon(x) \equiv U\left(\min_{\gamma < \max\{g(x), \delta_0\}} T(\varepsilon, \beta, \gamma)\right).$$

It follows (from results of Jensen and Karp) that  $\phi_\varepsilon$  is  $\alpha$ -Prim.

(ii) We prove the existence of a  $g \in \text{Prim}_0$  and  $f \in C_g^\Phi$  where  $f \notin \text{Prim}_0$ . Define

$$f(x) = \begin{cases} x, & x \neq 3, \\ \omega + 17, & x = 3, \end{cases}$$

and let  $g(x)$  be the  $\text{Prim}_0$  function  $\langle 1, x, \text{GN}(“f(x) = x”) \rangle$ . Since  $\text{Prim}_0$

functions map  $\omega$  into  $\omega$ ,  $f$  cannot be  $\text{Prim}_0$ ; however, it is the case that  $f \in C_g^\Phi$ .  $\square$

One of the more well-known properties of the  $\omega$ -Prim functions was first discovered by Grzegorzczuk [5]. Namely, the existence of an  $\omega$ -hierarchy,  $\bigcup_{n < \omega} \mathcal{E}^n$ , for  $\omega$ -Prim. The Grzegorzczuk hierarchy is based on the existence of a strictly increasing r.e. sequence of  $\omega$ -Prim functions  $\{f_e(x) | e < \omega\}$  which majorize the  $\omega$ -Prim functions. The class  $\mathcal{E}^n$ ,  $n < \omega$ , is defined as containing successor, zero, projections,  $f_n$  and closed under operations of composition and a limited or bounded recursion. As a consequence,

5.4 COROLLARY.  $\omega$ -Prim  $\models$  Proposition C.

Since  $\omega$ -Prim satisfies all three propositions, by Theorem 5.1,

5.5 COROLLARY.  $\omega$ -Prim  $= \bigcup_{e < \omega} C_{f_e}^\Phi = C_t^\Phi$  for some  $\omega$ -recursive  $t$ .

Our next result implies that for  $\alpha > \omega$  an  $\alpha$ -hierarchy constructed along the lines of Grzegorzczuk is impossible for  $\text{Prim}_0 | \alpha$ .

5.6 LEMMA. Let  $\alpha$  be any admissible  $> \omega$ . Then there cannot exist a sequence  $\{f_e | e < \alpha\}$  of strictly increasing  $\text{Prim}_0 | \alpha$  functions.

PROOF. Suppose  $\{f_e | e < \alpha\}$  is such a sequence. Let  $n_0 \in \omega$  and define  $m: \alpha \rightarrow \omega$  by  $m(\beta) = f_\beta(n_0)$ . Since each  $f_\beta \in \text{Prim}_0$ ,  $f_\beta(n_0) \in \omega$ . By the increasingness of  $\{f_e\}$ ,  $m$  is a strictly increasing (hence, one-one) map of  $\alpha$  into  $\omega$ . Since  $\omega < \alpha$ ,  $m[\omega] < m_0 < \omega$ , implying the cardinality of  $\omega$  is finite.  $\square$

We have an immediate

5.7 COROLLARY.  $\text{Prim}_0 \not\models$  Proposition C.  $\square$

Since  $\text{Prim}_0$  is a model of Proposition A, but not of B nor C, Theorem 5.1 cannot be used to obtain an  $\alpha$ -hierarchy,  $\bigcup_{e < \alpha} C_{f_e}^\Phi = \text{Prim}_0 = C_t^\Phi$  for  $\alpha$ -recursive  $t$  and Kripke measure. We, therefore, leave as open the question of the existence of such. Namely, for all admissible  $\alpha$  and arbitrary  $\alpha$ -complexity measure  $\Phi$  (not necessarily that of Kripke) do there exist (1) complexity hierarchies for  $\text{Prim}_0$  based upon  $\Phi$  and (2) an  $\alpha$ -recursive  $t$  such that  $\text{Prim}_0$  is the  $\Phi$  complexity class of  $t$ ? If negative answers arise, we would naturally be interested in necessary and sufficient characterizations of  $\alpha$  where (1) and (2) hold.

Another open problem is the question of whether or not  $\alpha$ -Prim is a model for Proposition C. Namely, does there exist an  $\alpha$ -r.e. strictly increasing sequence of  $\alpha$ -Prim functions having type  $\alpha$  such that each  $\alpha$ -Prim function is majorized? If the answer is yes, Theorem 5.1 tells us we have an  $\alpha$ -hierarchy  $\bigcup_{e < \alpha} C_{f_e}^\Phi = \alpha\text{-Prim} = C_t$  for  $\alpha$ -recursive  $t$ .

An obvious approach to the above would be the construction of  $\alpha$ -analogues to Grzegorzczuk's bounding functions. However, difficulties arise in demonstrating that each  $f_\varepsilon$ ,  $\varepsilon < \alpha$ , is  $\alpha$ -Prim. In particular, when  $\varepsilon = \lambda$  is a limit, any generalized  $f_\lambda$ , in some way, incorporates a clause

$$f_\lambda(0, y) = \bigcup_{\delta < \lambda} f_\delta(y + 1, y + 1).$$

The inherent difficulty lies in the fact that an infinite union is being taken where the universal  $m(\varepsilon, x, y) = f_\varepsilon(x, y)$  is not  $\alpha$ -Prim (in  $\varepsilon, x$  and  $y$ ).

A similar problem arises if one attempts to build the sequence of Proposition C from an arbitrary  $\alpha$ -r.e. sequence  $\{p_\tau \mid \tau < \alpha\}$  for  $\alpha$ -Prim functions. For instance, the maximizing sequence  $\{m_\varepsilon(x)\} = \{\sup_{\tau < \varepsilon} p_\tau(x)\}$  is clearly  $\alpha$ -r.e., strictly increasing, and  $\alpha$ -Prim majorizing. However, the problem occurs in showing  $m_\lambda(x)$   $\alpha$ -Prim for  $\lambda$  a limit. In the  $\omega$ -case each  $m_\varepsilon$  is  $\omega$ -Prim since it has a finite definition obtainable by incorporating  $\omega$ -Prim definitions of  $m_t$ ,  $t < \varepsilon$ . Upon passing to the infinite this argument is no longer valid. Consequently, one is again dependent upon a non- $\alpha$ -Prim universal function for the  $p_\tau$ .

#### REFERENCES

1. M. Blum, *A machine-independent theory of the complexity of recursive functions*, J. Assoc. Comput. Mach. **14** (1967), 322–336. MR **38** #4213.
2. R. L. Constable, *The operator gap*, J. Assoc. Comput. Mach. **19** (1972), 175–183. MR **47** #3159.
3. M. Davis, *Computability and unsolvability*, McGraw-Hill, New York, 1958. MR **23** #A1525.
4. K. Gödel, *The consistency of the continuum hypothesis*, Ann. of Math. Studies, no. 3, Princeton Univ. Press, Princeton, N. J., 1940. MR **2**, 66.
5. A. Grzegorzczuk, *Some classes of recursive functions*, Rozprawy Math. **4** (1953), 1–45. MR **15**, 667.
6. Leo Harrington, *Some contributions to the theory of recursion on higher types*, Ph.D. Thesis, Massachusetts Institute of Technology, 1972.
7. B. E. Jacobs,  *$\alpha$ -computational complexity*, Ph.D. Thesis; Tech. Report IMM-408, Courant Inst. Math. Sci., New York Univ., New York, 1975.
8. \_\_\_\_\_, *On generalized computational complexity*, J. Symbolic Logic (to appear).
9. R. B. Jensen and C. Karp, *Primitive recursive set functions*, Axiomatic Set Theory (Proc. Sympos. Pure Math., Vol. 13, Part I), Amer. Math. Soc., Providence, R. I., 1971. MR **43** #7317.
10. S. C. Kleene, *Introduction to metamathematics*, North-Holland, Amsterdam, 1971.
11. G. Kreisel, *Set theoretic problems suggested by the notion of potential totality*, Infinitistic Methods (Proc. Sympos. Foundations of Math., Warsaw, 1959), Pergamon Press, Oxford, 1961, pp. 103–140. MR **26** #3599.
12. G. Kreisel and G. E. Sacks, *Metarecursive sets*, J. Symbolic Logic **30** (1965), 318–338. MR **35** #4097.
13. S. Kripke, *Transfinite recursion on admissible ordinals*. I, II (abstracts), J. Symbolic Logic **29** (1964), 161–162.
14. \_\_\_\_\_, *The theory of transfinite recursions*, Class Notes by A. Thomas Tymoczko (unpublished).
15. A. Leggett and R. Shore, *Types of simple  $\alpha$ -recursively enumerable sets* (to appear).

16. M. Lerman, *Least upper bounds for pairs of  $\alpha$ -r.e.  $\alpha$ -degrees*, J. Symbolic Logic **39** (1974), 49–56.
17. M. Lerman and G. E. Sacks, *Some minimal pairs of  $\alpha$ -recursively enumerable degrees*, Ann. Math. Logic **4** (1972), 415–442.
18. M. Machover, *The theory of transfinite recursion*, Bull. Amer. Math. Soc. **67** (1961), 575–578. MR **26** #17.
19. E. McCreight and A. Meyer, *Classes of computable functions defined by bounds on computation*, Preliminary report, Proc. ACM Sympos. on Theory of Computing, 1969, pp. 79–88.
20. R. Platek, *Foundations of recursion theory*, Ph.D. Thesis, Stanford Univ., 1966.
21. G. E. Sacks and S. G. Simpson, *The  $\alpha$ -finite injury method*, Ann. Math. Logic **4** (1972), 343–367. MR **51** #5277.
22. R. A. Shore,  *$\Sigma_n$  sets which are  $\Delta_n$  incomparable (uniformly)*, J. Symbolic Logic **39** (1974), 295–304.
23. \_\_\_\_\_, *Splitting an  $\alpha$ -recursively enumerable set*, Trans. Amer. Math. Soc. **204** (1975), 65–77. MR **52** #60.
24. \_\_\_\_\_, *The recursively enumerable  $\alpha$ -degrees are dense*, Ann. Math. Logic (to appear).
25. \_\_\_\_\_,  *$\alpha$ -recursion theory* (to appear).
26. G. Takeuti, *Construction of the set theory from the theory of ordinal numbers*, J. Math. Soc. Japan **6** (1954), 196–220. MR **16**, 783.
27. \_\_\_\_\_, *On the theory of ordinal numbers*, J. Math. Soc. Japan **9** (1957), 93–113. MR **19**, 237.
28. T. Tugué, *On the partial recursive functions of ordinal numbers*, J. Math. Soc. Japan **16** (1964), 1–31. MR **29** #4677.
29. P. Young, *Easy constructions in complexity theory: Gap and speed-up theorems*, Proc. Amer. Math. Soc. **37** (1973), 555–563. MR **47** #1323.

DEPARTMENT OF COMPUTER SCIENCE, QUEENS COLLEGE (CUNY), FLUSHING, NEW YORK 11367