

**CONDITIONALLY COMPACT
 SEMITOPOLOGICAL ONE-PARAMETER
 INVERSE SEMIGROUPS
 OF PARTIAL ISOMETRIES**

M. O. BERTMAN

ABSTRACT. The algebraic structure of one-parameter inverse semigroups has been completely described. Furthermore, if B is the bicyclic semigroup and if B is contained in any semitopological semigroup, the relative topology on B is discrete. We show that if F is an inverse semigroup generated by an element and its inverse, and F is contained in a compact semitopological semigroup, then the relative topology is discrete; in fact, if F is any one-parameter inverse semigroup contained in a compact semitopological semigroup, then the multiplication on F is jointly continuous if and only if the inversion is continuous on F , and we describe \bar{F} in that case. We also show that if $\{J_i\}$ is a one-parameter semigroup of bounded linear operators on a (separable) Hilbert space, then $\{J_i\} \cup \{J_i^*\}$ generates a one-parameter inverse semigroup T with $J_i^{-1} = J_i^*$ if and only if $\{J_i\}$ is a one-parameter semigroup of partial isometries, and we describe the weak operator closure of T in that case.

1. Introduction. An *inverse semigroup* is a semigroup in which each element x has a unique inverse x^{-1} with the properties that $xx^{-1}x = x$ and $x^{-1}xx^{-1} = x^{-1}$. If G is any subgroup of the positive real numbers under ordinary multiplication, let $P = G \cap [1, \infty)$. Let $F_P = \{(x, y, z) \in P^3: y \geq x \text{ and } y \geq z\}$ together with the following multiplication:

$$(x, y, z)(r, s, t) = \left(\frac{xzr}{y \wedge zr}, \frac{yzrs}{(y \wedge zr)(zr \wedge s)}, \frac{zrt}{zr \wedge s} \right)$$

where $x \wedge y = \min(x, y)$. Any inverse semigroup generated by a homomorphic image of P is a homomorphic image of F_P [6] and is called a *one-parameter inverse semigroup*. We mention at this point one homomorphic image of F_P . Let $B_P = P \times P$ together with this multiplication:

$$(x, y)(z, w) = \left(\frac{xz}{y \wedge z}, \frac{yw}{y \wedge z} \right).$$

Received by the editors March 1, 1976.

AMS (MOS) subject classifications (1970). Primary 47D05, 22A10.

Key words and phrases. Partial isometries, one-parameter semigroups, inverse semigroups, compact semitopological semigroups.


If $P = \{1, x, x^2, \dots\}$ where $x > 1$, then B_P is the *bicyclic semigroup* whose properties are discussed in [4].

Suppose that Y is a semigroup with a Hausdorff topology in which the maps $x \rightarrow xy$ and $x \rightarrow yx$ of $Y \rightarrow Y$ are continuous for each y in Y . Then the multiplication on Y is said to be *separately continuous* and Y is called a *semitopological semigroup*. If the map $(x, y) \rightarrow xy$ of $Y \times Y \rightarrow Y$ is continuous, then the multiplication on Y is *jointly continuous* (or just *continuous*) and Y is called a *topological semigroup*.

Let X be a one-parameter inverse semigroup generated by the image of P under a continuous homomorphism $f: P \rightarrow X$. Let $j: P \rightarrow F_P$ be the map $x \rightarrow (x, x, x)$. Then j is an isomorphism of P into F_P and we can identify P with its image in F_P . Then $x^{-1} = (1, x, 1)$ and hence $(x, y, z) = xy^{-1}z$. Furthermore, if $\bar{f}: F_P \rightarrow X$ is defined by

$$\bar{f}(xy^{-1}z) = f(x)f(y)^{-1}f(z),$$

then \bar{f} is a homomorphism and the following diagram commutes [6], [1]:



If F_P is endowed with the relative product topology which it inherits from R^3 , then the multiplication on F_P is jointly continuous. We shall show that if X is embedded densely in a compact semitopological semigroup and that if the inversion on $f(P) \cup f(P)^{-1}$ is continuous, then in fact \bar{f} is a *continuous* homomorphism of F_P onto X .

One-parameter inverse semigroups occur naturally in $B(H)$, the semigroup of bounded linear operators on Hilbert space. In the weak operator topology, the closed unit ball is a compact semitopological semigroup, and the map $T \rightarrow T^*$ is continuous where T^* is the adjoint of the operator T . A partial isometry is an operator which is isometric on the orthogonal complement of its kernel; it is known [8, p. 99] that T is a partial isometry if and only if $TT^*T = T$. We will show that if $J = \{J_t; t \in [0, \infty)\}$ is a one-parameter semigroup of partial isometries, then $J \cup J^*$ generates a one-parameter inverse semigroup.

This example shows that it is not unreasonable to ask that the map $x \rightarrow x^{-1}$ be continuous on X , and in that case we are able to give a description of X .

The author is grateful to D. B. McAlister and to T. T. West for many helpful suggestions. In particular, Professor McAlister pointed out (2.2) as a generalization of (2.3).

1.1. REMARK. The closure of an inverse semigroup in a compact topological semigroup is an inverse semigroup, but Brown and Moran [3, Proposition 1] give an example of a unitary group whose weak closure is not an inverse semigroup.

2. The one-parameter inverse semigroups. In this section we summarize the possible congruence relations on F_P (and hence its homomorphic images), as given in [6, 3.10]. We then point out that every one-parameter semigroup J of partial isometries generates a one-parameter inverse semigroup X , and that each of the possibilities for X can be realized as an inverse semigroup generated by a partial isometry.

If $1 < t \in P$, let $I_t = \{xy^{-1}z : y \succ t\} \subseteq F_P$ and $I_t^0 = \{xy^{-1}z \in F_P; y \succ t\}$. I_t and I_t^0 are the only ideals of F_P . Let α and β be the congruences on F_P defined so that $xy^{-1}z \alpha rs^{-1}t$ if and only if $x = r$ and $yt = sz$ and $xy^{-1}z \beta rs^{-1}t$ if and only if $z = t$ and $yr = xs$. Then $F_P/\alpha \approx B_P \approx F_P/\beta$. For each subgroup N of G , define σ_N to be the group congruence on F_P defined so that $xy^{-1}z \sigma_N rs^{-1}t$ if and only if $xzs/rty \in N$. Then if I is an ideal of F_P and $\delta = \alpha, \beta$, or σ_N , define the relation $I\delta$ by $xy^{-1}z I\delta rs^{-1}t$ if and only if either

- (a) $xy^{-1}z$ and $rs^{-1}t$ are in I and $xy^{-1}z \delta rs^{-1}t$ or
- (b) $xy^{-1}z = rs^{-1}t \notin I$.

2.1. PROPOSITION (EBERHART AND SELDEN). *Every congruence on F_P is $I\delta$ for some ideal I and congruence $\delta = \alpha, \beta$, or σ_N . Hence the only possible homomorphic images of F_P are $(F_P \setminus I) \cup B_P$, $(F_P \setminus I) \cup (G/N)$ or F_P [6, Theorem 3.10].*

2.2. PROPOSITION. *Let T be a semigroup with an involution $*$, i.e., a mapping $u \rightarrow u^*$ such that $(uv)^* = v^*u^*$ and $(u^*)^* = u$, and let $R = \{R_t; t \in P\}$ be a one-parameter subsemigroup of T . Then $\langle R, R^* \rangle$, the semigroup generated by $R \cup R^*$, is an inverse semigroup if and only if*

- (i) $R_t R_t^* R_t = R_t$ for each t in P and
- (ii) $R_t R_t^* R_s^* R_s = R_s^* R_s R_t R_t^*$ for each s and t in P .

PROOF. The conditions are clearly necessary. Now if R is as above, and $s \succ t$, we have

$$\begin{aligned} R_s R_t^* &= R_{s-t} R_t R_t^* = R_{s-t} R_{s-t}^* R_{s-t} R_t R_t^* \\ &= R_{s-t} R_t R_t^* R_{s-t}^* R_{s-t} = R_s R_s^* R_{s-t}. \end{aligned}$$

Similarly, $R_t^* R_s = R_{s-t} R_s^* R_s$, and it follows that any "word" from $\langle R, R^* \rangle$ can be expressed as a triple $R_x R_y^* R_z$ where $y \succ x$ and $y \succ z$. The arguments

of [6, pp. 56-57] can be used to show that the map $h: F_P \rightarrow \langle R, R^* \rangle$ defined by $h(xy^{-1}z) = R_x R_y^* R_z$ is a homomorphism, and since F_P is an inverse semigroup, so is its image $\langle R, R^* \rangle$.

Now let $J = \{J_t\}_{t=0}^\infty$ be a one-parameter semigroup of partial isometries on a (separable) Hilbert space. By [7, Theorem B],

$$J_t = U_t \oplus K_t \oplus S_t \oplus T_t,$$

where U_t is unitary, K_t is purely isometric, S_t is purely coisometric, and T_t is a nilpotent and hence, by [12], the direct integral of truncated shifts. Recall that a *truncated shift* is an operator which is unitarily equivalent to R_t , where for f in $\mathcal{L}^2[K, a]$ (if K is a separable Hilbert space and $a > 0$, $\mathcal{L}^2[K, a]$ is the Hilbert space of measurable K -valued functions on $[0, a]$ with square-integrable K -norm):

$$R_t f(x) = \begin{cases} 0 & \text{if } x < t \\ f(x-t) & \text{if } t \leq x < a \end{cases}$$

and $R_t = 0$ if $t > a$. This a is called the *index* of R_t . Since

$$R_t^* f(x) = \begin{cases} 0 & \text{if } a-t < x < a \\ f(x+t) & \text{if } 0 < x < a-t \end{cases}$$

and $R_t^* = 0$ if $t > a$, we have the following.

2.3. PROPOSITION. *If $R = \{R_t\}$ is a one-parameter semigroup of truncated shifts of index a , then $\langle R, R^* \rangle$ is isomorphic to F_P/I_a .*

PROOF. Since, for x, y , and z in P , with $y < a$,

$$R_x R_y^* R_z f(t) = \begin{cases} 0 & \text{if } t < x \text{ or } t > a - y + x \\ f(t - x + y - z) & \text{otherwise,} \end{cases}$$

it is easy to see that R satisfies the conditions of 2.2 and hence $\langle R, R^* \rangle$ is an inverse semigroup. Furthermore, one sees that $R_x R_y^* R_z = 0$ if and only if $y > a$, and hence by 2.1, $\langle R, R^* \rangle \approx F_P/I_a$.

It now follows that if $T = \{T_t\}$ is a one-parameter semigroup of nilpotent partial isometries of index a (and hence a direct integral of truncated shifts of index less than or equal to a), then $\langle T, T^* \rangle$ is isomorphic to $(F_P \setminus I_a) \cup \{0\}$.

Proposition 2.2 also gives the following.

2.4. PROPOSITION. *If $J = \{J_t\} = \{U_t \oplus K_t \oplus S_t \oplus T_t\}$ is a one-parameter semigroup of partial isometries, then $\langle J, J^* \rangle$ is a one-parameter inverse semigroup. Furthermore, if $\{U_t\} \neq I$ is a semigroup of unitary operators, $\{K_t\}$ of nonunitary isometries, $\{S_t\}$ of nonunitary coisometries, and $\{T_t\}$ of nilpotents of index a , then*

- (1) $\langle \{U_t \oplus S_t\}, \{U_t^* \oplus S_t^*\} \rangle \approx (F_P \setminus I_a) \cup R^+$;
- (2) $\langle \{K_t \oplus T_t\}, \{K_t^* \oplus T_t^*\} \rangle \approx (F_P \setminus I_a) \cup B_P$;

$$(3) \langle \{K_t \oplus S_t\}, \{K_t^* \oplus S_t^*\} \rangle \approx F_P;$$

$$(4) \langle \{K_t\}, \{K_t^*\} \rangle \approx B_P.$$

3. Conditionally compact one-parameter inverse semigroups. We now assume that X is a one-parameter inverse semigroup densely embedded in a compact semitopological semigroup Y . In case $P = \{1, x, x^2, \dots\}$ where $x > 1$, we show that X has the discrete topology as a subspace of Y . Furthermore, if P is dense in $[1, \infty)$, and the inversion $x \rightarrow x^{-1}$ on $P \cup P^{-1}$ is continuous, then the multiplication is continuous on X . We then assume, from 3.2.7 on, that inversion is continuous on X , and describe \bar{X} .

Since semigroups of the type B_P are somewhat more accessible than those of the more general type, and some information is known about them in case $P = \{1, x, x^2, \dots\}$ for $x > 1$, we devote a short section to these semigroups.

3.1. BICYCLIC SEMIGROUPS. As we have mentioned before, if $P = \{1, x, x^2, \dots\}$ where $x > 1$, then B_P is the familiar bicyclic semigroup, B , so-called because it is generated by two mutually inverse elements p and q subject to the generating relation $pq = 1$; we summarize the results of [2] on this subject:

(a) B is a discrete subspace of \bar{B} in the relative topology. (Originally proved in [5], as was (b).)

(b) $\bar{B} \setminus B$ is a closed two-sided ideal of \bar{B} .

(c) The minimal idempotent e of the monothetic semigroup $\Gamma = \{q, q^2, \dots\}$ is the minimal idempotent of \bar{B} , and the minimal ideal K of Γ , which is a compact topological group, with identity e , in the relative topology from \bar{B} , is the minimal ideal of \bar{B} . Furthermore $K = e\bar{B} = \bar{B}e$.

(d) The decreasing sequence $q^n p^n$ of idempotents converges to an idempotent w ; wB is a cyclic group with identity $w\bar{B} = \bar{w}\bar{B} = \bar{B} \setminus B$.


(e) If the map $x \rightarrow x^*$ is a continuous involution, then e and w are selfadjoint, and for each $x \in \bar{B}$, $(ex)^* = ex^* = (ex)^{-1}$ in K .

(f) If $B = \langle T, T^* \rangle$ where T is an isometry, then w is the Wold idempotent which decomposes H into $H_1 \oplus H_2$ where $T|H_1$ is unitary and $T|H_2$ is a unilateral shift. The minimal idempotent e decomposes H into $H_3 \oplus H_4$, where $T|H_3$ is unitary, and H_4 is the closed linear span of the eigenvalues of T (and T^*) corresponding to its unimodular eigenvalues.

These results show that we have no more (nor less) information about the closure of B than we have about the closure of a group, which examples in [3] and [12] show may be arbitrarily pathological.

If P is a dense subsemigroup of $[1, \infty)$, the existence of the idempotent w follows exactly as in [2], as does the fact that wB_P is a group. That $\bar{B}_P \setminus B_P = \bar{w}B_P$ turns out to be equivalent to the continuity of the inversion on B_P as we see for F_P in 3.2.6, and from 3.2.7 on, we shall assume this condition.

3.2. F_P . Suppose $X = F_P$. Then X may be pictured as below:



The idempotents of F_P are precisely those of the form $x(xy)^{-1}y$ where x and y are elements of P . [6, 2.13]. Now the idempotents $\{xx^{-1}: x \in P\}$ and $\{x^{-1}x: x \in P\}$ form two decreasing nets as $x \rightarrow \infty$, and hence there exist two idempotents E and F such that $xx^{-1} \rightarrow E$ and $x^{-1}x \rightarrow F$ as $x \rightarrow \infty$. Since xx^{-1} is decreasing, $E \neq xx^{-1}$ for any x in P , and similarly, $F \neq x^{-1}x$ for any x in P . Furthermore, $xx^{-1}E = E = Exx^{-1}$ for every x in P and $x^{-1}xF = F = Fx^{-1}x$ for every x in P .

PROPOSITION 3.2.1. *E and F commute with every element $xy^{-1}z$ of F_P , and hence with every element of \overline{F}_P .*

PROOF. As $x \rightarrow \infty$, $xx^{-1} \rightarrow E$, so if $r \in P$, $rxx^{-1} = r(r^{-1}r)(xx^{-1}) = r(xx^{-1})(r^{-1}r)$ (since idempotents commute) $= (rx)(rx)^{-1}r \rightarrow Er$; but $rxx^{-1} \rightarrow rE$, so $Er = rE$. On the other hand, for each r in P , $xx^{-1}r^{-1} = xx^{-1}r^{-1}rr^{-1} = r^{-1}rxx^{-1}r^{-1} = r^{-1}(rx)(rx)^{-1} \rightarrow r^{-1}E$; but $xx^{-1}r^{-1} \rightarrow Er^{-1}$, so $Er^{-1} = r^{-1}E$ for each r in P . Hence if $xy^{-1}z \in F_P$, $Exy^{-1}z = xEy^{-1}z = xy^{-1}Ez = xy^{-1}zE$. Similarly, F commutes with each element of F_P and hence with each element of \overline{F}_P .

PROPOSITION 3.2.2. *If $\{z_\alpha\}$ and $\{y_\alpha\}$ are unbounded increasing nets from P , then $\{x_\alpha x_\alpha^{-1} y_\alpha^{-1} y_\alpha\}$ converges to EF .*

PROOF. Let $\{x_{\alpha_\beta} x_{\alpha_\beta}^{-1} y_{\alpha_\beta}^{-1} y_{\alpha_\beta}\}$ be a subnet which converges, say to Z . Then

$$EFx_{\alpha_\beta} x_{\alpha_\beta}^{-1} y_{\alpha_\beta}^{-1} y_{\alpha_\beta} = Ex_{\alpha_\beta} x_{\alpha_\beta}^{-1} Fy_{\alpha_\beta}^{-1} y_{\alpha_\beta} = EF,$$

but this net converges to EFZ , so $EF = EFZ$. Now let $x \in P$, and assume $x_{\alpha_\beta} > x$. Then

$$xx^{-1}x_{\alpha_\beta} x_{\alpha_\beta}^{-1} y_{\alpha_\beta}^{-1} y_{\alpha_\beta} = x_{\alpha_\beta} x_{\alpha_\beta}^{-1} y_{\alpha_\beta}^{-1} y_{\alpha_\beta} \rightarrow Z,$$

so $xx^{-1}Z = Z$ for each x in P . Similarly $Zx^{-1}x = Z$ for each x in P , and

hence, letting x approach ∞ , $EZ = Z = ZF$, so $EFZ = EZF = EZ = Z$, so $EF = EFZ = Z$. It now follows that $x_\alpha x_\alpha^{-1} y_\alpha^{-1} y_\alpha \rightarrow EF$.

PROPOSITION 3.2.3. $EF_P \cap F_P = \emptyset$; $FF_P \cap F_P = \emptyset$ and $EFF_P \cap F_P = \emptyset$.

PROOF. Suppose $E \in F_P$; then $E = xx^{-1}y^{-1}y$ for some x and y in P . But if $k > x$, $E = kk^{-1}E = kk^{-1}xx^{-1}y^{-1}y = kk^{-1}y^{-1}y \neq xx^{-1}y^{-1}y = E$. Thus E is not in F_P , and a similar maneuver shows that F and EF are not in F_P . Now we show that $EF_P \cap F_P = \emptyset$. Suppose that there exists x in P such that $Ex^{-1} \in F_P$. Then $E = Exx^{-1} = xEx^{-1} \in F_P$. So there is no such x . Now if $xy^{-1}z$ is any element of F_P such that $Exy^{-1}z \in F_P$, then

$$\begin{aligned} E(y/x)^{-1} &= Exx^{-1}(y/x)^{-1} = Exy^{-1} \\ &= xy^{-1}Ezz^{-1} = (Exy^{-1}z)z^{-1} \in F_P, \end{aligned}$$

and this contradiction proves that there is no such $xy^{-1}z$. The proof for F and EF is similar.

PROPOSITION 3.2.4. *If $\{x_\alpha y_\alpha^{-1} z_\alpha\}$ is a net from F_P such that $\{y_\alpha\}$ is unbounded, then if $x_\alpha y_\alpha^{-1} z_\alpha \rightarrow u$, $u \notin F_P$. In fact, $u \in \overline{EF_P} \cup \overline{FF_P}$.*

PROOF. Note that if e is any idempotent and x any element of $\overline{F_P}$, then $ex = x$ if and only if $x \in \overline{eF_P}$. Now if $\{x_\alpha\}$ is unbounded, for any k in P , x_α is eventually larger than k , so $kk^{-1}x_\alpha y_\alpha^{-1} z_\alpha = x_\alpha y_\alpha^{-1} z_\alpha \rightarrow u$, so $kk^{-1}u = u$, and hence $Eu = u$. If u were in F_P , u would be in EF_P , so $u \notin F_P$. Similarly, if $\{z_\alpha\}$ is unbounded, u would be in $FF_P \setminus F_P$. If $\{x_\alpha\}$ and $\{z_\alpha\}$ are both bounded nets and $\{y_\alpha\}$ is unbounded, then for every $x > \sup\{x_\alpha\}$,

$$\begin{aligned} x^{-1}x x_\alpha y_\alpha^{-1} z_\alpha &= x^{-1}x x_\alpha x_\alpha^{-1} (y_\alpha/x_\alpha)^{-1} z_\alpha \\ &= x_\alpha x_\alpha^{-1} x^{-1} x (y_\alpha/x_\alpha)^{-1} z_\alpha \\ &= x_\alpha x_\alpha^{-1} (y_\alpha/x_\alpha)^{-1} z_\alpha = x_\alpha y_\alpha^{-1} z_\alpha \rightarrow u. \end{aligned}$$

But $x^{-1}x x_\alpha y_\alpha^{-1} z_\alpha \rightarrow x^{-1}xu$, so $x^{-1}xu = u$, and again $u \in FF_P \setminus F_P$.

PROPOSITION 3.2.5. *If $P = \{1, x, x^2, x^3, \dots\}$, then F_P has the discrete topology as a subspace of any compact semitopological semigroup.*

PROOF. Suppose $x_\alpha y_\alpha^{-1} z_\alpha \rightarrow xy^{-1}z$. Then by 3.2.4, $\{y_\alpha\}$, $\{x_\alpha\}$ and $\{z_\alpha\}$ are bounded nets in P . Hence we can find x_0, y_0, z_0 and subnets $x_{\alpha_0}, y_{\alpha_0}, z_{\alpha_0}$ such that $x_{\alpha_0} \rightarrow x_0, y_{\alpha_0} \rightarrow y_0$, and $z_{\alpha_0} \rightarrow z_0$. Thus eventually $x_{\alpha_0} = x_0, y_{\alpha_0} = y_0$, and $z_{\alpha_0} = z_0$, and hence eventually $x_{\alpha_0} y_{\alpha_0}^{-1} z_{\alpha_0} = x_0 y_0^{-1} z_0$. Thus since

$$x_{\alpha_0} y_{\alpha_0}^{-1} z_{\alpha_0} \rightarrow xy^{-1}z, \quad x_0 y_0^{-1} z_0 = xy^{-1}z.$$

This shows that the only convergent nets are eventually constant, and so the relative topology on F_P is the discrete.

Proposition 3.2.5 assures us that if P is $\{1, x, x^2, \dots\}$, then both the

multiplication on F_P and the inversion map on F_P are continuous functions. Moreover, since every convergent net which is bounded in its second coordinate converges to a point of F_P , and every convergent net which is unbounded in its second coordinate converges to a point of $\overline{EP_P}$ or $\overline{FF_P}$, we see that $\overline{EF_P} \cup \overline{FF_P}$ is an ideal of $\overline{F_P}$ and $\overline{F_P} \setminus F_P = \overline{EF_P} \cup \overline{FF_P}$. We next show that all these related conditions are equivalent.

PROPOSITION 3.2.6. *If F_P is contained densely in a compact semitopological semigroup, and $P \subseteq F_P$ is a continuous embedding, then the following are equivalent:*

- (a) *Inversion is continuous on $P \cup P^{-1}$;*
 - (b) *$P^{-1} \subseteq F_P$ is a continuous embedding;*
 - (c) *The topology on F_P is the product topology;*
- that is, $x_\alpha y_\alpha^{-1} z_\alpha \rightarrow xy^{-1}z$ if and only if $x_\alpha \rightarrow x, y_\alpha \rightarrow y$, and $z_\alpha \rightarrow z$;
- (d) *If $\{y_\alpha\}$ is a bounded net from P , and if $x_\alpha y_\alpha^{-1} z_\alpha \rightarrow u, u \in F_P$;*
 - (e) *$\overline{F_P} \setminus F_P = \overline{EF_P} \cup \overline{FF_P}$.*
 - (f) *$\overline{F_P} \setminus F_P$ is an ideal of F_P , if $\overline{F_P} \setminus F_P \neq \emptyset$.*

PROOF. The implications (a) \Leftrightarrow (b) and (c) \Rightarrow (d) are easy. We show (a) \Rightarrow (c) and (d) \Rightarrow (a), and (d) \Rightarrow (e) \Rightarrow (f) \Rightarrow (d).

(a) \Rightarrow (c). Suppose inversion is continuous in $P \cup P^{-1}$. We note that if $x_\alpha y_\alpha^{-1} z_\alpha \rightarrow u$ where $x_\alpha \rightarrow x$ and $z_\alpha \rightarrow z$, and if $x > 1$, and $1 < r < x$, then since $r < x_\alpha$ eventually, $rr^{-1}x_\alpha y_\alpha^{-1} z_\alpha = x_\alpha y_\alpha^{-1} z_\alpha$ eventually, so $rr^{-1}u = u$. (Similarly $ut^{-1}t = u$ if $t < z$.) Now if $x_\alpha \rightarrow x$, we show that $x_\alpha x_\alpha^{-1} \rightarrow xx^{-1}$. We can assume that $x_\alpha x_\alpha^{-1} \rightarrow u$ for some $u \in \overline{F_P}$. If $r > x$, then eventually $r > x_\alpha$, so $r^{-1}x_\alpha x_\alpha^{-1} = r^{-1}$, and so $r^{-1} = r^{-1}u$; letting $r \rightarrow x, r^{-1} \rightarrow x^{-1}$ by assumption and thus $x^{-1} = x^{-1}u$, and $xx^{-1} = xx^{-1}u$. If $x_\alpha \geq x$ frequently, then

$$xx^{-1}x_\alpha x_\alpha^{-1} = x_\alpha x_\alpha^{-1}$$

frequently, so $xx^{-1}u = u$. On the other hand, if $x_\alpha < x$ eventually, then since $xx_\alpha^{-1} \rightarrow xx^{-1}$ we have

$$xx_\alpha^{-1}u = (x/x_\alpha)x_\alpha x_\alpha^{-1}u = x/x_\alpha u$$

by the above, so $xx^{-1}u = u$, so in either case, $u = xx^{-1}u = xx^{-1}$. Similarly $x_\alpha^{-1}x_\alpha \rightarrow x^{-1}x$. Now if $x_\alpha \rightarrow x, y_\alpha \rightarrow y, z_\alpha \rightarrow z$, we can assume $x_\alpha y_\alpha^{-1} z_\alpha \rightarrow u$. Let $r > x$ and $t > z$; then

$$rr^{-1}x_\alpha y_\alpha^{-1} z_\alpha t^{-1}t = r \left(\frac{ry_\alpha t}{x_\alpha z_\alpha} \right)^{-1} t,$$

so

$$rr^{-1}ut^{-1}t = r \left(\frac{ryt}{xz} \right)^{-1} t = rr^{-1}(yt/xz)t.$$

Letting $r \rightarrow x$, we have $xx^{-1}ut^{-1}t = xx^{-1}(yt/xz)^{-1}z = x(yt/z)^{-1}t = x(y/z)^{-1}t^{-1}t$, and letting $t \rightarrow z$,

$$u = xx^{-1}uz^{-1}z = x(y/z)^{-1}z^{-1}z = xy^{-1}z.$$

It follows that $x_\alpha y_\alpha^{-1} z_\alpha \rightarrow xy^{-1}z$ if and only if $x_\alpha \rightarrow x, y_\alpha \rightarrow y$, and $z_\alpha \rightarrow z$.

(d) \Rightarrow (a). Let $x_\alpha \rightarrow x$ in P , and suppose $x_\alpha^{-1} \rightarrow rs^{-1}t$ in F_P . We show $r \leq x$, and $t \leq x$. For suppose $r > x$. Then $r > x_\alpha$ eventually, so since

$$rr^{-1}\left(\frac{r}{x_\alpha}\right) = rx_\alpha^{-1} \rightarrow rrs^{-1}t = r^2\left(\frac{r^2s}{r^2 \wedge s}\right) \frac{r^2t}{r^2t}$$

we have

$$rr^{-1}r/x = r^2\left(\frac{r^2s}{r^2 \wedge s}\right)^{-1} \frac{r^2t}{r^2 \wedge s}$$

and hence $r = 1$, and this contradiction shows that $r \leq x$; similarly $t \leq x$. Now suppose that $s < x$. Then since $s < x_\alpha$ eventually, and $x_\alpha^{-1} = x_\alpha^{-1}ss^{-1} \rightarrow rs^{-1}tss^{-1} = rt(st)^{-1}t$, we have $t = 1$. Similarly, we can show $r = 1$. Thus $x_\alpha^{-1} \rightarrow s^{-1}$ where $s < x$. Now if $x_\alpha < x$ frequently, the fact that

$$(x/x)x^{-1} = x_\alpha^{-1}xx^{-1} \rightarrow s^{-1}xx^{-1} = (x/s)x^{-1}$$

implies that $x = s$. If $x_\alpha > x$ frequently, then

$$s^{-1} = \lim_\alpha x_\alpha^{-1} = \lim_\alpha (x_\alpha^{-1}xx^{-1}) = s^{-1}xx^{-1} = (x/s)x^{-1},$$

so $x = s$. Hence if $x_\alpha^{-1} \rightarrow rs^{-1}t$, $r \leq x$, $t \leq x$, and $s \geq x$. Suppose $s > x$. Then $s > x_\alpha$ eventually, so since $(s/x_\alpha)s^{-1} = x_\alpha^{-1}ss^{-1} \rightarrow rs^{-1}tss^{-1} = rt(st)^{-1}t$, $t = 1$, and $s/x = r$. Similarly, $r = 1$, so $s = x$. Therefore, if $x_\alpha \rightarrow x$, then $x_\alpha^{-1} \rightarrow x^{-1}$.

(d) \Rightarrow (e). Let $u \in \overline{F_P} \setminus F_P$. Then $u = \lim_\alpha x_\alpha y_\alpha^{-1} z_\alpha$, where $\{y_\alpha\}$ is unbounded. As in 3.2.4, either $Eu = u$ or $Fu = u$. Hence $u \in \overline{EF_P} \cup \overline{FF_P}$. If $v \in \overline{EF_P} \cup \overline{FF_P}$, then $Ev = v$ or $Fv = v$, so by 3.2.3, $v \in \overline{F_P} \setminus F_P$.

(e) \Rightarrow (f). Suppose $\overline{F_P} \setminus F_P = \overline{EF_P} \cup \overline{FF_P}$. If $u \in \overline{F_P} \setminus F_P$, then either $Eu = u$ or $Fu = u$, so if $v \in \overline{F_P}$, $uv = Eu v \in \overline{EF_P}$, or else $uv = Fuv \in \overline{FF_P}$, so $uv \in \overline{F_P} \setminus F_P$.

(f) \Rightarrow (d). We first show that if $\{y_\alpha\}$ is bounded and $y_\alpha^{-1} \rightarrow u$, then $u \in F_P$. Let $r > y_\alpha$ for each α . We can assume y_α converges, say to y in P , and hence $r^{-1}ru = r^{-1}r/y$, and since $r^{-1}ru \in F_P$, $u \in F_P$. Now let $x_\alpha y_\alpha^{-1} z_\alpha \rightarrow v$. Let $r > x_\alpha$ and $t > z_\alpha$ for each α . Then

$$rr^{-1}x_\alpha y_\alpha^{-1} z_\alpha t^{-1}t = r\left(\frac{ry_\alpha t}{x_\alpha z_\alpha}\right)^{-1} t;$$

we can assume $ry_\alpha t^{-1}/x_\alpha z_\alpha$ converges to q , hence $q \in F_P$; so $rr^{-1}vt^{-1}t = rqt \in F_P$ and hence $v \in F_P$.

3.3. For the remainder of this paper, we will assume X is a homomorphic image of F_P embedded densely in a compact semitopological semigroup, and that the inversion on X is a continuous mapping.

PROPOSITION 3.3.1. *If $X = F_P/\delta$ and if δ^h is the natural map of F_P to X , then, under the assumption that inversion is continuous on X , δ^h is a continuous homomorphism from F_P with the product topology onto X .*

PROOF. It follows as in 3.2.6 that δ^h is continuous on $F_P \setminus I$ where $\delta = I\sigma$. Now since δ^h must be continuous on P , we know $I = I_a$ for some $a > 1$ and if $\sigma = \sigma_N$ for some subgroup N of G , N must be a closed subgroup of G . We can assume P is dense in $[1, \infty)$. We prove the proposition for $F_P/I_a\alpha$ and point out that the proofs for the remaining cases are similar.

Suppose $X = F_P/I_a\alpha$. Then we note that $X \approx (F_P \setminus I_a) \cup B_P$, where the multiplication is as follows: If both u and v are in F_P then uv is as usual. If either u or v is in B_P , then $uv = \alpha^h(u)\alpha^h(v)$. For the sake of simplifying the notation, we equate $xy^{-1}z$ with $(x, y/z)$ if $y > a$.

Let $x_\alpha \rightarrow a$ from $[1, a)$, and suppose $x_\alpha x_\alpha^{-1} \rightarrow e$. Since for each $r < a$, $x_\alpha x_\alpha^{-1} r r^{-1} = x_\alpha x_\alpha^{-1}$ eventually, $err^{-1} = e$. Now $x_\alpha x_\alpha^{-1}(a, a) = (a, a)$, so $e(a, a) = (a, a)$. But $e(a/x_\alpha)^{-1} \rightarrow e$ as $x_\alpha \rightarrow a$, and

$$e(a/x_\alpha)^{-1} = ex_\alpha x_\alpha^{-1}(a/x_\alpha)^{-1} = e(x_\alpha, a) \rightarrow e(a, a) = (a, a),$$

so $e = (a, a)$. Similarly, if $x_\alpha^{-1}x_\alpha \rightarrow f, f(1, 1) = (1, 1)$, so since

$$f(a/x_\alpha) = fx_\alpha^{-1}x_\alpha(a/x_\alpha) = fx_\alpha^{-1}(a, 1) = f(a/x_\alpha, 1) \rightarrow f(1, 1)$$

and $f(a/x_\alpha) \rightarrow f$, we have $f = f(1, 1) = (1, 1)$. Now suppose that $x_\alpha \rightarrow x$, and $y_\alpha \rightarrow b > a$; we assume $x_\alpha y_\alpha^{-1} \rightarrow u$. Then $y_\alpha > a$ eventually, so $x_\alpha y_\alpha^{-1} = (x_\alpha, y_\alpha)$ eventually, and $x_\alpha y_\alpha^{-1} \rightarrow (x, b)$. If $y_\alpha \rightarrow a$, we can assume $y_\alpha < a$ eventually. As above, if $r < a$, $x_\alpha y_\alpha^{-1} = x_\alpha y_\alpha^{-1} r r^{-1} \rightarrow u r r^{-1}$ so $u = u r r^{-1}$ and hence, letting $r \rightarrow a$, $u = u(a, a)$. But $x_\alpha y_\alpha^{-1}(a, a) \rightarrow u(a, a)$ and $x_\alpha y_\alpha^{-1}(a, a) = (x_\alpha a/y_\alpha, a) \rightarrow (x, a)$ so $u = (x, a)$. If $z_\alpha \rightarrow z$, and $y_\alpha \rightarrow b > a$, then since $y_\alpha^{-1}z_\alpha = (1, y_\alpha/z_\alpha)$ eventually, $y_\alpha^{-1}z_\alpha \rightarrow (1, b/z)$. If $y_\alpha \rightarrow a$, assume $y_\alpha^{-1}z_\alpha \rightarrow u$, and $y_\alpha < a$ eventually. Then if $r < a$, $r^{-1}ru = u$, and, letting $r \rightarrow a$, $(1, 1)u = u$ by the above; but $(1, 1)(y_\alpha^{-1}z_\alpha) = (1, y_\alpha/z_\alpha) \rightarrow (1, a/z)$ so $u = (1, a/z)$. Now we show in general that if $x_\alpha \rightarrow x, y_\alpha \rightarrow y$, and $z_\alpha \rightarrow z$, then $x_\alpha y_\alpha^{-1}z_\alpha \rightarrow xy^{-1}z$. We can assume $x_\alpha y_\alpha^{-1}z_\alpha \rightarrow u$. If $y > a$, then $x_\alpha y_\alpha^{-1}z_\alpha = (x_\alpha, y_\alpha/z_\alpha)$ eventually and we are done. If $y = a$ and $x > 1$, suppose $x < a$, and let $x < r < a$. Then eventually $rr^{-1}x_\alpha y_\alpha^{-1}z_\alpha = r(ry_\alpha/x_\alpha)^{-1}z_\alpha \rightarrow (r, ry/xz)$ so $rr^{-1}u = (r, ry/xz)$, and, letting $r \rightarrow x$, $xx^{-1}u = (x, y/z)$ but as before $xx^{-1}u = u$, so $u = (x, y/z)$. Finally, if $x = a$, then $u = (a, a)u = (a, a/z)$. Thus if $\delta^h(x_\alpha) \rightarrow \delta^h(x)$, $\delta^h(y_\alpha) \rightarrow \delta^h(y)$, and $\delta^h(z_\alpha) \rightarrow \delta^h(z)$, then $\delta^h(x_\alpha y_\alpha^{-1}z_\alpha) \rightarrow \delta^h(xy^{-1}z)$, and it follows that δ^h is a continuous homomorphism of F_P onto X . The proof is exactly analogous if $X = F_P/I_a\sigma_N$.

Now recall that for each X , the decreasing nets $\{\delta^{\sharp}(x)\delta^{\flat}(x)^{-1}: x \in P\}$ and $\{\delta^{\sharp}(x)^{-1}\delta^{\flat}(x): x \in P\}$ converge to idempotents E and F respectively as $x \rightarrow \infty$. Furthermore, in each case, E and F commute with X , and 3.3.1 allows us to conclude that in fact, $\bar{X} = X \cup \overline{EX} \cup \overline{FX}$. We point out that if $X = F_P/I_a\sigma_N$, $E = F = \delta^{\sharp}(aa^{-1})$ is the identity element of the group which is the minimal ideal of X . If $X = F_P/I_a\alpha$, $F = (1, 1) = \delta^{\sharp}(a^{-1}a)$ but $E \notin X$. In this case, $\bar{X} \setminus X = \overline{EX}$. Similarly, if $X = F_P/I_a\beta$, then $\bar{X} \setminus X = \overline{FX}$.

PROPOSITION 3.3.3. *If X is any homomorphic image of F_P , then $EX \approx B_P$ or EX is a commutative group; EX is a group if and only if $E = EF$, and these remarks are true if E and F are interchanged. Furthermore, $\bar{X} = X \cup \overline{EX} \cup \overline{FX}$.*

PROOF. Since E commutes with each element of X , the map $u \rightarrow Eu$ is a homomorphism of X to EX . Hence $EX = F_P/I_b\sigma$ for some $b \in P$ and congruence σ on FP . Since $Exx^{-1} = E$ for each x in $(I_b\sigma)^{\sharp}(P)$, $b = 1$, and hence $EX = F_P/\beta$ or F_P/σ_N . In the first case, $EX \approx B_P$ with E as identity and

$$E\delta^{\sharp}(xy^{-1}z) = E(y/x)^{-1}z \approx (y/x, z),$$

and the idempotents $Ex^{-1}x$ converge to EF as $x \rightarrow \infty$. It is easy to see that EFX is a commutative group in this case. In the second case, EX is a group and since $Ex^{-1}x = E$, $E = EF$. The discussion in the previous paragraph together with 3.2.6 implies that $\bar{X} = X \cup \overline{EX} \cup \overline{FX}$.

The idempotents E and F are thus maximal among idempotents of $\bar{X} \setminus X$, if $\bar{X} \setminus X \neq \emptyset$. We now turn to a discussion of minimal idempotents of \bar{X} . The one-parameter semigroups $\overline{\delta^{\sharp}(P)}$ and $\overline{\delta^{\flat}(P^{-1})}$ possess as kernels (minimal ideals) commutative compact topological groups K and K' , respectively, with identities e_0 and f_0 . If $X = B_P$, it follows as in [2] that $e_0 = f_0$ and hence $K = K'$ is the minimal ideal for \bar{X} ; furthermore $K \subseteq \overline{EX}$ and $K = \overline{EX}$ if and only if $e_0 = E = w$. These facts are easy to prove if X is any of the other images of F_P ; we discuss the case where $X = F_P$.

If $\{x_{\alpha}\}$ is a net from P which converges to e_0 , by 3.2.6, $\{x_{\alpha}\}$ is unbounded. If $x \in P$, $x < x_{\alpha}$ frequently, so $xx^{-1}x_{\alpha} = x_{\alpha}$ frequently and hence $xx^{-1}e_0 = e_0 = Ee_0 = e_0E$. Similarly $e_0F = e_0 = Fe_0$. Thus $EFe_0 = e_0$ so $e_0 \in \overline{EFX}$, and hence e_0 commutes with each element of \bar{X} . Now if $y \in P$, $x_{\alpha}y^{-1} \rightarrow e_0y^{-1}$. But since $x_{\alpha} > y$ frequently, $x_{\alpha}y^{-1} = x_{\alpha}x_{\alpha}^{-1}x_{\alpha}/y$, and $e_0x_{\alpha}y^{-1} \rightarrow e_0y^{-1}$. Thus since $e_0x_{\alpha}y^{-1} = e_0x_{\alpha}x_{\alpha}^{-1}x_{\alpha}/y = e_0x_{\alpha}/y$, e_0y^{-1} is in $\bar{K} = K$. Similarly $y^{-1}e_0$ is in K . Thus $e_0X = Xe_0 \subseteq K$, and $e_0\bar{X} = \bar{X}e_0 \subseteq K$. Now if e is any idempotent of \bar{X} , $(e_0e)(e_0e) = e_0e_0ee = e_0e$, and since $e_0e \in K$, $e_0e = e$. Hence e_0 is the minimal idempotent for \bar{X} . It follows that $e_0 = f_0$ and as in [2, Proposition 2] that K is the kernel for \bar{X} and that $K = e_0\bar{X}$. We summarize this discussion in the following proposition.

PROPOSITION 3.3.4. *If $X = F_P/\delta$, then the kernel of \overline{X} is K , the compact topological group which is the kernel of $\delta^{\sharp}(P)$. If e_0 is the identity element of K , then $K = e_0\overline{X}$.*

We now derive some corollaries and give some examples from the space of bounded linear operators on complex Hilbert space. This one is proved as in [2, Proposition 5].

COROLLARY 3.3.5. *If \overline{X} is a compact semitopological semigroup with a continuous involution $t \rightarrow t^*$, all idempotents of X , EX and FX are selfadjoint, as are EF and e_0 . If $x \in K$, then $x^* \in K$, and furthermore, $x^*x = e = xx^*$ for every x in K .*

The following corollary was proved in [1, Theorem 4].

COROLLARY 3.3.6. *If \overline{X} is a compact topological semigroup, then since $EX \neq B_P \neq FX$, $E = F$ and hence $\overline{X} = X \cup \overline{EX}$.*

EXAMPLE 3.3.7. If U is unitary, and T and S are unilateral shifts, let $J = U \oplus T \oplus S^*$. Then $J^n = U^n \oplus T^n \oplus (S^*)^n$; let $X = \langle J, J^* \rangle$. We have $E = I_1 \oplus O_2 \oplus I_3$, $F = I_1 \oplus I_2 \oplus O_3$, and $EF = I_1 \oplus O_2 \oplus O_3$. EX is the bicyclic semigroup generated by $U \oplus O_2 \oplus S^* = p$ and $U^* \oplus O_2 \oplus S = q$ with $1 = pq = E$, and FX is the bicyclic semigroup $\langle U \oplus T \oplus O_3, U^* \oplus T^* \oplus O_3 \rangle$. It is easy to see that the structure of \overline{EFX} is totally dependent on U .

REMARK 3.3.9. If $\{J_i\} = J$ is a one-parameter semigroup of partial isometries on a finite-dimensional complex space, then $J_i = U_i \oplus T_i$, where U_i is unitary and T_i is a truncated shift.

PROOF. The uniform and weak topologies coincide, so $\overline{X} = \langle J \cup J^* \rangle$ is a compact topological semigroup. If $J_i = U_i \oplus S_i \oplus C_i^* \oplus T_i$, where either S_i or C_i is a nonunitary isometry, then either EX or FX would be bicyclic. Since this cannot happen in a compact topological semigroup, S_i and C_i^* are both unitary.

REFERENCES

1. M. Bertman, *Free topological inverse semigroups*, Semigroup Forum 8 (1974), 226–270.
2. M. Bertman and T. T. West, *Conditionally compact bicyclic semitopological semigroups*, Proc. Roy. Irish Acad. Sect. A 76 (1975), 219–226.
3. G. Brown and W. Moran, *Idempotents of compact monothetic semigroups*, Proc. London Math. Soc. 22 (1971), 203–216.
4. A. H. Clifford and G. Preston, *The algebraic theory of semigroups*, vol. I, Math. Surveys, no. 7, Amer. Math. Soc., Providence, R.I., 1961.
5. C. Eberhart and J. Selden, *On the closure of the bicyclic semigroup*, Trans. Amer. Math. Soc. 144 (1969), 115–126.
6. ———, *One-parameter inverse semigroups*, Trans. Amer. Math. Soc. 168 (1972), 53–66.
7. M. Embry, A. Lambert and L. Wallen, *A simplified treatment of the structure of semigroups of partial isometries*, Michigan Math. J. 22 (1975), 175–179.
8. P. R. Halmos, *A Hilbert space problem book*, Van Nostrand, Princeton, N. J., 1967.

9. K. H. Hofmann and P. S. Mostert, *Elements of compact semigroups*, Merrill, Columbus, Ohio, 1966.
10. K. DeLeeuw and I. Glicksberg, *Applications of almost periodic compactifications*, *Acta Math.* **105** (1961), 63–98.
11. D. B. McAllister, *A homomorphism theorem for semigroups*, *J. London Math. Soc.* **43** (1968), 355–366.
12. L. J. Wallen, *Decomposition of semi-groups of partial isometries*, *Indiana Univ. Math. J.* **20** (1970), 207–212.
13. T. T. West, *Weakly compact monothetic semigroups of operators in Banach spaces*, *Proc. Roy. Irish Acad. Sect. A* **67** (1968), 27–37.

DEPARTMENT OF MATHEMATICS, CLARKSON COLLEGE, POTSDAM, NEW YORK 13676