

GENERIC COHOMOLOGY FOR TWISTED GROUPS

BY

GEORGE S. AVRUNIN

ABSTRACT. Let G be a simple algebraic group defined and split over $k_0 = \mathbb{F}_p$, and let σ be a surjective endomorphism of G with finite fixed-point set G_σ . We give conditions under which cohomology groups of G are isomorphic to cohomology groups of G_σ .

Let G be a simple algebraic group defined and split over $k_0 = \mathbb{F}_p$, and, for $q = p^m$, let $G(q)$ be the subgroup of \mathbb{F}_q -rational points. For a finite dimensional rational G -module V and a nonnegative integer e , let $V(e)$ be the G -module obtained by twisting the original G -action on V by the Frobenius endomorphism $x \rightarrow x^{[p^e]}$ of G . Cline, Parshall, Scott, and van der Kallen proved in [2] that, for sufficiently large q and e (depending on V and n), the restriction map induces an isomorphism from the rational cohomology group $H^n(G, V(e))$ to $H^n(G(q), V(e))$. This implies that, as q increases, the groups $H^n(G(q), V(e))$ have a stable or generic value $H_{\text{gen}}^n(G, V)$.

In this paper, we prove an analogous theorem with $G(q)$ replaced by G_σ for a surjective endomorphism σ of G having finite fixed point set. The first section of the paper summarizes the basic results on endomorphisms of algebraic groups required for the proof. Some arithmetic facts are established in the second section, and the main theorem is proved in the third section.

The author is grateful to Leonard L. Scott for several helpful conversations, and to the referee for comments on an earlier version of this paper.

1. Endomorphisms of algebraic groups. We briefly review here some results on endomorphisms of algebraic groups which will be needed later. We refer the reader to [4] and [5] for more details.

Let k be the algebraic closure of the prime field $k_0 = \mathbb{F}_p$ and let G be a simple algebraic group defined and split over k_0 . If σ is a surjective rational endomorphism of G having finite fixed-point set G_σ , σ stabilizes a Borel subgroup B and a maximal torus $T \leq B$. All such pairs (B, T) are G_σ -conjugate.

Assume G_σ is finite and fix a σ -stable pair (B, T) . Let Σ be the root system of T in G , with fundamental system $\Delta \leq \Sigma^+$ defined by B . The comorphism σ^* of $\sigma|_T$ determines a permutation ρ of Σ stabilizing Σ^+ and Δ , and powers $q(\alpha)$ of p such that $\sigma^*\rho(\alpha) = q(\alpha)\alpha$. If $U_\alpha \leq G$ is the root subgroup determined by α and $x_\alpha: {}_k\text{Add} \rightarrow U_\alpha$ is a T -equivariant isomorphism, then $\sigma x_\alpha(u) = x_{\rho\alpha}(c_\alpha u^{q(\alpha)})$ for some

Received by the editors May 5, 1980 and, in revised form, November 12, 1980.

1980 *Mathematics Subject Classification.* Primary 20G10.

Key words and phrases. Cohomology of algebraic groups, finite groups of Lie type.

© 1981 American Mathematical Society
0002-9947/81/0000-0513/\$02.75

$c_\alpha \in k$. Adjusting the isomorphisms x_α as necessary, the c_α can be taken to be ± 1 , with $c_\alpha = 1$ if α or $-\alpha$ is fundamental.

If ρ is the identity, all $q(\alpha)$ are equal, G_σ is a finite Chevalley group, and the results of [2] apply. We are primarily concerned with the cases in which ρ is a nontrivial permutation. These are listed below.

I. G is of type A_n , ρ is the nontrivial graph automorphism, and $q(\alpha) = p^f$ for all α . G_σ is ${}^2A_n(p^f)$.

II. G is of type D_n , ρ is a graph automorphism of order 2, and $q(\alpha) = p^f$ for all α . G_σ is ${}^2D_n(p^f)$.

III. G is of type D_4 , ρ is a graph automorphism of order 3, and $q(\alpha) = p^f$ for all α . G_σ is ${}^3D_4(p^f)$.

IV. G is of type E_6 , ρ is the nontrivial graph automorphism, and $q(\alpha) = p^f$ for all α . G_σ is ${}^2E_6(p^f)$.

V. G is of type C_2 , ρ interchanges long and short roots, $p = 2$, and $q(\alpha) = 2^f$ if α is long, while $q(\alpha) = 2^{f+1}$ if α is short. G_σ is ${}^2C_2(2^f)$.

VI. G is of type F_4 . If the successive nodes of the Dynkin diagram are $\alpha_1, \alpha_2, \alpha_3$ and α_4 with α_1 and α_2 long, ρ interchanges α_1 and α_4 and interchanges α_2 and α_3 . $p = 2$ and $q(\alpha)$ is as in case V. G_σ is ${}^2F_4(2^f)$.

VII. G is of type G_2 , ρ interchanges long and short roots, $p = 3$, and $q(\alpha) = 3^f$ if α is long, while $q(\alpha) = 3^{f+1}$ if α is short. G_σ is ${}^2G_2(3^f)$.

2. Arithmetic results. In the next section, we will show that certain sufficient conditions for the restriction map $H^n(G, V) \rightarrow H^n(G_\sigma, V)$ to be an injection or an isomorphism reduce to conditions on particular systems of equations. Here we establish some results concerning these systems of equations. Our arguments are based on Theorem 1 below, which follows from Lemma 6.4 and the discussion at the beginning of §6 of [2].

THEOREM 1. *Let α, β, x and d be integers. Suppose*

$$\sum_{j=1}^{\alpha} p^j = p^\alpha x + d(p^\beta - 1)$$

with $\alpha \leq n$ and $0 \leq j < \beta$ for each j . Let $[\]$ denote the greatest integer function. If $\alpha \geq [(n - 1)/(p - 1)]$ and $\beta \geq [\log_p(|x| + 1)] + \alpha + 2$, then $d = 0$.

THEOREM 2. *Let e, f, x, y, c, d, d' and n be integers with $0 \leq e$ and $0 \leq c < p^{2f} - 1$. Suppose*

$$\sum_{j=1}^{\alpha} p^j = p^e x + p^f c + d(p^{2f} - 1)$$

and

$$\sum_{l=1}^{\alpha'} p^{k_l} = p^e y - c + d'(p^{2f} - 1)$$

where $0 \leq \alpha, \alpha' \leq n$ and $0 \leq j, k_l < f$ for each j, l .

If $e \geq [(2n - 1)/(p - 1)]$ and $f \geq [\log_p(\max(|x|, |y|) + 1)] + e + 2$, then $c = d = d' = 0$.

PROOF. Multiplying the second equation by p^f and adding gives

$$\sum_{j=1}^a p^j + p^f \sum_{l=1}^{a'} p^{k_l} = p^3(x + p^f y) + (d + p^f d')(p^{2f} - 1).$$

The hypotheses of the theorem imply that $2f \geq [\log_p(|x + p^f y| + 1)] + e + 2$ and $e \geq [(a + a' - 1)/(p - 1)]$. Theorem 1 then gives $d + p^f d' = 0$. Multiplying the resulting equation by p^f and rewriting, we have

$$p^f \sum_{j=1}^a p^j + \sum_{l=1}^{a'} p^{k_l} = p^e(p^f x + y) + \left(p^e y - \sum_{l=1}^{a'} p^{k_l} \right) (p^{2f} - 1).$$

Another application of Theorem 1 yields $p^e y - \sum_{l=1}^{a'} p^{k_l} = 0$ from which it follows that $c = d = d' = 0$.

THEOREM 3. *Let e, f, x, y, c, d, d' and n be integers with $0 < e$ and $0 < c < p^{2f+1} - 1$. Suppose*

$$\sum_{j=1}^a p^j = p^e x + p^f c + d(p^{2f+1} - 1)$$

and

$$\sum_{l=1}^{a'} p^{k_l} = p^e y - c + d'(p^{2f+1} - 1),$$

where $0 < a, a' \leq n$ and $0 < i_j, k_l < f$ for each j, l . If $e \geq [(2n - 1)/(p - 1)]$ and $f \geq [\log_p(\max(|x|, |y|) + 1)] + e + 2$, then $c = d = d' = 0$ and $i_j, k_l < f$ for each j, l .

PROOF. Arguing as before, we apply Theorem 1 with $\beta = 2f + 1$ to obtain $\sum_{j=1}^a p^j + p^f \sum_{l=1}^{a'} p^{k_l} = p^e x + p^{e+f} y$. The hypothesis on f implies that the right side is less than p^{2f} , so each $k_l < f$. Then after multiplying by p^f and rewriting as in the previous proof, we may apply Theorem 1 with $\beta = 2f$ to conclude that $\sum p^{k_l} = p^e y$. Then $\sum p^j = p^e x$, so $c = d = d' = 0$, and, since $|p^e x| < p^f$, each $i_j < f$.

THEOREM 4. *Let $e, f, x, y, z, c, c', d, d', d''$ and n be integers with $0 < e$ and $0 < c, c' < p^{3f} - 1$. Suppose*

$$\sum_{j=1}^a p^j = p^e x + p^f c + p^{2f} c' + d(p^{3f} - 1),$$

$$\sum_{l=1}^{a'} p^{k_l} = p^e y - c + d'(p^{3f} - 1),$$

and

$$\sum_{s=1}^{a''} p^{r_s} = p^e z - c' + d''(p^{3f} - 1),$$

where $0 < a, a', a'' \leq n$ and $0 < i_j, k_l, r_s < f$ for each j, l, s . Then if $e \geq [(3n - 1)/(p - 1)]$ and $f \geq [\log_p(\max(|x|, |y|, |z|) + 1)] + e + 2$, $c = c' = d = d' = d'' = 0$.

PROOF. Combining the three equations gives

$$\begin{aligned} & \sum_{j=1}^a p^j + p^f \sum_{l=1}^{a'} p^{k_l} + p^{2f} \sum_{s=1}^{a''} p^{r_s} \\ & = p^e(x + p^f y + p^{2f} z) + (d + p^f d' + p^{2f} d'')(p^{3f} - 1). \end{aligned}$$

Theorem 1 implies that $d + p^f d' + p^{2f} d'' = 0$. Multiplying by p^f , we have

$$p^f \sum_{j=1}^a p^j + p^{2f} \sum_{l=1}^{a'} p^{k_l} + \sum_{s=1}^{a''} p^{r_s} = p^e(p^f x + p^{2f} y + z) + \left(p^e z - \sum_{s=1}^{a''} p^{r_s} \right) (p^{3f} - 1).$$

This time Theorem 1 gives $p^e z = \sum p^{r_s}$, so $c' = d'' = 0$. Again multiplying by p^f and applying Theorem 1 yields $p^e y = \sum p^{k_l}$ and $p^e x = \sum p^j$, which completes the proof.

3. The main theorem. Let G be a simple algebraic group defined and split over k_0 and let σ be a surjective rational endomorphism of G having finite fixed-point set G_σ . Fix a σ -stable Borel subgroup B and maximal torus $T \leq B$.

Let U be the unipotent subgroup of B and let P_1, \dots, P_l be the orbits of ρ on Σ^+ . Renumbering the P_i if necessary, we can find a σ -stable central series $U = U_1 > \dots > U_l > U_{l+1} = 1$ with $\bar{U}_i = U_i / U_{i+1} = \prod_{\alpha \in P_i} U_\alpha$ [1]. T_σ acts on the factor group $(U_i)_\sigma / (U_{i+1})_\sigma$ of the central series $U_\sigma = (U_1)_\sigma > \dots > (U_l)_\sigma > 1$ with weight $\alpha_i|_{T_\sigma}$. We will write $(\bar{U}_i)_\sigma$ for $(U_i)_\sigma / (U_{i+1})_\sigma$.

Let V be a rational G -module. By Theorem 2.1 of [2], $H^n(G, V) \cong H^n(B, V)$. We are interested in conditions which insure that the restriction map $H^n(G, V) \rightarrow H^n(G_\sigma, V)$ is an isomorphism or an injection. Since the index of B_σ in G_σ is prime to p , restriction from $H^n(G_\sigma, V)$ to $H^n(B_\sigma, V)$ is always injective, and it is then easy to see that if restriction from $H^n(B, V)$ to $H^n(B_\sigma, V)$ is an isomorphism (or an injection), so is restriction from $H^n(G, V)$ to $H^n(G_\sigma, V)$.

By the spectral sequence argument of [2, §5], restriction $H^n(B, V) \rightarrow H^n(B_\sigma, V)$ will be an isomorphism for $n \leq m$ and an injection for $n = m + 1$, if, for each weight λ in V , restriction induces an isomorphism

$$\begin{aligned} & (H^{s_1}(\bar{U}_1, k) \otimes \dots \otimes H^{s_l}(\bar{U}_l, k))_{-\lambda} \\ (3.1) \quad & \rightarrow (H^{s_1}((\bar{U}_1)_\sigma, k) \otimes \dots \otimes H^{s_l}((\bar{U}_l)_\sigma, k))_{(-\lambda|_{T_\sigma})} \end{aligned}$$

whenever $s_1 + \dots + s_l \leq m$.

The cohomology ring $H^*(\bar{U}_i, k)$ is isomorphic to $\otimes_{\alpha \in P_i} H^*(U_\alpha, k)$. Let V_α be the k -vector space with basis $a(-p^i \alpha)$, $i = 0, 1, 2, \dots$, and let T act on V_α by $ta(\mu) = \mu(t)a(\mu)$. Similarly let W_α be the k -vector space with basis $b(-p^i \alpha)$, $i = 1, 2, \dots$, and T -action given by $tb(\mu) = \mu(t)b(\mu)$. For a vector space V , let $S(V)$ and $\Lambda(V)$ denote the symmetric and exterior algebras of V respectively. Then [2, Theorem 4.1] as graded T -algebras

$$H^*(U_\alpha, k) \simeq \begin{cases} S(V_\alpha) & \text{if } p = 2, \\ \Lambda(V_\alpha) \otimes S(W_\alpha) & \text{if } p \neq 2, \end{cases}$$

where V_α has degree 1 and W_α has degree 2.

Choose a representative $\alpha_i \in P_i$ for each orbit, taking α_i long if there are two root lengths, and put $q_i = \prod_{\alpha \in P_i} q(\alpha)$. Let $V_\alpha(q_i)$ be the k -vector space with basis $\bar{a}(-p^j\alpha_i), j = 0, 1, \dots, \log_p q_i - 1$, and let $W_\alpha(q_i)$ be the k -vector space with basis $\bar{b}(-p^j\alpha_i), j = 1, 2, \dots, \log_p q_i$. We define T_σ -actions on $V_\alpha(q_i)$ and $W_\alpha(q_i)$ in the same way as the T -actions on V_α and W_α above. Exactly as in [2, Theorem 4.1], we see that, as graded T_σ -algebras

$$H^*((\bar{U}_i)_\sigma, k) \simeq \begin{cases} S(V_\alpha(q_i)) & \text{if } p = 2, \\ \Lambda(V_\alpha(q_i)) \otimes S(W_\alpha(q_i)) & \text{if } p \neq 2, \end{cases}$$

where $V_\alpha(q_i)$ has degree 1 and $W_\alpha(q_i)$ has degree 2.

Let $f = f(\sigma) = \min_{\alpha \in \Sigma^+} \log_p q(\alpha)$, as in §1. The proof of the preceding isomorphism shows that, for $p^j < p^f$, the restriction map sends $a(-p^j\alpha_i)$ to $\bar{a}(-p^j\alpha_i)$. If $\rho\alpha_i \neq \alpha_i$, the restriction of $a(-p^j\rho\alpha_i)$ is $\bar{a}(-p^{j+f}\alpha_i)$, and, if $\rho^2\alpha_i \neq \alpha_i$, the restriction of $a(-p^j\rho^2\alpha_i)$ is $\bar{a}(-p^{j+2f}\alpha_i)$. Similarly, if $1 < p^j \leq p^f$, $b(-p^j\alpha_i)$ restricts to $\bar{b}(-p^j\alpha_i)$, when $\rho\alpha_i \neq \alpha_i$ $b(-p^j\rho\alpha_i)$ restricts to $\bar{b}(-p^{j+f}\alpha_i)$, and when $\rho^2\alpha_i \neq \alpha_i$ $b(-p^j\rho^2\alpha_i)$ restricts to $\bar{b}(-p^{j+2f}\alpha_i)$.

Then we have the following sufficient conditions for the map of (3.1) to be an injection or an isomorphism.

Injectivity condition for $p = 2$. In every equation $\sum_{j=1}^n p^j\alpha_j = \lambda$, with $0 \leq i_j$ and $\alpha_j \in \Sigma^+$ for each j , we have each $i_j < f$.

Isomorphism condition for $p = 2$. If $\sum_{j=1}^n p^j\alpha_j|_{T_\sigma} = \lambda|_{T_\sigma}$ with $0 \leq i_j < f$ ($0 \leq i_j < f$ if there are two root lengths) and $\alpha_j \in \Sigma^+$ for each j , then $\sum_{j=1}^n p^j\alpha_j = \lambda$.

Injectivity condition for $p \neq 2$. In every equation $\sum_{j=1}^{n_1} p^j\alpha_j + \sum_{l=1}^{n_2} p^{k_l}\alpha_l = \lambda$, with $0 \leq i_j, 1 \leq k_l, \alpha_j, \alpha_l \in \Sigma^+$ for each j and l , and $n_1 + 2n_2 = n$, we have $i_j < f$ and $k_l < f$ for each j and l .

Isomorphism condition for $p \neq 2$. If $\sum_{j=1}^{n_1} p^j\alpha_j|_{T_\sigma} + \sum_{l=1}^{n_2} p^{k_l}\alpha_l|_{T_\sigma} = \lambda|_{T_\sigma}$ with $0 \leq i_j < f, 0 < k_l \leq f$ ($0 \leq i_j < f, 0 < k_l \leq f + 1$ if there are two root lengths), $\alpha_j, \alpha_l \in \Sigma^+$ for each j and l , and $n_1 + 2n_2 = n$, the $\sum_{j=1}^{n_1} p^j\alpha_j + \sum_{l=1}^{n_2} p^{k_l}\alpha_l = \lambda$.

Let $\omega = \sum_{\delta \in \Delta} n_\delta \delta$ be the maximal root and put $b = \max n_\delta$. For ζ in the root lattice Q , write $\zeta = \sum_{\delta \in \Delta} m_\delta \delta$ and put $b(\zeta) = \max |m_\delta|$. For λ in the weight lattice Λ , let $t(\lambda)$ be the order of λ in Λ/Q , with $t_p(\lambda)$ the p -part of $t(\lambda)$, and let t be the exponent of Λ/Q . We write $\bar{\lambda} = t\lambda$.

For integers b, n and e , let

$$e_\sigma(n) = \begin{cases} [(n - 1)/(p - 1)] & \text{if } \rho = 1, \\ [(2n - 1)/(p - 1)] & \text{if } \rho \neq 1 \text{ and } G_\sigma \text{ is not } {}^3D_4, \\ [(3n - 1)/(p - 1)] & \text{if } G_\sigma \text{ is } {}^3D_4 \end{cases}$$

and $f(b, e) = [\log_p(|b| + 1)] + e + 2$.

THEOREM 5. *Let V be a finite dimensional rational G -module and m a nonnegative integer. Let e be a nonnegative integer with $e \geq e_\sigma(mbt)$ and suppose $f = f(\sigma) \geq f(b(\bar{\lambda}), e)$ for all weights λ in V . If $p \neq 2$, assume further that*

$$e \geq (m + p - 1)bt_p(\lambda)/(p - 1).$$

Then the restriction map $H^n(G, V(e)) \rightarrow H^n(G_\sigma, V(e))$ is an isomorphism for $n < m$ and injection for $n = m + 1$.

PROOF. First assume $p = 2$.

To check injectivity, assume $\sum_{j=1}^{m+1} p^j \alpha_j = p^e \lambda$. If some $i_j > f$, multiply both sides of the equation by t . Examining the coefficients of α_i on both sides, we get $tp^f \leq p^e b(\bar{\lambda}) < p^f$, a contradiction.

To check the isomorphism condition assume that $\sum_{j=1}^n p^j \alpha_j|_{T_\sigma} = p^e \lambda|_{T_\sigma}$, with $0 \leq i_j < f$ ($0 \leq i_j \leq f$ if there are two root lengths) and $\alpha_i \in \Sigma^+$ for each j . Multiplying both sides by t , we have $\sum_{v=1}^n p^{u_v} \alpha_v|_{T_\sigma} = p^e \bar{\lambda}|_{T_\sigma}$ where $0 \leq u_v < f$ ($0 \leq u_v \leq f$ if there are two roots lengths) and $\alpha_v \in \Sigma^+$ for each v .

Consider the map on the root lattice given by restricting a weight from T to T_σ . t times the kernel of this map is contained in the group generated by the elements $p^f \alpha_i - \rho \alpha_i$ (which may be $(p^f - 1)\alpha_i$), and $p^{2f} \alpha_i - p^2 \alpha_i$ if $G_\sigma = {}^3D_4$ for the orbit representatives α_i in Δ , and the elements $p^{2f} \delta_i - \delta_i$ if G_σ is ${}^2A_n, {}^2D_n, {}^2E_6, p^{3f} \delta_i - \delta_i$ if G_σ is 3D_4 , and $p^{2f+1} \delta_i - \delta_i$ if G_σ is 2C_2 or 2F_4 , where δ_i ranges over Δ (cf. [3, §4a]).

Thus $\sum_{v=1}^n p^{u_v} \alpha_v$ and $p^e \bar{\lambda}$ differ by an element of this group. Considering the coefficients of elements of a ρ -orbit on Δ leads to systems of equations of the forms discussed §2, and the hypotheses of the theorem imply that the appropriate conditions on e and f in the theorems of that section are satisfied. Then the results of §2 imply that $\sum_{v=1}^n p^{u_v} \alpha_v = p^e \bar{\lambda}$, so $\sum_{j=1}^n p^j \alpha_j = p^e \lambda$.

Now suppose p is odd. The injectivity condition is proved exactly as when $p = 2$. For the isomorphism condition, suppose $\sum_{j=1}^{n_1} p^j \alpha_j|_{T_\sigma} + \sum_{l=2}^{n_2} p^k \alpha_l|_{T_\sigma} = p^e \lambda|_{T_\sigma}$ with $0 \leq i_j < f$ ($0 \leq i_j \leq f$ if there are two root lengths), $0 < k_l \leq f$ ($0 < k_l \leq f + 1$ if there are two root lengths) and $\alpha_j, \alpha_l \in \Sigma^+$ for each j and l , and $n_1 + 2n_2 \leq m$. We want to reduce this to the arithmetic problems of §2, so we have to eliminate those $p^{k_l} = f$ ($= f + 1$ in the case of two root lengths). But $p^f \alpha|_{T_\sigma} = (\rho \alpha)|_{T_\sigma}$ if Σ has one root length and $p^{f+1} \alpha|_{T_\sigma} = p(\rho \alpha)|_{T_\sigma}$ if Σ has two root lengths and α is long, while $p^{f+1} \alpha|_{T_\sigma} = (\rho \alpha)|_{T_\sigma}$ if there are two root lengths α is short.

Making the appropriate replacements of $p^{k_l} \alpha_l$ by terms with the same restriction to T_σ we get $\sum_{j=1}^{n_1} p^j \alpha_j|_{T_\sigma} + \sum_{l=2}^{n_2} p^{k_l} \alpha'_l|_{T_\sigma} = p^e \lambda|_{T_\sigma}$ with $0 \leq k'_l < f$ ($0 \leq k'_l < f + 1$ if there are two root lengths). The argument in the $p = 2$ case shows that $\sum_{j=1}^{n_1} p^j \alpha_j + \sum_{l=2}^{n_2} p^{k_l} \alpha'_l = p^e \lambda$.

Now suppose $\alpha'_l \neq \alpha_l$ for some l , and multiply the preceding equation by $t_p(\lambda)$. Subtract a simple root δ appearing in α'_l from both sides, and consider the coefficients of δ . On the left side we have at most $(m - 1)bt_p(\lambda)$ terms of the form $p^k \delta$, and one term of the form $t_p(\lambda)m_\delta(\alpha'_l) - 1$ if there is one root length or there are two root lengths and α'_l is long, or of the form $t_p(\lambda)pm_\delta(\alpha'_l) - 1$ if α'_l is short. On the right side, the coefficient of δ is $p^e m_\delta(t_p(\lambda)\lambda) - 1$, where $m_\delta(t_p(\lambda)\lambda) > 1$.

Then the sum of the p -adic digits of the coefficient of δ on the left is at most $(m - 1)bt_p(\lambda) + pt_p(\lambda)b - 1 = (m + p - 1)bt_p(\lambda) - 1$. The sum of the p -adic digits of the coefficient on the right is at least $e(p - 1) \geq (m + p - 1)bt_p(\lambda)$, which is a contradiction.

Thus $\alpha'_l = \alpha_l$ for all l , and the proof is complete.

The theorem says that if e is sufficiently large (with respect to n) and $f(\sigma)$ is sufficiently large (with respect to V and e), then $H^n(G, V(e)) \simeq H^n(G_\sigma, V(e))$. Since $H^n(G_\sigma, V) \simeq H^n(G_\sigma, V(e))$, this implies that for large values of $f(\sigma)$, $H^n(G_\sigma, v)$ has a stable or generic value $H^n_{\text{gen}}(G, V)$.

It is also natural to consider twisting the module V by σ . If $r: G \rightarrow GL(V)$ is the morphism of algebraic groups associated with V , let $V(\sigma)$ denote the module with associated morphism $r \circ \sigma$. Since the weights of T in $V(\sigma)$ are all divisible by $p^{f(\sigma)}$, the arguments above imply that, for $f(\sigma)$ sufficiently large with respect to n , $H^n(G, V(\sigma)) \simeq H_{\text{gen}}^n(G, V)$. Thus, the generic cohomology group $H_{\text{gen}}^n(G, V)$ may be computed as $H^n(G_\sigma, V)$ if $f(\sigma)$ is sufficiently large with respect to n and V , or as $H^n(G, V(\sigma))$ if $f(\sigma)$ is sufficiently large with respect to n .

REFERENCES

1. G. S. Avrunin, *A vanishing theorem for second degree cohomology*, J. Algebra **53** (1978), 382–388.
2. E. Cline, B. Parshall, L. Scott and W. van der Kallen, *Rational and generic cohomology*, Invent. Math. **39** (1977), 143–169.
3. W. Jones and B. Parshall, *On the 1-cohomology of finite groups of Lie type*, Proc. Conf. Finite Groups (W. R. Scott and F. Gross, eds.), Academic Press, New York, 1976.
4. R. Steinberg, *Endomorphisms of linear algebraic groups*, Mem. Amer. Math. Soc., no. 80, Amer. Math. Soc., Providence, R. I., 1968.
5. _____, *Lectures on Chevalley groups*, Yale University notes, 1968.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MASSACHUSETTS, AMHERST, MASSACHUSETTS 01003 (Current address)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VIRGINIA 22903