

THE HOMOLOGY AND HIGHER REPRESENTATIONS OF THE AUTOMORPHISM GROUP OF A RIEMANN SURFACE

S. A. BROUGHTON

ABSTRACT. The representations of the automorphism group of a compact Riemann surface on the first homology group and the spaces of q -differentials are decomposed into irreducibles. As an application it is shown that M_{24} is not a Hurwitz group.

1. Introduction. Let G be a finite group of orientation-preserving homeomorphisms of a Riemann surface S of genus $\sigma \geq 2$. We then have a representation of G on the first homology group $H_1(S) = H_1(S, \mathbb{C})$. If S has a conformal structure which is preserved under the G -action, then there are also representations of G on the various spaces of q -differentials $\mathcal{H}^q(S)$ ($\mathcal{H}^q(S)$ = holomorphic sections of $T^*(S) \otimes \cdots \otimes T^*(S)$ (q times), $T^*(S)$ = cotangent bundle). In this note we give formulae (Propositions 1–2) for the decompositions of these representations into irreducibles.

The decompositions for $H_1(S) \simeq \mathcal{H}^1(S) \oplus \mathcal{H}^1(S)^*$ and $\mathcal{H}^2(S)$ may be applied to the study of surfaces of genus σ . From the decomposition of the homology representation it follows that the characters of G must satisfy certain inequalities (see (13) below). This is useful in showing that certain groups cannot occur as automorphism groups of a surface of a given genus σ . In [S] L. L. Scott has given a formula equivalent to (13), though derived by a purely group-theoretic argument.

The decompositions of $\mathcal{H}^2(S)$ may be used to locally describe the action of the Teichmüller modular group Mod_σ on Teichmüller space, \mathcal{T}_σ (see [R]). This was used by J. Lewittes [L] to compute the dimensions of the branch loci of the action of Mod_σ on \mathcal{T}_σ .

The decompositions are derived in §2 from the Eichler Trace Formula and the Lefschetz Fixed Point Formula, using a simple character theory argument. In §3 we give an application showing that the Mathieu group M_{24} is not a Hurwitz group.

2. The decomposition formulae and their derivations. First we recall some facts about actions of a finite group G on a surface S (cf. [H, T]). The space $T = S/G$ is a surface T of genus τ , and $\pi: S \rightarrow T$ is branched over $Q_1, \dots, Q_t \in T$ with branching orders n_1, \dots, n_t . Call $(\tau; n_1, \dots, n_t)$ the *branching data* of G (write (n_1, \dots, n_t) if $\tau = 0$). The Riemann-Hurwitz formula [FK, p. 243] gives

$$(1) \quad (2\sigma - 2)/|G| = 2\tau - 2 + \sum_{i=1}^t (1 - 1/n_i).$$

Received by the editors July 23, 1985 and, in revised form, March 21, 1986.
1980 *Mathematics Subject Classification* (1985 Revision). Primary 30F30.

©1987 American Mathematical Society
0002-9947/87 \$1.00 + \$.25 per page

We denote the right-hand side by κ . There are elements $a_1, \dots, a_\tau, b_1, \dots, b_\tau, c_1, \dots, c_t$, generating G , such that

$$(2) \quad \prod_{i=1}^\tau [a_i, b_i] \prod_{j=1}^t c_j = 1,$$

and

$$(3) \quad o(c_i) = n_i.$$

If $P \in S$ is a point fixed by $g \in G$, then the induced map of tangent spaces $dg^{-1}: T_P(S) \rightarrow T_P(S)$ is multiplication by an $o(g)$ th root of unity, denoted by $\varepsilon(P, g)$. It is easy to show that we may pick the c_i and $P_i \in \pi^{-1}(Q_i)$ such that $G_{P_i} = \{g \in G \mid gP_i = P_i\} = \langle c_i \rangle$ and

$$(4) \quad \varepsilon(P_i, c_i) = \exp(2\pi\sqrt{-1}/n_i).$$

Let $U_n \subseteq S^1$ be the group of n th roots of unity and let $\varphi_k: S^1 \rightarrow S^1$ be the character $z \rightarrow z^k, k \in \mathbf{Z}$. Let c_1, \dots, c_t be as defined above and let $\nu_i: \langle c_i \rangle \rightarrow U_{n_i}$ be the isomorphism defined by $c_i \rightarrow \exp(2\pi\sqrt{-1}/n_i)$. Let χ_0, \dots, χ_t be the irreducible characters of G with $\chi_0 =$ principal character. Each χ_j defines a character of U_{n_i} by means of the isomorphism ν_i . Define $m_i^k(\chi_j), 0 \leq k \leq n_i - 1$, by

$$(5) \quad \chi_j|_{U_{n_i}} = \sum_{k=0}^{n_i-1} m_i^k(\chi_j) \varphi_k|_{U_{n_i}}$$

and define $m_i^k(\chi_j)$ for all $k \in \mathbf{Z}$ by periodicity: $m_i^k(\chi_j) = m_i^{k+n_i}(\chi_j)$. Let $\text{ch}_{\mathcal{H}^q(S)}$ be the character of the representation of G on $\mathcal{H}^q(S)$, and write

$$\text{ch}_{\mathcal{H}^q(S)} = \mu_q^0 \chi_0 + \dots + \mu_q^t \chi_t.$$

Define the Poincaré series $P_{\chi_j}(z)$ by

$$P_{\chi_j}(z) = \sum_{q=0}^\infty \mu_q^j z^q.$$

We have the following propositions.

PROPOSITION 1. *Let G be a group of conformal automorphisms of a Riemann surface S of genus ≥ 2 and let all notation be as above. Then:*

- (i) $P_{\chi_0}(z) = 1 + z + zR_{\chi_0}(z)$,
- (ii) $P_{\chi_j} = zR_{\chi_j}(z), j \neq 0$, where
- (iii)

$$R_{\chi_j} = \frac{(1 - \tau)\chi_j(1)}{1 - z} + \frac{\kappa\chi_j(1)}{(1 - z)^2} - \sum_{i=1}^t \frac{1}{n_i} \cdot \frac{e_i^0(j) + e_i^1(j)z + \dots + e_i^{n_i-1}(j)z^{n_i-1}}{1 - z^{n_i}}$$

and

$$e_i^r(j) = \sum_{k=0}^{n_i-1} k \cdot m_i^{1+r+k}(\chi_j).$$

PROPOSITION 2. *Let G be a finite group of homeomorphisms of a Riemann surface S , $\text{ch}_{H_1(S)}$ the character of the homology representation, and other notation as above. Then the multiplicity of χ_j in $\text{ch}_{H_1(S)}, \langle \chi_j, \text{ch}_{H_1(S)} \rangle$, is given by*

- (i) $\langle \chi_0, \text{ch}_{H_1(S)} \rangle = 2\tau,$
- (ii)

$$\langle \chi_j, \text{ch}_{H_1(S)} \rangle = (2\tau - 2 + t)\chi_j(1) - \sum_{i=1}^t m_i^0(\chi_j), \quad j \neq 0.$$

Let ρ be the regular representation of G and ρ_i the permutation character determined by G acting on the coset space $G/\langle c_i \rangle$. Then (i) and (iii) may be rewritten:

- (iii)

$$\text{ch}_{H_1(S)} = 2\chi_0 + (2\tau - 2 + t)\rho - \sum_{i=1}^t \rho_i.$$

Before proving Propositions 1–2 we recall the Eichler Trace Formula and the Lefschetz Fixed Point Formula. Let $\eta: G \rightarrow \mathbf{Z}$ be the class function on G obtained by setting $\eta(g)$ equal to the negative of the Euler characteristic of the fixed point subset S^g of g , i.e.

$$\eta(1) = 2\sigma - 2, \quad \eta(g) = -|S^g|, \quad g \neq 1.$$

By the Lefschetz Fixed Point Formula,

$$(6) \quad \text{ch}_{H_1(S)}(g) = 2 + \eta(g), \quad g \in G.$$

Define $\lambda_q: G \rightarrow \mathbf{C}, q \geq 0$, as follows:

$$\begin{aligned} \lambda_0(g) &= 1, & g \in G, \\ \lambda_q(1) &= (\sigma - 1)(2q - 1), & q \geq 1, \\ \lambda_q(g) &= \sum_{P \in S^g} \frac{(\varepsilon(P, g))^q}{1 - \varepsilon(P, g)}, & q \geq 1, \end{aligned}$$

where the last sum is zero if S^g is empty. The Riemann-Roch Theorem and the Eichler Trace Formula state that the characters $\text{ch}_{\mathcal{H}^q(S)}$ are given by

$$(7) \quad \begin{aligned} \text{ch}_{\mathcal{H}^q(S)}(g) &= \lambda_q(g), & q \neq 1, \\ \text{ch}_{\mathcal{H}^1(S)}(g) &= 1 + \lambda_q(g). \end{aligned}$$

For proofs of (6)–(7) see [FK]. Observe [FK] that $\eta(g) = 2 \text{Re } \lambda_1(g)$.

Write

$$\eta = \eta^0 + \dots + \eta^t, \quad \lambda_q = \lambda_q^0 + \dots + \lambda_q^t,$$

where

$$\begin{aligned} \eta^0(1) &= 2\sigma - 2, & \eta^0(g) &= 0, & g \neq 1, \\ \eta^i(1) &= 0, & \eta^i(g) &= -|S^g \cap \pi^{-1}(Q_i)|, & i > 0, \\ \lambda_q^0(1) &= (\sigma - 1)(2q - 1), & \lambda_q^0(g) &= 0, & g \neq 1, q \geq 1, \\ \lambda_q^i(1) &= 0, & \lambda_q^i(g) &= \sum_{P \in S^g \cap \pi^{-1}(Q_i)} \frac{(\varepsilon(P, g))^q}{1 - \varepsilon(P, g)}, & i > 0, q \geq 1. \end{aligned}$$

For $1 \neq g \in G$, $S^g \cap \pi^{-1}(Q_i) \neq \emptyset$ if and only if the conjugacy class of g , $\text{Cl}(g)$, meets $\langle c_i \rangle$. Assume $g \in \langle c_i \rangle$, then since G_{P_i} is cyclic, $S^g \cap \pi^{-1}(Q_i)$ is in 1-1 correspondence with $N_G(\langle g \rangle) / \langle c_i \rangle$ by $h \rightarrow h \cdot P_i$. Furthermore, $N_G(\langle g \rangle) / \text{Cent}(g)$ is in 1-1 correspondence with $\text{Cl}(g) \cap \langle c_i \rangle$ by $h \rightarrow hgh^{-1}$. From (4) and the definition of ν_i , $\varepsilon(P_i, g) = \nu_i(g)$, $g \in \langle c_i \rangle$. It easily follows for $i > 0$ that

$$(8) \quad \lambda^i_q(g) = \frac{|\text{Cent}(g)|}{n_i} \cdot \sum_{h \in \text{Cl}(g) \cap \langle c_i \rangle} \frac{(\nu_i(h))^q}{1 - \nu_i(h)}.$$

Since λ^i_q is a class function, this holds for all $1 \neq g \in G$. Similarly, for $1 \neq g \in G$,

$$(9) \quad \eta^i(g) = -\frac{|\text{Cent}(g)|}{n_i} |\text{Cl}(g) \cap \langle c_i \rangle|.$$

We now give proofs of the decompositions, first Proposition 2. Let $1 = g_0, \dots, g_t$ be a set of representatives of conjugacy classes of G . For $j = 0, 1, \dots, t$:

$$(10) \quad \begin{aligned} \langle \eta, \chi_j \rangle &= \sum_{i=0}^t \langle \eta^i, \chi_j \rangle = \sum_{i=0}^t \frac{1}{|G|} \sum_{g \in G} \eta^i(g) \bar{\chi}_j(g) \\ &= \sum_{i=0}^t \sum_{k=0}^t \frac{\eta^i(g_k) \bar{\chi}_j(g_k)}{|\text{Cent}(g_k)|} \\ &= \frac{2\sigma - 2}{|G|} \chi_j(1) - \sum_{i=1}^t \frac{1}{n_i} \sum_{1 \neq g \in \langle c_i \rangle} \overline{\chi_j(g)}, \end{aligned}$$

from (9) above. By the Riemann-Hurwitz Formula (1), (10) may be rewritten as

$$(2\tau - 2 + t) \chi_j(1) - \sum_{i=1}^t \frac{1}{n_i} \sum_{g \in \langle c_i \rangle} \bar{\chi}_j(g) = (2\tau - 2 + t) \chi_j(1) - \sum_{i=1}^t m_i^0(\chi_j).$$

Since $\text{ch}_{H_1(S)} = 2\chi_0 + \eta$, (i) and (ii) of Proposition 2 follow immediately; (iii) follows from (i)-(ii) and Frobenius reciprocity.

Let $R_g(z) = \sum_{q=1}^\infty \lambda_q(g) z^{q-1}$. To prove Proposition 1 it suffices by (7) to prove

$$(11) \quad R_{\chi_j}(z) = \frac{1}{|G|} \sum_{g \in G} R_g(z) \overline{\chi_j(g)}.$$

Using (8) and arguing as above, the right-hand side of (11) equals

(12)

$$\begin{aligned} &\sum_{q=1}^\infty \frac{(\sigma - 1) \chi_j(1)}{|G|} (2q - 1) z^{q-1} + \sum_{i=1}^t \sum_{q=1}^\infty \sum_{1 \neq g \in \langle c_i \rangle} \frac{1}{n_i} \frac{(\nu_i(g))^q}{1 - \nu_i(g)} \bar{\chi}_j(g) z^{q-1} \\ &= \kappa \chi_j(1) (1 - z)^{-2} \frac{\kappa \chi_j(1) (1 - z)^{-1}}{2} \\ &\quad + \sum_{i=1}^t \sum_{r=0}^{n_i-1} \frac{1}{n_i} \sum_{1 \neq \omega \in U_{n_i}} \frac{\omega^{r+1}}{1 - \omega} \bar{\chi}_j(\omega) z^r (1 - z)^{-n_i}. \end{aligned}$$

We calculate

$$\begin{aligned} \sum_{1 \neq \omega \in U_n} \frac{\omega^s}{1 - \omega} \bar{\chi}(\omega) &= \lim_{x \rightarrow 1} \sum_{1 \neq \omega \in U_n} \frac{\omega^s}{1 - x\omega} \bar{\chi}(\omega) \\ &= \lim_{x \rightarrow 1} \left(\sum_{q=0}^{\infty} \sum_{\omega \in U_n} \omega^{q+s} \bar{\chi}(\omega) x^q - \sum_{q=0}^{\infty} \chi(1) x^q \right) \\ &= \lim_{x \rightarrow 1} \left(n \frac{L_s(x)}{1 - x^n} - \frac{\chi(1)}{1 - x} \right), \end{aligned}$$

where $L_s(x) = a_0 + a_1x + \dots + a_{n-1}x^{n-1}$ and $a_k = (1/n) \sum_{\omega \in U_n} \omega^{k+s} \bar{\chi}(\omega)$. The limit is easily calculated by l'Hôpital's rule and equals $(n - 1)\chi(1)/2 - L'_s(1)$. Setting $n = n_j$, $s = r + 1$, $\chi = \chi_j$, then $a_k = m_i^{1+k+r}(\chi_j)$ and (11) now follows easily from (12) and the definition of R_{χ_j} .

3. Application. If $\tau = 0$, then G is generated by c_1, \dots, c_t with $c_1 \cdot c_2 \cdot \dots \cdot c_t = 1$, and from (ii) of Proposition 2 it follows that for a nonprincipal character χ_j

$$(13) \quad (t - 2)\chi_j(1) \geq \sum_{i=1}^t m_i^0(\chi_j).$$

This is a reformulation of the inequality that L. L. Scott obtains in [S] by purely group theoretic means for arbitrary characteristic. The G -module he constructs on p. 475 of [S] may be identified with $H_1(S)$. The inequality (13) may sometimes be used as a "Brauer trick" to show that a given group cannot occur as the automorphism group of a surface of given genus.

As an example of this let us verify that the Mathieu group M_{24} is not a Hurwitz group. The group G is a Hurwitz group if it occurs as the automorphism group of a surface S of genus σ with $|G| = 84(\sigma - 1)$, Hurwitz' upper bound for the order of an automorphism group. If G acts on S as above then the branching data is $(2, 3, 7)$ and G has a generating $(2, 3, 7)$ -vector (c_1, c_2, c_3) . In Table 1 we have copied a portion of the character table of M_{24} [Fr, p. 346], giving, for selected characters, the character values of all elements of order 1, 2, 3, or 7. The classes are given in cycle notation, M_{24} being realized as a permutation group of degree 24.

TABLE 1

	1^{24}	$1^{82}8$	2^{12}	1^63^6	3^8	$1^37^3_+$	$1^37^3_-$
χ_1	45	-3	5	0	3	$(-1 + \sqrt{-7})/2$	$(-1 - \sqrt{-7})/2$
χ_2	252	28	12	9	0	0	0

For c_i chosen from the classes in Table 1 all the nonidentity elements of $\langle c_i \rangle$ are conjugate in M_{24} except for $\langle c_3 \rangle$, where half lie in $1^37^3_+$ and the other half lie in $1^37^3_-$. Since $\kappa = 1/42$, we obtain from (10), for any nonprincipal character χ of M_{24} ,

$$\frac{1}{42}(\chi(1) - 21\chi(c_1) - 28\chi(c_2) - 36 \operatorname{Re} \chi(c_3)) = \langle \eta, \chi \rangle \geq 0,$$

or

$$\chi(1) \geq 21\chi(c_1) + 28\chi(c_2) + 36 \operatorname{Re} \chi(c_3).$$

(This is equivalent to (13) but slightly more convenient.) There is no possible choice of c_1, c_2, c_3 for which this inequality holds for both the characters χ_1, χ_2 above. It is interesting to note that for $c_1 \in 2^{12}, c_2 \in 3^8, c_3 \in 1^3 7_+^3$, or $1^3 7_-^3$, χ_2 and its conjugate $\bar{\chi}_2$ are the only irreducible characters for which (13) fails, and that the standard Brauer trick [I, p. 70] applied to any pair of $\langle c_1 \rangle, \langle c_2 \rangle$, or $\langle c_3 \rangle$ fails.

REFERENCES

- [FK] H. M. Farkas and I. Kra, *Riemann surfaces*, Graduate Texts in Math., no. 71, Springer-Verlag, Berlin and New York, 1979.
- [Fr] F. G. Frobenius, *Gesammelte Abhandlungen*, Vol. 3, Springer-Verlag, Berlin and New York, 1968.
- [H] J. Harvey, *On branch loci in Teichmüller space*, Trans. Amer. Math. Soc. **153** (1971), 387–399.
- [I] I. M. Isaacs, *Character theory of finite groups*, Academic Press, New York, 1976.
- [L] J. Lewittes, *Invariant quadratic differentials*, Bull. Amer. Math. Soc. **68** (1962), 320–322.
- [R] H. E. Rauch, *A transcendental view of the space of algebraic Riemann surfaces*, Bull. Amer. Math. Soc. **21** (1965), 1–39.
- [S] L. L. Scott, *Matrices and cohomology*, Ann. of Math. (2) **105** (1977), 473–492.
- [T] T. Tucker, *Finite groups acting on surfaces and the genus of a group*. J. Combin. Theory Ser. B **34** (1983), 82–98.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN - MADISON, MADISON, WISCONSIN 53706

Current address: Department of Mathematics, Cleveland State University, Cleveland, Ohio 44115