

RINGS OF DIFFERENTIAL OPERATORS ON INVARIANT RINGS OF TORI

IAN M. MUSSON

ABSTRACT. Let k be an algebraically closed field of characteristic zero and G a torus acting diagonally on k^s . For a subset β of $s = \{1, 2, \dots, s\}$, set $U_\beta = \{u \in k^s \mid u_j \neq 0 \text{ if } j \in \beta\}$. Then G acts on $\mathcal{O}(U_\beta)$, the ring of regular functions on U_β , and we study the ring $D(\mathcal{O}(U_\beta)^G)$ of all differential operators on the invariant ring.

More generally suppose that Δ is a set of subsets of s , such that each invariant ring $\mathcal{O}(U_\beta)^G$, $\beta \in \Delta$, has the same quotient field. We prove that $\bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta)^G)$ is Noetherian and finitely generated as a k -algebra.

Now G acts on each $D(\mathcal{O}(U_\beta))$ and there is a natural map

$$\theta: \bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta))^G \rightarrow \bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta)^G) = D(Y_\Delta/G)$$

obtained by restriction of the differential operators. We find necessary and sufficient conditions for θ to be surjective and describe the kernel of θ .

The algebras $\bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta))^G$ and $\bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta)^G)$ carry a natural filtration given by the order of the differential operators. We show that the associated graded rings are finitely generated commutative algebras and are Gorenstein rings. We also determine the centers of $\bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta))^G$ and $\bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta)^G)$.

Introduction. Throughout this paper k will be an algebraically closed field of characteristic zero. If K is a commutative k -algebra we denote by $D_0(K)$ the set of k -linear maps $K \rightarrow K$ and if $p \geq 1$ a k -linear map $f: K \rightarrow K$ belongs to $D_p(K)$ provided the map $[f, r]$ defined by $[f, r](s) = f(rs) - rf(s)$ for $s \in K$ belongs to $D_{p-1}(K)$ for all $r \in K$. The set $D(K) = \bigcup_{p \geq 0} D_p(K)$ forms a subring of $\text{End}_k(K)$ called the *ring of differential operators on K* . We consider K as a left $D(K)$ -module where $f \cdot r = f(r)$ for all $f \in D(K)$, $r \in K$.

The ring of differential operators on the invariant ring of a finite group acting on a polynomial ring has been studied in [Ka, L1 and L2]. We carry out a similar study for the invariant ring of a torus.

We assume that the torus G acts on k^s as a group of diagonal matrices. If β is a subset of $s = \{1, 2, \dots, s\}$, then G acts on $U_\beta = \{u \in k^s \mid u_j \neq 0 \text{ if } j \in \beta\}$. Also G acts on $\mathcal{O}(U_\beta)$ via $(gf)(u) = f(g^{-1}u)$ for $g \in G$, $f \in \mathcal{O}(U_\beta)$, $u \in U_\beta$. More generally suppose that for some set of subsets Δ of s we have $Y = \bigcup_{\beta \in \Delta} U_\beta$, where each invariant ring $\mathcal{O}(U_\beta)^G$ has the same quotient field F . Then each $D(\mathcal{O}(U_\beta)^G)$ is a

Received by the editors June 18, 1986 and, in revised form, December 11, 1986.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 16A33; Secondary 13N05, 14L30.

©1987 American Mathematical Society
 0002-9947/87 \$1.00 + \$.25 per page

subring of $D(F)$ and we define

$$D(Y_\Delta/G) = \bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta)^G).$$

THEOREM A. *With the above notation $D(Y_\Delta/G)$ is Noetherian and finitely generated as a k -algebra.*

If Δ consists of a single subset $\Delta = \{\beta\}$ then $D(Y_\Delta/G) = D(\mathcal{O}(U_\beta)^G)$. However the ring $\mathcal{O}(U_\beta)^G$ does not in general reflect the geometry of the orbit space U_β/G very well. A simple example of this is where $G = \text{GL}(1)$ acts on $U_\phi = k^s$ by scalar multiplication. Then the origin O is the only closed orbit, and $(U_\phi - O)/G \cong \mathbf{P}^{s-1}$ but $\mathcal{O}(U_\phi)^G = k$.

However, there are many nonconstant global differential operators on \mathbf{P}^{s-1} ; see [BB, Example 3.10a]. The definition of $D(Y_\Delta/G)$ given above is designed to accommodate such examples.

We shall write $D(U_\beta)$ and $D(Y)$ in place of $D(\mathcal{O}(U_\beta))$ and $D(\mathcal{O}(Y))$.

The group G acts on $D(U_\beta)$ via $(g \cdot d)(f) = g(d(g^{-1}f))$ for $f \in \mathcal{O}(U_\beta)$, $d \in D(U_\beta)$, and $g \in G$. Moreover there is a natural map $\theta_\beta: D(U_\beta)^G \rightarrow D(\mathcal{O}(U_\beta)^G)$ which is obtained by restriction of the differential operators. Taking intersections over $\beta \in \Delta$, we obtain a map

$$\theta: D(Y)^G = \bigcap_{\beta \in \Delta} D(U_\beta)^G \rightarrow \bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta)^G) = D(Y_\Delta/G).$$

In Theorem B below we obtain necessary and sufficient conditions for θ to be surjective and describe the kernel of θ . The corresponding result for the action of a finite group on a polynomial ring is proved in [Ka, Chapter III, Théorème 4 and L1, Théorème 5].

The statement of Theorem B is given at the beginning of §2. It is formulated in terms of certain convex rational polyhedral cones which are related to the invariant rings of a torus. For other results on invariant rings of tori which use this approach see [Ho and Ke].

This paper is organized as follows. In §1 we assemble the results we need on rings of invariants of tori acting on $\mathcal{O}(U_\beta)$ and convex cones. Theorem B is proved in §2. In §3 we describe the centers of $D(Y)^G$ and $D(Y_\Delta/G)$ (see Theorem C and Corollary 3.4) and in Corollary 3.6, obtain an important special case of Theorem B. In §4 we filter the rings $D(Y)^G$ and $D(Y_\Delta/G)$ by the order of the differential operators, and show that the associated graded rings are finitely generated commutative algebras. Theorem A follows easily from this. We also show that these graded rings are Gorenstein rings (Theorem D). This allows us to deduce homological information about modules over $D(Y)^G$ and $D(Y_\Delta/G)$; see [L2, Théorème 4.4].

In our companion paper [M] we give some examples and indicate some applications to enveloping algebras of some semisimple Lie algebras.

Index of notation. \mathbf{Z}_+ , \mathbf{Q}_+ refer to the set of nonnegative integers and rational numbers, respectively.

Other symbols used frequently in the text are given below in order of appearance.

Introduction. $D_p(K)$, $D(K)$, s , U_β , Δ , $Y = \bigcup_{\beta \in \Delta} U_\beta$, $D(Y)^G$, $D(Y_\Delta/G)$, θ_β , θ .

1.1. H_f , H_f^0

1.3. $\mathbf{Q}\Lambda$, ϕ , ψ , ψ^* , x_i , f_i

1.4. T_i , m_i , Λ_β , V , H_i , Γ , M , N

2.0. \mathcal{H}_i , α , J

2.2. Q_i , P_i , A , $I(\lambda)$

2.3. $A_{\mathbf{Q}}$, $\Omega(\lambda)$

2.4. Σ

2.5. Λ_b , $\overline{\Omega(\lambda)}$, $\overline{\Omega_{\beta,i}(\lambda)}$, $S_{\lambda,\beta,i}$, S_λ

2.6. t_i , q_i , p_i , B , $B_{\mathbf{Q}}$, $\omega_\beta(\lambda)$, $\omega_{\beta,i}(\lambda)$

2.8. I

3.1. $Z(R)$, $Q(R)$, $GK(R)$

3.5. I_i

4.1. \mathcal{E}_i , \mathcal{D}_i , $\text{Gr}(R)$, E_i , D_i , Π_1 , Π_2

4.9. $\text{Kd}(M)$, $d(M)$, $\text{grade}(M)$

4.10. $\text{tr deg}_k(R)$

1. Convex cones and rings of invariants of tori.

1.1. If $f \in (\mathbf{Q}^n)^*$ a subset of the form $H_f = \{\lambda \in \mathbf{Q}^n \mid f(\lambda) \geq 0\}$ is called a *halfspace* in \mathbf{Q}^n . We sometimes write H_f^0 for $\ker f$. Given a finite set $S = \{v_1, \dots, v_s\}$ in \mathbf{Q}^n , the *convex rational polyhedral cone*, $\text{cone}(S)$ spanned by S is the set

$$\text{cone}(S) = \sum_{i=1}^s \mathbf{Q}^+ v_i.$$

THEOREM. (a) *If S is a finite subset of \mathbf{Q}^n , then $\text{cone}(S)$ is a finite intersection of halfspaces. Moreover, if S spans \mathbf{Q}^n , then $\text{cone}(S) = \bigcap_{i \in I} H_i$ where the halfspaces H_i , $i \in I$, are determined as those halfspaces H such that H^0 contains $n - 1$ linearly independent elements of S .*

(b) *Conversely any finite intersection of halfspaces in \mathbf{Q}^n has the form $\text{cone}(S)$ for some finite set S .*

PROOF. This is the analogue for cones of the fact that a nonempty subset of \mathbf{Q}^n is a convex polytope if and only if it is a bounded polyhedral set [B, Theorem 9.2].

When S spans \mathbf{Q}^n we call the halfspaces H_i of Theorem 1.1(a) the *bounding halfspaces* of $\text{cone}(S)$. In the terminology of [B] these are the supporting halfspaces which intersect $\text{cone}(S)$ in a facet. If H is a bounding halfspace of $\text{cone}(S)$ then H^0 will be called a *bounding hyperplane*.

1.2. **LEMMA.** *Suppose $C = H_1 \cap H_2 \cap \dots \cap H_r$ is an intersection of halfspaces in \mathbf{Q}^n and that C spans \mathbf{Q}^n . Then any bounding halfspace of C is equal to some H_i .*

PROOF. Suppose $H = \{v \in \mathbf{Q}^n \mid f(v) \geq 0\}$ is a bounding halfspace different from each $H_i = \{v \in \mathbf{Q}^n \mid f_i(v) \geq 0\}$. Since $H^0 \cap C$ contains $n - 1$ linearly independent elements, there exists $v \in H^0 \cap C$ such that $v \notin H_i^0$. Adding the elements obtained in this way we obtain u with $f(u) = 0$ and $f_i(u) > 0$ for $i = 1, \dots, r$. The idea now is to perturb u slightly to obtain an element u' of C which does not lie in H .

There exists $w \in \mathbf{Q}^n$ such that $f(w) = -1$. Let $f_i(u) = \varepsilon_i > 0$ and $f_i(w) = \delta_i$. If $\delta_i \geq 0$ for $1 \leq i \leq r$, then $u' = u + w$ is the desired element. If $\delta_i < 0$ for some i , let $\lambda = \min\{-\varepsilon_i/\delta_i \mid 1 \leq i \leq r \text{ and } \delta_i < 0\}$, so $\lambda > 0$.

We set $u' = u + \lambda w$. Then it is easily checked that $f_i(u') \geq 0$ for $1 \leq i \leq r$, so $u' \in C$, but $f(u') = -\lambda < 0$ so $u' \notin H$ and H is not a bounding halfspace of C .

1.3. We next characterize certain semigroups which we shall be interested in. An additive abelian semigroup Λ is *normal* if it is finitely generated, cancellative, and whenever $p_1, p_2, p_3 \in \Lambda$ with $p_1 + np_2 = np_3$ for a positive integer n , there exists $p_4 \in \Lambda$ with $p_1 = np_4$. In this situation Λ can be embedded in a finitely generated abelian group ($\mathbf{Z}\Lambda$ for example). However the definition of normality is independent of the embedding. For Λ a subsemigroup of \mathbf{Z}^n , we write $k\Lambda = \text{span}\{t^\lambda \mid \lambda \in \mathbf{Z}^n\}$.

PROPOSITION. *For a semigroup Λ the following are equivalent:*

- (1) Λ is normal.
- (2) For any field k , $k\Lambda$ is an integrally closed Noetherian domain.
- (3) For some $s \geq t \geq 0$, Λ is isomorphic to a semigroup of the form $(\mathbf{Z}'_+ \times \mathbf{Z}^{s-t}) \cap V$, where V is a subspace of \mathbf{Q}^s and under this isomorphism $\dim \mathbf{Q}\Lambda = n$.
- (4) For some $t, n \geq 0$, Λ is isomorphic to a semigroup of the form $\mathbf{Z}^n \cap H_1 \cap \dots \cap H_t$ where the H_i are halfspaces in \mathbf{Q}^n , and under this isomorphism $\mathbf{Z}\Lambda = \mathbf{Z}^n$.

In the proof that (4) \Rightarrow (3) we may take $s = t + n$.

PROOF. The proof of the equivalence of (1)–(3) can be adapted from the proof of [Ho, Proposition 1].

(4) \Rightarrow (3). Assume $\Lambda = \mathbf{Z}^n \cap H_1 \cap \dots \cap H_t$ where $H_i = \{v \in \mathbf{Q}^n \mid f_i(v) \geq 0\}$. Define $y_i = f_i(x_1, \dots, x_n)$ for $i = 1, \dots, t$. Then $y_i \in \mathbf{Z}^+$ for $(x_1, \dots, x_n) \in \Lambda$. The map $(x_1, \dots, x_n) \rightarrow (y_1, \dots, y_t, x_1, \dots, x_n)$ is an isomorphism of $\mathbf{Q}\Lambda$ onto the subspace V of \mathbf{Q}^{t+n} defined by the equations $Y_i = f_i(X_1, \dots, X_n)$. Clearly the restriction of this map to Λ maps Λ isomorphically onto $(\mathbf{Z}'_+ \times \mathbf{Z}^n) \cap V$.

(3) \Rightarrow (4). We assume $\Lambda = (\mathbf{Z}'_+ \times \mathbf{Z}^{s-t}) \cap V$ and show Λ is isomorphic to a semigroup of the form $\mathbf{Z}^n \cap H_1 \cap \dots \cap H_t$ as in part (4). We can choose a basis v_1, \dots, v_n for $\mathbf{Q}\Lambda$ such that v_1, \dots, v_n generate $\mathbf{Z}\Lambda$ as a group. Then we define an isomorphism $\phi: \mathbf{Q}\Lambda \rightarrow \mathbf{Q}^n$ by $\phi(\sum_{i=1}^n \lambda_i v_i) = (\lambda_1, \dots, \lambda_n)$. Clearly ϕ maps $\mathbf{Z}\Lambda$ isomorphically onto \mathbf{Z}^n . Let ψ be the composite $\mathbf{Q}^n \xrightarrow{\phi^{-1}} \mathbf{Q}\Lambda \subseteq \mathbf{Q}^s$. Define $\psi^*: (\mathbf{Q}^s)^* \rightarrow (\mathbf{Q}^n)^*$ by $(\psi^*g)(v) = g(\psi(v))$ for $g \in (\mathbf{Q}^s)^*$, $v \in \mathbf{Q}^n$. Let $x_i \in (\mathbf{Q}^s)^*$ be the coordinate functions defined by $x_i(\mu) = \mu_i$ for $\mu = (\mu_1, \dots, \mu_s) \in \mathbf{Q}^s$ and for $i = 1, \dots, t$ define $f_i = \psi^*x_i \in (\mathbf{Q}^n)^*$. Let H_i be the halfspace $H_i = \{u \in \mathbf{Q}^n \mid f_i(v) \geq 0\}$. We claim that ϕ maps Λ isomorphically onto $\mathbf{Z}^n \cap H_1 \cap \dots \cap H_t$.

Certainly $\phi(\Lambda) \subseteq \mathbf{Z}^n$, since $\phi(\mathbf{Z}\Lambda) = \mathbf{Z}^n$. If $\lambda \in \Lambda$, then $f_i(\phi(\lambda)) = (\psi^*x_i)(\phi(\lambda)) = x_i(\psi(\phi(\lambda))) = x_i(\lambda) \geq 0$ since $\psi \circ \phi$ is just the inclusion $\mathbf{Q}\Lambda \subseteq \mathbf{Q}^s$. Hence $\phi(\lambda) \in H_i$ and $\phi(\Lambda) \subseteq \mathbf{Z}^n \cap H_1 \cap \dots \cap H_t$.

Finally, given $\mu \in \mathbf{Z}^n \cap H_1 \cap \dots \cap H_t$ there exists $\lambda \in \mathbf{Z}\Lambda$ with $\phi(\lambda) = \mu$, since $\phi: \mathbf{Z}\Lambda \rightarrow \mathbf{Z}^n$ is an isomorphism. The computation above shows that $x_i(\lambda) \geq 0$ for $i = 1, \dots, t$ so $\lambda \in \Lambda$ and $\phi: \Lambda \rightarrow \mathbf{Z}^n \cap H_1 \cap \dots \cap H_t$ is the desired isomorphism.

The maps ϕ, ψ , and ψ^* constructed above will play an important role in §2.

1.4. We now interpret our work in terms of rings of invariants of tori. Assume that $G = \text{GL}(1, k)^r$ is a torus acting on $\mathcal{O}(U_s) = k[T_1^{\pm 1}, \dots, T_s^{\pm 1}]$ so that for $g = (g_1, \dots, g_r) \in G$,

$$(1) \quad g \cdot T_j = \prod_{i=1}^r g_i^{a_{ij}} T_j \quad \text{where } a_{ij} \in \mathbf{Z}.$$

It is easily checked that a monomial T^λ is fixed by G if and only if $m_i(\lambda) = 0$ for $1 \leq i \leq r$ where

$$(2) \quad m_i = \sum_{j=1}^s a_{ij} x_j \in (\mathbf{Q}^s)^*.$$

It follows that $\mathcal{O}(U_\beta)^G = k\Lambda_\beta$, a semigroup algebra, where

$$\Lambda_\beta = \{ \lambda \in \mathbf{Z}^s \mid m_i(\lambda) = 0 \text{ for } 1 \leq i \leq r, \lambda_j \geq 0 \text{ if } j \notin \beta \}.$$

Let $V = \{ \lambda \in \mathbf{Q}^s \mid m_i(\lambda) = 0 \text{ for } 1 \leq i \leq r \}$, and $H_i = \{ \lambda \in \mathbf{Q}^s \mid \lambda_i \geq 0 \}$. Then $\Lambda_\beta = \mathbf{Z}^s \cap V \cap \bigcap_{j \notin \beta} H_j$ is a semigroup of the type described in statement (3) of Proposition 1.3, and $\mathcal{O}(U_s)^G = k\Lambda_s$ where $\Lambda_s = \mathbf{Z}^s \cap V$.

Conversely given Λ_β, Λ_s of this form, we may assume the vector space V is the solution space of a system of linear equations as in (2), define an action of a torus G on $\mathcal{O}(U_s)$ as in (1), and then we obtain $\mathcal{O}(U_s)^G = k\Lambda_s$ and $\mathcal{O}(U_\beta)^G = k\Lambda_\beta$.

It is often useful to regard our torus G as a subtorus of the torus $\Gamma = \text{GL}(1)^s$ which consists of all invertible linear transformations represented by diagonal matrices on the basis T_1, \dots, T_s . For $g = (g_1, \dots, g_s) \in \Gamma$ we have $gT_i = g_i T_i$, so if we define $T_i(g) = g_i$ then T_1, \dots, T_s generate the character group $M = M(\Gamma)$ of Γ and

$$g \cdot T^\lambda = T^\lambda(g) T^\lambda \quad \text{for } g \in \Gamma, \lambda \in \mathbf{Z}^s.$$

The correspondence $\lambda \leftrightarrow T^\lambda$ allows us to switch between additive and multiplicative notation for M . We obtain an induced action of Γ on \mathbf{Z}^s by

$$g \cdot \lambda = T^\lambda(g) \lambda \quad \text{for } g \in \Gamma, \lambda \in \mathbf{Z}^s.$$

If $N = N(\Gamma)$ is the group of one parameter subgroups of Γ there is a pairing $(,) : N \times M \rightarrow \mathbf{Z}$ defined by

$$T^\lambda(x(g)) = g^{(x, \lambda)} \quad \text{for } g \in k, x \in N, \lambda \in M.$$

Let x_1, \dots, x_s be the basis for N such that $T_i(x_j(g)) = g^{\delta_{ij}}$, for all $g \in k$, where δ_{ij} is the Kronecker delta. We also write $x(\lambda) = (x, \lambda)$. Then $x_i(\lambda) = \lambda_i$ and x_1, \dots, x_s can be viewed as coordinate functions on \mathbf{Z}^s .

LEMMA. *There exists a subset β' of s containing β such that $\mathcal{O}(U_\beta)^G = \mathcal{O}(U_{\beta'})^G$ and the distinct bounding halfspaces of $\mathbf{Q}^+ \Lambda_\beta = \mathbf{Q}^+ \Lambda_{\beta'}$ are precisely the halfspaces $\mathbf{Q} \Lambda_\beta \cap H_j$ with $j \notin \beta'$.*

PROOF. By Lemma 1.2 any bounding halfspace of $\mathbf{Q}^+ \Lambda_\beta$ has the form $\mathbf{Q} \Lambda_\beta \cap H_j$ where $j \notin \beta$. Simply choose a subset β' containing β so the distinct bounding halfspaces have the form $\mathbf{Q} \Lambda_\beta \cap H_j$ where $j \notin \beta'$.

1.5. In the situations described above where $\mathcal{O}(U_s)^G = k\Lambda_s$, $\mathcal{O}(U_\beta)^G = k\Lambda_\beta$, it is possible to have $\mathbf{Z}\Lambda_\beta$ strictly contained in the group $\Lambda_s = \mathbf{Z}^s \cap V$. However if we are given $\Lambda_\beta = \mathbf{Z}^s \cap V \cap \bigcap_{j \in \beta} H_j$ for some subspace V of $\mathbf{Q}\Lambda_\beta$ then since $\Lambda_\beta \subseteq \mathbf{Q}\Lambda_\beta \subseteq V$ we obtain $\Lambda_\beta \subseteq \mathbf{Z}^s \cap \mathbf{Q}\Lambda_\beta \cap \bigcap_{j \in \beta} H_j \subseteq \mathbf{Z}^s \cap V \cap \bigcap_{j \in \beta} H_j = \Lambda_\beta$. Hence $\Lambda_\beta = \mathbf{Z}^s \cap \mathbf{Q}\Lambda_\beta \cap \bigcap_{j \in \beta} H_j$, and we could replace V by $\mathbf{Q}\Lambda_\beta$ in the definition of Λ_β . We also note

LEMMA. $\mathbf{Z}\Lambda_\beta = \mathbf{Z}^s \cap \mathbf{Q}\Lambda_\beta$.

PROOF. If $\lambda \in \mathbf{Z}^s \cap \mathbf{Q}\Lambda_\beta$, there is an integer $p \geq 1$ such that $p\lambda \in \mathbf{Z}\Lambda_\beta$. Hence $p\lambda = \lambda_1 - \lambda_2$ for $\lambda_1, \lambda_2 \in \Lambda_\beta$, so $p\lambda = p\lambda_1 - \lambda_3$ where $\lambda_3 = \lambda_2 + (p - 1)\lambda_1 \in \Lambda_\beta \cap p\mathbf{Z}^s = p\Lambda_\beta$. Therefore $\lambda_3 = p\lambda_4$ for some $\lambda_4 \in \Lambda_\beta$ and $\lambda = \lambda_1 - \lambda_4 \in \mathbf{Z}\Lambda_\beta$. Since the inclusion $\mathbf{Z}\Lambda_\beta \subseteq \mathbf{Z}^s \cap \mathbf{Q}\Lambda_\beta$ is obvious this proves the result.

1.6. Applying the procedure outlined after Proposition 1.3 to Λ_β and $\mathbf{Z}\Lambda_\beta$ we obtain

COROLLARY. Given an action of a torus G on $\mathcal{O}(U_s)$ we may extend the action of G to a larger torus G' so that $\mathcal{O}(U_\beta)^{G'} = k\Lambda_\beta$ and $\mathcal{O}(U_s)^{G'} = k\mathbf{Z}\Lambda_\beta$.

There is no loss of generality in assuming that the torus G acts faithfully on $\mathcal{O}(U_s)$ and we shall always do so. This is equivalent to requiring that the equations $m_j(x)$, $j = 1, \dots, r$, are linearly independent. If $\mathcal{O}(U_\beta)^G = k\Lambda_\beta$, then the Krull dimension of $\mathcal{O}(U_\beta)^G$ is equal to the rank of $\mathbf{Z}\Lambda_\beta$. Thus by linear algebra

$$\dim \mathcal{O}(U_\beta) - \dim G = s - r = \dim \mathcal{O}(U_s)^G \geq \dim \mathcal{O}(U_\beta)^G.$$

Here $\dim G = r$ is the dimension of G as an algebraic group, and the dimensions of the rings are their Krull dimensions. We say that G satisfies the *dimension equality on $\mathcal{O}(U_\beta)$* provided $\dim \mathcal{O}(U_\beta)^G = \dim \mathcal{O}(U_\beta) - \dim G$. By Lemma 1.5 this is equivalent to the requirement that $\mathbf{Z}\Lambda_\beta = \Lambda_s$. By Corollary 1.6 we may assume the dimension equality holds by embedding G in a larger torus without changing $\mathcal{O}(U_\beta)^G$.

1.7. LEMMA. Suppose $\Lambda = (\mathbf{Z}'_+ \times \mathbf{Z}^{s-r}) \cap V$ and that $\mathbf{Q}\Lambda \cap \ker x_1$ is a bounding hyperplane of $\mathbf{Q}^+\Lambda$. If $x_1(\mathbf{Z}\Lambda) = a\mathbf{Z}$ for some $a > 0$ then there exists $\lambda \in \Lambda$ such that $x_1(\lambda) = a$.

PROOF. By assumption $x_1(\lambda') = a$ for some $\lambda' \in \mathbf{Z}\Lambda$. Let $H_j = \{\lambda \in \mathbf{Q}^s \mid x_j(\lambda) \geq 0\}$. For $2 \leq j \leq r$ we consider three cases which may arise. If $\mathbf{Q}\Lambda \subseteq H_j^0$ then $x_j(\lambda') = 0$; in this case let $u_j = 0$. In the other cases $\mathbf{Q}\Lambda \cap H_j^0$ has codimension one in $\mathbf{Q}\Lambda$.

Suppose next that $H_1^0 \cap \mathbf{Q}\Lambda = H_j^0 \cap \mathbf{Q}\Lambda$. Then $H_1 \cap \mathbf{Q}\Lambda = H_j \cap \mathbf{Q}\Lambda$ or $H_1 \cap \mathbf{Q}\Lambda = (-H_j) \cap \mathbf{Q}\Lambda$. Since $\mathbf{Q}\Lambda \not\subseteq H_1^0$ there exists $\mu \in \Lambda$ with $x_1(\mu) > 0$, and so $\mu \in H_1 \cap \mathbf{Q}\Lambda$ but $\mu \notin (-H_j) \cap \mathbf{Q}\Lambda$. Therefore $H_1 \cap \mathbf{Q}\Lambda = H_j \cap \mathbf{Q}\Lambda$, so $x_j(\lambda') > 0$ in this case and we set $u_j = 0$.

Finally suppose that $H_1^0 \cap \mathbf{Q}\Lambda$ and $H_j^0 \cap \mathbf{Q}\Lambda$ are distinct subspaces of codimension one in $\mathbf{Q}\Lambda$. Then $H_1^0 \cap \mathbf{Q}\Lambda \not\subseteq H_j^0$ and since $H_1^0 \cap \Lambda$ spans $H_1^0 \cap \mathbf{Q}\Lambda$, there exists $u_j \in H_1^0 \cap \Lambda$ with $u_j \notin H_j^0$. Therefore $x_1(u_j) = 0$ and $x_j(u_j) > 0$.

Set $u = u_2 + \dots + u_r \in \Lambda$. Then $x_1(u) = 0$ and in the first two cases $x_j(\lambda') \geq 0$ and $x_j(u) \geq 0$, while in the last case $x_j(u) > 0$. If we set $\lambda = \lambda' + Nu$ for a sufficiently large positive integer N , then λ will satisfy the conditions of the lemma.

1.8. We make one further reduction. Suppose $\Lambda_\beta = \mathbf{Z}^s \cap V \cap \bigcap_{j \notin \beta} H_j$ and the groups $\mathbf{Z}\Lambda_\beta$ are equal for all $\beta \in \Delta$. For all j such that $\mathbf{Q}\Lambda \not\subseteq \ker x_j$, let $x_j(\mathbf{Z}\Lambda_\beta) = a_j\mathbf{Z}$ where $a_j \geq 1$. We define an isomorphism $\phi: \mathbf{Q}^s \rightarrow \mathbf{Q}^s$ by $\phi(\lambda_1, \dots, \lambda_s) = (\mu_1, \dots, \mu_s)$ where $\mu_j = \lambda_j$ if $\mathbf{Q}\Lambda \subset \ker x_j$ and $\mu_j = \lambda_j/a_j$ otherwise. Let $\psi = \phi^{-1}$ and as in the proof of Proposition 1.3 define $\psi^*: (\mathbf{Q}^s)^* \rightarrow (\mathbf{Q}^s)^*$ by $(\psi^*f)(v) = f(\psi(v))$. Let $V' \subseteq \mathbf{Q}^s$ be the solution space to the equations $m'_j = \psi^*m_j$, $1 \leq j \leq r$. Then ϕ maps Λ_β isomorphically onto the semigroup $\Lambda'_\beta = \mathbf{Z}^s \cap V' \cap \bigcap_{j \notin \beta} H_j$ and $x_j(\mathbf{Z}\Lambda'_\beta) = \mathbf{Z}$ for all $j \notin \beta$. In terms of fixed rings of tori we have shown the following lemma.

LEMMA. *There exists an action of $G' \cong G$ on k^s such that for all $\beta \in \Delta$, $\mathcal{O}(U_\beta)^{G'} = k\Lambda'_\beta \cong k\Lambda_\beta$ and for all $j \notin \beta$, $x_j(\mathbf{Z}\Lambda'_\beta) = \mathbf{Z}$.*

1.9. If G satisfies the dimension equality, the reduction achieved in the previous lemma has an interesting interpretation in terms of pseudoreflections.

LEMMA. *Assume G satisfies the dimension equality on $\mathcal{O}(U_\beta)$. If $\mathcal{O}(U_\beta)^G = k\Lambda_\beta$ then $x_j(\mathbf{Z}\Lambda_\beta) = \mathbf{Z}$ if and only if G contains no nonidentity pseudoreflection fixing the hyperplane defined by $x_j = 0$.*

PROOF. Without loss of generality $j = 1$. If G contains a pseudoreflection of the form $(\omega, 1, \dots, 1)$ where ω is a primitive m th root of unity, $m > 1$, then $k\Lambda_\beta = \mathcal{O}(U_\beta)^G \subseteq \mathcal{O}(U_s)^G \subseteq k[T_1^{\pm m}, T_2^{\pm 1}, \dots, T_s^{\pm 1}]$ and $x_1(\mathbf{Z}\Lambda_\beta) \subseteq m\mathbf{Z}$.

Conversely, assume $x_1(\mathbf{Z}\Lambda_\beta) = m\mathbf{Z}$ where $m > 1$. For $g \in \Gamma$, $\lambda \in \mathbf{Z}^s$ we have $g \cdot T^\lambda = T^\lambda(g)T^\lambda$. Hence $\mathcal{O}(U_s)^G = \text{Span}\{T^\lambda \mid \lambda \in \mathbf{Z}^s, G \subseteq \text{Ker } T^\lambda\}$. Since $x_1(\mathbf{Z}\Lambda_\beta) = m\mathbf{Z}$ and $\mathbf{Z}\Lambda_\beta = \Lambda_s$ by the dimension equality, we have $\mathcal{O}(U_s)^G \subseteq k[T_1^{\pm m}, T_2^{\pm 1}, \dots, T_s^{\pm 1}]$. Hence the pseudoreflection $g = (\omega, 1, \dots, 1)$ fixes $\mathcal{O}(U_s)^G$ where ω is a primitive m th root of unity. Therefore $g \in \text{Ker } T^\lambda$ for all $\lambda \in \mathbf{Z}^s$ such that $G \subseteq \text{Ker } T^\lambda$. However by [Hu, Proposition 16.1] $G = \bigcap \text{Ker } T^\lambda$ where the intersection is over those kernels of characters which contain G . Hence $g \in G$ as required.

1.10. EXAMPLE. If G does not satisfy the dimension equality on $\mathcal{O}(U_\beta)$ the above result may fail to hold.

Let $G = \text{GL}(1, k)^2$ and let G act on $K = k[T_1, T_2, T_3, T_4]$ by $g \cdot T_j = \prod_i g_i^{a_{ij}} T_j$ where

$$(a_{ij}) = \begin{pmatrix} 1 & -1 & 0 & -2 \\ -1 & 0 & -1 & 2 \end{pmatrix}.$$

Then $T^\lambda \in K^G$ if and only if λ satisfies

$$\begin{aligned} \lambda_1 - \lambda_2 & & -2\lambda_4 & = 0 & \text{and } \lambda_i & \geq 0, \\ -\lambda_1 & & -\lambda_3 + 2\lambda_4 & = 0. \end{aligned}$$

Adding these equations we find $\lambda_2 + \lambda_3 = 0$. Therefore $\lambda_2 = \lambda_3 = 0$, and $K^G = k\Lambda$ where $\Lambda = \mathbf{Z}_+(2, 0, 0, 1)$. Hence $x_1(\mathbf{Z}\Lambda) = 2\mathbf{Z}$. However if $g \in G$ is a pseudoreflection fixing the hyperplane defined by $x_1 = 0$, then from $g \cdot T_2 = T_2$ and $g \cdot T_3 = T_3$, it follows that $g_1 = g_2 = 1$ so $g = 1$.

2. Rings of differential operators.

2.0. We explain the notation for halfspaces and hyperplanes which we will be using from now on. Let x_i be the coordinate functions on \mathbf{Q}^s , $\mathcal{H}_i = \{\lambda \in \mathbf{Q}^s \mid x_i(\lambda) \geq 0\}$, and $\mathcal{H}_i^0 = \ker x_i$. Let f_i be the restriction of x_i to $\mathbf{Q}\Lambda$, $H_i = \mathcal{H}_i \cap \mathbf{Q}\Lambda$, and $H_i^0 = \mathcal{H}_i^0 \cap \mathbf{Q}\Lambda = \ker f_i$. Identifying $\mathbf{Q}\Lambda$ with \mathbf{Q}^r by means of the isomorphism ϕ , we have $f_i = \psi^*x_i$ as in Proposition 1.3.

We are now able to formulate Theorem B. Suppose the torus $G = \text{GL}(1)^r$ acts on $Y = \bigcup_{\beta \in \Delta} U_\beta$ as in the introduction. For each $\beta \in \Delta$, $\mathcal{O}(U_\beta)^G = k\Lambda_\beta$ where

$$\Lambda_\beta = \{ \lambda \in \mathbf{Z}^s \mid m_i(\lambda) = 0 \text{ for } 1 \leq i \leq r, \lambda_j \geq 0 \text{ if } j \notin \beta \}$$

as described in §1. Since for each $\beta \in \Delta$, $k\Lambda_\beta$ has the same quotient field, the groups $\mathbf{Z}\Lambda_\beta$ are equal for $\beta \in \Delta$, and we write $\mathbf{Z}\Lambda = \mathbf{Z}\Lambda_\beta$ and similarly $\mathbf{Q}\Lambda = \mathbf{Q}\Lambda_\beta$ for any $\beta \in \Delta$.

The conditions for θ to be surjective are expressed in terms of the relationship between the halfspaces in \mathbf{Q}^s determined by the coordinate hyperplanes and their restriction to the subspace $\mathbf{Q}\Lambda$. By Lemma 1.2 if $\beta \in \Delta$, any bounding halfspace of $\mathbf{Q}^+ \Lambda_\beta$ has the form $\mathbf{Q}\Lambda \cap \mathcal{H}_j$ where $\mathbf{Q}\Lambda \not\subseteq \mathcal{H}_j^0$ and $j \notin \beta$. We call the set of such halfspaces as β runs over Δ , the induced halfspaces of $\mathbf{Q}\Lambda$. That is set $\alpha = \{ j \in \mathbf{s} \mid \mathbf{Q}\Lambda \not\subseteq \mathcal{H}_j^0, \text{ and } j \notin \bigcap_{\beta \in \Delta} \beta \}$. Then an induced halfspace of $\mathbf{Q}\Lambda$ is of the form $\mathbf{Q}\Lambda \cap \mathcal{H}_j$ with $j \in \alpha$. We say that a differential operator $d \in D(U_s)$ is homogeneous if for every monomial $f \in \mathcal{O}(U_s)$ we have $d \cdot f = \lambda f$ for some $\lambda \in k$, depending on f .

THEOREM B. (1) *The map $\theta: D(Y)^G \rightarrow D(Y_\Delta/G)$ is surjective if and only if*

- (a) *The induced halfspaces of $\mathbf{Q}\Lambda$ are distinct.*
 - (b) *Every induced halfspace of $\mathbf{Q}\Lambda$ is a bounding halfspace of $\mathbf{Q}^+ \Lambda_\beta$ for some $\beta \in \Delta$.*
 - (c) *If $\mathbf{Q}\Lambda \cap \mathcal{H}_i$ is an induced halfspace of $\mathbf{Q}\Lambda$ then $x_i(\mathbf{Z}\Lambda) = \mathbf{Z}$.*
- (2) *Let J be the left ideal of $D(U_s)^G$ generated by all first order homogeneous differential operators which vanish on $k\mathbf{Z}\Lambda$. Then $\text{Ker } \theta = J \cap D(Y)^G$.*

We give some examples to illustrate conditions (a)–(c) of Theorem B part (1) at the end of this section. We also show that the definition of $D(Y_\Delta/G)$ may depend on Δ and not just on Y and G .

In Proposition 3.7, we show how conditions (a)–(c) of (1) can be achieved without changing $D(Y_\Delta/G)$ up to isomorphism.

To prove Theorem B we shall obtain explicit descriptions of the rings $D(Y)^G$, $D(Y_\Delta/G)$, and the map θ .

2.1. **LEMMA.** *If K is a commutative k -algebra and \mathcal{C} a multiplicatively closed subset of K consisting of regular elements, then any differential operator on K has a unique extension to $K_\mathcal{C}$. In addition $D(K) = \{ x \in D(K_\mathcal{C}) \mid x \cdot K \subseteq K \}$.*

PROOF. See [H, Lemma 2].

2.2. Let $k[Q_1, \dots, Q_s, P_1, \dots, P_s]$ be the s th Weyl algebra over k with generators Q_j, P_i satisfying $[P_i, Q_j] = \delta_{ij}$ where $[a, b] = ab - ba$. We regard this ring as the ring of differential operators on $k[T_1, \dots, T_s]$ where Q_i acts as multiplication by T_i

and P_i as $\partial/\partial T_i$. Since $\mathcal{O}(Y) = \bigcap_{\beta \in \Delta} \mathcal{O}(U_\beta) = k[T_1, \dots, T_s, T_i^{-1}; i \in \bigcap_{\beta \in \Delta} \beta]$ we can regard $D(Y)$ as a subalgebra of $D(U_s) = k[Q_1^{\pm 1}, \dots, Q_s^{\pm 1}, P_1, \dots, P_s]$ by Lemma 2.1. It is easy to show that $D(U_s) = \bigoplus_{\lambda \in \mathbf{Z}^s} D(U_s)(\lambda)$ where $D(U_s)(\lambda) = \{x \in D(U_s) \mid [Q_i P_i, x] = \lambda_i x \text{ for } 1 \leq i \leq s\}$. We denote by A the subring $A = D(U_s)(0) = k[Q_1 P_1, \dots, Q_s P_s]$ of $D(U_s)$. Then A is exactly the set of homogeneous differential operators on $\mathcal{O}(U_s)$.

LEMMA. *Let R be a subring of $D(U_s)$ containing A . If I is an ideal of R then $I = \bigoplus_{\lambda \in \mathbf{Z}^s} I(\lambda)$ where $I(\lambda) = \{x \in I \mid [Q_i P_i, x] = \lambda_i x \text{ for } 1 \leq i \leq s\}$.*

PROOF. Let $\Gamma = \text{GL}(1)^s$. Then Γ acts rationally on $\mathcal{O}(\Gamma) = \mathcal{O}(U_s)$ and $D(\Gamma) = D(U_s)$. Furthermore the Lie algebra $\text{Lie } \Gamma$ of Γ can be identified, as differential operators on $\mathcal{O}(\Gamma)$, with $kQ_1 P_1 + \dots + kQ_s P_s$. The action of Γ on $D(U_s)$ gives the following action of $\text{Lie } \Gamma$:

$$\xi \cdot d = [\xi, d] \quad \text{for } \xi \in \text{Lie } \Gamma, d \in D(U_s).$$

Therefore, if R is a subring of $D(U_s)$ containing A , and I an ideal of R , then I and R are stable under $\text{Lie } \Gamma$ and so are Γ -submodules of $D(U_s)$. Identifying the character group of Γ with \mathbf{Z}^s , we have $I = \bigoplus_{\lambda \in \mathbf{Z}^s} I(\lambda)$ with $I(\lambda)$ as above, by standard results.

We note also that $R(\lambda) \subseteq Q^\lambda A$ and $R(0) = A$.

2.3. We next obtain a description of $D(k\Lambda)$ where Λ is an arbitrary subsemigroup of \mathbf{Z}^s . There is no loss of generality in assuming $\mathbf{Z}\Lambda = \mathbf{Z}^s$. In $k\Lambda$ set $\mathcal{C} = \{T^\lambda \mid \lambda \in \Lambda\}$. Then $k\Lambda_{\mathcal{C}} = \mathcal{O}(U_s) = k[T_1^{\pm 1}, \dots, T_s^{\pm 1}]$. We view $D(k\Lambda)$ as a subring of $D(U_s)$. Note that for $\mu \in \mathbf{Z}^s$ we have $Q_i P_i \cdot T^\mu = \mu_i T^\mu$.

Let $A_{\mathbf{Q}} = \mathbf{Q}[Q_1 P_1, \dots, Q_s P_s]$ so $A = A_{\mathbf{Q}} \otimes_{\mathbf{Q}} k$. Elements of A define polynomial functions from $\mathbf{Z}^s \subseteq \mathbf{Q}^s$ to k by the rule

$$Q_i P_i(\mu) = \mu_i \quad \text{for } \mu \in \mathbf{Z}^s.$$

With this definition we have for $f \in A$

$$f(Q_1 P_1, \dots, Q_s P_s) \cdot T^\mu = f(\mu) \cdot T^\mu \quad \text{for } \mu \in \mathbf{Z}^s.$$

Since we have identified $A_{\mathbf{Q}}$ with the symmetric algebra on $(\mathbf{Q}^s)^*$, we can now regard an element $f = f(x_1, \dots, x_s) \in (\mathbf{Q}^s)^*$ as an element $f = f(Q_1 P_1, \dots, Q_s P_s)$ of $A_{\mathbf{Q}}$.

For a subset Ω of \mathbf{Q}^s we write $\text{ann}_A \mathbf{Q} = \{f \in A \mid f(w) = 0 \text{ for all } w \in \Omega\}$. Clearly $\text{ann}_A \Omega = \text{ann}_A \bar{\Omega}$ where $\bar{\Omega}$ denotes the Zariski closure of Ω .

THEOREM. *Let Λ be a subsemigroup of \mathbf{Z}^s with $\mathbf{Z}\Lambda = \mathbf{Z}^s$. Then*

$$D(k\Lambda) = \bigoplus_{\lambda \in \mathbf{Z}^s} D(k\Lambda)(\lambda)$$

where $D(k\Lambda)(\lambda) = Q^\lambda \text{ann}_A \Omega(\lambda)$ and $\Omega(\lambda) = \{\mu \in \Lambda \mid \mu + \lambda \notin \Lambda\}$.

PROOF. The direct sum decomposition is immediate from Lemma 2.2. It is also clear that $D(k\Lambda)(\lambda) \subseteq Q^\lambda A$. If $f \in A$, then $Q^\lambda f \cdot T^\mu = f(\mu) T^{\mu+\lambda}$. By Lemma 2.1, $Q^\lambda f \in A$ if and only if f vanishes on $\Omega(\lambda)$. This proves the result.

2.4. We can regard $\mathcal{O}(Y) = k[T_1, \dots, T_s, T_i^{-1} : i \in \cap_{\beta \in \Delta} \beta]$ as a semigroup algebra $k\Sigma$ where Σ is the semigroup $\Sigma = \{\sigma \in \mathbf{Z}^s \mid \sigma_i \geq 0 \text{ if } i \notin \cap_{\beta \in \Delta} \beta\}$. Let $\Omega(\lambda) = \{\mu \in \Sigma \mid \lambda + \mu \notin \Sigma\}$. Then we have $D(Y) = \bigoplus_{\lambda \in \mathbf{Z}^s} D(Y)(\lambda)$ where $D(Y)(\lambda) = Q^\lambda \text{ann}_A \Omega(\lambda)$.

Now $D(U_s)$ is the twisted group algebra of the group $M = \langle Q_1^{\pm 1}, \dots, Q_s^{\pm 1} \rangle$ over the subring $D(U_s)(0) = A$. As explained in the introduction the action of G on $\mathcal{O}(U_s)$ extends to an action of G on $D(U_s)$ by the rule

$$(g \cdot d)(f) = g(d(g^{-1}f))$$

for $f \in \mathcal{O}(U_s)$, $d \in D(U_s)$, and $g \in G$. A simple computation shows that for $g = (g_1, \dots, g_r) \in G$ we have

$$g \cdot Q_j = \prod_{i=1}^r g_i^{a_{ij}} Q_j$$

and

$$g \cdot P_j = \prod_{i=1}^r g_i^{-a_{ij}} P_j.$$

It follows from this that the subring A is invariant under G and that $D(U_s)^G$ is the twisted group algebra of the subgroup M_1 of G -fixed elements of M over A . It is easy to see that $Q^\lambda \in M$ is G -fixed if and only if $\lambda \in \mathbf{Z}^s \cap V$, and thus we obtain

$$D(U_s)^G = \bigoplus_{\lambda \in \mathbf{Z}^s \cap V} D(U_s)(\lambda).$$

Since $D(Y)^G = D(Y) \cap D(U_s)^G$ we obtain the following result.

THEOREM.

$$D(Y)^G = \bigoplus_{\lambda \in \mathbf{Z}^s \cap V} D(Y)^G(\lambda)$$

where $D(Y)^G(\lambda) = Q^\lambda \text{ann}_A \overline{\Omega(\lambda)}$, and $\Omega(\lambda) = \{\mu \in \Sigma \mid \mu + \lambda \notin \Sigma\}$.

2.5. By identifying $\mathbf{Z}\Lambda$ with \mathbf{Z}^n we have $k\mathbf{Z}\Lambda = k[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$. Let $k[q_1, \dots, q_n, p_1, \dots, p_n]$ be the n th Weyl algebra with generators q_i, p_j satisfying $[p_i, q_j] = \delta_{ij}$. We regard this as the ring of differential operators on $k[t_1, \dots, t_n]$ where q_i acts as multiplication by t_i and p_j as $\partial/\partial t_j$. Let $B_{\mathbf{Q}} = \mathbf{Q}[q_1 p_1, \dots, q_n p_n]$ and $B = B_{\mathbf{Q}} \otimes_{\mathbf{Q}} k$. We may identify $B_{\mathbf{Q}}$ with the symmetric algebra on $(\mathbf{Q}^n)^*$. The next lemma will be used to find the Zariski closures of certain sets.

LEMMA. *Let Λ be a subsemigroup of \mathbf{Q}^n with $\mathbf{Q}\Lambda = \mathbf{Q}^n$. For $g \in (\mathbf{Q}^n)^*$ and $b \in \mathbf{Q}$ set $\Lambda_b = \{\lambda \in \Lambda \mid g(\lambda) = b\}$. Suppose that*

- (i) $\Lambda_b \neq \emptyset$,
- (ii) $\dim \mathbf{Q}\Lambda_0 = n - 1$.

Then $\overline{\Lambda}_b = \{\lambda \in \mathbf{Q}^n \mid g(\lambda) = b\}$.

PROOF. It suffices to show that if $f \in B_{\mathbf{Q}}$ and $f(\Lambda_b) = 0$, then $f(\lambda) = 0$ for all $\lambda \in \mathbf{Q}^n$ such that $g(\lambda) = b$.

We first prove the case $b = 0$. Write $X_1 = q_1 p_1, \dots, X_n = q_n p_n$ and introduce new indeterminates Y_1, \dots, Y_n, T . Suppose

$$f(X + TY) = f(X_1 + TY_1, \dots, X_n + TY_n) = \sum_{j=0}^m T^j g_j(X, Y)$$

where $g_j(X, Y) \in \mathbf{Q}[X_1, \dots, X_n, Y_1, \dots, Y_n]$.

For $\lambda, \mu \in \Lambda_0$ and t a positive integer we have $0 = f(\lambda + t\mu) = \sum_{j=0}^m t^j g_j(\lambda, \mu)$. An argument using Vandermonde determinants shows that $g_j(\lambda, \mu) = 0$ for each j . Taking $\lambda = 0, \mu \in \Lambda_0$, and $t \in \mathbf{Q}^+$ we find that $f(\mathbf{Q}^+ \Lambda_0) = 0$ and so $g_j(\lambda, \mu) = 0$ for $\lambda, \mu \in \mathbf{Q}^+ \Lambda_0$. Then taking $t = -1$ we find that $f(\mathbf{Q} \Lambda_0) = 0$. Since $\dim \mathbf{Q} \Lambda_0 = n - 1$, this proves the case $b = 0$.

For the general case, there exists $\mu \in \Lambda$ such that $g(\mu) = b$, by (i). Define $f^* \in B_{\mathbf{Q}}$ by $f^*(\nu) = f(\nu + \mu)$. Then $f^*(\Lambda_0) = 0$ so by the case $b = 0$, $f^*(\mathbf{Q} \Lambda_0) = 0$. If now $\lambda \in \mathbf{Q}^n$ and $g(\lambda) = b$ then $g(\lambda - \mu) = 0$. Therefore $f(\lambda) = f^*(\lambda - \mu) = 0$ as required.

We can now describe the Zariski closed sets $\overline{\Omega(\lambda)}$ of Theorem 2.4 as a union of hyperplanes. For $\beta \in \Delta$ and $i \notin \beta$ define $\overline{\Omega_{\beta,i}(\lambda)} = \{\mu \in \mathbf{Q}^s \mid 0 \leq \mu_i < -\lambda_i, \mu_i \in \mathbf{Z}\}$. Then using Lemma 2.5 $\overline{\Omega(\lambda)} = \bigcup_{\beta \in \Delta} \bigcup_{i \notin \beta} \overline{\Omega_{\beta,i}(\lambda)}$. To describe $\text{ann}_A \overline{\Omega(\lambda)}$, set $S_{\lambda,\beta,i} = \prod_{j=0}^{-\lambda_i} (Q_j P_i - j)$ if $\beta \in \Delta, i \notin \beta, \lambda_i < 0$, and $S_{\lambda,\beta,i} = 1$ otherwise. Then $\text{ann}_A \overline{\Omega(\lambda)} = S_{\lambda} A$ where $S_{\lambda} = \prod_{\beta \in \Delta} \prod_{i \notin \beta} S_{\lambda,\beta,i}$.

2.6. We next give a description of the ring $D(k\Lambda_{\beta})$ where Λ_{β} is a semigroup of the form $\Lambda_{\beta} = \mathbf{Z}^n \cap \bigcap_{i \notin \beta} H_i$ and $\mathbf{Z} \Lambda_{\beta} = \mathbf{Z}^n$. We may assume that the bounding halfspaces of $\mathbf{Q}^+ \Lambda_{\beta}$ are the H_i with $i \notin \beta$. Then by Theorem 2.3, $D(k\Lambda_{\beta}) = \bigoplus_{\lambda \in \mathbf{Z}^n} q^{\lambda} \text{ann}_B \omega_{\beta}(\lambda)$, where $\omega_{\beta}(\lambda) = \{\mu \in \Lambda_{\beta} \mid \mu + \lambda \notin \Lambda_{\beta}\}$. It is easily seen that $\omega_{\beta}(\lambda) = \bigcup_{i \notin \beta} \omega_{\beta,i}(\lambda)$ where $\omega_{\beta,i}(\lambda) = \{\mu \in \Lambda_{\beta} \mid f_i(\lambda + \mu) < 0\}$.

The next lemma shows that $\overline{\omega_{\beta,i}(\lambda)}$ is a finite union of hyperplanes parallel to H_i^0 .

LEMMA. Suppose $i \notin \beta, H_i$ is a bounding halfspace of $\mathbf{Q} \Lambda_{\beta}$, and $f_i(\lambda) < 0$. Then

$$\overline{\omega_{\beta,i}(\lambda)} = \{\mu \in \mathbf{Q}^n \mid f_i(\mu) \in f_i(\mathbf{Z}^n) \text{ and } 0 \leq f_i(\mu) < -f_i(\lambda)\}.$$

PROOF. The left side is contained in the right since f_i is linear and the right side is Zariski closed in \mathbf{Q}^n .

We apply Lemma 2.5 with $g = f_i$. Suppose $f_i(\mu) = b \in f_i(\mathbf{Z}^n)$ and $0 \leq f_i(\mu) < -f_i(\lambda)$. By Lemma 1.7, there is a $\nu \in \Lambda_{\beta}$ such that $f_i(\nu) = b$, so condition (i) of Lemma 2.5 holds. Since H_i^0 is a bounding hyperplane of $\mathbf{Q} \Lambda_{\beta}$, condition (ii) holds. We conclude that $\mu \in \overline{\omega_{\beta,i}(\lambda)}$.

2.7. By taking intersections over all $\beta \in \Delta$, we can obtain the following description of $D(Y_{\Delta}/G)$.

THEOREM. Suppose that for all $\beta \in \Delta, \mathbf{Z} \Lambda_{\beta} = \mathbf{Z}^n$ and that for $i \notin \beta, H_i$ is a bounding halfspace of $\mathbf{Q}^+ \Lambda_{\beta}$. Then

$$D(Y_{\Delta}/G) = \bigoplus_{\mu \in \mathbf{Z}^n} D(Y_{\Delta}/G)(\mu)$$

where $D(Y_{\Delta}/G)(\mu) = q^{\lambda} \text{ann}_B \overline{\omega(\lambda)}$, and $\overline{\omega(\lambda)} = \bigcup_{\beta \in \Delta} \bigcup_{i \notin \beta} \overline{\omega_{\beta,i}(\lambda)}$, and $\overline{\omega_{\beta,i}(\lambda)}$ is a union of hyperplanes as described in Lemma 2.6.

2.8. We next investigate the map $\theta: D(Y)^G \rightarrow D(Y_\Delta/G)$. Let I be the ideal of functions of A which vanish on $\mathbf{Q}\Lambda$. Then I is generated by linear polynomials, that is by first order homogeneous differential operators.

LEMMA. Assume $Q^\lambda g \in D(U_\beta)^G$ where $\lambda \in \mathbf{Z}^s \cap V$, $g \in A$. Then

- (i) $Q^\lambda g \in \text{Ker } \theta_\beta$ if and only if $g \in I$,
- (ii) if $\lambda \notin \mathbf{Z}\Lambda$ then $Q^\lambda g \in \text{Ker } \theta_\beta$.

PROOF. (i) $Q^\lambda g \in \text{Ker } \theta_\beta$ if and only if for all $\mu \in \Lambda_\beta$, $0 = Q^\lambda g \cdot T^\mu = g(\mu)T^{\lambda+\mu}$.

This occurs if and only if $g(\Lambda_\beta) = 0$. By an argument used in the proof of Lemma 2.5 this is equivalent to $g(\mathbf{Q}\Lambda) = 0$. Hence $Q^\lambda g \in \text{Ker } \theta_\beta$ if and only if $g \in I$.

(ii) If $\lambda \notin \mathbf{Z}\Lambda$ and $\mu \in \Lambda_\beta$ then $\lambda + \mu \notin \Lambda_\beta$. Therefore $g(\Lambda_\beta) = 0$ and $Q^\lambda g \in \text{Ker } \theta_\beta$ as in (i).

If $Q^\lambda g \in D(U_\beta)^G$, and $\lambda \in \mathbf{Z}\Lambda$, then the unique extension of $\theta_\beta(Q^\lambda g)$ to a differential operator on $k\mathbf{Z}\Lambda$ is given by

$$\theta_\beta(Q^\lambda g)_t^{\phi(\mu)} = g(\mu)_t^{\phi(\lambda+\mu)} \quad \text{for } \mu \in \mathbf{Z}\Lambda.$$

Here $\phi: \mathbf{Z}\Lambda \rightarrow \mathbf{Z}^n$ is the isomorphism of Proposition 1.3. It follows that if $x \in D(U_\beta)^G \cap D(U_\gamma)^G$ then $\theta_\beta(x) = \theta_\gamma(x)$ as elements of $D(k\mathbf{Z}\Lambda)$. The map $\theta: D(Y)^G = \cap D(U_\beta)^G \rightarrow \cap D(\mathcal{O}(U_\beta)^G) = D(Y_\Delta/G)$ is obtained as the restriction of any θ_β , $\beta \in \Delta$.

PROOF OF THEOREM B. The second part of the theorem follows from Lemmas 2.2 and 2.8 since $\text{Ker } \theta = D(Y)^G \cap \text{Ker } \theta_\beta$.

Using Corollary 1.6 the action of G on $\mathcal{O}(U_s)$ may be extended to an action of a larger torus G' such that $\mathcal{O}(U_\beta)^{G'} = \mathcal{O}(U_\beta)^G = k\Lambda_\beta$ for all $\beta \in \Delta$, and $\mathcal{O}(U_s)^{G'} = k\mathbf{Z}\Lambda$. By Theorem 2.4 we have $D(Y)^{G'} = \bigoplus_{\lambda \in \mathbf{Z}\Lambda} D(Y)^{G'}(\lambda)$ where $D(Y)^{G'}(\lambda) = D(Y)^G(\lambda)$ for $\lambda \in \mathbf{Z}\Lambda$. By Lemma 2.8 $\theta(D(Y)^{G'}(\lambda)) = 0$ unless $\lambda \in \mathbf{Z}\Lambda$. Hence in studying surjectivity we may replace $D(Y)^G$ with the subring $D(Y)^{G'}$. In other words we can replace G by G' and assume $\mathbf{Z}\Lambda_\beta = \mathbf{Z}^s \cap V$ for all $\beta \in \Delta$.

Let $\bar{\theta}: D(U_s)^G \rightarrow D(\mathcal{O}(U_s)^G)$ be the map obtained from restriction of the differential operators. For any $x \in D(U_\beta)^G$, $\bar{\theta}(x) \in D(\mathcal{O}(U_s)^G)$ extends $\theta_\beta(x) \in D(\mathcal{O}(U_\beta)^G)$. The ring $D(U_s)^G$ is a skew group ring of the group $M = \{Q^\lambda \mid \lambda \in \mathbf{Z}\Lambda\}$ over $B = k[Q_1P_1, \dots, Q_sP_s]$. Therefore $\bar{\theta}$ is completely determined by its restriction to M and B . We note that $\mathcal{O}(U_s)^G$ is embedded in $D(U_s)^G$ as the subring spanned by $\{Q^\lambda \mid \lambda \in \mathbf{Z}\Lambda\}$ and in $D(\mathcal{O}(U_s)^G)$ as the subring $\{q^{\phi(\lambda)} \mid \lambda \in \mathbf{Z}\Lambda\}$. Hence, we have $\bar{\theta}(Q^\lambda) = q^{\phi(\lambda)}$ for all $\lambda \in \mathbf{Z}\Lambda$. We claim the restriction of $\bar{\theta}$ to B is given by

$$\bar{\theta}(Q_iP_i) = f_i(q_1p_1, \dots, q_n p_n) \quad \text{for } 1 \leq i \leq s.$$

Indeed if $\lambda \in \mathbf{Z}\Lambda$ we have

$$\begin{aligned} \bar{\theta}(Q_iP_i) \cdot t^{\phi(\lambda)} &= \lambda_i t^{\phi(\lambda)} = (x_i \phi^{-1})(\phi(\lambda))_t^{\phi(\lambda)} \\ &= (\psi^* x_i)(\phi(\lambda))_t^{\phi(\lambda)} = f_i(\phi(\lambda))_t^{\phi(\lambda)} \\ &= f_i(q_1p_1, \dots, q_n p_n) \cdot t^{\phi(\lambda)}. \end{aligned}$$

We note that the restriction of θ to A gives a surjective map from A to B . To see this note that the linear map $\psi: \mathbf{Q}^n \rightarrow \mathbf{Q}^s$ has rank n , and so $\psi^*: (\mathbf{Q}^s)^* \rightarrow (\mathbf{Q}^n)^*$ has rank n also. Hence the linear functionals $f_i = \psi^* x_i$, $1 \leq i \leq s$ span $(\mathbf{Q}^n)^*$ and so the linear polynomials $f_i = \theta(Q_iP_i)$ generate B .

To prove the theorem observe that $\theta: D(Y)^G \rightarrow D(Y_\Delta/G)$ will be surjective if and only if for all $\lambda \in \mathbf{Z}\Lambda$, the restriction of θ to $D(Y)^G(\lambda) \rightarrow D(Y_\Delta/G)(\phi(\lambda))$ is surjective. Using Theorems 2.4 and 2.7 we see that θ is surjective if and only if for all $\lambda \in \mathbf{Z}\Lambda$ we have $\theta(\text{ann}_A \overline{\Omega(\lambda)}) = \text{ann}_B \overline{\omega(\phi(\lambda))}$.

We first show that condition (a) is necessary for surjectivity. If (a) fails then for some $i, j \in \alpha, i \neq j$, we have $cf_i = f_j$ for some $c > 0$. Choose $\lambda \in \mathbf{Z}\Lambda$ with $\lambda_i < 0$. Then $\overline{\Omega(\lambda)}$ contains the hyperplanes \mathcal{H}_i^0 and \mathcal{H}_j^0 . Therefore $f_i^2 \in \theta(\text{ann}_A \overline{\Omega(\lambda)})$, but $f_j \notin \theta(\text{ann}_A \overline{\Omega(\lambda)})$. Therefore θ is not surjective, since $\text{ann}_B \overline{\omega(\phi(\lambda))}$ is a semiprime ideal.

Henceforth we assume that (a) holds. We have

$$\theta(\text{ann}_A \overline{\Omega(\lambda)}) = \text{ann}_B(\overline{\Omega(\lambda)} \cap \mathbf{Q}\Lambda),$$

since now distinct hyperplanes in $\overline{\Omega(\lambda)}$ now have distinct intersections with $\mathbf{Q}\Lambda$.

Since we always have $\theta(\text{ann}_A \overline{\Omega(\lambda)}) \subset \text{ann}_B \overline{\omega(\phi(\lambda))}$ it now suffices to show that conditions (b) and (c) hold if and only if for all $\lambda \in \mathbf{Z}\Lambda$ we have $\overline{\Omega(\lambda)} \cap \mathbf{Q}\Lambda \subset \overline{\omega(\phi(\lambda))}$.

Suppose that (b) and (c) hold. If H is a hyperplane contained in $\overline{\Omega(\lambda)} \cap \mathbf{Q}\Lambda$, there exists $i \notin \bigcap_{\beta \in \Delta} \beta$ with $x_i(\lambda) < 0$ and an integer b with $0 \leq b < -x_i(\lambda)$ such that

$$H = \left\{ \mu \in \mathbf{Q}\Lambda \mid x_j(\mu) \geq 0 \text{ for } j \notin \bigcap_{\beta \in \Delta} \beta \text{ and } x_i(\mu) = b \right\}.$$

We have $i \in \alpha$, so by (b) $\ker f_i = \ker x_i \cap \mathbf{Q}\Lambda$ is a bounding hyperplane of $\mathbf{Q}\Lambda_\beta$ for some $\beta \in \Delta$. Since $f_i(\mathbf{Z}\Lambda) = \mathbf{Z}$, we obtain $H \subset \overline{\omega_{\beta,i}(\phi(\lambda))} \subset \overline{\omega_\beta(\phi(\lambda))}$ as required.

Conversely, suppose that some induced hyperplane $\mathbf{Q}\Lambda \cap \ker x_i$ is not a bounding hyperplane of $\mathbf{Q}^+ \Lambda_\beta$ for any $\beta \in \Delta$. Choose $\lambda \in \mathbf{Z}\Lambda$ with $f_i(\lambda) < 0$. Then the hyperplane $\{\mu \in \mathbf{Q}\Lambda \mid f_i(\mu) = 0\}$ is contained in $\overline{\Omega(\lambda)} \cap \mathbf{Q}\Lambda$ but not in $\overline{\omega_\beta(\phi(\lambda))}$ for any β . Hence condition (b) is necessary for surjectivity.

Finally assume conditions (a) and (b) hold, but that (c) fails. Then for some induced hyperplane $\mathbf{Q}\Lambda \cap \ker x_i$ of $\mathbf{Q}^+ \Lambda_\beta, \beta \in \Delta$, we have $x_i(\mathbf{Z}\Lambda) = a\mathbf{Z}$ with $a > 1$. Choose $\lambda \in \mathbf{Z}\Lambda$ with $f_i(\lambda) < 0$. Then the hyperplane $\{\mu \in \mathbf{Q}\Lambda \mid f_i(\mu) = 1\}$ is contained in $\overline{\Omega(\lambda)} \cap \mathbf{Q}\Lambda$ but not in $\overline{\omega_\beta(\phi(\lambda))}$ for any β . This shows conditions (a)–(c) are necessary for surjectivity and completes the proof of Theorem B.

2.9. EXAMPLES. We first give some examples to illustrate the independence of conditions (a)–(c) of Theorem B, part 1. We suppose that $G = \text{GL}(1)^r$ acts on $\mathcal{O}(U_s) = k[T_1^\pm, \dots, T_s^\pm]$ by the rule $g \cdot T_j = \prod_i g_i^{a_{ij}} T_j$ where (a_{ij}) is an $r \times s$ matrix with integer entries.

(a) Suppose $r = 1, s = 2, (a_{ij}) = (1 \ -1)$, and $\Delta = \{\phi\}$. Then $T^\lambda \in \mathcal{O}(U_\phi) = k[T_1, T_2]$ is G -invariant if and only if $\lambda \in \mathbf{Z}^+(1, 1) = \Lambda$. Clearly the halfspaces in \mathbf{Q}^2 defined by $x_1 \geq 0$ and $x_2 \geq 0$ give the same induced halfspaces of $\mathbf{Q}\Lambda$. Hence condition (a) of Theorem B fails. It is easily verified that conditions (b) and (c) hold.

(b) Suppose $r = 1, s = 3, (a_{ij}) = (1 \ 1 \ -2)$, and $\Delta = \{\phi\}$. Then $T^\lambda \in \mathcal{O}(U_\phi) = k[T_1, T_2, T_3]$ is G -invariant if and only if λ belongs to the semigroup Λ generated by $(2, 0, 1), (0, 2, 1)$ and $(1, 1, 1)$. We note that the induced halfspace of $\mathbf{Q}\Lambda$ defined

by $x_3 \geq 0$ is not a bounding halfspace of $\mathbf{Q}^+ \Lambda$. Therefore condition (b) of Theorem B fails. It is easily checked that conditions (a), (c) of Theorem B hold. Note that if we let $\Delta = \{\{3\}\}$ instead, then $\mathcal{O}(U_{\{3\}}) = k[T_1, T_2, T_3^{\pm 1}]$ and the halfspace of $\mathbf{Q} \Lambda$ defined by $x_3 \geq 0$ is no longer an induced halfspace. Hence conditions (a)–(c) of Theorem B hold in this case. Also we have $\mathcal{O}(U_\phi)^G = \mathcal{O}(U_{\{3\}})^G$.

(c) Suppose $r = 1, s = 3, (a_{ij}) = \begin{pmatrix} 1 & 2 & -4 \end{pmatrix}$ and $\Delta = \{\{3\}\}$. Then $\mathcal{O}(U_{\{3\}})^G = k\Lambda$ where Λ is the semigroup generated by $(4, 0, 1), (2, 1, 1),$ and $(0, 2, 1)$. Conditions (a), (b) of Theorem B hold, but $x_1 \geq 0$ defines an induced halfspace of $\mathbf{Q}^+ \Lambda$ and $x_1(\mathbf{Z}\Lambda) = 2\mathbf{Z}$. Note also that G satisfies the dimension equality and contains the pseudoreflection $(-1, 1, 1)$ which fixes the hyperplane defined by $x_1 = 0$.

(d) Finally, we give an example to show that the definition of $D(Y_\Delta/G)$ depends on Δ . Let $r = 1, s = 3, (a_{ij}) = \begin{pmatrix} 1 & 1 & -2 \end{pmatrix}$. We have seen in (b) that $\mathcal{O}(U_\phi)^G = \mathcal{O}(U_{\{3\}})^G$ and the map $D(\mathcal{O}(U_{\{3\}}))^G \rightarrow D(\mathcal{O}(U_\phi))^G$ is surjective. Hence $P_1 P_2 Q_3^{-1} \in D(\mathcal{O}(U_{\{3\}}))^G$ induces a differential operator on $\mathcal{O}(U_\phi)^G$. On the other hand we have $\mathcal{O}(U_{\{1\}})^G = k[T_1^{-1} T_2, T_1^2 T_3]$ and $P_1 P_2 Q_3^{-1} \cdot T_1^{-1} T_2 = -T_1^{-2} T_3^{-1} \notin \mathcal{O}(U_{\{1\}})^G$. Hence $D(\mathcal{O}(U_\phi)^G) \not\subseteq D(\mathcal{O}(U_{\{1\}})^G)$. Now we can write $U_\phi = U_\phi \cup U_{\{1\}}$. Thus with $\Delta = \{\phi\}$ and $\Delta' = \{\phi, \{1\}\}$, we have $Y = \bigcup_{\beta \in \Delta} U_\beta = \bigcup_{\beta \in \Delta'} U_\beta$ and $D(Y_\Delta/G) \not\subseteq D(Y_{\Delta'}/G)$.

3. Centers.

3.1. We denote the center of a ring R by $Z(R)$, and the Ore quotient ring of R by $Q(R)$ when it exists. Let R be a subring of $D(U_s) = k[Q_1^{\pm 1}, \dots, Q_s^{\pm 1}, P_1, \dots, P_s]$ containing $A = k[Q_1 P_1, \dots, Q_s P_s]$. By Lemma 2.2, $R = \bigoplus_{\lambda \in \mathbf{Z}^s} R(\lambda)$ where $R(\lambda) = \{x \in R \mid [Q_i P_i, x] = \lambda_i x \text{ for } 1 \leq i \leq s\}$. Let $\Lambda = \{\lambda \in \mathbf{Z}^s \mid R(\lambda) \neq 0\}$, and $S = k[Q_1 P_1, \dots, Q_s P_s, Q^\lambda; \lambda \in \mathbf{Z}\Lambda]$. We denote by $\text{GK}(R)$ the Gelfand-Kirillov dimension of R ; see [KL].

LEMMA. *With the above notation $\text{GK}(R) = s + \text{rank}(\mathbf{Z}\Lambda)$.*

PROOF. We may assume that R is finitely generated as a k -algebra. Since $R \subset S$, we have $\text{GK}(R) \leq \text{GK}(S)$. A straightforward calculation shows that $\text{GK}(S) = s + \text{rank}(\mathbf{Z}\Lambda)$.

Suppose $\lambda_1, \dots, \lambda_r$ are \mathbf{Z} -linearly independent elements of Λ , where $r = \text{rank}(\mathbf{Z}\Lambda)$. Choose a finite-dimensional subspace W of R containing $1, Q_1 P_1, \dots, Q_s P_s$ and such that $W \cap R(\lambda_i) \neq 0$ for $i = 1, \dots, r$. Let $R_m(\lambda) = W^m \cap R(\lambda)$. For $\lambda = \sum a_i \lambda_i \in \sum \mathbf{Z}_+ \lambda_i$ define $|\lambda| = \sum a_i$. Clearly $R_{|\lambda|}(\lambda) \neq 0$ for all $\lambda \in \sum \mathbf{Z}_+ \lambda_i$. Since the number of monomials of degree at most d in s indeterminates is $\binom{s+d}{s}$ we see that $\dim R_m(\lambda) \geq \binom{s+m-|\lambda|}{s}$, for $m \geq |\lambda|$. Therefore $\dim W^m \geq \sum \binom{s+m-|\lambda|}{s}$, where the sum is over all $\lambda \in \sum \mathbf{Z}_+ \lambda_i$ with $|\lambda| \leq m$. By induction on r it is possible to show that the right side is bounded below by a polynomial in m of degree $s + r$. Hence $\text{GK}(R) \geq s + r$ and this proves the lemma.

3.2. COROLLARY. (i) *Suppose $G = \text{GL}(1)^r$ acts faithfully on k^s . Then $\text{GK}(D(Y)^G) = 2s - r$.*

(ii) *Let $Y = \bigcup U_\beta$ and suppose the common quotient field of the rings $\mathcal{O}(U_\beta)^G$ has transcendence degree n over k . Then $\text{GK}(D(Y_\Delta/G)) = 2n$.*

PROOF. This follows immediately from Lemma 3.1, and Theorems 2.4 and 2.7. It follows from [KL, Theorem 4.12(a)] and Lemma 3.1 that $Q(R)$ exists.

3.3. LEMMA. *With the above notation $Q(R) = Q(S)$.*

PROOF. Since $R \subset S$, and S is a domain, it suffices to show $S \subset Q(R)$. If $\lambda \in \Lambda$, there exists $\alpha \in A$, $\alpha \neq 0$ such that $Q^\lambda \alpha \in R$. Since $A \subset R$, it follows that $Q^\lambda \in Q(R)$. Hence $Q^\lambda \in Q(R)$ for $\lambda \in \mathbf{Z}\Lambda$ and $S \subset Q(R)$.

Now suppose g_1, \dots, g_r form a basis for the kernel of the restriction map $(\mathbf{Q}^s)^* \rightarrow (\mathbf{Q}\Lambda)^*$, and choose $h_1, \dots, h_t \in (\mathbf{Q}^s)^*$, $t = s - r$ so that $g_1, \dots, g_r, h_1, \dots, h_t$ form a basis of $(\mathbf{Q}^s)^*$. We regard the elements g_j and h_i as elements of $A = k[Q_1P_1, \dots, Q_sP_s]$. Then A is a polynomial ring in h_1, \dots, h_t over $C = k[g_1, \dots, g_r]$.

THEOREM C. *With the above notation $Z(R) = C$.*

PROOF. We show first that $C \subseteq Z(S)$. Since S is the twisted group algebra of $\{Q^\lambda | \lambda \in \mathbf{Z}\Lambda\}$ over A , it is enough to show that $[Q^\lambda, g_j] = 0$ for $\lambda \in \mathbf{Z}\Lambda$ and $1 \leq j \leq r$. Now

$$\begin{aligned} [Q^\lambda, g_j] &= Q^\lambda \{ g_j(Q_1P_1, \dots, Q_sP_s) - g_j(Q_1P_1 + \lambda_1, \dots, Q_sP_s + \lambda_s) \} \\ &= Q^\lambda \{ g_j(Q_1P_1, \dots, Q_sP_s) - g_j(Q_1P_1, \dots, Q_sP_s) - g_j(\lambda) \} = 0, \end{aligned}$$

as g_j is a linear polynomial and $g_j(\lambda) = 0$.

Since $C \subseteq R \subseteq S$, it is now sufficient to prove that $Z(R) \subseteq C$. Note that if $x \in Z(R)$, then $[\xi, x] = 0$ for all $\xi \in \text{Lie } \Gamma = kQ_1P_1 + \dots + kQ_sP_s$. Hence $x \in R(0) = A$.

We let \leq denote the natural partial order on $(\mathbf{Z}_+)^t$, that is for $\alpha, \beta \in (\mathbf{Z}_+)^t$, $\alpha \leq \beta$ if and only if $\alpha_i \leq \beta_i$ for $1 \leq i \leq t$.

Suppose $x = \sum_{\beta \in (\mathbf{Z}_+)^t} z_\beta h^\beta \in A$ is central in R where $z_\beta \in C$. If $x \notin C$, then $z_\beta \neq 0$ for some $\beta > 0$. We choose an element $\alpha \in (\mathbf{Z}_+)^t$ maximal under the partial order \leq such that $z_\alpha \neq 0$. Suppose $\alpha_i \neq 0$. Since the restrictions of h_1, \dots, h_t form a basis for $(\mathbf{Q}\Lambda)^*$, there exists $\lambda \in \mathbf{Z}\Lambda$ such that $h_j(\lambda) = 0$ if $j \neq i$ and $h_i(\lambda) \neq 0$. Since $x \in Z(R)$ and $S \subset Q(R)$ we have

$$0 = [x, Q^\lambda] = Q^\lambda \left\{ \sum_{\beta} z_\beta (h^\beta(Q_1P_1 + \lambda_1, \dots, Q_sP_s + \lambda_s) - h^\beta(Q_1P_1, \dots, Q_sP_s)) \right\}.$$

For any β we have

$$\begin{aligned} h^\beta(Q_1P_1 + \lambda_1, \dots, Q_sP_s + \lambda_s) - h^\beta(Q_1P_1, \dots, Q_sP_s) \\ = \beta_i h_i(\lambda) h^{\beta'}(Q_1P_1, \dots, Q_sP_s) + \text{lower degree terms} \end{aligned}$$

where $\beta' = (\beta_1, \dots, \beta_{i-1}, \beta_i - 1, \beta_{i+1}, \dots, \beta_t)$.

For $\beta \neq \alpha$, none of the above terms involves $h^{\alpha'}$ and it follows that the coefficient of $h^{\alpha'}$ in the above sum is $\alpha_i h_i(\lambda) z_\alpha \neq 0$. This contradiction shows that $Z(R) \subseteq C$ which proves the theorem.

3.4. We now specialize Theorem C to the rings $D(Y)^G$ and $D(Y_\Delta/G)$. The fixed ring $\mathcal{O}(U_s)^G$ is the group algebra of $\Lambda_s = \{\lambda \in \mathbf{Z}^s \mid m_i(\lambda) = 0\}$ where the $m_i \in (\mathbf{Q}^s)^*$ are given by equation (2) of §1. In the notation of Theorem 2.4, $\Lambda_s = \mathbf{Z}^s \cap V$. This gives the first part of the following corollary.

COROLLARY. *With the above notation*

- (i) $Z(D(Y)^G) = k[m_1, \dots, m_r]$.
- (ii) $Z(D(Y_\Delta/G)) = k$.

PROOF. Part (ii) follows from Theorem C and Theorem 2.7.

3.5. Recall that for $\lambda \in \mathbf{Z}^s \cap V$, $\text{ann}_A \overline{\Omega(\lambda)} = S_\lambda A$, where S_λ is defined after Lemma 2.5.

LEMMA. *Assume G satisfies the dimension equality on $\mathcal{O}(U_\beta)$ for $\beta \in \Delta$. For $1 \leq l \leq r$, let $I_l = \sum_{k=1}^l m_k A$. Then $S_\lambda A \cap I_l = S_\lambda I_l$.*

PROOF. If $Q_i P_i - j$ is a factor of S_λ then for some $\beta \in \Delta$ and $i \notin \beta$ we have $\lambda_i < 0$ and $0 \leq j < -\lambda_i$. Since $\lambda \in Q\Lambda$ and $x_i(\lambda) = \lambda_i$ we have $Q\Lambda \not\subseteq \ker x_i$, and clearly $Q\Lambda \not\subseteq \ker(x_i - j)$ if $j \neq 0$. Hence $(Q_i P_i - j) \notin I_r$ which is the ideal of functions of A that vanish on $Q\Lambda = V$. Therefore $(Q_i P_i - j) \notin I_l$. Since I_l is a prime ideal of A it follows that $S_\lambda \notin I_l$ and $S_\lambda A \cap I_l = S_\lambda I_l$.

3.6. Note that if G satisfies the dimension equality on $\mathcal{O}(U_\beta)$, $\beta \in \Delta$, then $I_r = I$ is the ideal used in the proof of Theorem B. We can now reformulate Theorem B in this case.

COROLLARY. *Assume G satisfies the dimension equality on $\mathcal{O}(U_\beta)$, $\beta \in \Delta$. Let $\theta: D(Y)^G \rightarrow D(Y_\Delta/G)$ be the natural map.*

- (1) θ is surjective if and only if
 - (a) *The induced halfspaces of $Q\Lambda$ are distinct and are precisely the bounding halfspaces of the convex cones $Q^+ \Lambda_\beta$, $\beta \in \Delta$.*
 - (b) *If $Q\Lambda \cap \mathcal{H}_j$ is a bounding halfspace of $Q^+ \Lambda_\beta$, then G contains no nonidentity pseudoreflection fixing the hyperplane defined by $x_j = 0$.*
- (2) $\text{Ker } \theta = D(Y)^G I = D(Y)^G (\text{Ker } \theta \cap Z(D(Y)^G))$.

PROOF. (1) follows immediately from Theorem B and Lemma 1.9.

(2) Theorem B gives $\text{Ker } \theta = D(U_s)^G I \cap D(Y)^G$. In particular $I \subseteq \text{Ker } \theta$. Also $I \subseteq Z(D(Y)^G)$ by Corollary 3.4. If $g \in A$ and $Q^\lambda g \in (\text{Ker } \theta)(\lambda)$, we have $g \in S_\lambda A \cap I = S_\lambda I$. Therefore $Q^\lambda g \in Q^\lambda S_\lambda I \subseteq D(Y)^G I$. Together with Lemma 2.2 this proves (2).

3.7. In later applications we shall often require the conditions of the corollary to be satisfied. It is important to know that we can do this without changing the ring $D(Y_\Delta/G)$ up to isomorphism.

PROPOSITION. *Suppose the torus G acts on $Y = \bigcup_{\beta \in \Delta} U_\beta$. Then there is a set of subsets Σ of s such that if $X = \bigcup_{\beta \in \Sigma} U_\beta$, then there is an action of a possibly larger torus G' on X such that*

- (i) $D(X_\Sigma/G') \cong D(Y_\Delta/G)$.
- (ii) G' satisfies the dimension equality on $\mathcal{O}(U_\beta)$, $\beta \in \Sigma$.
- (iii) The map $\theta: D(X)^G \rightarrow D(X_\Sigma/G')$ is surjective.

PROOF. We define an equivalence relation \sim on $\alpha = \alpha(\Delta)$ by $i \sim j$ for $i, j \in \alpha$ if and only if $f_i = af_j$ for some $a > 0$. Let γ be a set of representatives for the equivalence classes of \sim . For $\beta \in \Delta$, we can choose a set β' containing β such that the bounding halfspaces of $\mathbf{Q}^+ \Lambda_\beta = \mathbf{Q}^+ \Lambda_{\beta'}$ have the form $\{\mathbf{Q}\Lambda \cap \mathcal{H}_j \mid j \in \mathfrak{s} \setminus \beta'\}$ and $(\mathfrak{s} \setminus \beta') \subseteq \gamma$. Let $\Sigma = \{\beta' \mid \beta \in \Delta\}$. If $X = \bigcup_{\beta' \in \Sigma} U_{\beta'}$, then G acts on X , conditions 1(a) and (b) of Theorem B now hold, and we have $\mathcal{O}(U_{\beta'})^G = \mathcal{O}(U_\beta)^G$ for all β . By embedding G in a larger torus, we may assume the dimension equality holds on $\mathcal{O}(U_\beta)$ for all $\beta \in \Sigma$. Since these reductions do not change the fixed rings we have $\bigcap_{\beta \in \Delta} D(\mathcal{O}(U_\beta)^G) = \bigcap_{\beta' \in \Sigma} D(\mathcal{O}(U_{\beta'})^G)$.

To obtain condition 1(c) of Theorem B, we apply Lemma 1.8. Since we replace $\mathcal{O}(U_{\beta'})^G$ by an isomorphic ring some care must be taken. Suppose $\phi: \mathbf{Q}^s \rightarrow \mathbf{Q}^s$ is the isomorphism of Lemma 1.8. We obtain an isomorphism, also denoted by ϕ from $k\mathbf{Z}\Lambda$ to $k\phi\mathbf{Z}\Lambda$. The map $\phi^*: D(k\mathbf{Z}\Lambda) \rightarrow D(k\phi\mathbf{Z}\Lambda)$ defined by $(\phi^*d)(f) = \phi(d(\phi^{-1}f))$ is an isomorphism, and using Lemma 2.1, we can show that ϕ^* maps $\bigcap_{\beta' \in \Sigma} D(k\Lambda_{\beta'})$ onto $\bigcap_{\beta' \in \Sigma} D(k\phi\Lambda_{\beta'})$.

4. Graded rings.

4.1. We filter the rings $D(Y)^G$ and $D(Y_\Delta/G)$ by the order of the differential operators. In $D(U_s)$ we let \mathcal{E}_t denote the span of all differential operators of order at most t , and $\mathcal{D}_t = \mathcal{E}_t \cap D(Y)$. Then \mathcal{E}_t is the free $\mathcal{O}(U_s)$ -module generated by monomials in Q_1P_1, \dots, Q_sP_s of degree at most t . We also set $\mathcal{D}_{-1} = \mathcal{E}_{-1} = 0$. Then

$$\text{Gr} D(Y) = \bigoplus_{t \geq 0} \frac{\mathcal{D}_t}{\mathcal{D}_{t-1}} \subseteq \text{Gr} D(U_s) = \bigoplus_{t \geq 0} \frac{\mathcal{E}_t}{\mathcal{E}_{t-1}}.$$

The action of G preserves the filtrations $\{\mathcal{D}_t\}$ and $\{\mathcal{E}_t\}$, and hence G acts on $\text{Gr} D(Y)$ by

$$g \cdot (a + \mathcal{D}_{t-1}) = (g \cdot a + \mathcal{D}_{t-1}) \quad \text{for } a \in \mathcal{D}_t.$$

We may filter $D(Y)^G$ by the subspaces $\mathcal{D}_t \cap D(Y)^G$, and it follows that

$$\begin{aligned} \text{Gr}(D(Y)^G) &= \bigoplus_{t \geq 0} \frac{D(Y)^G \cap \mathcal{D}_t}{D(Y)^G \cap \mathcal{D}_{t-1}} \cong \bigoplus_{t \geq 0} \frac{(D(Y)^G \cap \mathcal{D}_t) + \mathcal{D}_{t-1}}{\mathcal{D}_{t-1}} \\ &= (\text{Gr} D(Y))^G. \end{aligned}$$

In the ring $D(k\mathbf{Z}\Lambda) = k[q_1^{\pm 1}, \dots, q_n^{\pm 1}, p_1, \dots, p_n]$ we let E_t denote the span of all differential operators of order at most t , and $D_t = E_t \cap D(Y_\Delta/G)$. Then E_t is the free $k\mathbf{Z}\Lambda$ -module generated by monomials in q_1p_1, \dots, q_np_n of degree at most t . As before

$$\text{Gr}(D(Y_\Delta/G)) = \bigoplus D_t/D_{t-1} \subseteq \bigoplus E_t/E_{t-1} = \text{Gr}(D(k\mathbf{Z}\Lambda)).$$

It follows from the proof of Theorem B that the map $\theta: D(Y)^G \rightarrow D(Y_\Delta/G)$ preserves the filtrations defined above, that is $\theta(\mathcal{D}_t \cap D(Y)^G) \subseteq D_t$ for all $t \geq 0$. Hence there is an induced map $\text{Gr}\theta: \text{Gr}(D(Y)^G) \rightarrow \text{Gr}(D(Y_\Delta/G))$ defined by $\text{Gr}\theta(a + (\mathcal{D}_{t-1} \cap D(Y)^G)) = \theta(a) + D_{t-1}$ for $a \in \mathcal{D}_t \cap D(Y)^G$. It is routine to

check that $\text{Gr } \theta$ is surjective if and only if θ is surjective and that the diagram

$$\begin{CD} D(Y)^G @>\theta>> D(Y_{\Delta}/G) \\ @V\Pi_1VV @VV\Pi_2V \\ \text{Gr } D(Y)^G @>\text{Gr } \theta>> \text{Gr } D(Y_{\Delta}/G) \end{CD}$$

commutes. Here if $a \in (\mathcal{D}_t \cap D(Y)^G) \setminus (\mathcal{D}_{t-1} \cap D(Y)^G)$ we define $\Pi_1(a) = a + (\mathcal{D}_{t-1} \cap D(Y)^G)$. The map Π_2 is defined similarly.

THEOREM. *The graded rings $\text{Gr}(D(Y)^G) = (\text{Gr } D(Y))^G$ and $\text{Gr}(D(Y_{\Delta}/G))$ are finitely generated commutative k -algebras.*

PROOF. The graded ring of $\text{Gr } D(Y) \cong k[X_1, \dots, X_s, Y_1, \dots, Y_s, X_i^{-1}; i \in \bigcap_{\beta \in \Delta} \beta]$ is a localization of a commutative polynomial ring at some of the indeterminates, and $G = \text{GL}(1)^r$ acts linearly on the subspace spanned by $X_1, \dots, X_s, Y_1, \dots, Y_s$. A well-known result from invariant theory, [Hu, §14, Exercise 2] implies that the fixed ring $(\text{Gr } D(Y))^G$ is finitely generated as a k -algebra.

To show that $\text{Gr } D(Y_{\Delta}/G)$ is finitely generated, we may assume the map θ is surjective by Proposition 3.7. Hence $\text{Gr } \theta$ is also surjective and the result follows.

4.2. The following is well known; see [V, I, Proposition 7] for example.

PROPOSITION. *Let R be a k -algebra which has a filtration $k \subseteq R_0 \subseteq R_1 \subseteq \dots \subseteq R_n \subseteq \dots \subseteq \bigcup_{n \geq 0} R_n = R$ such that the associated graded ring $\text{Gr } R = \bigoplus_{n \geq 0} R_n/R_{n-1}$ is a finitely generated commutative k -algebra. Then*

- (i) R is (left and right) Noetherian.
- (ii) R is finitely generated as a k -algebra.

Theorem A now follows immediately from Theorem 4.1 and Proposition 4.2.

4.3. We next show that the kernel of $\text{Gr } \theta$ is $\Pi_1(\text{Ker } \theta)$. For this we need the following lemma.

LEMMA. *Let $\psi: \mathbf{Q}^s \rightarrow \mathbf{Q}^n$ be an epimorphism of vector spaces, and extend ψ to a map $k[X_1, \dots, X_s] \rightarrow k[x_1, \dots, x_n]$, also denoted by ψ , where X_1, \dots, X_s and x_1, \dots, x_n are bases for \mathbf{Q}^s and \mathbf{Q}^n respectively. If $g = g(X_1, \dots, X_s)$ is a homogeneous polynomial of degree t , and $\deg \psi g < t$, then $\psi g = 0$.*

PROOF. We can by renumbering assume that $\psi X_1, \dots, \psi X_n$ form a basis for \mathbf{Q}^n . Then for $n + 1 \leq i \leq s$ we have

$$X_i - \sum_{j=1}^n a_{ij} X_j \in \text{Ker } \psi \quad \text{for some } a_{ij} \in k.$$

Hence we may write

$$\begin{aligned} g(X_1, \dots, X_s) &= g\left(X_1, \dots, X_n, \sum_{j=1}^n a_{n+1j} X_j, \dots, \sum_{j=1}^n a_{sj} X_j\right) + g_2(X_1, \dots, X_s) \\ &= g_1(X_1, \dots, X_n) + g_2(X_1, \dots, X_s) \end{aligned}$$

where $g_2 \in \ker \psi$ and g_1 is either zero or a homogeneous polynomial of degree t in X_1, \dots, X_n . The result follows since the restriction of ψ to $k[X_1, \dots, X_n]$ is an isomorphism which preserves degrees.

4.4. THEOREM. $\text{Ker Gr } \theta = \Pi_1(\text{Ker } \theta)$.

PROOF. It is clear that $\Pi_1(\text{Ker } \theta) \subseteq \text{Ker Gr } \theta$. Suppose $y \in \text{Ker Gr } \theta$. We may assume that $y = \Pi_1(x)$, where $x = \sum_{\lambda \in \mathbf{Z}^s \cap V} Q^\lambda g_\lambda \in D(Y)^G$ and each g_λ is a homogeneous polynomial of degree t in $Q_1 P_1, \dots, Q_s P_s$. Then $\theta(Q^\lambda g_\lambda) = 0$ for $\lambda \notin \mathbf{Z}\Lambda$, so $\theta(\sum_{\lambda \in \mathbf{Z}^s \cap V} Q^\lambda g_\lambda) = \sum_{\lambda \in \mathbf{Z}\Lambda} q^{\phi(\lambda)} \theta(g_\lambda)$. Since $y \in \text{Ker Gr } \theta$ we have $\theta(x) \in D_{t-1}$, so each polynomial $\theta(g_\lambda)$ has degree $\leq t - 1$ in the variables $q_1 p_1, \dots, q_n p_n$. By the proof of Theorem B, the restriction of θ to a map from A to B is a map of the type described in Lemma 4.3. It follows that $\theta(g_\lambda) = 0$ for each λ and so $x \in \text{Ker } \theta$ and $y \in \Pi_1(\text{Ker } \theta)$.

4.5. We draw a further consequence from this. Recall that a sequence of elements x_1, \dots, x_r in a commutative ring R is an R -sequence provided $\sum_{i=1}^r x_i R \neq R$ and for each $i = 1, \dots, r$ the image of x_i in $R/\sum_{j=1}^{i-1} x_j R$ is a nonzero divisor (see [K, 3.1]).

LEMMA. Assume G satisfies the dimension equality on $\mathcal{O}(U_\beta)$ for $\beta \in \Delta$. Then $\text{Ker Gr } \theta$ is generated by an R -sequence.

PROOF. We can write $A = k[m_1, \dots, m_r, h_1, \dots, h_t]$ where m_1, \dots, m_r generate the ideal of A consisting of functions which vanish on $\mathbf{Q}\Lambda$. Then $D(U_s)^G = \bigoplus_{\lambda \in \mathbf{Z}\Lambda} D(U_s)^G(\lambda)$ where $D(U_s)^G(\lambda) = Q^\lambda A$. Let $X^\lambda = \Pi_1(Q^\lambda)$, $Z_i = \Pi_1(m_i)$ for $1 \leq i \leq r$, and $Z_i = \Pi_1(h_{i-r})$ for $r + 1 \leq i \leq r + t = s$. Then

$$\begin{aligned} \text{Gr } D(U_s)^G &= k[X^\lambda, Z_i, \lambda \in \mathbf{Z}\Lambda, 1 \leq i \leq s] \\ &= k[X^\lambda][Z_1, \dots, Z_s] \text{ a polynomial ring over } k[X^\lambda]. \end{aligned}$$

Now $D(Y)^G$ is a subring of $D(U_s)^G$ and we have

$$\text{Gr } D(Y)^G = \bigoplus_{\lambda \in \mathbf{Z}\Lambda} X^\lambda \Pi_1(S_\lambda) k[Z_1, \dots, Z_s].$$

By Theorem 4.4 and Corollary 3.4, $\text{Ker Gr } \theta$ is the ideal of $\text{Gr } D(Y)^G$ generated by Z_1, \dots, Z_r . By the proof of Lemma 3.5 we have

$$\Pi_1(S_\lambda) k[Z_1, \dots, Z_s] \cap \sum_{i=1}^l Z_i k[Z_1, \dots, Z_s] = \Pi_1(S_\lambda) \sum_{i=1}^l Z_i k[Z_1, \dots, Z_s]$$

for $1 \leq l \leq r$ and hence

$$\sum_{i=1}^l \text{Gr } D(U_s)^G Z_i \cap \text{Gr } D(Y)^G = \sum_{i=1}^l \text{Gr } D(Y)^G Z_i.$$

Since the elements Z_1, \dots, Z_r form an R -sequence in the ring $\text{Gr } D(U_s)^G$ it follows from the equality above that they form an R -sequence in $\text{Gr } D(Y)^G$.

4.6. Our next aim is to show that the graded rings $\text{Gr } D(Y)^G$ and $\text{Gr } D(Y_\Delta/G)$ are Gorenstein rings. We have

$$\text{Gr } D(Y) = k \left[X_1, \dots, X_s, Y_1, \dots, Y_s, X_i^{-1}; i \in \bigcap_{\beta \in \Delta} \beta \right]$$

and G acts on $V = \text{span}\{X_1, \dots, X_s, Y_1, \dots, Y_s\}$ as a subgroup of $\text{SL}(V)$. Hence the fact that $D(Y)^G$ is Gorenstein follows from the following which is a slight extension of a result of R. P. Stanley [S, p. 81].

THEOREM. *Assume the torus G acts as a diagonal subgroup of $\text{SL}(V)$ where $V = k^s$. For a fixed subset β of $\{1, 2, \dots, s\}$ let $U = \{v \in V \mid v_i \neq 0 \text{ if } i \in \beta\}$. Then $\mathcal{O}(U)^G$ is a Gorenstein ring.*

We can assume that $\mathcal{O}(U) = k[T_1^{\pm 1}, \dots, T_t^{\pm 1}, T_{t+1}, \dots, T_s]$, and that as explained in §1, $\mathcal{O}(U)^G$ is a semigroup algebra $k\Lambda$ where Λ has the form

$$\Lambda = \{ b \in \mathbf{Z}^t \times (\mathbf{Z}_+)^{s-t} \mid f_i(b) = 0 \text{ for } 1 \leq i \leq r \}$$

for certain linear equations $f_i(x_1, \dots, x_s) = \sum_{j=1}^s a_{ij}x_j \in (\mathbf{Q}^s)^*$.

A subsemigroup M of \mathbf{Z}_+^s will be called a *positive toroidal semigroup* if it is the set of solutions in nonnegative integers to a system of homogeneous linear equations with integer coefficients and there exists an element $m = (m_1, \dots, m_s) \in M$ with $m_i > 0$ for each i . If M is a positive toroidal semigroup, an element $m = (m_1, \dots, m_s) \in M$ is a *minimal positive element* provided $m_i > 0$ for each i , and whenever $n = (n_1, \dots, n_s) \in M$ with $n_i > 0$ for each i , we have $m_i \leq n_i$. Theorem 4.6 is deduced from the following result of Stanley.

4.7. THEOREM. *If M is a positive toroidal semigroup, then kM is a Gorenstein ring if and only if M has a unique minimal positive element.*

PROOF. [S, Theorem 6.7].

4.8. LEMMA. *If $\mathcal{O}(U)^G = k\Lambda$ where $G \subseteq \text{SL}(V)$ as in Theorem 4.6 then $k\Lambda = (kM)_{\mathcal{Q}}$ is a localization of a semigroup algebra kM , where M is isomorphic to a positive toroidal semigroup with a unique minimal positive element.*

Together with Theorem 4.7, and the fact that a localization of a Gorenstein ring is again a Gorenstein ring [K, Exercise 12, p. 164], this will prove Theorem 4.6.

PROOF OF LEMMA 4.8. The hypothesis $G \subseteq \text{SL}(V)$ is equivalent to the fact that $\sum_{j=1}^s a_{ij} = 0$ for $1 \leq i \leq r$. That is f_1, \dots, f_r belong to the subspace W of $(\mathbf{Q}^s)^*$ spanned by $x_1 - x_2, x_2 - x_3, \dots, x_{s-1} - x_s$. Let (b_{ij}) be the row reduced form of the matrix (a_{ij}) and $g_i(x) = \sum b_{ij}x_j$. Then $g_i \in W$ for each i , and Λ is the set of solutions in $\mathbf{Z}^t \times (\mathbf{Z}_+)^{s-t}$ to the equations $g_i(x) = 0$.

As usual the variables x_j corresponding to a column of (b_{ij}) containing a leading one are called leading variables, and the others are nonleading variables. We can assign the nonleading variables arbitrary values and solve for the leading variables. Moreover the value of a leading variable x_j is determined by the values of the nonleading variables x_k with $k \geq j$. Therefore we can assign a value zero to every variable x_j with $j \geq t + 1$ and assign the remaining nonleading variables arbitrary values.

For each nonleading variable x_j with $1 \leq j \leq t$, we obtain in this way a solution $\beta_j = (\beta_{j1}, \dots, \beta_{js}) \in \Lambda$ with β_{jj} a positive integer and $\beta_{jk} = 0$ if $k \geq t + 1$ or x_k is any nonleading variable with $k \neq j$.

Let M be the subsemigroup of Λ given by

$$M = \{ b \in \mathbf{Z}^t \times (\mathbf{Z}_+)^{s-t} \mid f_i(b) = 0, 1 \leq i \leq r \text{ and } b_j \geq 0 \}$$

if x_j is a nonleading variable with $1 \leq j \leq t$

and let \mathcal{C} be the multiplicatively closed subset of kM generated by the set $\{T^{B_j}\}$. Then it follows easily that $k\Lambda = (kM)_{\mathcal{C}}$.

It remains to show that M has the properties claimed. By changing notation, we may assume that

$$M = \{ b \in \mathbf{Z}^p \times (\mathbf{Z}_+)^{s-p} \mid b_i = f_i(b_{p+1}, \dots, b_s) \text{ for } 1 \leq i \leq p \}$$

and $g_j(b_{p+1}, \dots, b_s) = 0, j \in J$

where as before $x_i - f_i(x_{p+1}, \dots, x_s)$ and $g_j(x_{p+1}, \dots, x_s)$ belong to W . To see this first regroup all the variables which are constrained to be nonnegative at the end, and relabel them as x_{p+1}, \dots, x_s . Then relabel the leading variables x_j with $1 \leq j \leq t$ as x_1, \dots, x_p and express their dependence on x_{p+1}, \dots, x_s by the equations $x_i = f_i(x_{p+1}, \dots, x_s)$. Any further relations between the x_{p+1}, \dots, x_s are expressed by certain equations $g_j(x_{p+1}, \dots, x_s) = 0$, where j runs over some index set J .

Suppose $f_i(x_{p+1}, \dots, x_s) = \sum_{k=p+1}^s c_{ik} x_k$ where $c_{ik} \in \mathbf{Q}$ and choose $m \in \mathbf{Z}$ such that $m + c_{ik} > 0$ for each i and k . For $1 \leq i \leq p$ define $h_i(x) = x_i + m \sum_{k=p+1}^s x_k$ and let $\zeta: \mathbf{Q}^s \rightarrow \mathbf{Q}^s$ be the linear map defined by

$$\zeta(b_1, \dots, b_s) = (h_1(b), \dots, h_p(b), b_{p+1}, \dots, b_s).$$

It is easily checked that ζ is an isomorphism and that

$$\zeta(M) = \left\{ (c_1, \dots, c_s) \in \mathbf{Z}_+^s \mid c_i = f_i(c_{p+1}, \dots, c_s) + m \sum_{k=p+1}^s c_k \right.$$

for $1 \leq i \leq p$ and $g_j(c_{p+1}, \dots, c_s) = 0, j \in J$.

Hence M is isomorphic to the positive toroidal semigroup $\zeta(M)$. For $1 \leq i \leq p$, let $a_i = f_i(1, 1, \dots, 1) + m(s - p)$. Then since $g_j \in W$ it follows that $(a_1, \dots, a_p, 1, 1, \dots, 1)$ is the unique minimal positive element of $\zeta(M)$. This completes the proof.

THEOREM D. *The graded rings $\text{Gr } D(Y)^G$ and $\text{Gr } D(Y_{\Delta}/G)$ are Gorenstein rings.*

PROOF. The fact that $\text{Gr } D(Y)^G$ is Gorenstein now follows directly from Theorem 4.6. To prove $\text{Gr } D(Y_{\Delta}/G)$ is Gorenstein we may assume by Proposition 3.7 that G satisfies the dimension equality on $\mathcal{O}(U_{\beta})$ for $\beta \in \Delta$, and the map $\theta: D(Y)^G \rightarrow D(Y_{\Delta}/G)$ is surjective. Hence the map $\text{Gr } \theta: \text{Gr } D(Y)^G \rightarrow \text{Gr } D(Y_{\Delta}/G)$ is surjective, and by Lemma 4.5 the kernel of $\text{Gr } \theta$ is generated by an R -sequence. It follows that $\text{Gr } D(Y_{\Delta}/G)$ is Gorenstein by [K, Exercise 13, p. 164].

4.9. The fact that these graded rings are Gorenstein can be used to apply a result of Levasseur, which we explain next.

Let A be a filtered ring such that the associated graded ring B is commutative Noetherian with $\text{Kd}(B) = \omega$, where Kd denotes Krull dimension. If M is a finitely generated A -module, we may filter M in such a way that the associated graded module $\text{Gr}(M)$ is a finitely generated $B = \text{Gr}(A)$ -module, and we set $d(M) = \text{Kd}({}_B\text{Gr}(M))$. We set $\text{grade}(M) = \inf\{i \mid \text{Ext}_A^i(M, A) \neq 0\}$. We say that B is of *pure dimension* ω if for every maximal ideal m of B , $\text{Kd}(B_m) = \omega$. The next result is [L2, Théorème 4.4].

THEOREM. *Suppose that B is a Gorenstein ring of pure dimension ω . Then for any finitely generated, nonzero A -module M , we have $d(M) + \text{grade}(M) = \omega$.*

In addition the left and right injective dimensions of A are finite and equal to $\mu \leq \omega$.

4.10. We note that if B is a finitely generated domain over a field k , and $\text{Kd}(B) = \omega$, then B has pure dimension ω . This follows from Noether's normalization lemma and the going up and going down theorems.

It is therefore of interest to compute the Krull dimensions of the graded rings of Theorem D. More generally in the next lemma we assume that R is a subring of $D(U_s) = k[Q_1^{\pm 1}, \dots, Q_s^{\pm 1}, P_1, \dots, P_s]$ containing $A = k[Q_1P_1, \dots, Q_sP_s]$. By Lemma 2.2, $R = \bigoplus_{\lambda \in \mathbf{Z}^s} R(\lambda)$ where $R(\lambda) = \{x \in R \mid [Q_i, P_i, x] = \lambda_i x\}$. Let $\Lambda = \{\lambda \in \mathbf{Z}^s \mid R(\lambda) \neq 0\}$, and suppose $\text{rank}(\mathbf{Z}\Lambda) = r$.

Let \mathcal{E}_t be the span of all differential operators of order at most t in $D(U_s)$ and $\text{Gr}(D(U_s)) = \bigoplus \mathcal{E}_t/\mathcal{E}_{t-1}$. If $x \in \mathcal{E}_t - \mathcal{E}_{t-1}$ we define $\Pi(x) = (x + \mathcal{E}_t)/\mathcal{E}_{t-1}$. Then $\text{Gr}(R) = \Pi(R)$ is a subring of $\text{Gr}(D(U_s))$.

If $\text{Gr}(R)$ is a finitely generated k -algebra, $\text{Kd}(\text{Gr}(R)) = \text{tr deg}_k(\text{Gr}(R))$, the transcendence degree of $\text{Gr}(R)$ over k . We compute this next.

LEMMA. *With the above notation $\text{tr deg}_k(\text{Gr}(R)) = s + r$.*

PROOF. Let $S = k[Q_1P_1, \dots, Q_nP_n, Q^\lambda, \lambda \in \mathbf{Z}\Lambda]$. Then $R \subset S$, $\text{Gr}(R) \subset \text{Gr}(S)$, and $\text{tr deg}_k(\text{Gr}(S)) = s + r$. It suffices to show that $\text{Gr}(R)$ and $\text{Gr}(S)$ have the same quotient fields, and this is proved in the same way as Lemma 3.3.

4.11. **COROLLARY.** (i) *Suppose $G = \text{GL}(1)^r$ acts faithfully on k^s . Then $\text{Kd Gr } D(Y)^G = 2s - r$.*

(ii) *Let $Y = \bigcup U_\beta$ and suppose the common quotient field of the rings $\mathcal{O}(U_\beta)^G$ has transcendence degree n over k . Then $\text{Kd Gr } D(Y_\Delta/G) = 2n$.*

PROOF. Since $\text{Gr } D(Y)^G$ and $\text{Gr } D(Y_\Delta/G)$ are finitely generated k -algebras by Theorem 4.1, this follows from Lemma 4.10 and Theorems 2.4 and 2.7.

REFERENCES

[BB] W. Borho and J.-L. Brylinski, *Differential operators on homogeneous spaces*. I, *Invent. Math.* **69** (1982), 437–476.
 [B] A. Brondsted, *An introduction to convex polytopes*, Graduate Texts in Math., no. 90, Springer-Verlag, Berlin, Heidelberg and New York, 1982.
 [H] R. Hart, *Differential operators on affine algebras*, *J. London Math. Soc.* (2) **28** (1983), 470–476.

[Ho] M. Hochster, *Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes*, Ann. of Math. (2) **96** (1972), 318–337.

[Hu] J. E. Humphreys, *Linear algebraic groups*, Graduate Texts in Math., no. 21, Springer-Verlag, Berlin, Heidelberg, and New York, 1975.

[Ka] J. M. Kantor, *Formes et opérateurs différentiels sur les espaces analytiques complexes*, Bull. Soc. Math. France **53** (1977), 5–80.

[K] I. Kaplansky, *Commutative rings*, Univ. of Chicago Press, Chicago and London, 1974.

[Ke] G. Kempf et al., *Toroidal embeddings. I*, Lecture Notes in Math., no. 339, Springer-Verlag, Berlin, Heidelberg, and New York, 1973.

[KL] G. R. Krause and T. H. Lenagan, *Growth of algebras and Gelfand-Kirillov dimension*, Research Notes in Math., No. 116, Pitman, Boston.

[L1] T. Levasseur, *Anneaux d'opérateurs différentiels*, Séminaire d'Algèbre Paul Dubreil et Marie-Paule Malliavin, Lecture Notes in Math., no. 867, Springer-Verlag, Berlin, Heidelberg, and New York, 1981, pp. 157–173.

[L2] _____, *Complexe bidualisant en algèbre non-commutative*, Séminaire d'Algèbre Paul Dubreil et Marie-Paule Malliavin, Lecture Notes in Math., no. 1146, Springer-Verlag, Berlin, Heidelberg, and New York, 1985, pp. 397–415.

[M] I. M. Musson, *Actions of tori on Weyl algebras*, Comm. Algebra (to appear).

[S] R. P. Stanley, *Hilbert functions of graded algebras*, Adv. in Math. **28** (1978), 57–83.

[V] J. P. Vigué, *Opérateurs différentiels sur les espaces analytiques*, Invent. Math. **20** (1973), 313–336.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MILWAUKEE, MILWAUKEE, WISCONSIN 53201