

## THE BLOW-UP SURFACE FOR NONLINEAR WAVE EQUATIONS WITH SMALL SPATIAL VELOCITY

AVNER FRIEDMAN AND LUC OSWALD

ABSTRACT. Consider the Cauchy problem for  $u_{tt} - \varepsilon^2 \Delta u = f(u)$  in space dimension  $\leq 3$  where  $f(u)$  is superlinear and nonnegative. The solution blows up on a surface  $t = \phi_\varepsilon(x)$ . Denote by  $t = \phi(x)$  the blow-up surface corresponding to  $v'' = f(v)$ . It is proved that  $|\phi_\varepsilon(x) - \phi(x)| \leq C\varepsilon^2$ ,  $|\nabla(\phi_\varepsilon(x) - \phi(x))| \leq C\varepsilon^2$  in a neighborhood of any point  $x_0$  where  $\phi(x_0) < \infty$ .

### 1. The main results. Let

$$\Delta = \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2}, \quad \square_\varepsilon = \frac{\partial^2}{\partial t^2} - \varepsilon^2 \Delta \quad (\varepsilon > 0)$$

and consider the Cauchy problem

$$(1.1) \quad \square_\varepsilon u_\varepsilon = f(u_\varepsilon) \quad \text{in } \mathbf{R}^N \times (0, \infty),$$

$$(1.2) \quad u_\varepsilon(x, 0) = g(x), \quad x \in \mathbf{R}^N,$$

$$(1.3) \quad \frac{\partial}{\partial t} u_\varepsilon(x, 0) = h(x), \quad x \in \mathbf{R}^N.$$

Here  $f(u)$  is a superlinear function such as  $(u^+)^p$ ; more generally we shall assume that  $f \geq 0$ ,  $f \in C^4(\mathbf{R})$ ; there exists a  $u_0 \geq 0$  such that

$$(1.4) \quad \begin{aligned} & f(u) > 0, \quad f'(u) \geq 0, \quad f''(u) \geq 0 \quad \text{if } u \geq u_0; \\ & (f(u)/u^p) \rightarrow 1 \quad \text{if } u \rightarrow \infty, \quad p > 1; \\ & \limsup_{u \rightarrow \infty} (f'(u)/u^{p-1}) < p + (p-1)/2; \\ & \liminf_{u \rightarrow \infty} (f'(u)/u^{p-1}) > 0; \\ & |f^{(j)}(u)| \leq Cu^{p-j} \quad \text{if } u \geq u_0, \quad 2 \leq j \leq 4. \end{aligned}$$

We also assume that

$$(1.5) \quad N \leq 3$$

and

$$(1.6) \quad \begin{aligned} & g \in C^5(\mathbf{R}^N), \quad h \in C^4(\mathbf{R}^N) \quad \text{if } N = 2, 3; \\ & g, h \in C^4(\mathbf{R}^1) \quad \text{if } N = 1. \end{aligned}$$

---

Received by the editors March 30, 1987 and, in revised form, June 8, 1987.  
 1980 *Mathematics Subject Classification* (1985 Revision). Primary 35L70; Secondary 35L05, 35L67.

This work is partially supported by National Science Foundation Grants DMS-8420896 and DMS-8501397.

Sufficient conditions for nonexistence of global solutions of (1.1)–(1.3) are given in [3–7]. In this paper we are interested in the behavior of (blowing-up) solutions  $u_\varepsilon$  of (1.1)–(1.3) as  $\varepsilon \rightarrow 0$ . This is naturally related to the behavior of the solutions of the ordinary differential equation

$$(1.7) \quad \frac{d^2u}{dt^2} = f(u) \quad \text{for } t > 0$$

under the Cauchy conditions

$$(1.8) \quad u(x, 0) = g(x), \quad u_t(x, 0) = h(x).$$

For each fixed  $x$  the solution of (1.7), (1.8) exists for  $0 < t < \phi(x)$  where either  $\phi(x) = \infty$  or  $\phi(x) < \infty$ ; in the latter case it can be shown (see §2) that  $u(x, t) \rightarrow \infty$  if  $t \rightarrow \phi(x)$ , and we say that  $u(x, t)$  *blows up* at time  $t = \phi(x)$ . The surface  $\{t = \phi(x)\}$  is called the *blow up surface* for  $u$ .

Caffarelli and Friedman [1] proved that if  $N = 1$  then there exists a unique classical solution of (1.1)–(1.3) for all  $0 < t < \phi_\varepsilon(x)$  where either  $\phi_\varepsilon(x) \equiv \infty$  (no blow-up) or else  $\phi_\varepsilon(x) < \infty$  for all  $x \in \mathbf{R}^1$  and  $\phi_\varepsilon \in C^1$ ,  $|\phi'_\varepsilon(x)| < 1/\varepsilon$ ; further,  $u_\varepsilon(x, t) \rightarrow \infty$  if  $t \rightarrow \phi_\varepsilon(x)$ . In [2] they extended these results to  $N = 2, 3$  under some restrictions on the Cauchy data (in addition to (1.6)). We shall recall a slightly simplified version of their result in case  $\varepsilon = 1$ ; this will be needed in the sequel.

Introduce the sets

$$\begin{aligned} K^\varepsilon(x_0, t_0) &= \{(x, t); |x - x_0| \leq \varepsilon(t_0 - t), 0 \leq t < t_0\}, \\ B_R(x_0) &= \{|x - x_0| < R\}, \quad B_R = B_R(0), \\ K_{R,T}^\varepsilon &= \bigcup_{x \in B_R} K^\varepsilon(x, T). \end{aligned}$$

We shall assume, in addition to (1.5), (1.6), the following conditions:

(1.9) the solution  $w$  of  $w''(t) = f(w)$ ,  $t > 0$  with  $w(0) = w'(0) = \gamma$  blows up in finite time  $T$ , where  $\gamma > u_0, T > 0$ ;

$$(1.10) \quad \begin{aligned} g(x) &\geq 2\gamma, \quad h(x) \geq 2\gamma \quad \text{in } B_{R+T}, \\ |\nabla g| + |\nabla^2 g| + |\nabla h| &< \eta \quad \text{in } B_{R+T}, \quad n > 0. \end{aligned}$$

**THEOREM 1.1 [2].** *If  $\eta$  is sufficiently small, depending on  $R, \gamma, T$ , then there exists a classical solution  $u_1(x, t)$  of (1.1)–(1.3) with  $\varepsilon = 1$  in  $K_{R,T}^1 \cap \Omega$  where  $\Omega = \{(x, t); x \in B_{R+T}, 0 < t < \phi_1(x)\}$ , and it satisfies*

- (i)  $0 < \phi_1(x) < T$ ,
- (ii)  $u_1(x, t) \rightarrow \infty$  if  $t \rightarrow \phi_1(x) - 0$ ,
- (iii)  $\phi_1 \in C^1(B_{R+T})$  and  $|\nabla \phi_1(x)| < 1$ . The solution is unique in  $K_{R,T}^1$  and it belongs to  $C^{3,1}$ .

The proof of existence of  $u_1$  begins by constructing a sequence of finite valued solutions  $U_n$  where  $U_0 = 0$  and

$$(1.11) \quad \begin{aligned} \square_1 U_{n+1} &= f(U_n) \quad \text{in } \mathbf{R}^N \times (0, \infty), \\ U_{n+1}(x, 0) &= g(x), \quad \frac{\partial}{\partial t} U_{n+1}(x, 0) = h(x) \quad (x \in \mathbf{R}^N). \end{aligned}$$

One shows that

$$(1.12) \quad U_n(x, t) \leq U_{n+1}(x, t) \quad \text{in } K_{R,T}^1$$

and that  $U_{n+1}(x, t) \rightarrow u_1(x, t)$  as  $n \rightarrow \infty$ , where  $u_1$  satisfies the properties asserted in Theorem 1.1.

Consider the case

$$(1.13) \quad \phi(0) < \infty.$$

In §2 we shall prove

LEMMA 1.2. *If (1.13) holds then there exists an  $R' > 0$  such that*

$$(1.14) \quad 0 < \phi(x) < \infty \quad \text{if } x \in B_{R'},$$

$$(1.15) \quad \phi \in C^1(B_{R'}).$$

Actually  $\phi$  belongs to  $C^4(B_{R'})$ , but this fact will not be needed.

In §3 we shall prove

LEMMA 1.3. *Fix any  $T$  such that  $\phi(0) < T < \infty$ . Then there exist  $R > 0$ ,  $C > 0$  and  $\varepsilon_0 > 0$  such that if  $0 < \varepsilon < \varepsilon_0$  then there exists a unique solution  $u_\varepsilon$  of (1.1)–(1.3) in  $K_{R,T}^\varepsilon \cap \Omega_{R,T}^\varepsilon$  where  $\Omega_{R,T}^\varepsilon = \{(x, t); x \in B_{R+T}, 0 < t < \phi_\varepsilon(x)\}$  and*

- (i)  $0 < \phi_\varepsilon(x) < T$ ,
- (ii)  $u_\varepsilon(x, t) \rightarrow \infty$  if  $t \rightarrow \phi_\varepsilon(x) - 0$ ,  $x \in B_R$ ,
- (iii)  $\phi_\varepsilon \in C^1(B_{R+T})$  and  $|\nabla\phi_\varepsilon(x)| \leq C$ ; the solution belongs to  $C^{3,1}$ .

We can now state the main results of this paper in case  $\phi(0) < \infty$ .

THEOREM 1.4. *If (1.4)–(1.6) hold and  $\phi(0) < \infty$  then there exist positive constants  $R, C$  such that, for all  $\varepsilon$  sufficiently small,*

$$(1.16) \quad \sup_{B_R} |\phi_\varepsilon(x) - \phi(x)| \leq C\varepsilon^2,$$

$$(1.17) \quad \sup_{B_R} |\nabla(\phi_\varepsilon(x) - \phi(x))| \leq C\varepsilon^2.$$

Theorem 1.4 will be proved in §§4–6.

Observe that Lemma 1.2 implies that the set  $D = \{x \in \mathbf{R}^N, \phi(x) < \infty\}$  is open, and Theorem 1.4 implies that the solution  $u_\varepsilon$  exists for  $0 < t < \phi_\varepsilon(x)$  and  $x$  in any compact subset  $D_0$  of  $D$ ; further

$$\begin{aligned} |\phi_\varepsilon(x) - \phi(x)| &\leq C\varepsilon^2 \quad \forall x \in D_0, \\ |\nabla(\phi_\varepsilon(x) - \phi(x))| &\leq C\varepsilon^2 \quad \forall x \in D_0. \end{aligned}$$

In §7 we shall consider the case  $\phi(0) = \infty$  and derive growth rates for  $\phi_\varepsilon(0)$  as  $\varepsilon \rightarrow 0$ .

**2. The equation  $u'' = f(u)$ .** Throughout §§2–6 we assume that (1.13) holds. Set

$$F(u) = \int_0^u f(s) ds.$$

From (1.7), (1.8) we obtain

$$(2.1) \quad u_t^2 - h^2(x) = 2[F(u) - F(g(x))] \quad \text{if } t < \phi(x),$$

and then also

$$(2.2) \quad \phi(x) = \int_{g(x)}^{\infty} \frac{du}{[2F(u) - 2F(g(x)) + h^2(x)]^{1/2}}$$

provided, say,  $g(x) > u_0$  (so that the denominator in the integrand is well defined).

Set

$$(2.3) \quad T(\gamma, \delta) = \int_{\gamma}^{\infty} \frac{du}{[2F(u) - 2F(\gamma) + \delta^2]^{1/2}}$$

for  $\gamma > u_0, \delta > 0$ . Then  $T(\gamma, \delta)$  is the blow-up time of (1.7) subject to  $u(0) = \gamma, u'(0) = \delta$ . Using (1.4) we can easily show that  $u'(t) > 0, u''(t) > 0$  if  $t > 0$  and then, from the differential equation for  $\partial u / \partial \gamma, \partial u / \partial \delta$  remains positive for all  $0 < t < T(\gamma, \delta)$ . It follows that  $\partial T / \partial \gamma \leq 0$ . Since

$$\frac{\partial T}{\partial \gamma} = -\frac{1}{\delta} + f(\gamma) \int_{\gamma}^{\infty} \frac{du}{[2F(u) - 2F(\gamma) + \delta^2]^{3/2}},$$

we deduce that

$$(2.4) \quad |\partial T / \partial \gamma| \leq 1/\delta.$$

Next

$$\frac{\partial T}{\partial \delta} = -\delta \int_{\gamma}^{\infty} \frac{du}{[2F(u) - 2F(\gamma) + \delta^2]^{3/2}}$$

so that

$$(2.5) \quad |\partial T / \partial \delta| \leq T/\delta.$$

The assumption  $\phi(0) < \infty$  implies that

$$(2.6) \quad u(0, t) \rightarrow \infty \quad \text{if } t \rightarrow \phi(0).$$

Indeed, if  $u(0, t_n)$  remains bounded for a sequence  $t_n \rightarrow \phi(0)$ , then, by (2.1), also  $u_t(0, t_n)$  remains bounded. But then the solution  $u(0, t)$  of  $u_{tt} = f(u)$  can be extended to  $t_n < t < t_n + \delta$  with  $\delta$  positive and independent of  $n$ , which is a contradiction if  $n$  is large enough.

From (2.6) and (2.1) we get

$$(2.7) \quad u_t(0, t) \rightarrow \infty \quad \text{if } t \rightarrow \phi(0).$$

Consequently, for any  $\gamma > u_0$  there exists a  $t_0 \in (0, \phi(0))$  such that  $u(0, t_0) > \gamma, u_t(0, t_0) > \gamma$  and, by continuity

$$(2.8) \quad u(x, t_0) > \gamma, \quad u_t(x, t_0) > \gamma \quad \text{if } x \in B_{R_0}$$

for some  $R_0 > 0$ . Using (1.4) we easily deduce that  $u(x, t)$  blows up in finite time  $\phi(x)$  for any  $x \in B_{R_0}$ . Further, analogously to (2.2), we have

$$(2.9) \quad \phi(x) = t_0 + T(u(x, t_0), u_t(x, t_0)), \quad x \in B_{R_0}.$$

Since  $u(x, t_0)$  and  $u_t(x, t_0)$  vary smoothly with  $x$  and since (2.4), (2.5) hold, we conclude:

LEMMA 2.1.  $\phi \in C^1(B_{R_0})$ .

Set

$$\Omega_\rho = \{(x, t); x \in B_\rho, 0 \leq t < \phi(x), \rho < R_0\}.$$

LEMMA 2.2. For any  $0 < R < R_0$  there exist positive constants  $C$  and  $c$  such that

$$(2.10) \quad |D^\alpha u(x, t)| \leq C(\phi(x) - t)^{-(pq+|\alpha|-2)} \quad \text{in } \Omega_R$$

where  $q = 2/(p - 1)$ ,  $0 \leq |\alpha| \leq 2$ , and

$$(2.11) \quad c(\phi(x) - t)^{-(pq+j-2)} \leq D_t^j u(x, t) \leq C(\phi(x) - t)^{-(pq+j-2)} \quad \text{in } \Omega_R$$

for  $0 \leq j \leq 3$ .

PROOF. From (2.1), by integration,

$$\int_{u(x,t)}^\infty \frac{du}{[h^2(x) + 2F(u) - 2F(g(x))]^{1/2}} = \phi(x) - t.$$

Since  $F(u) \sim u^{p+1}/(p + 1)$  as  $u \rightarrow \infty$ , the estimate (2.11) for  $u$  readily follows. Next using (2.1) we can establish (2.11) for  $D_t u$ , and using (1.7) we can further establish (2.11) for  $j = 2$  and then (from  $u_{ttt} = f'(u)u_t$ ) for  $j = 3$ .

To prove (2.10) we introduce (cf. [2]) the functions

$$\begin{aligned} J_1 &= C_1 u_t \pm D^\alpha u, & |\alpha| &= 1, \\ J_2 &= C_2 u_{tt} \pm D^\alpha u, & |\alpha| &= 2, \end{aligned}$$

with  $C_1, C_2$  positive constants. For any  $x_0 \in B_R$  we can choose  $\delta > 0$  and  $t_1 \in (0, \phi(x_0))$  such that  $u(x, t_1) > u_0$  and  $D_t^j u(x, t_1) > 1$  if  $x \in B_\delta(x_0)$  ( $0 \leq j \leq 3$ ). Hence, if  $C_1$  is large enough then  $J_1(x, t_1) > 0$  and  $J_{1,t}(x, t_1) > 0$  for  $x \in B_\delta(x_0)$ . Since

$$d^2 J_1/dt^2 = f'(u)J_1,$$

we can easily deduce by a continuity argument that  $J_1(x, t)$  remains positive for  $t_1 < t < \phi(x)$ , if  $x \in B_\delta(x_0)$ .

Next we choose  $C_2$  such that  $J_2(x, t_1) > 0$  and  $J_{2,t}(x, t_1) > 0$  for  $x \in B_\delta(x_0)$ . We have

$$\frac{d^2 J_2}{dt^2} = f'(u)J_2 + f''(u)(C_2 u_t^2 \pm D^{\beta_1} u D^{\beta_2} u)$$

where  $\beta_1 + \beta_2 = \alpha$ . Since  $J_1 > 0$ , if  $C_2$  is large enough then the coefficient of  $f''(u)$  is positive. Hence, by a continuity argument,  $J_2(x, t) > 0$  if  $x \in B_\delta(x_0)$ ,  $t_1 < t < \phi(x)$ . Combining the positivity of  $J_1, J_2$  with (2.11), the estimate (2.10) follows.

**3. Proof of Lemma 1.3.** In the sequence we shall need an integral representation for solutions of the inhomogeneous wave equation. The formula has a different form depending on the space dimension  $N$ . We shall consider only the case  $N = 3$ ; the cases  $N = 1, 2$  can be treated in a similar way.

For  $N = 3$  we have

$$(3.1) \quad \begin{aligned} w(x, t) &= \frac{t}{4\pi} \int_{|\xi|=1} w_1(x + \varepsilon t \xi) d\omega_\xi + \frac{\partial}{\partial t} \frac{t}{4\pi} \int_{|\xi|=1} w_0(x + \varepsilon t \xi) d\omega_\xi \\ &\quad + \frac{1}{4\pi} \int_0^t (t - s) ds \int_{|\eta|=1} h(x + \varepsilon(t - s)\eta, s) d\omega_\eta \end{aligned}$$

where

$$w_0(y) = w(y, 0), \quad w_1(y) = w_t(y, 0), \quad h(y, t) = \square_\varepsilon w(y, t).$$

For any  $R_1 > 0$ ,  $0 < \varepsilon < 1$  we can construct a solution  $u_\varepsilon$  of (1.1)–(1.3) in  $K_{R_1, \sigma_1}^\varepsilon$  provided  $\sigma_1$  is sufficiently small, independently of  $\varepsilon$ . In fact, define an operator  $S$  by

$$(3.2) \quad (Sw)(x, t) = G(x, t) + \frac{1}{4\pi} \int_0^t (t-s) ds \int_{|\eta|=1} f(w(x + \varepsilon(t-s)\eta, s)) d\omega_\eta,$$

where

$$(3.3) \quad G(x, t) = \frac{t}{4\pi} \int_{|\xi|=1} h(x + \varepsilon t\xi) dw_\xi + \frac{\partial}{\partial t} \frac{t}{4\pi} \int_{|\xi|=1} g(x + \varepsilon t\xi) dw_\xi.$$

The domain of  $S$  is taken to be

$$X_{\sigma_1, M_1} = \{w \in C^0(K_{R_1, \sigma_1}^\varepsilon); \sup |w| \leq M_1\}$$

where

$$M_1 = 1 + \sup_{K_{R_1, \tau}^\varepsilon} |G|.$$

Then, if  $\sigma_1$  is sufficiently small,  $S$  maps  $X_{\sigma_1, M_1}$  into itself and it is a contraction. It follows that  $S$  has a fixed point, which is clearly a solution  $u_\varepsilon$  of (1.1)–(1.3). If  $\tilde{u}_\varepsilon$  is another solution in  $K_{R_1, \sigma_2}$  then we easily deduce that

$$\|S\tilde{u}_\varepsilon - Su_\varepsilon\| < \frac{1}{2} \|\tilde{u}_\varepsilon - u_\varepsilon\|$$

where the norm is the supremum norm in  $K_{R_1, \sigma}$  with  $\sigma$  small enough, depending on  $\|\tilde{u}_\varepsilon\|$ . It follows that  $\tilde{u}_\varepsilon = u_\varepsilon$  if  $0 < t < \sigma$ , and by a step-by-step argument, also if  $0 < t < \min(\sigma_1, \sigma_2)$ .

Since  $\square_\varepsilon(DU_\varepsilon) = f'(u_\varepsilon)(Du_\varepsilon)$ , we can apply (3.1) to  $Du_\varepsilon$  and, by estimating successively the right-hand side, we find that

$$|Du_\varepsilon| \leq M'_1 \quad \text{in } K_{R_1, \sigma_1}^\varepsilon$$

provided  $\sigma_1$  is small enough;  $\sigma_1$  and  $M'_1$  are independent of  $\varepsilon$ . Similarly

$$(3.4) \quad |D^\alpha u_\varepsilon| \leq M'_1 \quad \text{in } K_{R_1, \sigma_1}^\varepsilon, \quad |\alpha| \leq 3,$$

with another constant  $M'_1$ .

From (3.4) it follows that any sequence  $\varepsilon \rightarrow 0$  has a subsequence such that

$$(3.5) \quad D^\alpha u_\varepsilon \rightarrow D^\alpha v \text{ in } (L^\infty(K_{R_1, \sigma_1}^\varepsilon))^* \text{ weakly; } 0 \leq |\alpha| \leq 3, \text{ and, therefore}$$

$$(3.6) \quad D^\alpha u_\varepsilon \rightarrow D^\alpha v \text{ uniformly in compact subsets of } \bigcap_{\varepsilon'} K_{R_1, \sigma_1}^{\varepsilon'}; \quad 0 \leq |\alpha| \leq 2.$$

Hence

$$(3.7) \quad v(x, t) = u(x, t)$$

where  $u$  is the solution of (1.7), (1.8) and (3.5), (3.6) hold for all  $\varepsilon \rightarrow 0$ .

We wish to extend the solution  $u_\varepsilon$  beyond  $t = \sigma_1$ . To do this we repeat the previous proof, considering  $S$  in the space

$$X_{\sigma_2, M_2} = \{w \in C^0(K_{R_2, \sigma_2}^\varepsilon), w \equiv u_\varepsilon \text{ in } K_{R_2, \sigma_2}^\varepsilon \cap \{t < \sigma_1\}, \sup |w| \leq M_1\}$$

for any  $R_2 < R_1$ ,  $M_2 = M_1 + 1$  and some  $\sigma_2 > \sigma_1$ . Then  $S$  is a contraction if  $\sigma_2 - \sigma_1$  is sufficiently small, depending on  $M_2$ . As before we can establish (3.5)–(3.7) in  $K_{R_2, \sigma_2}^\epsilon$ .

We can carry out the above procedure with  $\sigma_3, R_3, M_3, \sigma_4, R_4, M_4$ , etc.; however, the numbers  $\sigma_{j+1} - \sigma_j$  are decreasing since the  $M_j$  are increasing. Let

$$(3.8) \quad \tilde{\sigma} < \inf_{B_{R_1}} \phi(x), \quad 0 < \tilde{R} < R_1, \quad R_1 \text{ small.}$$

We claim that the above procedure yields, in a finite number  $j_0$  of steps ( $j_0$  independent of  $\epsilon$ ) a solution  $u_\epsilon$  in  $K_{\tilde{R}, \tilde{\sigma}}$ . Indeed, from (3.8) we have that

$$|D^\alpha u(x, t)| \leq C \quad \text{in } B_{R_1} \times [0, \tilde{\sigma}] \quad (|\alpha| \leq 3).$$

Hence, in view of (3.5)–(3.7) we may repeat the previous construction of  $u_\epsilon$  but with the following modifications: at each step  $j$  we must take  $\epsilon \leq \epsilon_j$  so that for  $|\alpha| \leq 2$  we have  $|D^\alpha u_\epsilon| < C + 1$  in  $K_{R_j, \sigma_j}$ . Hence we obtain the bound  $C + 2$  instead of  $M_j$  for  $|D^\alpha u_\epsilon|$  in  $K_{R_{j+1}, \sigma_{j+1}}$  ( $|\alpha| \leq 2$ ). Next, by Gronwall’s inequality we can derive a bound  $M_{j+1}$ , independent of  $\epsilon$ , for  $|D^\alpha u_\epsilon|$  in  $K_{R_{j+1}, \sigma_{j+1}}$  where  $|\alpha| = 3$ . We have  $\epsilon_1 > \epsilon_2 > \dots$ . However, the differences  $\sigma_{j+1} - \sigma_j$  remain uniformly positive independently of the choice of the  $R_j$ ; say  $\sigma_{j+1} - \sigma_j \geq \delta > 0$  ( $\epsilon_j$  depends on  $R_j$ ). Choosing  $j_0 = [\tilde{\sigma}/\delta] + 1$ , and  $R_j = R_1 - (R_1 - \tilde{R})/j_0$ , we obtain the desired solution  $u_\epsilon$  in  $K_{\tilde{R}, \tilde{\sigma}}^\epsilon$ . Further,

$$(3.9) \quad D^\alpha u_\epsilon \rightarrow D^\alpha u \text{ uniformly in compact subsets of } B_{\tilde{R}} \times [0, \tilde{\sigma}], \quad 0 \leq |\alpha| \leq 2.$$

Choosing  $R_1$  sufficiently small, we can take  $\tilde{\sigma}$  sufficiently close to  $\phi(0)$ . Hence in view of (2.8), (3.9) we have

$$(3.10) \quad u_\epsilon(x, t_0) > 2\gamma, \quad u_{\epsilon, t}(x, t_0) > 2\gamma \quad \text{if } x \in B_{2R},$$

provided  $2R < \tilde{R}$ , where  $\gamma > u_0$  and  $t_0$  is some point in  $(0, \tilde{\sigma})$ .

We now introduce the scaled functions

$$U_\epsilon(x, t) = u_\epsilon(\epsilon x, t) \quad \text{for } t \geq t_0.$$

Then  $\square_1 U_\epsilon = f(U_\epsilon)$ . Setting  $g_\epsilon(x) = U_\epsilon(x, t_0)$ ,  $h_\epsilon(x) = U_{\epsilon, t}(x, t_0)$ , we have

$$(3.11) \quad |\nabla g_\epsilon| + |\nabla h_\epsilon| \leq C\epsilon, \quad |\nabla^2 g_\epsilon| \leq C\epsilon^2.$$

Using (3.10), (3.11) we can now apply the proof of Theorem 1.4 (as given in [2]) in order to establish the existence of a unique solution  $U_\epsilon$  in  $K_{(R/\epsilon), T}^1 \cap \{t \geq t_0\} \cap \{t < \tilde{\phi}_\epsilon(x)\}$  for all  $\epsilon$  small enough, and the estimate

$$(3.12) \quad U_{\epsilon, t} \geq (c_0/\epsilon)|\nabla_x U_\epsilon|, \quad c_0 > 0:$$

the function  $U_\epsilon(x/\epsilon, t)$  is then the unique extension of the solution  $u_\epsilon$  to  $\{t < \phi_\epsilon(x)\}$ , where  $\phi_\epsilon(x) = \tilde{\phi}(\epsilon x)$ . It will be shown below that

$$(3.13) \quad \phi_\epsilon(x) < T.$$

Then, the assertion (ii) of Lemma 1.3 follows (from the proof of Theorem 1.4 for  $U_\epsilon$ ), and  $\phi_\epsilon \in C^1$ . Further, from (3.12) we deduce that  $|\nabla \phi_\epsilon| \leq 1/c_0$  and thus Lemma 1.3 follows.

To prove (3.13) let  $W_\delta(t)$  ( $\delta$  positive and small) be the solution of

$$(3.14) \quad \begin{aligned} W_\delta'' &= f(W_\delta) \quad \text{if } t > t_0, \\ W_\delta(t_0) &= u(0, t_0) - 2\delta, \\ W_\delta'(t_0) &= u_t(0, t_0) - 2\delta. \end{aligned}$$

We shall compare  $W_\delta$  with  $U_\varepsilon$  in  $K_{(R/\varepsilon), T}$ . By (3.9), if  $R$  is small enough, depending on  $\delta$ , then

$$(3.15) \quad \begin{aligned} W_\delta(t_0) &\leq U_\varepsilon(x, t_0) - \delta, \\ W_\delta'(t_0) &\leq U_{\varepsilon, t}(x, t_0) - \delta \quad \text{in } B_{(R/\varepsilon)+T} \end{aligned}$$

provided  $\varepsilon$  is sufficiently small. Also,

$$(3.16) \quad |\nabla U_\varepsilon(x, t_0)| \leq C\varepsilon < \delta \quad \text{if } \varepsilon < \delta/C.$$

Hence, by a comparison argument based on the representation (3.1), (3.2) (cf. the proof of Lemma 2.3 in [2]), it follows that

$$W_\delta(t) \leq U_\varepsilon(x, t) \quad \text{in } K_{(R/\varepsilon), T},$$

and thus  $\phi_\varepsilon(x) < T_\delta$  where  $T_\delta$  is the blow-up time for  $W_\delta$ . By the results of §2 (cf. (2.4), (2.5)),  $|T_\delta - \phi(x)| < C_1\delta$ . Consequently

$$(3.17) \quad \phi_\varepsilon(x) \leq \phi(x) + C_1\delta,$$

and (3.13) follows.

REMARK 3.1. If we replace  $-\delta$  by  $+\delta$  in (3.14) then the previous argument yields the estimate

$$(3.18) \quad \phi_\varepsilon(x) \geq \phi(x) - C_1\delta.$$

#### 4. Estimating $u_\varepsilon - u$ .

LEMMA 4.1. *If  $R$  is sufficiently small then for any compact subset  $D_0$  of*

$$\Omega_R \equiv \{(x, t); x \in B_R, 0 \leq t < \phi(x)\}$$

*there exists a constant  $C$  such that*

$$(4.1) \quad |u_\varepsilon - u| \leq C\varepsilon^2 \quad \text{in } D_0,$$

$$(4.2) \quad |u_{\varepsilon, t} - u_t| \leq C\varepsilon^2 \quad \text{in } D_0$$

*if  $\varepsilon$  is sufficiently small.*

PROOF. By the estimates of §3 we deduce that if  $\rho$  is sufficiently small then

$$(4.3) \quad |D^\alpha u_\varepsilon| \leq C \quad \text{for } 0 \leq |\alpha| \leq 3$$

provided  $(x, t) \in D_0$  and  $|x| < \rho$ . Similarly, for any  $x_0 \in B_R$  the estimate (4.3) holds on  $\{(x, t) \in D_0, x \in B_\rho(x_0)\}$ , where  $C$  and  $\rho$  can be taken independently of  $x_0$ . It follows that (4.3) holds in  $D_0$ .

We can then write

$$(4.4) \quad \frac{\partial^2}{\partial t^2} u_\varepsilon = f(u_\varepsilon) + h_\varepsilon, \quad |h_\varepsilon| = |\varepsilon^2 \Delta u_\varepsilon| \leq C\varepsilon^2.$$

Representing  $u$  in the form

$$(4.5) \quad u(x, t) = g(x) + th(x) + \int_0^t (t - \tau)f(u(x, \tau)) d\tau$$

and, similarly,

$$(4.6) \quad \begin{aligned} u_\epsilon(x, t) &= g(x) + th(x) + \int_0^t (t - \tau)f(u_\epsilon(x, \tau)) d\tau \\ &+ \int_0^t (t - \tau)h_\epsilon(x, \tau) d\tau \end{aligned}$$

and taking the difference, we get

$$|u_\epsilon(x, t) - u(x, t)| \leq C \int_0^t |u_\epsilon(x, \tau) - u(x, \tau)| + C\epsilon^2$$

provided  $D_0$  is taken to be a subgraph in the  $t$ -direction, and (4.1) follows.

Similarly

$$\begin{aligned} \frac{\partial^2}{\partial t^2} u_{\epsilon,t} &= f'(u_\epsilon)u_{\epsilon,t} + h_1, & |h_1| &= |\epsilon^2 \Delta u_{\epsilon,t}| \leq C\epsilon^2 \\ \frac{\partial^2}{\partial t^2} u_t &= f'(u)u_t, \end{aligned}$$

and

$$\begin{aligned} u_{\epsilon,t}(x, 0) &= h(x) = u_t(x, 0), \\ u_{\epsilon,tt}(x, 0) &= \epsilon^2 \Delta g + f(g) = u_{tt}(x, 0) + \epsilon^2 \Delta g \end{aligned}$$

and the previous argument coupled with the estimate (4.1) yields the assertion (4.2).

**5. Estimating  $\phi_\epsilon - \phi$  and estimating  $D^\alpha(u_\epsilon - u)$  near  $\{t = \phi\}$ .**

LEMMA 5.1. *If  $R$  is sufficiently small then there exists a constant  $C$  such that*

$$(5.1) \quad \sup_{B_R} |\phi_\epsilon(x) - \phi(x)| \leq C\epsilon$$

for all  $\epsilon$  sufficiently small.

PROOF. We re-examine the proof of (3.17), (3.18). It is easy to see that (3.15) holds with  $\delta = A\epsilon$  provided  $A$  is a sufficiently large positive number. Recalling also (3.16), we deduce as before that (3.17) holds if  $\delta = C\epsilon$  and  $\epsilon$  is sufficiently small. The proof of (3.18) with  $\delta = C\epsilon$  is similar.

In the sequel we shall need some estimates on  $D^\alpha u_\epsilon$  and  $D^\alpha(u_\epsilon - u)$  near the blow-up surface. We begin with

LEMMA 5.2. *There exist  $t_1 \in (0, \phi(0))$  and  $R > 0$  such that, for all  $\epsilon$  sufficiently small,*

$$(5.2) \quad c(\phi_\epsilon(x) - t)^{(pq+j-2)} \leq D_t^j u_\epsilon(x, t) \leq C(\phi_\epsilon(x) - t)^{-(pq+j-2)} \quad (0 \leq j \leq 2),$$

$$(5.3) \quad |D^\alpha u_\epsilon(x, t)| \leq C(\phi_\epsilon(x) - t)^{-(pq+|\alpha|-2)} \quad (0 \leq |\alpha| \leq 2)$$

for  $x \in B_R$ ,  $t_1 < t < \phi_\epsilon(x)$ , where  $c, C$  are positive constants, and  $q = 2/(p - 1)$ .

PROOF. To prove (5.2) we work with  $U_\varepsilon(x, t) = u_\varepsilon(\varepsilon x, t)$  and establish for  $D_t^j U_\varepsilon$  estimates (as in (5.2) with  $\phi_\varepsilon(x)$  replaced by  $\phi_\varepsilon(\varepsilon x)$ ) by the method of [2]. Since  $D_t^j u_\varepsilon(x, t) = D_t^j U_\varepsilon(x/\varepsilon, t)$ , the inequalities in (5.2) follow.

Consider next

$$\begin{aligned} J_1^\varepsilon &= C_1 u_{\varepsilon,t} \pm D_x^\alpha u_\varepsilon \quad (|\alpha| = 1), \\ J_2^\varepsilon &= C_2 u_{\varepsilon,tt} \pm D_x^\alpha u_\varepsilon \quad (|\alpha| = 2). \end{aligned}$$

If  $C_1, C_2$  are positive and sufficiently large, then

$$J_i^0(x, t_1) > 0, \quad J_{i,t}^0(x, t_1) > 0 \quad \text{if } x \in B_R,$$

where  $J_i^0$  is  $J_i^\varepsilon$  with  $u_\varepsilon$  replaced by  $u$ . Consequently, by (3.9),

$$(5.4) \quad J_i^\varepsilon(x, t_1) > 0, \quad J_{i,t}^\varepsilon(x, t_1) > 0 \quad \text{if } x \in B_R$$

provided  $\varepsilon$  is sufficiently small. We can now proceed by a comparison argument as in [2] (cf. also the proof of Lemma 2.2) to show that if  $\varepsilon$  is small enough (so that also  $|\nabla_x J_1^\varepsilon(\varepsilon x, t_1)| < \varepsilon J_{1,t}^\varepsilon(\varepsilon x, t_1)$ ,  $\varepsilon x \in B_R$ ) then  $J_1^\varepsilon(x, t) > 0$  in the set  $(\bigcup_{x_0 \in B_r} K^\varepsilon(x_0, \phi_\varepsilon(x_0))) \cap \{t > t_1\}$ . This yields the assertion (5.3) for  $|\alpha| = 1$  (with a different  $R$ ). Using this information we can next establish by comparison that  $J_2^\varepsilon(x, t) > 0$  in the same domain as before provided  $C_2$  is suitably chosen, and (5.3) thus follows for  $|\alpha| = 2$ .

By (5.1) we know that

$$|\phi_\varepsilon(x) - \phi(x)| \leq C_0 \varepsilon \quad \text{if } x \in B_R.$$

We shall choose a constant  $M$  such that  $M > 2C_0$ . Then

$$(5.5) \quad \tilde{c} \leq \frac{\phi_\varepsilon(x) - t}{\phi(x) - t} < \frac{1}{\tilde{c}} \quad \text{if } x \in B_R, \quad 0 < t < \phi(x) - M\varepsilon$$

where  $\tilde{c}$  is a positive constant independent of  $\varepsilon$ .

LEMMA 5.3. *The following estimates hold for  $t_1 < t < \phi(x) - M\varepsilon$ ,  $x \in B_R$ :*

$$(5.6) \quad |D^\alpha(u_\varepsilon - u)(x, t)| \leq C\varepsilon^2(\phi(x) - t)^{-(pq+|\alpha|-2)} \quad (0 \leq |\alpha| \leq 2)$$

where  $C$  is a positive constant independent of  $\varepsilon$ .

PROOF. We proceed as in Lemma 4.1 but use the estimates of Lemma 5.2. From the integral representation of  $u_\varepsilon$  and  $u$  in (4.6) and (4.5) we obtain, by taking the difference,

$$|u_\varepsilon(x, t) - u(x, t)| \leq C\varepsilon^2 \int_{t_1}^t \int_{t_1}^{\tilde{t}} |\Delta u_\varepsilon| + \int_{t_1}^t \int_{t_1}^{\tilde{t}} f'(\tilde{u})|u_\varepsilon - u| + C\varepsilon^2$$

where, in each integral, both integrations are in  $t$ , and  $\tilde{u}$  lies between  $u$  and  $u_\varepsilon$ . Setting  $\lambda = \phi(x) - t$  ( $x$  fixed) and using Lemma 5.2, we obtain

$$\begin{aligned} |u_\varepsilon| &\leq C\varepsilon^2 \int_{t_*}^\lambda \int_{t_*}^{\tilde{\lambda}} \frac{1}{\tilde{\lambda}^{pq}} + C \int_{t_*}^\lambda \int_{t_*}^{\tilde{\lambda}} \frac{|u_\varepsilon - u|}{\tilde{\lambda}^{(pq-2)(p-1)}} + C\varepsilon^2 \\ &\leq \frac{C\varepsilon^2}{\lambda^{pq-2}} + C \int_{t_*}^\lambda \int_{t_*}^{\tilde{\lambda}} \frac{|u_\varepsilon - u|}{\tilde{\lambda}^2} \\ &\leq \frac{C\varepsilon^2}{\lambda^{pq-2}} + \frac{C}{\lambda} \int_{t_*}^\lambda |u_\varepsilon - u| \quad (\phi(x) - t_1 = t_*). \end{aligned}$$

Hence the function

$$z = \int_{t_*}^\lambda |u_\epsilon - u|$$

satisfies

$$z' - Cz/\lambda \leq C\epsilon^2/\lambda^{pq-2}, \quad z(t_*) = 0,$$

from which we deduce that  $z \leq C\epsilon^2/\lambda^{pq-3}$ . Hence

$$(5.7) \quad |u_\epsilon - u| \leq C\epsilon^2/(\phi(x) - t)^{pq-2}.$$

This establishes (5.6) for  $|\alpha| = 0$ .

To consider the case  $|\alpha| = 1$  we first take  $D^\alpha = D_t$ . By differentiating the integral representation of  $u_\epsilon$  and  $u$  with respect to  $t$  and taking the difference, we get

$$|u_{\epsilon,t} - u_t| \leq c\epsilon \int_{t_1}^t |\Delta u_\epsilon| + \int_{t_1}^t f'(\tilde{u})|u_\epsilon - u|.$$

Using (5.7) and Lemma 5.2, we easily estimate the right-hand side, thereby deriving (5.6).

To estimate  $D_x^\alpha(u_\epsilon - u)$  for  $|\alpha| = 1$  we apply  $D_x^\alpha$  to the integral representation of  $u_\epsilon - u$  and obtain

$$|D_x^\alpha(u_\epsilon - u)| \leq C\epsilon^2 \iint |D_x^\alpha \Delta u_\epsilon| + \iint |f'(u_\epsilon)D_x^\alpha u_\epsilon - f'(u)D_x^\alpha u|.$$

Estimating

$$|[f'(u_\epsilon) - f'(u)]D_x^\alpha u_\epsilon|$$

by (5.7) and Lemma 5.2 we find that

$$|D_x^\alpha(u_\epsilon - u)| \leq \frac{C\epsilon^2}{\lambda^{pq-1}} + C \int_{t_*}^\lambda \int_{t_*}^{\tilde{\lambda}} \frac{|D_x^\alpha(u_\epsilon - u)|}{\tilde{\lambda}^{(pq-2)(p-1)}}.$$

We can now proceed as before to establish (5.6) (with  $|\alpha| = 1$ ). Finally, the proof of (5.6) for  $|\alpha| = 2$  is similar; we argue separately in the cases  $D_t^2$ ,  $D_t D_x^\alpha$  ( $|\alpha| = 1$ ) and  $D_x^\alpha$  ( $|\alpha| = 2$ ).

Using Lemmas 5.2 and 5.3 we shall now complete the proof of (1.16).

LEMMA 5.4. *If  $R$  is sufficiently small then there exists a constant  $C$  such that*

$$(5.8) \quad \sup_{B_R} |\phi_\epsilon(x) - \phi(x)| \leq C\epsilon^2$$

for all  $\epsilon$  sufficiently small.

PROOF. We repeat the proof of Lemma 5.1 choosing, in the comparison argument (3.14),

$$t_0 = \phi(0) - 3M\epsilon$$

and working in the cone  $K$  with base  $B_{5M\epsilon^2}(0)$  on  $t = t_0$  and vertex  $(0, \phi(0) + 2M\epsilon)$ . Set  $d = 3M\epsilon$ . By Lemmas 5.2 and 5.3,

$$(5.9) \quad \begin{aligned} \gamma_\epsilon &\equiv u(0, t_0) + C_0\epsilon^2/d^{pq-1} \geq u_\epsilon(x, t_0) & \text{if } x \in B_{5M\epsilon^2}(0), \\ \delta_\epsilon &\equiv u_t(0, t_0) + C_1\epsilon^2/d^{pq} \geq u_{\epsilon,t}(x, t_0) & \text{if } x \in B_{5M\epsilon^2}(0), \end{aligned}$$

if  $C_0, C_1$  are sufficiently large positive constants. Let  $W_\varepsilon(t)$  be the solution of

$$\begin{aligned} W''_\varepsilon &= f(W) \quad \text{if } t > t_0, \\ W_\varepsilon(t_0) &= \gamma_\varepsilon, \quad W'_\varepsilon(t_0) = \delta_\varepsilon. \end{aligned}$$

Set  $U_\varepsilon(x, t) = u_\varepsilon(\varepsilon x, t)$ . From the integral representation (3.1) and a comparison argument (as in [2]) we see that if

$$(5.10) \quad \begin{aligned} W'_\varepsilon(t_0) + (t - t_0)W_\varepsilon(t_0) &> U_{\varepsilon,t}(x, t_0) + (t - t_0)U_\varepsilon(x, t_0) \\ &+ (t - t_0)|\nabla_x U_\varepsilon(x, t_0)| \end{aligned}$$

then

$$W_\varepsilon(t) > u_\varepsilon(x, t) \quad \text{in } K$$

and consequently

$$(5.11) \quad \phi_\varepsilon(0) > T_\varepsilon$$

where  $T_\varepsilon$  is the blow-up time of  $W_\varepsilon(t)$ . Since  $t - t_0 \leq 5M\varepsilon$  and  $|\nabla_x U(x, t)| = \varepsilon|\nabla u_\varepsilon(\varepsilon x, t)|$ , (5.10) is a consequence of Lemma 5.2 provided we choose  $C_1$  to be sufficiently large (independently of  $\varepsilon$ ).

Setting  $\gamma = u(0, t_0)$ ,  $\delta = u_t(0, t_0)$  and using (2.3), we compute that

$$\begin{aligned} \phi(0) - T_\varepsilon &= \int_\gamma^\infty [2F(u) - 2F(\gamma) + \delta^2]^{-1/2} du \\ &\quad - \int_{\gamma_\varepsilon}^\infty [2F(u) - 2F(\gamma_\varepsilon) + \delta_\varepsilon^2]^{-1/2} du \\ &\leq \int_\gamma^{\gamma_\varepsilon} [2F(u) - 2F(\delta) + \delta^2]^{1/2} du \\ &\quad + \int_{\gamma_\varepsilon}^\infty \frac{C\varepsilon^2 d^{-(3p+1)/(p-1)}}{[2F(u) - 2F(\gamma) + \delta^2]^{3/2}} \equiv I_1 + I_2. \end{aligned}$$

Clearly

$$I_1 \leq C(\gamma_\varepsilon - \gamma)/\delta \leq C\varepsilon^2.$$

Next, substituting  $u = \gamma v$  into  $I_2$ , we get

$$I_2 \leq \frac{C\gamma}{\gamma^{(p+1)3/2}} \frac{\varepsilon^2}{d^{(3p+1)/(p-1)}} \leq C\varepsilon^2.$$

We have thus proved that  $\phi(0) - T_\varepsilon \leq C\varepsilon^2$ . Combining this with (5.10), it follows that

$$\phi(0) - \phi_\varepsilon(0) \leq C\varepsilon^2.$$

Similarly one proves that  $\phi(0) - \phi_\varepsilon(0) \geq -C\varepsilon^2$ , and thus  $|\phi(0) - \phi_\varepsilon(0)| \leq C\varepsilon^2$ . Since the above proof applies about each point  $x$  in some neighborhood of  $x = 0$ , (5.8) follows.

**6. Estimating  $\nabla(\phi_\varepsilon - \phi)$ .** Denote by  $\phi_m(x)$  and  $\psi_m(x)$  the solutions of

$$u(x, \phi_m(x)) = m, \quad u_t(x, \psi_m(x)) = m;$$

in view of Lemma 2.2,  $\phi_m(x)$  and  $\psi_m(x)$  are well defined for  $x \in B_R$ , provided  $m$  is sufficiently large;  $R$  is as usual a small enough positive number. Denote by

$N(x)$ ,  $N_m(x)$  and  $M_m(x)$  the unit normals in the positive  $t$ -direction of the surfaces  $\{t = \phi(x)\}$ ,  $\{t = \phi_m(x)\}$  and  $\{t = \psi_m(x)\}$  respectively. Thus, for instance,

$$N_m(x) = [1 + |\nabla\phi_m(x)|^2]^{-1/2}(-\nabla\phi_m(x), 1).$$

For any  $\eta > 0$ , denote by  $S_\eta(x)$  the set of all unit vectors  $\tau = \tau(x)$  with

$$(6.2) \quad \tau \cdot N(x) \geq \eta.$$

We claim

$$(6.3) \quad |\nabla(\phi_m(x) - \phi(x))| \leq Cm^{-(3p+1)/2} \quad \text{if } x \in B_R.$$

Indeed, analogously to (2.9) ( $t_0$  is taken close to  $\phi(0)$ )  $\phi_m(x)$  is given by

$$\phi_m(x) = t_0 + \int_{u(x,t_0)}^m \frac{du}{[2F(u) - 2F(u(x,t_0)) + (u_t(x,t_0))^2]^{1/2}}.$$

Therefore

$$\begin{aligned} \nabla\phi_m(x) = & -\frac{\nabla u(x,t_0)}{u_t(x,t_0)} \\ & + \int_{u(x,t_0)}^m \frac{[f(u(x,t_0))\nabla u(x,t_0) - u_t(x,t_0)\nabla u_t(x,t_0)] du}{[2F(u) - 2F(u(x,t_0)) + (u_t(x,t_0))^2]^{3/2}}. \end{aligned}$$

Since  $\phi(x) = \phi_\infty(x)$ , we deduce that

$$\nabla(\phi(x) - \phi_m(x)) = \int_m^\infty [\dots] du$$

with the same integrand as in the preceding integral. Hence

$$|\nabla(\phi_m(x) - \phi(x))| \leq C \int_m^\infty \frac{du}{u^{(p+1)3/2}},$$

and (6.3) follows.

We shall next prove that

$$(6.4) \quad |\nabla(\psi_m(x) - \phi(x))| \leq Cm^{-(3p+1)/(p+1)} \quad \text{if } x \in B_R.$$

Indeed, we have

$$u_t^2 - u_t^2(x,t_0) + 2F(u(x,t_0)) = 2F(u)$$

and  $F(u)$  has an inverse  $G = F^{-1}$ , well defined and smooth, for all  $u = u(x,t)$  with  $t > t_0$  (by (2.6), (2.7)). We can then write

$$u = G(\frac{1}{2}u_t^2 - \frac{1}{2}u_t^2(x,t_0) + F(u(x,t_0))),$$

and (1.7) takes the form

$$u_{tt} = f(G(\frac{1}{2}u_t^2 - \frac{1}{2}u_t^2(x,t_0) + F(u(x,t_0)))).$$

By integration we then obtain

$$\psi_m(x) = t_0 + \int_{u_t(x,t_0)}^m \frac{dv}{f(G(\frac{1}{2}v^2 + a(x)))}$$

where

$$a(x) = -\frac{1}{2}u_t^2(x,t_0) + F(u(x,t_0)).$$

The same formula holds for  $\phi(x)$  with  $m = \infty$ . Taking the gradient of the difference, we get

$$|\nabla(\psi(x) - \psi_m(x))| \leq C \int_m^\infty \frac{dv}{(\frac{1}{2}v^2 + a(x))^{1+p/(p+1)}}$$

and (6.4) follows.

LEMMA 6.1. *There exist a positive constant  $c$  such that, for any  $\eta \in (0, 1)$ , if*

$$(6.5) \quad \phi(x) - t < c\eta^{(p-1)/(3p+1)}$$

then

$$(6.6) \quad \tau \cdot N_{u(x,t)}(x) > \eta/3,$$

$$(6.7) \quad \tau \cdot M_{u_t(x,t)}(x) > \eta/3$$

for any  $x \in B_R$ ,  $\tau \in S_{2\eta/3}(x)$ .

PROOF. From (6.3), (6.4) and Lemma 2.2,

$$(6.8) \quad |\nabla(\phi_m - \phi)| \leq C/u^{(3p+1)/2} \leq C(\phi - t)^{(3p+1)/(p-1)}, \quad m = u(x, t),$$

$$(6.9) \quad |\nabla(\psi_m - \phi)| \leq C/u_t^{(3p+1)/(p+1)} \leq C(\phi - t)^{(3p+1)/(p-1)}, \quad m = u_t(x, t).$$

From (6.1) we have

$$|N(x) - N_{u(x,t)}(x)| \leq C|\nabla(\phi - \psi_m)(x)|.$$

Using (6.8) and (6.5) we get

$$|N(x) - N_{u(x,t)}(x)| \leq C(\phi(x) - t)^{(3p+1)/(p-1)} \leq Cc\eta < \eta/3$$

if  $c < 1/(3C)$ ; thus (6.6) follows. The proof of (6.7) follows similarly, making use of (6.9).

LEMMA 6.2. *There exist positive constants  $c_0, c_1, C_0$  such that for any  $\eta \in (0, 1)$  such that*

$$(6.10) \quad \varepsilon^2 < c_0\eta,$$

the following is true: if

$$(6.11) \quad 2M\varepsilon \leq \phi(x) - t < c\eta^{(p-1)/(3p+1)},$$

$$(6.12) \quad \tau \in S_{2\eta/3}(x), \quad x \in B_R,$$

then

$$(6.13) \quad \frac{\partial u_\varepsilon(x, t)}{\partial \tau} \geq \frac{c_1\eta}{(\phi(x) - t)^{pq-1}},$$

$$(6.14) \quad \frac{\partial}{\partial t} \frac{\partial u_\varepsilon(x, t)}{\partial \tau} \geq \frac{c_1\eta}{(\phi(x) - t)^{pq}},$$

$$(6.15) \quad \left| \nabla \frac{\partial u_\varepsilon(x, t)}{\partial \tau} \right| \leq \frac{C_0}{(\phi(x) - t)^{pq}}$$

provided  $\varepsilon$  is sufficiently small.

PROOF. The estimate (6.15) follows from Lemma 5.2. Next, from Lemma 6.1 we have, if  $\tau \in S_{2\eta/3}(x)$ ,

$$\frac{\partial u(x, t)}{\partial \tau} \geq \frac{\eta}{3} \frac{\partial u}{\partial N_u}, \quad \frac{\partial u}{\partial t} \frac{\partial u(x, t)}{\partial \tau} \geq \frac{\eta}{3} \frac{\partial u_t}{\partial M_{u_t}}.$$

We also clearly have

$$\frac{\partial u}{\partial t} = \alpha \frac{\partial u}{\partial N_u}, \quad \frac{\partial^2 u}{\partial t^2} = \beta \frac{\partial u_t}{\partial M_{u_t}}$$

where

$$\begin{aligned} \alpha &= \alpha(x, t) = [1 + |\nabla \phi_m(x)|^2]^{-1/2}, & m &= u(x, t), \\ \beta &= \beta(x, t) = [1 + |\nabla \psi_m(x)|^2]^{-1/2}, & m &= u_t(x, t). \end{aligned}$$

Recalling the estimates in (2.11), we conclude that

$$\frac{\partial u}{\partial \tau} \geq \frac{c\eta}{(\phi - t)^{pq-1}}, \quad \frac{\partial}{\partial t} \frac{\partial u}{\partial \tau} \geq \frac{c\eta}{(\phi - t)^{pq}} \quad (c > 0).$$

If we now make use of Lemma 5.3, we obtain from the last two inequalities the inequalities (6.13), (6.14) provided  $\varepsilon^2/\eta$  is bounded by a sufficiently small constant, i.e., provided (6.10) holds with  $c_0$  small enough (independently of  $\eta, \varepsilon$ ). This completes the proof of the lemma.

We now proceed to establish (1.17). Fix a point  $y$  in  $B_R$  ( $R$  small) and let  $t_1 = \phi(y) - 3M\varepsilon$ . Denote by  $K$  the cone with base

$$B \equiv \{(x, t_1); |x - y| < 5M\varepsilon^2\}$$

and vertex  $(y, \phi(y) + 2M\varepsilon)$ .

If  $\tau \in S_\eta(y)$  then (since  $\phi$  is smooth)  $\tau \in S_{2\eta/3}(x)$  for any  $x \in B_{5M\varepsilon^2}(y)$  provided

$$(6.16) \quad \varepsilon^2 \leq \eta/C$$

and  $C$  is a sufficiently large positive constant. It follows that (6.13)–(6.15) hold on  $B$ . We can therefore apply a comparison argument to  $U_\varepsilon(x, t) = \partial u_\varepsilon(y + \varepsilon x, t)/\partial \tau$  (as in [2]; cf. also the proof of Lemma 5.4) and deduce that

$$(6.17) \quad \partial u_\varepsilon / \partial \tau > 0 \quad \text{in } K \cap \{t < \phi\}$$

provided (6.16) holds with  $C$  large enough. Since  $K$  contains  $(y, \phi_\varepsilon(y))$  in its interior, we see from (6.17) that  $u_\varepsilon$  is increasing along any direction  $\tau (\tau \in S_\eta(y))$ , in some neighborhood of  $(y, \phi_\varepsilon(y))$ . This means that the direction of  $\nabla \phi_\varepsilon(y)$  and  $\nabla \phi(y)$  differ by at most  $\eta$ . Thus

$$|\nabla(\phi_\varepsilon - \phi)| \leq \eta \quad \text{at } y.$$

The constant  $\eta$  was subject only to the constraints (6.10), (6.11) for any  $x \in B_{5M\varepsilon^2}(y)$  with  $\phi(y) - t = 3M\varepsilon$ , and (6.16). Thus we can choose  $\eta = C\varepsilon^2$ , where  $C$  is a sufficiently large positive constant, and then (1.17) follows.

**REMARK 6.1.** Using Lemma 5.4 we can extend the estimates of Lemma 5.3 to  $t_1 < t < \phi(x) - M\varepsilon^2$ , and of Lemma 6.2 under the condition  $2M\varepsilon^2 < \phi(x) - t < c\eta^{(p-1)/(3p+1)}$  instead of (6.11). If we now follow the proofs of (1.16), (1.17) with these improved lemmas (using, for instance, in the proof of (1.17), cones with base  $B$  on  $t = t_1 \equiv \phi(y) - 3M\varepsilon^2$  given by  $B = \{(x, t_1); |x - y| < 5M\varepsilon^3\}$ ), we do not get any improvements of the estimates (1.16), (1.17).

**7. The case  $\phi(0) = \infty$ .** In this section we consider the case where  $\phi(0) = \infty$  but  $\phi(x) < \infty$  if  $x \in B_R \setminus \{0\}$  for some  $R > 0$ . By Lemma 1.3 for any  $x_0 \in B_R \setminus \{0\}$  there exists  $\delta_0 = \delta_0(|x_0|)$  and  $\varepsilon_0 = \varepsilon_0(\delta_0, |x_0|)$  positive such that there is a unique solution  $u_\varepsilon(x, t)$  with blow-up surface  $t = \phi_\varepsilon(x)$  if  $x \in B_\delta(x, 0)$  and  $0 < \varepsilon < \varepsilon_0$ ;

however  $\varepsilon_0$  may go to zero if  $x_0 \rightarrow 0$ . On the other hand, for  $N = 1$ ,  $\phi_\varepsilon$  is finite for all  $x \in B_R$  or even for all  $x \in \mathbf{R}^1$  with  $|\phi'_\varepsilon(x)| < 1/\varepsilon$  (by [1]).

For simplicity we take

$$(7.1) \quad f(u) = (u^+)^p, \quad p > 1,$$

and consider first the case where

$$(7.2) \quad g(x) > 0 \text{ in } B_R \setminus \{0\}, \quad g(x) = \sum_{i=1}^N \alpha_i x_i^2 + O(|x|^3) \quad \text{where } \alpha_i > 0 \forall i,$$

$$(7.3) \quad h(x) > 0 \text{ in } B_R \setminus \{0\}, \quad h(x) = \sum_{i=1}^N \beta_i x_i^2 + O(|x|^3) \quad \text{where } \beta_i > 0 \forall i.$$

We further assume that if  $N = 2$  or  $N = 3$  then the solution  $u_\varepsilon(x, t)$  with finite-valued blow-up surface  $t = \phi_\varepsilon(x)$  exist for all  $x \in B_R$ .

**THEOREM 7.1.** *Under the foregoing assumptions, there exist positive constants  $c_0, c_1$  such that*

$$(7.4) \quad c_0 \varepsilon^{-\sigma} \leq \phi_\varepsilon(x) \leq c_1 \varepsilon^{-\sigma} \quad \text{where } \sigma = 2(p-1)/3p-1.$$

**PROOF.** We shall construct a subsolution  $v_\varepsilon(x, t)$  with blow-up surface  $t = \psi_\varepsilon(x)$  such that

$$(7.5) \quad \psi_\varepsilon(0) = c_1 \varepsilon^{-\sigma}.$$

Similarly we shall construct a supersolution  $w_\varepsilon(t)$  with blow-up time  $T_\varepsilon$ ,

$$(7.6) \quad T_\varepsilon \geq c_0 \varepsilon^{-\sigma}.$$

Since  $w_\varepsilon \geq u_\varepsilon > v_\varepsilon$ ,

$$(7.7) \quad \psi_\varepsilon(0) \geq T_\varepsilon \geq c_0 \varepsilon^{-\sigma},$$

and the proof of (6.4) will be completed.

Observe that the domain of dependence of a point  $(0, T)$  with  $T \sim c\varepsilon^{-\sigma}$  is a cone whose base on  $\{t = 0\}$  is  $B_{R_0}$  where  $R_0 \sim c\varepsilon^{1-\sigma}$ . Therefore, in constructing  $v_\varepsilon$  and  $w_\varepsilon$  we need to define their Cauchy data only in a ball  $\{|x| < C\varepsilon^{1-\sigma}\}$  with an appropriate positive constant  $C$ .

We take

$$(7.8) \quad v_\varepsilon(x, t) = (1+t)\delta|x|^2 + k(t)$$

where  $\delta$  is a small positive constant and  $k$  is the solution of

$$(7.9) \quad k'' = k^p + 2\delta N \varepsilon^2 t \quad \text{if } t > 0, \quad k(0) = k'(0) = 0.$$

It is easy to check that  $\square_\varepsilon v_\varepsilon \leq v_\varepsilon^p$ . In order to compare the Cauchy data, we work with  $U_\varepsilon(x, t) = u_\varepsilon(\varepsilon x, t)$  and  $V_\varepsilon(x, t) = v_\varepsilon(\varepsilon x, t)$ . Denote by  $G_\varepsilon$  the function defined by (3.3) with  $g = g(\varepsilon x)$ ,  $h = h(\varepsilon x)$ , and denote by  $\tilde{G}$  the function defined by (3.3) with  $g = h = \delta\varepsilon^2|x|^2$ . Using the integral representations (3.1) for  $U_\varepsilon$  and  $V_\varepsilon$ , and noting that  $G \geq \tilde{G}$  if  $\delta$  is small enough, we deduce by comparison (cf. [2]) that  $U_\varepsilon \geq V_\varepsilon$ , i.e.,  $u_\varepsilon \geq v_\varepsilon$ .

The function  $K(t) = \lambda^{2/(1-p)}k(\lambda t)$  where  $\lambda = \varepsilon^{-\sigma}$  satisfies

$$K'' = K^p + 2\delta Nt, \quad K(0) = K'(0) = 0$$

and it blows up in some finite time  $c_1$ . Therefore  $k(t)$  blows up in time  $c_1\varepsilon^{-\sigma}$ , and (7.5) follows.

We next construct a supersolution  $w_\varepsilon(t)$ ,

$$(7.10) \quad \begin{aligned} w''_\varepsilon &= w^p_2 \quad \text{for } t > 0, \\ w_\varepsilon(0) &= w'_\varepsilon(0) = A\varepsilon^{2(1-\sigma)}, \quad A > 0. \end{aligned}$$

If  $A$  is sufficiently large then, using the representation (3.1) for  $U_\varepsilon(x, t)$  and  $w_\varepsilon(t)$  we deduce that  $w_\varepsilon \geq u_\varepsilon$  in  $K^\varepsilon(0, T)$  for  $T = c_1\varepsilon^{-\sigma}$ . Thus it remains to prove that  $w_\varepsilon$  blows up in time  $T_\varepsilon \geq c_0\varepsilon^{-\sigma}$  for some  $0 < c_0 < c_1$ .

Now, from (7.10) we get, by integration,

$$\frac{1}{2}w^2_{\varepsilon,t} = \frac{w^{p+1}_\varepsilon}{p+1} + \frac{A^2}{2}\varepsilon^{4(1-\sigma)}(1 + o(1))$$

where  $o(1) \rightarrow 0$  if  $\varepsilon \rightarrow 0$ ; hence

$$T_\varepsilon = \int_{A\varepsilon^{2(1-\sigma)}}^\infty \frac{dw}{[2w^{p+1}/(p+1) + A^2\varepsilon^{4(1-\sigma)}]^{1/2}} \geq c_0\varepsilon^{-\sigma}.$$

We shall next consider the case where  $g$  and  $h$  vanish to higher order at  $x = 0$ . We take  $N = 1$  and assume that

$$(7.11) \quad g(x) \geq 0, \quad g''(x) \geq 0 \quad \text{for } |x| \text{ near } 0,$$

and

$$(7.12) \quad \begin{aligned} h(x) &= \beta|x|^n + O(|x|^{n+1}), \\ h'(x) &= n\beta|x|^{n-2}x + O(|x|^n), \quad \beta > 0, \end{aligned}$$

where  $n$  is a positive number  $\geq 2$ .

Set

$$(7.13) \quad \sigma = \{1 + (p+1)/n(p-1)\}^{-1}.$$

**THEOREM 7.2.** *Let  $N = 1$  and let (7.1), (7.11), (7.12) hold. Then there exist positive constants  $c_0, c_1$  such that (7.4) holds with  $\sigma$  given by (7.13).*

Notice that for  $n = 2$  this is an improvement of Theorem 7.1 for  $N = 1$ , since the condition (7.11) is weaker than (7.2).

**PROOF.** We represent  $u_\varepsilon$  in the form

$$(7.14) \quad \begin{aligned} u_\varepsilon(x, t) &= \frac{1}{2}[g(x + \varepsilon t) + g(x - \varepsilon t)] + \frac{1}{2\varepsilon} \int_{x-\varepsilon t}^{x+\varepsilon t} h(y) dy \\ &\quad + \frac{1}{2\varepsilon} \int_0^t \int_{x-\varepsilon(t-\tau)}^{x+\varepsilon(t-\tau)} (u^+)^p(y, \tau) dy d\tau. \end{aligned}$$

By differentiation,

$$(7.15) \quad \begin{aligned} u_{\varepsilon,t}(x, t) &= \frac{\varepsilon}{2}[g'(x + \varepsilon t) - g'(x - \varepsilon t)] + \frac{1}{2}[h(x + \varepsilon t) + h(x - \varepsilon t)] \\ &\quad + \frac{1}{2} \int_0^t [(u^+)^p(x + \varepsilon(t - \tau), \tau) + (u^+)^p(x - \varepsilon(t - \tau), \tau)] d\tau. \end{aligned}$$

Using (7.14) and the assumption  $g(x) \geq 0$  for  $x$  near 0, we get

$$(7.16) \quad u_\varepsilon(x, \varepsilon^{-\sigma}) \geq \frac{1}{2\varepsilon} \int_{x-\varepsilon^{1-\sigma}}^{x+\varepsilon^{1-\sigma}} h(y) dy \geq \beta \varepsilon^{-\sigma} \varepsilon^{n(1-\sigma)} \quad \text{if } |x| < R$$

provided  $R$  is small;  $\alpha$  is some positive constant.

Next, since  $g'' \geq 0$ , we obtain from (7.15),

$$(7.17) \quad u_{\varepsilon,t}(x, \varepsilon^{-\sigma}) \geq \beta \varepsilon^{n(1-\sigma)}/4 \quad \text{if } |x| \leq R.$$

We now compare  $u_\varepsilon(x, \varepsilon^{-\sigma} + t)$  with the solution  $v_\varepsilon(t)$  of

$$v_\varepsilon'' = v_\varepsilon^p, \quad t > 0, \\ v_\varepsilon(0) = v_\varepsilon'(0) = \gamma_0 \varepsilon^{n(1-\sigma)}, \quad \gamma_0 = \beta/4.$$

This solution blows up at time  $T(\gamma_0 \varepsilon^{n(1-\sigma)}, \gamma_0 \varepsilon^{n(1-\sigma)})$  where  $T(\gamma, \delta)$  is defined by (2.3). It is easily computed that the blow-up time is bounded from above by

$$C(\varepsilon^{-n(1-\sigma)})^{(p-1)/(p+1)} \quad (C \text{ positive constant}),$$

which is equal to  $C\varepsilon^{-\sigma}$  (with another positive constant  $C$ ), by the definition of  $\sigma$  in (7.13). Hence

$$(7.18) \quad \phi_\varepsilon(0) < \varepsilon^{-\sigma} + C_* \varepsilon^{-\sigma} = c_1 \varepsilon^{-\sigma} \quad (c_1 = 1 + C).$$

In order to estimate  $\phi_\varepsilon(0)$  from below it is sufficient (in view of (7.18)) to use the initial data only in the interval  $\{|x| < c_1 \varepsilon^{1-\sigma}\}$ . Thus we can use the same function  $w_\varepsilon$  as in (7.10) for a supersolution. Proceeding as in the argument following (7.10), we then derive the lower estimate in (7.4) with  $\sigma$  as in (7.13).

REMARK. The above proof extends to  $N = 2, 3$  and say

$$(7.19) \quad g(x) \sim \alpha|x|^n, \quad h(x) \sim \beta|x|^n \quad (\alpha > 0, \beta > 0)$$

provided we already know that

$$(7.20) \quad u_\varepsilon(x, t) \geq 0, \quad u_{\varepsilon,t}(x, t) \geq 0$$

for  $|x| \leq C\varepsilon^{1-\sigma}$ ,  $0 \leq t \leq \varepsilon^{-\sigma}$ ; we use here the representation (3.1) for both  $u_\varepsilon$  and  $u_{\varepsilon,t}$ . However the usual method for proving (7.20) (by using the representation (3.1), as in [2]) does not extend to the present situation where (7.19) holds, even for  $t$  small.

### REFERENCES

1. L. A. Caffarelli and A. Friedman, *Differentiability of the blow-up curve for one dimensional nonlinear wave equations*, Arch. Rational Mech. Anal. **91** (1985), 83–98.
2. ———, *The blow-up boundary for nonlinear wave equations*, Trans. Amer. Math. Soc. **297** (1986), 223–241.
3. R. Glassey, *Blow-up theorems for nonlinear wave equations*, Math. Z. **132** (1973), 183–203.
4. ———, *Finite-time blow-up for solutions of nonlinear wave equations*, Math. Z. **177** (1981), 323–340.
5. F. John, *Blow-up of solutions of nonlinear wave equations in three space dimensions*, Manuscripta Math. **28** (1979), 235–268.

6. H. Levine, *Instability and nonexistence of global solutions to nonlinear wave equations of the form  $Pu_{tt} = -Au + F(u)$* , Trans. Amer. Math. Soc. **192** (1974), 1-21.
7. T. Kato, *Blow-up of solutions of some nonlinear hyperbolic equations*, Comm. Pure Appl. Math. **32** (1980), 501-505.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55455

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA 47907