TRANSACTIONS OF THE
AMERICAN MATHEMATICAL SOCIETY
Volume 330, Number 1, March 1992

ERRATUM TO "CHARACTERIZATIONS OF NORMAL QUINTIC K-3 SURFACES"

JIN-GEN YANG

T. Urabe [1] kindly pointed out that the proof of Lemma 2.1 of [2] does not work. In fact, the statement of the lemma is false. There exist normal quintic surfaces with two triple points and one elliptic double point. For example, let S_0 be a quintic surface defined by the equation

$$(y-1)(y-x^2)(y^2-2y+x^2)+(y-x^2)z^2+\lambda yx^3+\mu z^4+\nu z^5=0,$$

where x, y, z are (affine) coordinates and λ, μ and ν are generic complex constants. It contains one elliptic double point $(0, 0, 0)$ and two triple points $(1, 1, 0)$ and $(-1, 1, 0)$. It can be checked that the minimal resolution S of S_0 contains three (-1)-curves and the minimal model of S is a $K3$ surface.

Therefore Lemma 2.1 and Theorem 1 should be deleted from [2].

References

1. T. Urabe, Private correspondence, 1991.
2. J.-G. Yang, *Characterizations of normal quintic K-3 surfaces*, Trans. Amer. Math. Soc. **313** (1989), 737–749.

Department of Mathematics, Fudan University, Shanghai, China

Received by the editors March 9, 1991.