

## THE WEIGHTED HARDY'S INEQUALITY FOR NONINCREASING FUNCTIONS

VLADIMIR D. STEPANOV

**ABSTRACT.** The purpose of this paper is to give an alternative proof of recent results of M. Arino and B. Muckenhoupt [1] and E. Sawyer [8], concerning Hardy's inequality for nonincreasing functions and related applications to the boundedness of some classical operators on general Lorentz spaces. Our approach will extend the results of [1, 8] to the values of the parameters which are inaccessible by the methods of these papers.

### 1. INTRODUCTION

The Hardy's inequality of the form

$$(1) \quad \left( \int_0^\infty \left( \int_0^x f(t) dt \right)^q u(x) dx \right)^{1/q} \leq C \left( \int_0^\infty f^p(x)v(x) dx \right)^{1/p}$$

is well known. The problem of determining weights  $u$  and  $v$  for fixed  $p$  and  $q$ ,  $1 \leq p, q \leq \infty$ , has been investigated by many authors [3, 5, 6, 12, 13]; furthermore, the important case  $0 < q < 1 < p < \infty$  has recently been studied by G. Sinnamon [9, 10] (see, also [11]). The final form of these results may be summarized as follows.

**Theorem 1.** *Necessary and sufficient conditions for the validity of the inequality (1) with weights  $u, v \geq 0$  for all  $f \geq 0$  such that the right side of (1) is finite, and constant  $C$  independent of  $f$  are*

(a) *Let  $1 \leq p \leq q \leq \infty$ ,  $1/p + 1/p' = 1$ . Then (1) holds iff*

$$(2) \quad A \equiv \sup_{t>0} \left( \int_t^\infty u(x) dx \right)^{1/q} \left( \int_0^t v^{1-p'}(x) dx \right)^{1/p'} < \infty.$$

Moreover  $A \leq C \leq \alpha(p, q)A$ .

(b) *Let  $0 < q < p < \infty$ ,  $p > 1$ ,  $1/r = 1/q - 1/p$ ,  $1/q + 1/q' = 1$ . Then (1) holds iff*

$$(3) \quad B \equiv \left( \int_0^\infty \left[ \left( \int_t^\infty u(x) dx \right)^{1/q} \left( \int_0^t v^{1-p'}(x) dx \right)^{1/q'} \right]^r v^{1-p'}(t) dt \right)^{1/r} < \infty$$

and  $\beta_1(p, q)B \leq C \leq \beta_2(p, q)B$ .

Received by the editors March 9, 1990 and, in revised form, March 22, 1991.  
 1980 *Mathematics Subject Classification* (1985 Revision). Primary 26D15.

The constants  $\alpha, \beta_1, \beta_2$  depend only on  $p, q$ .

Several authors [1, 7, 8] have recently considered inequality

$$(4) \quad \left( \int_0^\infty \left( \frac{1}{x} \int_0^x f(t) dt \right)^q w(x) dx \right)^{1/q} \leq C \left( \int_0^\infty f^p(x) v(x) dx \right)^{1/p}$$

where  $f$  is a nonnegative and nonincreasing function.

Necessary and sufficient conditions are known for (4) if  $1 \leq p = q < \infty$ ,  $w = v$ . These can be derived using the so called "dilation function." Specifically, let  $V(\tau) = \int_0^\tau v(x) dx$ . Then the inequality

$$(5) \quad \left( \int_0^\infty \left( \frac{1}{x} \int_0^x g(t) dt \right)^p v(x) dx \right)^{1/p} \leq C \left( \int_0^\infty g^p(x) v(x) dx \right)^{1/p}$$

is valid for nonincreasing functions  $g \geq 0$  iff for some  $\varepsilon > 0$

$$(6) \quad \sup_{s>0} \frac{V(\tau s)}{V(s)} = O(\tau^{\varepsilon-p}), \quad \tau \geq 1.$$

This was shown by D. W. Boyd [2]. S. G. Krein and E. M. Semenov [4, Chapter 2, Theorem 6.6].

Recently M. Arino and M. Muckenhoupt [1] have shown that (5) is equivalent to  $v$  satisfying the condition

$$(7) \quad \int_t^\infty x^{-p} v(x) dx \leq \frac{D}{t^p} \int_0^t v(x) dx$$

where the constant  $D$  does not depend on  $t \in (0, \infty)$ . They have also pointed out the application of this criterion to a similar problem for Hardy-Littlewood's maximal function on general Lorentz spaces.

E. Sawyer [8] has extended the above results to cases of different weights  $v, w$  and exponents  $1 < p, q < \infty$ . Sawyer's approach is quite general and reduces the problem to Hardy's inequality with arbitrary functions by means of duality. His method relies upon a proof of the reverse Hölder inequality of the form

$$\sup_{g \geq 0, g \downarrow} \frac{\int_0^\infty g f}{\left( \int_0^\infty g^p v \right)^{1/p}} \approx \left( \int_0^\infty \left( \int_0^x f \right)^{p'} \left( \frac{1}{V} \right)^{p'} v dx \right)^{1/p'} + \frac{\int_0^\infty f}{\left( \int_0^\infty v \right)^{1/p}}$$

where  $g \downarrow$  means that the function  $g$  is nonincreasing. If  $T$  is the operator defined by  $Tf(x) = \int_0^\infty k(x, y) f(y) dy$  where  $k(x, y)$  is a nonnegative kernel and if for simplicity we assume that  $\int_0^\infty v = \infty$ , one can apply this result to obtain the equivalence

$$\begin{aligned} \left( \int_0^\infty (Tg)^q w \right)^{1/q} &\leq C \left( \int_0^\infty g^p v \right)^{1/p} \quad \forall 0 \leq g \downarrow \\ \Leftrightarrow \left( \int_0^\infty \left( \int_0^x T^* f \right)^{p'} \left[ \frac{1}{V} \right]^{p'} v dx \right)^{1/p'} &\leq C \left( \int_0^\infty f^{q'} w^{1-q'} \right)^{1/q'} \quad \forall f \geq 0. \end{aligned}$$

For instance, if  $Tf(x) = \frac{1}{x} \int_0^x f(t) dt$ , then

$$(8) \quad \int_0^x T^* f = \int_0^x f + x \int_x^\infty f(t) \frac{dt}{t}.$$

Sawyer then proved the following theorem via the corresponding Hardy's inequalities for the operators on the right side of (8) using Theorem 1 and its dual.

**Theorem 2.** *Necessary and sufficient conditions for the validity of (4) with the weights  $w \geq 0$  and  $v \geq 0$  for all nonincreasing functions  $f \geq 0$  with finite right side of (4) and constant  $C$  independent of  $f$  are*

(a) *Let  $1 < p \leq q < \infty$ . Then (4) is valid iff*

$$(9) \quad A_0 \equiv \sup_{t>0} \left( \int_0^t w(x) dx \right)^{1/q} \left( \int_0^t v(x) dx \right)^{-1/p} < \infty,$$

$$(10) \quad A_1 \equiv \sup_{t>0} \left( \int_t^\infty x^{-q} w(x) dx \right)^{1/q} \left( \int_0^t x^{p'} V^{-p'}(x) v(x) dx \right)^{1/p'} < \infty$$

and  $\alpha_1(p, q)(A_0 + A_1) \leq C \leq \alpha_2(p, q)(A_0 + A_1)$ .

(b) *Let  $1 < q < p < \infty$ ,  $1/r = 1/q - 1/p$ . Then (4) is valid iff*

$$(11) \quad B_0 \equiv \left( \int_0^\infty \left[ \left( \int_0^t w(x) dx \right)^{1/p} \left( \int_0^t v(x) dx \right)^{-1/p} \right]^r w(t) dt \right)^{1/r} < \infty,$$

(12)

$$B_1 = \left( \int_0^\infty \left[ \left( \int_t^\infty x^{-q} w(x) dx \right)^{1/q} \cdot \left( \int_0^t x^{p'} V^{-p'}(x) v(x) dx \right)^{1/q'} \right]^r t^{p'} V^{-p'}(t) v(t) dt \right)^{1/r} < \infty$$

and also  $\beta_3(p, q)(B_0 + B_1) \leq C \leq \beta_4(p, q)(B_0 + B_1)$ . The constants  $\alpha_1, \alpha_2, \beta_3, \beta_4$  depend only on  $p, q$ .

In §2 we shall give an alternative proof of Theorem 2. The main new result of the paper is the following extension of Theorem 2 to the case  $0 < q < 1 < p < \infty$ , and  $0 < p \leq q < \infty, 0 < p < 1$ .

**Theorem 3.** *Let the assumptions of Theorem 2 hold. Then*

(a) *Assertion (b) of Theorem 2 holds if  $0 < q < 1 < p < \infty$ .*

(b) *Let  $0 < p \leq q < \infty, 0 < p < 1$ . Then (4) is valid iff  $A_0 < \infty$  and*

$$(13) \quad \mathcal{A}_1 = \sup_{t>0} t \left( \int_t^\infty x^{-q} w(x) dx \right)^{1/q} \left( \int_0^t v(x) dx \right)^{-1/p} < \infty$$

and  $\gamma_1(p, q)(A_0 + \mathcal{A}_1) \leq C \leq \gamma_2(p, q)(A_0 + \mathcal{A}_1)$ . The constants  $\gamma_1, \gamma_2$  depend only on  $p, q$ .

The proof of Theorem 3 is given in §3; at the end of this section we state and prove a new sufficient condition (Proposition 2) for (4) when  $0 < q < p < 1$ .

The following notations will be employed in the paper.

$$V(\tau) \equiv \int_0^\tau v(x) dx; \quad W(\tau) \equiv \int_0^\tau w(x) dx; \quad \Psi_0^{1-p'}(x) \equiv x^{p'} V^{-p'}(x) v(x).$$

$A \ll B$  means the inequality  $A \leq cB$  with the constant  $c$  depending on  $p, q$  only. Products and quotients of the forms  $0 \cdot \infty, \frac{\infty}{\infty}, \frac{0}{0}$  are taken to be 0.

The author expresses his deep gratitude to Professors H. P. Heinig and E. T. Sawyer for the opportunity to visit McMaster University and for fruitful discussions concerning these results. He also wishes to thank Professor R. C. Brown for editorial assistance in the preparation of the final English draft of the paper.

## 2. PROOF OF THEOREM 2

2.1. First we consider the characterization problem for the inequality

$$(14) \quad \left( \int_0^\infty f^q w \right)^{1/q} \leq C \left( \int_0^\infty f^p v \right)^{1/p}.$$

We need the following preliminary results.

**Lemma.** *If  $0 < q < p < \infty$ ,  $1/r = 1/q - 1/p$ , then  $A_0 \ll B_0$  and*

$$B_0 \equiv \left( \int_0^\infty W^{r/p} V^{-r/p} w \right)^{q/r} < \infty \Leftrightarrow \mathcal{B}_0 \equiv \left( \int_0^\infty W^{r/q} V^{-r/q} v \right)^{1/r} < \infty.$$

Moreover,  $B_0 \ll \mathcal{B}_0 \ll B_0$ .

*Proof.* First we show " $\Rightarrow$ ". Let  $B_0 < \infty$ , then

$$V^{-r/p}(x) W^{r/q}(x) = V^{-r/p} \int_0^x dW^{r/q} \leq \int_0^x V^{-r/p} dW^{r/q} \rightarrow 0, \quad x \rightarrow 0.$$

This implies that  $A_0 \ll B_0$  and integration by parts yields

$$\infty > B_0^r = \frac{q}{r} \int_0^\infty V^{-r/p} dW^{r/q} \geq \frac{q}{r} \int_0^\infty W^{r/q} d(-V^{-r/p}) = \frac{q}{r} \mathcal{B}_0^r.$$

The proof of " $\Leftarrow$ " is similar.

**Proposition 1.** *Necessary and sufficient conditions for the validity of (14) for all nonincreasing  $f \geq 0$  are as follows.*

- (a) *Let  $0 < p \leq q < \infty$ . Then (14) holds iff  $A_0 < \infty$ .*
- (b) *Let  $0 < q < p < \infty$ . Then (14) holds iff  $B_0 < \infty$ .*

*Remark.* For  $1 < p, q < \infty$  Proposition 1 is contained in Remark (i) of [8].

*Proof.* (a) The necessary part is trivial by substituting in the inequality  $f = f_I = \chi_{[0, I]}$ , where  $\chi_E(x) = 1$  for  $x \in E$  and  $\chi_E(x) = 0$  for  $x \notin E$ . To prove sufficiency we proceed as follows. Let  $h \geq 0$ ,  $\text{supp } h \subset (0, \infty)$ ,  $f(x) = \int_x^\infty h$  and  $\int_0^\infty f^p v < \infty$ . Integration by parts and Minkowski's inequality yields

$$\begin{aligned} \int_0^\infty f^q w &= \int_0^\infty W d(-f^q) \leq A_0^q \int_0^\infty V^{q/p} d(-f^q) \\ &\leq A_0^q \left( \int_0^\infty \left( \int_x^\infty d(-f^q) \right)^{p/q} v \right)^{q/p} = A_0^q \left( \int_0^\infty f^p v \right)^{q/p}. \end{aligned}$$

The proof is complete if  $\int_0^\infty v = \infty$ . If  $\int_0^\infty v < \infty$ , then we modify the argument by taking  $f(x) = c + \int_x^\infty h$ .

We next consider Proposition 1(b). Let us assume  $\int_0^\infty v = \infty$ . The necessary part proceeds as follows.

$$\mathcal{B}_0^r = \int_0^\infty W^{r/q} V^{-r/q} v = \int_0^\infty v(x) dx \int_x^\infty W^{r/q} V^{-(r/q)-1} v \equiv \int_0^\infty f_0^p v$$

where  $f_0$  is defined by the left side. Let us assume temporarily that  $B_0 < \infty$ , then using Lemma (14) with  $f = f_0$  yields

$$\begin{aligned} C^q B_0^{qr/p} &\geq C^q \left(\frac{q}{p}\right)^{q/p} \mathcal{B}_0^{qr/p} \geq \int_0^\infty w(x) \left(\int_x^\infty W^{r/q} V^{-(r/q)-1} v\right)^{q/p} dx \\ &\gg \int_0^\infty w W^{r/p} V^{-r/p} = B_0^r. \end{aligned}$$

This implies that  $C \gg B_0$  and the temporary assumption  $B_0 < \infty$  can be removed in the usual way.

To prove sufficiency, let  $h \geq 0$ ,  $\text{supp } h \subset (0, \infty)$ ,  $f(x) = \int_x^\infty h$  and  $\int_0^\infty f^p v < \infty$ . We have

$$\begin{aligned} \int_0^\infty f^q w &= \int_0^\infty \left(\int_s^\infty h\right)^q w(s) ds \\ &= \int_0^\infty \left(\left(\int_s^\infty h\right)^q \frac{w^{q/p}(s) ds}{\left(\int_s^\infty W^{r/p} V^{-r/p} v\right)^{q/r}}\right) \\ &\quad \cdot \left(w^{1-q/p}(s) \left(\int_s^\infty W^{r/p} V^{-r/q} v\right)^{q/r}\right) \end{aligned}$$

(apply Hölder's inequality with  $p/q, r/q$ ),

$$\begin{aligned} &\leq \left(\int_0^\infty \left(\int_s^\infty h\right)^p \frac{w(s) ds}{\left(\int_s^\infty W^{r/p} V^{-r/q} v\right)^{p/r}}\right)^{q/p} \left(\int_0^\infty w(s) ds \int_s^\infty W^{r/p} V^{-r/q} v\right)^{q/r} \\ &= \mathcal{B}_0^q \left(\int_0^\infty \left(\int_s^\infty h\right)^p \left(\int_s^\infty W^{r/p} V^{-r/q} v\right)^{-p/r} dW(s)\right)^{q/p} \end{aligned}$$

(integrating by parts)

$$\begin{aligned} &\ll \mathcal{B}_0^q \left(\int_0^\infty \left(\int_s^\infty h\right)^{p-1} h(s) \left(\int_s^\infty W^{r/p} V^{-r/q} v\right)^{-p/r} W(s) ds\right)^{q/p} \\ &\ll \mathcal{B}_0^q \left(\int_0^\infty \left(\int_s^\infty h\right)^{p-1} h(s) V(s) ds\right)^{q/p}, \end{aligned}$$

because

$$\left(\int_s^\infty W^{r/p} V^{-r/q} v\right)^{-p/r} \ll \frac{V(s)}{W(s)}.$$

Consequently

$$\int_0^\infty f^q w \ll \mathcal{B}_0^q \left(\int_0^\infty \left(\int_s^\infty h\right)^p v(s) ds\right)^{q/p} \ll B_0^q \left(\int_0^\infty f^p v\right)^{q/p}$$

and Proposition 1(b) is proved under the assumption  $\int_0^\infty v = \infty$ .

In the case  $\int_0^\infty v < \infty$  we have to change the weight  $v$  to  $v + \varepsilon$ ,  $\varepsilon > 0$ . The result follows from the previous case, by obtaining uniform estimates with respect to  $\varepsilon$  and then applying Fatou's lemma. For instance, if (14) holds and  $\int_0^\infty v < \infty$  then the inequality  $(\int_0^\infty f^q w)^{1/q} \leq C(\int_0^\infty f^p(v + \varepsilon))^{1/p}$  is also valid with the same constant  $C$ . It implies  $B_{0,\varepsilon} \ll C$ , where

$$B_{0,\varepsilon} = \left[ \int_0^\infty W^{r/p}(t) \left( \int_0^t v + \varepsilon \right)^{-r/p} w(t) dt \right]^{1/r}.$$

By Fatou's lemma, we get  $B_0 \ll C$ . Conversely, if  $B_0 < \infty$  and  $\int_0^\infty v < \infty$ , then  $B_{0,\varepsilon} \leq B_0$  and we get  $C \ll B_{0,\varepsilon} \leq B_0$  by the above arguments.

2.2. Next we prove the necessary part of Theorem 2. Let us assume that  $\int_0^\infty v = \infty$  and start with case (a),  $1 < p \leq q < \infty$ . Let  $0 < \tau < \infty$  and

$$f_\tau(s) = \left( \int_s^\tau x^{p'} V^{-p'-1}(x)v(x) dx \right)^{1/p} \chi_{[0,\tau]}(s),$$

then  $f_\tau$  is nonincreasing and

$$\int_0^\infty f_\tau^p v = \int_0^\tau x^{p'} V^{-p'-1}(x)v(x) dx.$$

Inequality (4) with  $f = f_\tau$  yields

$$\begin{aligned} C \left( \int_0^\tau x^{p'} V^{-p'-1}(x)v(x) dx \right)^{1/p} &\geq \left( \int_0^\infty \left( \frac{1}{x} \int_0^x \left( \int_s^\tau y^{p'} V^{-p'-1}(y)v(y) dy \right)^{1/p} \chi_{[0,\tau]}(s) ds \right)^q w(x) dx \right)^{1/q} \\ &\geq \left( \int_\tau^\infty x^{-q} w(x) dx \right)^{1/q} \int_0^\tau \left( \int_s^\tau y^{p'} V^{-p'-1}(y)v(y) dy \right)^{1/p} ds. \end{aligned}$$

Changing the order of integration we have

$$\begin{aligned} \int_0^\tau \left( \int_s^\tau y^{p'} V^{-p'-1}(y)v(y) dy \right)^{1/p} ds &\geq \int_0^\tau s^{p'/p} \left( \int_s^\tau V^{-p'-1}(y)v(y) dy \right)^{1/p} ds \\ &= p \int_0^\tau s^{p'/p} \int_s^\tau \left( \int_y^\tau V^{-p'-1}v \right)^{-1/p'} V^{-p'-1}(y)v(y) dy ds \\ &= p \int_0^\tau V^{-p'-1}(y)v(y) \left( \int_0^y s^{p'/p} ds \right) \left( \int_y^\tau V^{-p'-1}v \right)^{-1/p'} dy \\ &\geq (p-1) \int_0^\tau V^{-p'-1}(y)v(y)y^{p'} \left( \int_y^\infty V^{-p'-1}v \right)^{-1/p'} dy \\ &= (p')^{-1/p} p \int_0^\tau y^{p'} V^{-p'}(y)v(y) dy. \end{aligned}$$

This shows that  $C \gg A_1$  when  $\int_0^\infty v = \infty$ . This case  $\int_0^\infty v < \infty$  can be easily proved by the arguments at the end of the previous section.

2.3. Let us note before proving the necessary part of case (b) that the following conditions are equivalent  $B_1 < \infty \Leftrightarrow$

$$\mathcal{B}_1 \equiv \left( \int_0^\infty \left[ \left( \int_t^\infty x^{-q} w(x) dx \right)^{1/p} \left( \int_0^t \Psi_0^{1-p'}(x) dx \right)^{1/p'} \right]^r t^{-q} w(t) dt \right)^{1/r} < \infty$$

and, in particular,  $\mathcal{B}_1 \geq (q/p')^{1/r} B_1$ .

We write

$$\begin{aligned} B_1^r &= \int_0^\infty \left( \int_t^\infty x^{-q} w(x) dx \right)^{r/q} \left( \int_0^t \Psi_0^{1-p'} \right)^{r/q'} \Psi_0^{1-p'}(t) V^{-1}(t) V(t) dt \\ &= \int_0^\infty \left( \int_y^\infty \left( \int_t^\infty x^{-q} w(x) dx \right)^{r/q} \right. \\ &\quad \left. \cdot \left( \int_0^t \Psi_0^{1-p'} \right)^{r/q'} \Psi_0^{1-p'}(t) V^{-1}(t) dt \right) v(y) dy \\ &\equiv \int_0^\infty f_0^p v \end{aligned}$$

where the function  $f_0$  is again determined by the left side. Then  $f_0$  is nonincreasing and (4) yields

$$\begin{aligned} C B_1^{r/p} &= C \left( \int_0^\infty f_0^p v \right)^{1/p} \geq \left( \int_0^\infty \left( \frac{1}{s} \int_0^s f_0 \right)^q w(s) ds \right)^{1/q} \\ &\equiv \left( \int_0^\infty \Phi^q(s) s^{-q} w(s) ds \right)^{1/q}. \end{aligned}$$

As usual we assume that  $\int_0^\infty v = \infty$  and  $B_1 < \infty$  and will prove that  $C \gg B_1$ . We have

$$\begin{aligned} \Phi(s) &\geq \int_0^s \left( \int_y^s \left( \int_t^\infty x^{-q} w(x) dx \right)^{r/q} \left( \int_0^t \Psi_0^{1-p'} \right)^{r/q'} \Psi_0^{1-p'}(t) V^{-1}(t) dt \right)^{1/p} dy \\ &\geq \left( \int_s^\infty x^{-q} w(x) dx \right)^{r/qp} J(s), \end{aligned}$$

where

$$J(s) \equiv \int_0^s \left( \int_y^s \left( \int_0^t \Psi_0^{1-p'} \right)^{r/q'} \Psi_0^{1-p'}(t) V^{-1}(t) dt \right)^{1/p} dy.$$

Also we have for  $1 < q < p < \infty$  the lower bound.

$$\begin{aligned} J(s) &\geq \int_0^s y^{p'/p} \left( \int_y^s \left( \int_0^t \Psi_0^{1-p'} \right)^{r/q'} V^{-p'-1}(t) v(t) dt \right)^{1/p} dy \\ &= p \int_0^s y^{p'/p} dy \int_y^s \left( \int_\tau^s \left( \int_0^t \Psi_0^{1-p'} \right)^{r/q'} V^{-p'-1}(t) v(t) dt \right)^{-1/p'} \\ &\quad \cdot \left( \int_0^\tau \Psi_0^{1-p'} \right)^{r/q'} V^{-p'-1}(\tau) v(\tau) d\tau \geq p \left( \int_0^s \Psi_0^{1-p'} \right)^{-r/q'p'} J_1(s), \end{aligned}$$

where

$$J_1(s) \equiv \int_0^s y^{p'/p} dy \int_y^s \left( \int_\tau^s V^{-p'-1}(t)v(t) dt \right)^{-1/p'} \\ \times \left( \int_0^\tau \Psi_0^{1-p'} \right)^{r/q'} V^{-p'-1}(\tau)v(\tau) d\tau.$$

Changing the order of integration we get

$$J_1(s) = \int_0^s \left( \int_0^\tau \Psi_0^{1-p'} \right)^{r/q'} V^{-p'-1}(\tau)v(\tau) \\ \times \left( \int_0^\tau y^{p'/p} dy \right) \left( \int_\tau^s V^{-p'-1}(t)v(t) dt \right)^{-1/p'} d\tau \\ \geq \frac{1}{p'} \int_0^s \tau^{p'} \left( \int_0^\tau \Psi_0^{1-p'} \right)^{r/q'} V^{-p'-1}(\tau)v(\tau) \left( \int_\tau^\infty V^{-p'-1}(t)v(t) dt \right)^{-1/p'} d\tau \\ = (p')^{-1/p} \int_0^s \left( \int_0^\tau \Psi_0^{1-p'} \right)^{r/q'} \Psi_0^{1-p'}(\tau) d\tau = (p')^{1/p'} (1/r) \left( \int_0^s \Psi_0^{1-p'} \right)^{r/p'}.$$

Putting these lower bounds on  $J_1(s)$  and  $J(s)$  into the estimate for  $\Phi(s)$  we have

$$\Phi(s) \gg \left( \int_s^\infty x^{-q}w(x) dx \right)^{r/qp} \left( \int_0^s \Psi_0^{1-p'} \right)^{r/p'q}.$$

This implies

$$[CB_1^{r/p}]^q \gg \int_0^\infty \left( \int_s^\infty x^{-q}w(x) dx \right)^{r/p} \left( \int_0^s \Psi_0^{1-p'} \right)^{r/p'} s^{-q}w(s) ds \\ = \mathcal{B}_1^r \geq (q/p')B_1^r.$$

Hence  $C \gg B_1$  in the case  $1 < q < p < \infty$  and  $\int_0^\infty v = \infty$ . As before the case  $\int_0^\infty v < \infty$  follows by the arguments used at the end of §2.1.

2.4. Now we prove the sufficient part of Theorem 2. Let us assume at first that  $\int_0^\infty v = \infty$ ; we take as before

$$f(x) = \int_x^\infty h, \quad h \geq 0, \quad \text{supp } h \subset (0, \infty), \quad \int_0^\infty f^p v < \infty.$$

We will get an upper bound on the expression

$$F = \left( \int_0^\infty \left( \frac{1}{x} \int_0^x f(t) dt \right)^q w(x) dx \right)^{1/q}.$$

We have

$$\frac{1}{x} \int_0^x f(t) dt = \frac{1}{x} \int_0^x \left( \int_t^\infty h \right) dt = \int_x^\infty h + \frac{1}{x} \int_0^x sh(s) ds,$$

consequently

$$F \leq \left( \int_0^\infty f^q w \right)^{1/q} + F_1.$$



The first term on the right side has been bounded in the proof of the Proposition 1. Let us bound the second term. We have, setting  $H = hV$ ,

$$F_1^q \equiv \int_0^\infty \left( \frac{1}{x} \int_0^x sh(s) ds \right)^q w(x) dx = \int_0^\infty \left( \frac{1}{x} \int_0^x \frac{sH(s)}{V(s)} ds \right)^q w(x) dx.$$

Integrating by parts and setting  $\int_0^s H = G$ , we get

$$\begin{aligned} \int_0^x \frac{sH(s)}{V(s)} ds &= \int_0^x \frac{s}{V(s)} dG(s) = \frac{xG(x)}{V(x)} - \int_0^x \frac{G(s)}{V(s)} ds + \int_0^x \frac{sv(s)}{V^2(s)} G(s) ds \\ &\leq \frac{xG(x)}{V(x)} + \int_0^x \frac{sv(s)}{V^2(s)} G(s) ds. \end{aligned}$$

This implies

$$(16) \quad \begin{aligned} F_1^q &\ll \int_0^\infty \left( \frac{G}{V} \right)^q w + \int_0^\infty \left( \frac{1}{x} \int_0^x \frac{sv(s)}{V^2(s)} G(s) ds \right)^q w(x) dx \\ &\equiv F_2^q + F_3^q. \end{aligned}$$

Integrating by parts we have

$$(17) \quad F_2^q = \int_0^\infty \left( \frac{G}{V} \right)^q w = \int_0^\infty \left( \frac{G}{V} \right)^q dW \leq q \int_0^\infty \frac{G^q v}{V^{q+1}} W.$$

Now we have to consider separately the cases  $1 < p \leq q < \infty$  and  $1 < q < p < \infty$ . If  $1 < p \leq q < \infty$  then (9) yields

$$F_2^q \leq q A_0^q \int_0^\infty \frac{G^q v}{V^{q+1-q/p}}.$$

We have

$$G(s) = \int_0^s hV = \int_0^s v(x) \left( \int_x^s h \right) dx \leq \int_0^s fv.$$

Consequently

$$F_2^q \leq q A_0^q \int_0^\infty \left( \int_0^s fv \right)^q \frac{v(s) ds}{V^{q+1-q/p(s)}}.$$

It is easy to show that

$$\left( \int_t^\infty \frac{v(s) ds}{V^{q+1-q/p(s)}} \right)^{1/q} \left( \int_0^t v \right)^{1/p'} \leq (p'/q)^{1/q},$$

therefore applying Theorem 1 we get

$$(18) \quad F_2^q \leq \alpha^q(p, q) p' A_0^q \left( \int_0^\infty f^p v \right)^{q/p}.$$

Denoting  $sv(s)G(s)/V^2(s) = \Phi(s)$  and again applying Theorem 1 and (10) we

find that

$$\begin{aligned}
 F_3^q &= \int_0^\infty \left( \frac{1}{x} \int_0^x \Phi(t) dt \right)^q w(x) dx \leq \alpha^q(p, q) A_1^q \left( \int_0^\infty \Phi^p \Psi_0 \right)^{q/p} \\
 &= \alpha^q(p, q) A_1^q \left( \int_0^\infty \frac{G^p v}{V^p} \right)^{q/p} \\
 (19) \quad &\leq \alpha^q(p, q) A_1^q \left( \int_0^\infty \left( \int_0^s f v \right)^p \frac{v(s) ds}{V^p(s)} \right)^{q/p} \\
 &\leq \alpha^q(p, q) \alpha^q(p, p) (p'/p)^{q/p} A_1^q \left( \int_0^\infty f^p v \right)^{q/p}.
 \end{aligned}$$

Combining the estimates (15), (16), (18), and (19) we get the sufficient part of Theorem 2(a).

The proof of the sufficient part of Theorem 2(b) is similar. Let  $0 < q < p < \infty$ , then using (17) and Hölder's inequality with exponents  $p/q$  and  $r/q$  we get

$$\begin{aligned}
 F_2^q &\leq q \int_0^\infty \frac{G^q v}{V^{q+1}} W = q \int_0^\infty \left\{ \left( \frac{G}{V} \right)^q v^{1-q/r} \right\} \left\{ W V^{-1} v^{q/r} \right\} \\
 &\leq q \left( \int_0^\infty W^{r/q} V^{-r/q} v \right)^{q/r} \left( \int_0^\infty \left( \frac{G}{V} \right)^p v \right)^{q/p}.
 \end{aligned}$$

Applying the lemma and the last part of (19) we find that

$$F_2^q \leq q \alpha^q(p, p) (p'/p)^{q/p} (p/q)^{q/r} B_0^q \left( \int_0^\infty f^p v \right)^{q/p}.$$

The estimate of  $F_3^q$  in the case  $0 < q < p < \infty$ ,  $p > 1$  is produced using Theorem 1(b) exactly as above; we get

$$F_3^q \leq \beta_2^q(p, q) \alpha^q(p, p) (p'/p)^{q/p} B_1^q \left( \int_0^\infty f^p v \right)^{q/p}.$$

Thus the sufficient parts of Theorem 2(b) and Theorem 3 are proved if  $\int_0^\infty v = \infty$ .

If  $0 < \int_0^\infty v < \infty$ , then the estimate must be slightly corrected. In this case we have instead of (17)

$$F_2^q \leq \left[ \int_0^\infty h V \right]^q \left( \int_0^\infty v \right)^{-q} \int_0^\infty w + q \int_0^\infty \frac{G^q v}{V^{q+1}} W.$$

The first term on the right side of this inequality can be estimated trivially since

$$\int_0^\infty h V = \int_0^\infty f v \leq \left( \int_0^\infty f^p v \right)^{1/p} \left( \int_0^\infty v \right)^{1/p'};$$

hence we get again that

$$F_2^q \ll A_0^q \left( \int_0^\infty f^p v \right)^{q/p} \quad \text{for } 1 < p \leq q < \infty$$

or

$$F_2^q \ll B_0^q \left( \int_0^\infty f^p v \right)^{q/p} \quad \text{for } 0 < q < p < \infty, \quad p > 1,$$

because  $A_0 \ll B_0$ .

## 3. PROOF OF THEOREM 3

3.1. The sufficient part of Theorem 3(a) has been proved in §2.4 above. We will show the necessary part exactly by the method of §2.3. We only need to modify the estimate of the term denoting  $J(s)$ . If  $0 < q < 1 < p < \infty$ , then  $q' < 0$  and we have

$$\begin{aligned} J(s) &\equiv \int_0^s \left( \int_y^s \left( \int_0^t \Psi_0^{1-p'} \right)^{r/q'} \Psi_0^{1-p'}(t) V^{-1}(t) dt \right)^{1/p} dy \\ &\geq \left( \int_0^s \Psi_0^{1-p'} \right)^{r/q'p} J_2(s). \end{aligned}$$

It has already been shown in §2.2 that

$$J_2(s) = \int_0^s \left[ \int_y^s \Psi_0^{1-p'}(\tau) V^{-1}(\tau) d\tau \right]^{1/p} dy \geq p(p')^{-1/p} \int_0^s \Psi_0^{1-p'}.$$

Thus we get

$$J(s) \geq p(p')^{-1/p} \left( \int_0^s \Psi_0^{1-p'} \right)^{r/p'q}$$

and the rest of the reasoning is exactly the same as above in §2.3.

3.2. Now we consider Theorem 3(b), i.e.,  $0 < p \leq q < \infty$ ,  $0 < p < 1$ . The necessary part is trivial. The sufficient part is demonstrated as follows. As above we use the notation

$$f(x) = \int_x^\infty h, \quad h \geq 0, \quad \text{supp } h \subset (0, \infty), \quad \int_0^\infty f^p v < \infty.$$

If  $0 < p < 1$  then applying Minkowski's inequality we get

$$\begin{aligned} \int_0^x f(s) ds &= \int_0^x \left( \int_s^\infty h \right) ds = p^{1/p} \int_0^x \left[ \int_s^\infty \left( \int_y^\infty h \right)^{p-1} h(y) dy \right]^{1/p} ds \\ &\leq 2^{-1/p'} p^{1/p} \left\{ \int_0^x \left[ \int_s^x \left( \int_y^\infty h \right)^{p-1} h(y) dy \right]^{1/p} ds \right. \\ &\quad \left. + \int_0^x \left[ \int_x^\infty \left( \int_y^\infty h \right)^{p-1} h(y) dy \right]^{1/p} ds \right\} \\ &\leq 2^{-1/p'} p^{1/p} \left( \int_0^x \left( \int_y^\infty h \right)^{p-1} h(y) y^p dy \right)^{1/p} + 2^{-1/p'} x f(x). \end{aligned}$$

Now we have

$$\begin{aligned} I &= \left( \int_0^\infty \left( \frac{1}{x} \int_0^x f(t) dt \right)^q w(x) dx \right)^{1/q} \ll \left( \int_0^\infty f^q w \right)^{1/q} \\ &\quad + \left( \int_0^\infty \left( \int_0^x \left( \int_y^\infty h \right)^{p-1} h(y) y^p dy \right)^{q/p} x^{-q} w(x) dx \right)^{1/q} \\ &\equiv I_1 + I_2. \end{aligned}$$

Proposition 1(a) yields

$$I_1 \ll A_0 \left( \int_0^\infty f^p(x)v(x) dx \right)^{1/p}.$$

Applying Minkowski's inequality we find that

$$\begin{aligned} I_2 &\leq \left( \int_0^\infty \left( \int_y^\infty h \right)^{p-1} h(y)y^p dy \left( \int_y^\infty x^{-q}w(x) dx \right)^{p/q} dy \right)^{1/p} \\ &\leq \mathcal{A}_1 \left( \int_0^\infty \left( \int_y^\infty h \right)^{p-1} h(y)V(y) dy \right)^{1/p} \\ &= p^{-1/p} \mathcal{A}_1 \left( \int_0^\infty f^p(x)v(x) dx \right)^{1/p}. \end{aligned}$$

Combination of these upper bounds on  $I_1$  and  $I_2$  completes the proof.

3.3. The following result is a sufficient condition for (4) if  $0 < q < p < 1$ .

**Proposition 2.** *Let  $0 < q < p < 1$ . Then the inequality (4) is valid for all nonincreasing functions  $f \geq 0$ , if  $B_0 < \infty$  and*

$$D_1 \equiv \left( \int_0^\infty \left( \int_t^\infty x^{-q}w(x) dx \right)^{r/q} V^{-r/p}(t)t^{r-1} dt \right)^{1/r} < \infty.$$

*Proof.* Let us denote  $\int_t^\infty x^{-q}w(x) dx$  by  $\mathcal{W}_0(t)$  and suppose that  $f(x) = \int_x^\infty h$  as above. It has already been shown that

$$\int_0^x f(s)dx \leq 2^{-1/p'} p^{1/p} \left( \int_0^x \left( \int_y^\infty h \right)^{p-1} h(y)y^p dy \right)^{1/p} + 2^{-1/p'} x f(x).$$

Then we have as above

$$I = \left( \int_0^\infty \left( \frac{1}{x} \int_0^x f(t) dt \right)^q w(x) dx \right)^{1/q} \ll I_1 + I_2.$$

Applying Proposition 1(b) we get

$$I_1 \ll B_0 \left( \int_0^\infty f^p(x)v(x) dx \right)^{1/p}.$$

An upper bound on  $I_2$  is obtained as follows.

$$\begin{aligned} I_2^q &\ll \int_0^\infty \left( \int_0^x \left( \int_y^\infty h \right)^{p-1} h(y)y^p dy \right)^{q/p} x^{-q}w(x) dx \\ &= \int_0^\infty \left[ \left( \int_0^x \left( \int_y^\infty h \right)^{p-1} h(y)y^p dy \right)^{q/p} x^{-q^2/p} w^{q/p}(x) \right. \\ &\quad \cdot \left. \left( \int_0^x \mathcal{W}_0^{r/p}(t)V^{-r/p}(t)t^{r-1} dt \right)^{-q/r} \right] \\ &\quad \cdot \left[ x^{-q(1-q/p)} w^{1-q/p}(x) \left( \int_0^x \mathcal{W}_0^{r/p}(t)V^{-r/p}(t)t^{r-1} dt \right)^{q/r} \right] dx \end{aligned}$$

(applying Hölder's inequality with exponents  $p/q$ ,  $r/q$ ),

$$\begin{aligned} &\leq \left( \int_0^\infty \left( \int_0^x \left( \int_y^\infty h \right)^{p-1} h(y) y^p dy \right) x^{-q} w(x) \right. \\ &\quad \cdot \left. \left( \int_0^x \mathscr{W}_0^{r/p}(t) V^{-r/p}(t) t^{r-1} dt \right)^{-p/r} dx \right)^{q/p} \\ &\quad \cdot \left( \int_0^\infty x^{-q} w(x) \int_0^x \mathscr{W}_0^{r/p}(t) V^{-r/p}(t) t^{r-1} dt \right)^{q/r} \\ &= D_1^q \left( \int_0^\infty \left( \int_y^\infty h \right)^{p-1} h(y) y^p \int_y^\infty x^{-q} w(x) \right. \\ &\quad \cdot \left. \left( \int_0^x \mathscr{W}_0^{r/p}(t) V^{-r/p}(t) t^{r-1} dt \right)^{-p/r} dx dy \right)^{q/p}. \end{aligned}$$

Let us note that

$$\begin{aligned} &y^p \int_y^\infty x^{-q} w(x) \left( \int_0^x \mathscr{W}_0^{r/p}(t) V^{-r/p}(t) t^{r-1} dt \right)^{-p/r} dx \\ &\leq y^p \left( \int_0^y \mathscr{W}_0^{r/p}(t) V^{-r/p}(t) t^{r-1} dt \right)^{-p/r} \mathscr{W}_0(y) \leq y^p V(y) \left( \int_0^y t^{r-1} dt \right)^{-p/r} \\ &= r^{p/r} V(y). \end{aligned}$$

This implies the estimate

$$I_2^q \ll D_1^q \left( \int_0^\infty f^p v \right)^{q/p},$$

and ends the proof.

## REFERENCES

1. M. Arino and B. Muckenhoupt, *Maximal function on classical Lorentz spaces and Hardy's inequality with weights for nonincreasing functions*, Trans. Amer. Math. Soc. **320** (1990), 727–735.
2. D. W. Boyd, *The Hilbert transform on rearrangement-invariant spaces*, Canad. J. Math. **19** (1967), 599–616.
3. J. S. Bradley, *Hardy inequalities with mixed norms*, Canad. Math. Bull. **21** (1978), 405–408.
4. S. G. Krein, Yu. I. Petunin and E. M. Semenov, *Interpolation of linear operators*, Transl. Math. Monographs, vol. 54, Amer. Math. Soc., Providence, R.I., 1982.
5. V. G. Maz'ja, *Sobolev spaces*, Springer-Verlag, Berlin, 1985.
6. B. Muckenhoupt, *Hardy's inequality with weights*, Studia Math. **44** (1972), 31–38.
7. C. J. Neugebauer, *Weighted norm inequalities for averaging operators of monotone functions*, Publ. Math. (to appear).
8. E. T. Sawyer, *Boundedness of classical operators on classical Lorentz spaces*, Studia Math. **96** (1990), 145–158.
9. G. Sinnamon, *A weighted gradient inequality*, Proc. Roy. Soc. Edinburgh Sect. A **111** (1989), 329–335.
10. —, *Weighted Hardy and Opial-type inequalities*, J. Math. Anal. Appl. **160** (1991), 434–445.

11. V. D. Stepanov, *Weighted inequalities for a class of Volterra convolution operators*, J. London Math. Soc. (2) **45** (1992), 232–242.
12. G. Talenti, *Osservazione sopra una classe di disuguaglianze*, Rend. Sem. Mat. Fis. Milano **39** (1969), 171–185.
13. G. Tomaselli, *A class of inequalities*, Boll. Un. Mat. Ital. Ser. **4** (1969), 622–631.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA, TUSCALOOSA, ALABAMA 35487-0350

*Current address:* Institute for Applied Mathematics, Shelest 118–205, Khabarovsk 680042, Russia

*E-mail address:* vlads@hpicnit.khabarovsk.su