

RELATIVELY FREE INVARIANT ALGEBRAS OF FINITE REFLECTION GROUPS

MÁTYÁS DOMOKOS

ABSTRACT. Let G be a finite subgroup of $Gl_n(K)$ (K is a field of characteristic 0 and $n \geq 2$) acting by linear substitution on a relatively free algebra $K\langle x_1, \dots, x_n \rangle / I$ of a variety of unitary associative algebras. The algebra of invariants is relatively free if and only if G is a pseudo-reflection group and I contains the polynomial $[[x_2, x_1], x_1]$.

1. INTRODUCTION

Throughout the paper K is a field of characteristic 0 and $K\langle x_1, \dots, x_n \rangle$ is the unitary free associative K -algebra of rank n . The general linear group $Gl_n = Gl_n(K)$ acts on the free algebra by linear substitution. More explicitly, if $g = (g_{ij}) \in Gl_n$, then

$$g \cdot x_j = \sum_{i=1}^n g_{ij} x_i,$$

and for any $f(x_1, \dots, x_n) \in K\langle x_1, \dots, x_n \rangle$

$$g \cdot f = f(g \cdot x_1, \dots, g \cdot x_n).$$

An ideal I of the unitary free associative algebra $K\langle x_1, x_2, \dots \rangle$ of countable rank is called a T -ideal if I is invariant under any K -algebra endomorphism of the free algebra, that is, $f(x_1, \dots, x_n) \in I$ implies $f(u_1, \dots, u_n) \in I$ for any $u_1, \dots, u_n \in K\langle x_1, x_2, \dots \rangle$.

$I_n = I \cap K\langle x_1, \dots, x_n \rangle$ is invariant under the action of Gl_n , so

$$F_n(I) = K\langle x_1, \dots, x_n \rangle / I_n$$

inherits the Gl_n -structure. $F_n(I)$ is a relatively free algebra of rank n of the variety of unitary associative algebras satisfying all the identities $f = 0$ with $f \in I$. Denote by y_i the image of x_i under the natural homomorphism

$$K\langle x_1, \dots, x_n \rangle \rightarrow K\langle x_1, \dots, x_n \rangle / I_n.$$

Received by the editors September 7, 1994.

1991 *Mathematics Subject Classification*. Primary 16W20; Secondary 16R10.

Key words and phrases. Relatively free algebra, reflection group, invariants, Hilbert series.

This research was partially supported by Széchenyi István Scholarship Foundation and by Hungarian National Foundation for Scientific Research grant no. T4265.

Let G be a finite subgroup of Gl_n and denote by

$$F_n(I)^G = \{f \in F_n(I) \mid g \cdot f = f \forall g \in G\}$$

the algebra of invariants of G . We refer to the article of Formanek [8] for a survey of the results on the algebra of invariants of a finite linear group acting on a relatively free algebra. In the special case of the variety of all unitary commutative algebras, that is, when I is generated by the commutator $[x_1, x_2] = x_1x_2 - x_2x_1$, the relatively free algebra

$$F_n(I) = K[x_1, \dots, x_n]$$

is the n variable commutative polynomial algebra, and the study of $F_n(I)^G$ is the topic of classical invariant theory.

An element $g \in Gl_n(K)$ is called a *pseudo-reflection*, if it fixes an $n - 1$ dimensional subspace of K^n . If g is of finite order then g is a pseudo-reflection if and only if g has the eigenvalue 1 with multiplicity $n - 1$. A finite subgroup $G < Gl_n$ is called a *pseudo-reflection group* if it is generated by pseudo-reflections. Our starting point is the following famous result (Shephard-Todd [13], Chevalley [2]):

Theorem 1.1. *$K[x_1, \dots, x_n]^G$ is a polynomial algebra if and only if G is a pseudo-reflection group. Moreover, if G is a pseudo-reflection group, then there exist n algebraically independent homogeneous forms which generate $K[x_1, \dots, x_n]^G$. The degrees $d_1 \leq \dots \leq d_n$ of these forms are uniquely determined by G , and are called the degrees of G . We have the equality $|G| = \prod_{i=1}^n d_i$, and the number of pseudo-reflections in G is $\sum_{i=1}^n (d_i - 1)$.*

We shall extend the notion of algebraic independence to any variety of unitary associative K -algebras. We fix a T-ideal I , and denote by $\mathcal{V}(I)$ the variety defined by I . Let R be an algebra in $\mathcal{V}(I)$. We say that the elements $f_1, \dots, f_m \in R$ are *algebraically independent*, if $h(f_1, \dots, f_m) = 0$ for some $h \in K\langle x_1, \dots, x_m \rangle$ implies that $h \in I$. Note that this notion of algebraic independence depends on the T-ideal I and belongs to the variety $\mathcal{V}(I)$.

Guralnick proved in [9] that if $n \geq 2$, $k \geq 2$ and I is the T-ideal of the $k \times k$ matrix algebra over K , then $F_n(I)^G$ is not relatively free for any finite group G . As it was pointed out in [8, p. 105], this result implies easily that $F_n(I)^G$ is not relatively free if I is contained in the T-ideal of the 2×2 matrix algebra. In this paper we give a complete answer to the question of determining when $F_n(I)^G$ is relatively free. The answer was conjectured by Drensky [7].

The case $n = 1$ is trivial, because then $F_1(I) \cong K[x_1] \cong K\langle x_1 \rangle$ and G is a cyclic group acting by scalar multiplication, so the algebra of invariants is $K[x^m]$ for some positive integer m .

Theorem 1.2. *Let $G < Gl_n(K)$ ($n \geq 2$) be a finite group and let I be a T-ideal. Then $F_n(I)^G$ is generated by algebraically independent elements if and only if G is a pseudo-reflection group and I contains the polynomial $[[x_2, x_1], x_1]$.*

2. PRELIMINARIES

The algebra $K\langle x_1, \dots, x_n \rangle^G$ is always free and almost never finitely generated by [4] and [10]. However, working in a proper subvariety of the variety of all unitary associative K -algebras the notion of transcendence degree makes sense.

Proposition 2.1. *Fix a non-zero T-ideal I and consider the variety $\mathcal{V}(I)$. If R is an algebra in the variety $\mathcal{V}(I)$ generated by n elements, then it does not contain more than n algebraically independent elements.*

Proof. R is a homomorphic image of $F_n(I)$ and obviously the preimages of algebraically independent elements of R are also algebraically independent, so it suffices to prove the proposition for the relatively free algebra $F_n(I)$. Assume that $f_1, \dots, f_m \in F_n(I)$ are algebraically independent. Let $J \subseteq K\langle x_1, x_2, \dots \rangle$ be the radical of I , that is,

$$J = \{f \in K\langle x_1, x_2, \dots \rangle \mid f^N \in I \text{ for some } N\}.$$

It is well known (see for example [12, Theorems 1.5.32, 2.4.7 and 3.2.6]) that J is the set of identities of the $k \times k$ matrix algebra for some $k \geq 1$. We have the natural onto homomorphism

$$K\langle x_1, \dots, x_n \rangle / I_n \rightarrow K\langle x_1, \dots, x_n \rangle / J_n.$$

Denote by h_1, \dots, h_m the images of f_1, \dots, f_m . Suppose that for some $p \in K\langle x_1, \dots, x_m \rangle$ we have $p(h_1, \dots, h_m) = 0$. Then there exists an integer N such that $p^N(f_1, \dots, f_m) = 0$ in $F_n(I)$, and the algebraic independence of the f_i s implies that $p^N(x_1, \dots, x_m) \in I_m$. Hence $p(x_1, \dots, x_m) \in J_m$ by the definition of J_m , and this shows the algebraic independence of h_1, \dots, h_m in $F_n(J)$.

The center of $F_m(J)$ is a commutative domain and its transcendence degree is $d = (m - 1)k^2 + 1$ (see for example [8, p. 105]). Let

$$c_1 = c_1(y_1, \dots, y_m), \dots, c_d = c_d(y_1, \dots, y_m)$$

be a transcendence basis of the center of $F_m(J)$ (so the $c_i(x_1, \dots, x_m)$ are central polynomials for the $k \times k$ matrix algebra). The algebraic independence of h_1, \dots, h_m in $F_n(J)$ together with the algebraic independence (in the ordinary sense) of c_1, \dots, c_d in the center of $F_m(J)$ implies that $c_i(h_1, \dots, h_m)$ ($i = 1, \dots, d$) are algebraically independent in the center of $F_n(J)$. But this center has transcendence degree $(n - 1)k^2 + 1$, implying that $m \leq n$. \square

We use the following crucial argument from [8] or [9]. Any non-zero T-ideal I is contained in the T-ideal generated by $[x_1, x_2]$. So we have the natural onto homomorphism

$$F_n(I) \rightarrow K[x_1, \dots, x_n]$$

which commutes with the action of G , and by the complete reducibility of this action the homomorphism

$$F_n(I)^G \rightarrow K[x_1, \dots, x_n]^G$$

is also onto. Therefore the image of a generating set of $F_n(I)^G$ is a generating set of $K[x_1, \dots, x_n]^G$, hence $F_n(I)^G$ can not be generated by less than n elements. Thus if $F_n(I)^G$ is generated by algebraically independent elements, then by Proposition 2.1 it is generated by n elements.

Assume that f_1, \dots, f_n generate $F_n(I)^G$. Their images in $K[x_1, \dots, x_n]$ generate $K[x_1, \dots, x_n]^G$, hence G must be a pseudo-reflection group. We may assume by [9, Lemma 2] that f_1, \dots, f_n are homogeneous, therefore their degrees are the degrees of G . Moreover, the proof of [9, Lemma 2] shows that if $h_1, \dots, h_n \in F_n(I)^G$ such that their images generate $K[x_1, \dots, x_n]^G$, then $F_n(I)^G = K\langle h_1, \dots, h_n \rangle$.

By the above discussion Theorem 1.2 splits into the next two statements:

Theorem 2.2. *Let G be a pseudo-reflection group and let I be a T -ideal containing $[[x_2, x_1], x_1]$. If f_1, \dots, f_n are homogeneous elements in $F_n(I)^G$ such that their images generate $K[x_1, \dots, x_n]^G$, then f_1, \dots, f_n are algebraically independent in $F_n(I)$ and they generate $F_n(I)^G$. In particular, $F_n(I)^G \cong F_n(I)$.*

Theorem 2.3. *If the T -ideal I does not contain the polynomial $[[x_2, x_1], x_1]$, then $F_n(I)^G$ can not be generated by n elements for any $n \geq 2$ and finite group $G < Gl_n(K)$.*

We recall some basic facts about the T -ideals of the unitary free associative K -algebra. Let B denote the subalgebra of $K\langle x_1, x_2, \dots \rangle$ generated by all the commutators $[x_{i_1}, \dots, x_{i_r}]$ with $r \geq 2$, where $[x_1, \dots, x_r]$ is defined inductively by $[x_1, \dots, x_r] = [[x_1, \dots, x_{r-1}], x_r]$ for $r \geq 3$. The elements of B are called *proper* polynomials, and they can be characterised as the polynomials with zero partial derivatives. Clearly, $B_n = B \cap K\langle x_1, \dots, x_n \rangle$ is a Gl_n -submodule of $K\langle x_1, \dots, x_n \rangle$. Drensky showed (see [5, Theorem 2.2]) that any T -ideal I of the free unitary algebra is generated by the proper polynomials that it contains, and as Gl_n -modules

$$(2.1) \quad F_n(I) \cong K[x_1, \dots, x_n] \otimes (B_n/B_n \cap I).$$

Both I_n and B_n are multigraded subalgebras with respect to the usual multigrading on $K\langle x_1, \dots, x_n \rangle$, so $F_n(I)$ and $B_n/B_n \cap I$ inherit the multigrading.

For any $\alpha = (\alpha_1, \dots, \alpha_n)$ and multigraded algebra R we put

$$R^{(\alpha)} = \{f \in R \mid f \text{ is multihomogeneous of multidegree } \alpha\},$$

and for any \mathbb{N} -graded algebra R and non-negative integer d

$$R^{(d)} = \{f \in R \mid f \text{ is homogeneous of total degree } d\}.$$

Obviously $B_n^{(0)} = K$ and $B_n^{(1)} = \emptyset$. If $n \geq 2$, then $B_n^{(2)}$ is an irreducible Gl_n -module generated by $[x_1, x_2]$, and $B_n^{(3)}$ is an irreducible Gl_n -module generated by $[x_2, x_1, x_1]$. Therefore there exists a unique maximal T -ideal M among the T -ideals not containing $[x_2, x_1, x_1]$, M is generated by the proper polynomials of degree greater than 3 (in particular, M has no elements of degree less than 4). If I is a T -ideal not containing $[x_2, x_1, x_1]$, then $F_n(M)^G$ is a homomorphic image of $F_n(I)^G$, thus Theorem 2.3 is a consequence of the following more special statement:

Proposition 2.4. *Let M be the T -ideal generated by the proper polynomials of degree greater than 3, and let $G < Gl_n$ ($n \geq 2$) be a finite pseudo-reflection group. Then $F_n(M)^G$ can not be generated by n elements.*

The Hilbert series of a multigraded algebra R is an n variable formal power series defined by

$$H(R; t_1, \dots, t_n) = \sum_{\alpha=(\alpha_1, \dots, \alpha_n)} \dim_K(R^{(\alpha)}) t_1^{\alpha_1} \dots t_n^{\alpha_n},$$

and the Hilbert series of an \mathbb{N} -graded algebra R is the formal power series

$$H(R; t) = \sum_{d=0}^{\infty} \dim_K(R^{(d)}) t^d.$$

Let G be a finite subgroup of $Gl_n(K)$. For any element $g \in G$ denote by $\omega_1(g), \dots, \omega_n(d)$ the eigenvalues of g in the algebraic closure of K . $F_n(I)^G$ is an \mathbb{N} -graded algebra, and we have the noncommutative Molien-Weyl formula (see [8, Theorem 7]):

$$(2.2) \quad H(F_n(I)^G; t) = \frac{1}{|G|} \sum_{g \in G} H(F_n(I); \omega_1(g)t, \dots, \omega_n(g)t).$$

3. PROOF OF THEOREM 2.2

Denote by J the T-ideal generated by $[x_2, x_1, x_1]$, and let

$$s_d(x_1, \dots, x_d) = \sum_{\pi \in Sym(d)} (-1)^\pi x_{\pi(1)} \dots x_{\pi(d)}$$

be the *standard polynomial*. It is well known (see [6, 3.2.1. Theorem]) that $B_n^{(d)}/B_n^{(d)} \cap J = 0$, if d is odd or $d > n$, and it is an irreducible Gl_n -module generated by $s_d(y_1, \dots, y_d)$ if d is even and $d \leq n$. Let $J(m)$ be the T-ideal generated by $[x_2, x_1, x_1]$ and $s_{2m}(x_1, \dots, x_{2m})$, so the only T-ideals containing $[x_2, x_1, x_1]$ are J and $J(m)$ ($m = 1, 2, \dots$). We note that

$$K\langle x_1, \dots, x_n \rangle / J(m)_n = K\langle x_1, \dots, x_n \rangle / J_n \quad \text{if } 2m > n.$$

Let $f_1, \dots, f_n \in F_n(J(m))^G$ be homogeneous invariants such that their images in $K[x_1, \dots, x_n]$ generate $K[x_1, \dots, x_n]^G$. We prove by induction on m that f_1, \dots, f_n are algebraically independent in $F_n(J(m))$. In the case $m = 1$ there is nothing to prove. Suppose that the statement is true for $F_n(J(k))$, where $k \leq m$. We may assume that $2m \leq n$. The algebra $F_n(J(m+1))$ has a linear basis (see [6])

$$\{y_1^{\alpha_1} \dots y_n^{\alpha_n} s_{2k}(y_{i_1}, \dots, y_{i_{2k}}) \mid \alpha_1, \dots, \alpha_n \in \mathbb{N}, 0 \leq k \leq m, 1 \leq i_1 < \dots < i_{2k} \leq n\}.$$

Suppose that $h(f_1, \dots, f_n) = 0$ in $F_n(J(m+1))$ for some $h \in K\langle x_1, \dots, x_n \rangle$ with $h \notin J(m+1)_n$. Then we may assume that h is of the form

$$h = \sum_{\alpha=(\alpha_1, \dots, \alpha_n)} \sum_{k=0}^m \sum_{1 \leq i_1 < \dots < i_{2k} \leq n} a_{\alpha, k, i} x_1^{\alpha_1} \dots x_n^{\alpha_n} s_{2k}(x_{i_1}, \dots, x_{i_{2k}}),$$

where at least one of the coefficients $a_{\alpha, k, i}$ is non-zero. Let k_0 be the minimal k such that $a_{\alpha, i, k} \neq 0$ for some α, i . If $k_0 \leq m - 1$, then by the induction hypothesis

$$\sum_{\alpha, i} a_{\alpha, k_0, i} f_1^{\alpha_1} \dots f_n^{\alpha_n} s_{2k_0}(f_{i_1}, \dots, f_{i_{2k_0}}) \notin J_n(k_0 + 1) / J_n(m + 1).$$

On the other hand

$$\sum_{\alpha, k > k_0, i} a_{\alpha, k, i} f_1^{\alpha_1} \dots f_n^{\alpha_n} s_{2k}(f_{i_1}, \dots, f_{i_{2k}}) \equiv 0 \pmod{J_n(k_0 + 1) / J_n(m + 1)},$$

contradicting that $h(f_1, \dots, f_n) = 0$ in $F_n(J(m + 1))$. Thus $k_0 = m$ and we have

$$\sum_{\alpha, i} a_{\alpha, i} f_1^{\alpha_1} \dots f_n^{\alpha_n} s_{2m}(f_{i_1}, \dots, f_{i_{2m}}) = 0$$

in $F_n(J(m + 1))$.

Define the maps

$$\frac{\partial}{\partial y_i} : F_n(J(m + 1)) \rightarrow F_n(J(m + 1)) \quad (i = 1, \dots, n)$$

in the following way. If $f = f(y_1, \dots, y_n) \in F_n(J(m + 1))$ is multihomogeneous of multidegree $\alpha = (\alpha_1, \dots, \alpha_n)$, then let $\frac{\partial}{\partial y_i} f$ be the multihomogeneous component of multidegree $(\alpha_1, \dots, \alpha_i - 1, \dots, \alpha_n)$ of $f(y_1 + 1, \dots, y_n + 1)$ ($\frac{\partial}{\partial y_i} f = 0$ if $\alpha_i = 0$). Now extend $\frac{\partial}{\partial y_i}$ to $F_n(J(m + 1))$ by linearity. Clearly, $\frac{\partial}{\partial y_i}$ is a derivation.

Lemma 3.1. *For any $f, g \in F_n(J(m + 1))$ we have the equality*

$$[f, g] \equiv \sum_{1 \leq i, j \leq n} \frac{\partial f}{\partial y_i} \frac{\partial g}{\partial y_j} [y_i, y_j] \quad (\text{modulo } C^2),$$

where

$$C = F_n(J(m + 1))[F_n(J(m + 1)), F_n(J(m + 1))]F_n(J(m + 1))$$

is the commutator ideal of $F_n(J(m + 1))$.

Proof. By the multilinearity of the derivations and $[,]$ it suffices to prove the lemma when

$$f = y_{i_1} \dots y_{i_k} \text{ and } g = y_{j_1} \dots y_{j_l}$$

are monomials. We have

$$\begin{aligned} & (y_{i_1} \dots y_{i_k})(y_{j_1} \dots y_{j_l}) \\ &= y_{i_1} \dots y_{i_{k-1}} y_{j_1} y_{i_k} y_{j_2} \dots y_{j_l} + y_{i_1} \dots y_{i_{k-1}} [y_{i_k}, y_{j_1}] y_{j_2} \dots y_{j_l} \\ &= (y_{i_1} \dots y_{i_{k-1}})(y_{j_2} \dots y_{j_l}) [y_{i_k}, y_{j_1}] + y_{i_1} \dots y_{i_{k-1}} y_{j_1} y_{i_k} y_{j_2} \dots y_{j_l} \end{aligned}$$

(the second equality follows from the fact that any commutator lies in the center of $F_n(J(m + 1))$). Now exchange $y_{i_{k-1}}$ and y_{j_1} in the second term of the above sum, that is, replace this term by

$$y_{i_1} \dots y_{i_{k-2}} y_{j_1} y_{i_{k-1}} y_{i_k} y_{j_2} \dots y_{j_l} + (y_{i_1} \dots y_{i_{k-2}} y_{j_k})(y_{j_2} \dots y_{j_l}) [y_{i_{k-1}}, y_{j_1}].$$

Continuing this process we obtain

$$\begin{aligned} & (y_{i_1} \dots y_{i_k})(y_{j_1} \dots y_{j_l}) \\ &= y_{j_1} (y_{i_1} \dots y_{i_k})(y_{j_2} \dots y_{j_l}) + \sum_{r=1}^k (y_{i_1} \dots \hat{y}_{i_r} \dots y_{i_k})(y_{j_2} \dots y_{j_l}) [y_{i_r}, y_{j_1}], \end{aligned}$$

where the sign \hat{y}_{i_r} means that we delete y_{i_r} in the word $y_{i_1} \dots y_{i_k}$. On applying the same process to the word $(y_{i_1} \dots y_{i_k})(y_{j_2} \dots y_{j_l})$ we obtain that

$$\begin{aligned} (y_{i_1} \dots y_{i_k})(y_{j_1} \dots y_{j_l}) &= \sum_{r=1}^k (y_{i_1} \dots \hat{y}_{i_r} \dots y_{i_k})(y_{j_2} \dots y_{j_l})[y_{i_r}, y_{j_1}] \\ &\quad + \sum_{r=1}^k y_{j_1} (y_{i_1} \dots \hat{y}_{i_r} \dots y_{i_k})(y_{j_3} \dots y_{j_l})[y_{i_r}, y_{j_2}] \\ &\quad \quad + y_{j_1} y_{j_2} (y_{i_1} \dots y_{i_k})(y_{j_3} \dots y_{j_l}) \\ &\stackrel{(\text{mod } C^2)}{\equiv} \sum_{r=1}^k (y_{i_1} \dots \hat{y}_{i_r} \dots y_{i_k})(\hat{y}_{j_1} y_{j_2} \dots y_{j_l})[y_{i_r}, y_{j_1}] \\ &\quad + \sum_{r=1}^k (y_{i_1} \dots \hat{y}_{i_r} \dots y_{i_k})(y_{j_1} \hat{y}_{j_2} y_{j_3} \dots y_{j_l})[y_{i_r}, y_{j_2}] \\ &\quad \quad + y_{j_1} y_{j_2} (y_{i_1} \dots y_{i_k})(y_{j_3} \dots y_{j_l}) \\ &= \sum_{i=1}^n \left(\frac{\partial}{\partial y_i} (y_{i_1} \dots y_{i_k}) \right) (\hat{y}_{j_1} y_{j_2} \dots y_{j_l}) [y_i, y_{j_1}] \\ &\quad + \sum_{i=1}^n \left(\frac{\partial}{\partial y_i} (y_{i_1} \dots y_{i_k}) \right) (y_{j_1} \hat{y}_{j_2} \dots y_{j_l}) [y_i, y_{j_2}] \\ &\quad \quad + y_{j_1} y_{j_2} (y_{i_1} \dots y_{i_k})(y_{j_3} \dots y_{j_l}). \end{aligned}$$

Repeating this algorithm finally we get

$$\begin{aligned} &(y_{i_1} \dots y_{i_k})(y_{j_1} \dots y_{j_l}) \\ &\stackrel{(\text{mod } C^2)}{\equiv} (y_{j_1} \dots y_{j_l})(y_{i_1} \dots y_{i_k}) + \sum_{i=1}^n \sum_{r=1}^l \left(\frac{\partial}{\partial y_i} (y_{i_1} \dots y_{i_k}) \right) (y_{j_1} \dots \hat{y}_{j_r} \dots y_{j_l}) [y_i, y_{j_r}] \\ &= (y_{j_1} \dots y_{j_l})(y_{i_1} \dots y_{i_k}) + \sum_{i=1}^n \sum_{j=1}^n \left(\frac{\partial}{\partial y_i} (y_{i_1} \dots y_{i_k}) \right) \left(\frac{\partial}{\partial y_j} (y_{j_1} \dots y_{j_l}) \right) [y_i, y_j], \end{aligned}$$

which explicitly shows the claim. □

Now we use Lemma 3.1 to rewrite $A = s_{2m}(f_1, \dots, f_{2m})$. By the definition of s_{2m}

$$A = \frac{1}{2^m} \sum_{\pi \in \text{Sym}(2m)} (-1)^\pi [f_{\pi(1)}, f_{\pi(2)}] \dots [f_{\pi(2m-1)}, f_{\pi(2m)}].$$

We have $C^{m+1} = 0$ in $F_n(J(m+1))$, because the commutators are central elements of $F_n(J(m+1))$ and any proper polynomial of degree greater than $2m+1$ is contained in $J(m+1)$. Therefore in the right hand side of the above equality we may replace $[f_{\pi(2k-1)}, f_{\pi(2k)}]$ by something that is congruent with it modulo C^2 .

So by Lemma 3.1 we have

$$\begin{aligned}
 A &= \frac{1}{2^m} \sum_{\pi \in \text{Sym}(2m)} (-1)^\pi \sum_{(i_1, \dots, i_{2m})} \frac{\partial f_{\pi(1)}}{\partial y_{i_1}} \frac{\partial f_{\pi(2)}}{\partial y_{i_2}} [y_{i_1}, y_{i_2}] \\
 &\quad \dots \frac{\partial f_{\pi(2m-1)}}{\partial y_{i_{2m-1}}} \frac{\partial f_{\pi(2m)}}{\partial y_{i_{2m}}} [y_{i_{2m-1}}, y_{i_{2m}}] \\
 &= \frac{1}{2^m} \sum_{\pi \in \text{Sym}(2m)} (-1)^\pi \sum_{(i_1, \dots, i_{2m})} \frac{\partial f_{\pi(1)}}{\partial y_{i_1}} \dots \frac{\partial f_{\pi(2m)}}{\partial y_{i_{2m}}} [y_{i_1}, y_{i_2}] \dots [y_{i_{2m-1}}, y_{i_{2m}}].
 \end{aligned}$$

The polynomial $[x_1, x_2][x_1, x_3]$ is contained in J , hence if the i_1, \dots, i_{2m} are not pairwise different, then the corresponding term is zero in the above expression, implying that

$$\begin{aligned}
 A &= \frac{1}{2^m} \sum_{\pi \in \text{Sym}(2m)} (-1)^\pi \sum_{1 \leq i_1 < \dots < i_{2m} \leq n} \sum_{\rho \in \text{Sym}(2m)} \frac{\partial f_{\pi(1)}}{\partial y_{i_{\rho(1)}}} \\
 &\quad \dots \frac{\partial f_{\pi(2m)}}{\partial y_{i_{\rho(2m)}}} [y_{i_{\rho(1)}}, y_{i_{\rho(2)}}] \dots [y_{i_{\rho(2m-1)}}, y_{i_{\rho(2m)}}] \\
 &= \sum_{1 \leq i_1 < \dots < i_{2m} \leq n} \frac{1}{2^m} \sum_{\rho \in \text{Sym}(2m)} (-1)^\rho [y_{i_{\rho(1)}}, y_{i_{\rho(2)}}] \dots [y_{i_{\rho(2m-1)}}, y_{i_{\rho(2m)}}] \\
 &\quad \times \sum_{\pi \in \text{Sym}(2m)} (-1)^\pi (-1)^\rho \frac{\partial f_{\pi(1)}}{\partial y_{i_{\rho(1)}}} \dots \frac{\partial f_{\pi(2m)}}{\partial y_{i_{\rho(2m)}}}.
 \end{aligned}$$

Again by $C^{m+1} = 0$ we may permute $\frac{\partial f_{\pi(1)}}{\partial y_{i_{\rho(1)}}}, \dots, \frac{\partial f_{\pi(2m)}}{\partial y_{i_{\rho(2m)}}}$ among each other in the above expression, and we get

$$A = \sum_{1 \leq i_1 < \dots < i_{2m} \leq n} \sum_{\sigma \in \text{Sym}(2m)} (-1)^\sigma \frac{\partial f_1}{\partial y_{i_{\sigma(1)}}} \dots \frac{\partial f_{2m}}{\partial y_{i_{\sigma(2m)}}} s_{2m}(y_{i_1}, \dots, y_{i_{2m}}).$$

For any $1 \leq i_1 < \dots < i_{2m} \leq n$ and $1 \leq j_1 < \dots < j_{2m} \leq n$ we put

$$f_{j_1, \dots, j_{2m}}^{i_1, \dots, i_{2m}} = \sum_{\sigma \in \text{Sym}(2m)} (-1)^\sigma \frac{\partial f_{i_1}}{\partial y_{j_{\sigma(1)}}} \dots \frac{\partial f_{i_{2m}}}{\partial y_{j_{\sigma(2m)}}}.$$

Consider the natural homomorphism

$$\psi : F_n(J(m+1)) \rightarrow K[x_1, \dots, x_n].$$

The image of any $f \in F_n(J(m+1))$ has a normal form

$$\psi(f) = \sum_{\alpha=(\alpha_1, \dots, \alpha_n)} b_\alpha x_1^{\alpha_1} \dots x_n^{\alpha_n}.$$

Now let ϕ be the map $F_n(J(m+1)) \rightarrow F_n(J(m+1))$ defined by

$$\phi(f) = \sum_{\alpha} b_\alpha y_1^{\alpha_1} \dots y_n^{\alpha_n}.$$

Obviously, we have $\phi(f) \equiv f \pmod{C}$, hence $C^{m+1} = 0$ implies that

$$f s_{2m}(y_{i_1}, \dots, y_{i_{2m}}) = \phi(f) s_{2m}(y_{i_1}, \dots, y_{i_{2m}}).$$

Summarizing, we have

$$\begin{aligned} 0 &= h(f_1, \dots, f_n) \\ &= \sum_{\alpha=(\alpha_1, \dots, \alpha_n)} \sum_{1 \leq i_1 < \dots < i_{2m} \leq n} a_{\alpha,i} f_1^{\alpha_1} \dots f_n^{\alpha_n} s_{2m}(f_{i_1}, \dots, f_{i_{2m}}) \\ &= \sum_{1 \leq j_1 < \dots < j_{2m} \leq n} \phi \left(\sum_{\alpha,i} a_{\alpha,i} f_1^{\alpha_1} \dots f_n^{\alpha_n} f_{j_1, \dots, j_{2m}}^{i_1, \dots, i_{2m}} \right) s_{2m}(y_{j_1}, \dots, y_{j_{2m}}). \end{aligned}$$

Now since the elements $y_1^{\alpha_1} \dots y_n^{\alpha_n} s_{2m}(y_{i_1}, \dots, y_{i_{2m}})$ are linearly independent in $F_n(J(m+1))$,

$$\phi \left(\sum_{\alpha,i} a_{\alpha,i} f_1^{\alpha_1} \dots f_n^{\alpha_n} f_{j_1, \dots, j_{2m}}^{i_1, \dots, i_{2m}} \right)$$

must be zero for all j_1, \dots, j_{2m} , implying that

$$\psi \left(\sum_{\alpha,i} a_{\alpha,i} f_1^{\alpha_1} \dots f_n^{\alpha_n} f_{j_1, \dots, j_{2m}}^{i_1, \dots, i_{2m}} \right) = 0,$$

because the monomials $y_1^{\beta_1} \dots y_n^{\beta_n}$ are linearly independent in $F_n(J(m+1))$.

Introduce the symbols dx_1, \dots, dx_n , and let

$$E = K \langle dx_1, \dots, dx_n \mid dx_i dx_j = -dx_j dx_i, 1 \leq i, j \leq n \rangle$$

be the Grassmann algebra of an n dimensional linear space. Consider the map

$$d : K[x_1, \dots, x_n] \rightarrow K[x_1, \dots, x_n] \otimes E$$

defined by

$$df = d(f) = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i.$$

The commutative polynomials $h_i = \psi(f_i)$ ($i = 1, \dots, n$) form an algebraically independent generating set of $K[x_1, \dots, x_n]^G$. Direct computation shows that

$$\begin{aligned} &\sum_{\alpha,i} a_{\alpha,i} h_1^{\alpha_1} \dots h_n^{\alpha_n} dh_{i_1} \dots dh_{i_{2m}} \\ &= \sum_{1 \leq j_1 < \dots < j_{2m} \leq n} \psi \left(\sum_{\alpha,i} a_{\alpha,i} f_1^{\alpha_1} \dots f_n^{\alpha_n} f_{j_1, \dots, j_{2m}}^{i_1, \dots, i_{2m}} \right) dx_{j_1} \dots dx_{j_n}, \end{aligned}$$

and by our hypothesis the latter is zero. Now [14, Theorem] implies that

$$\{h_1^{\alpha_1} \dots h_n^{\alpha_n} dh_{i_1} \dots dh_{i_{2m}} \mid \alpha_1, \dots, \alpha_n \in \mathbb{N}, 1 \leq i_1 < \dots < i_{2m} \leq n\}$$

is a linearly independent subset of $K[x_1, \dots, x_n] \otimes E$, hence $a_{\alpha,i} = 0$ for every α, i , contradicting the hypothesis that $f_1, \dots, f_n \in F_n(J(m+1))$ are algebraically dependent.

Remark. The above argument did not use the fact that f_1, \dots, f_n are invariants. So we have proved for all $m = 1, 2, \dots$ that if $f_1, \dots, f_n \in F_n(J(m))$ are algebraically independent (in the ordinary sense) modulo the commutator ideal, then they are algebraically independent in $F_n(J(m))$.

We have to show that f_1, \dots, f_n generate $F_n(J(m))^G$. They are homogeneous of degree d_1, \dots, d_n , and as we pointed out earlier these are the degrees of G . We conclude from the formula (2.1) and the Gl_n -structure of $B_n/B_n \cap J(m)$ that the Hilbert series of $F_n(J(m))$ is

$$H(F_n(J(m)); t_1, \dots, t_n) = \frac{\sum_{i=0}^{m-1} \sigma_{2i}(t_1, \dots, t_n)}{\prod_{j=1}^n (1 - t_j)},$$

where σ_k is the k th elementary symmetric function ($\sigma_0 = 1$ and $\sigma_k = 0$ if $k > n$). We have the isomorphism

$$K\langle f_1, \dots, f_n \rangle \cong F_n(J(m)),$$

hence the \mathbb{N} -graded Hilbert series of the graded subalgebra $K\langle f_1, \dots, f_n \rangle$ of $F_n(J(m))^G$ is

$$H(F_n(J(m)); t^{d_1}, \dots, t^{d_n}) = \frac{\sum_{i=0}^{m-1} \sigma_{2i}(t^{d_1}, \dots, t^{d_n})}{\prod_{j=1}^n (1 - t^{d_j})}.$$

Using the noncommutative Molien-Weyl formula (2.2)

$$H(F_n(J(m))^G; t) = \frac{1}{|G|} \sum_{g \in G} \frac{\sum_{i=0}^{m-1} \sigma_{2i}(\omega_1(g)t, \dots, \omega_n(g)t)}{\prod_{j=1}^n (1 - \omega_j(g)t)}.$$

Solomon’s formula

$$(3.1) \quad \frac{1}{|G|} \sum_{g \in G} \frac{\sigma_p(\omega_1(g), \dots, \omega_n(g))}{\prod_{j=1}^n (1 - \omega_j(g)t)} = \frac{\sigma_p(t^{d_1-1}, \dots, t^{d_n-1})}{\prod_{j=1}^n (1 - t^{d_j})}$$

(see [14, formula (5)]) says that $F_n(J(m))^G$ has the same Hilbert series as its subalgebra $K\langle f_1, \dots, f_n \rangle$, therefore the two algebras must coincide. This finishes the proof of Theorem 2.2.

Remark. The polynomial $[x_2, x_1, x_1]$ appears in the theory of PI-algebras as a generator of the T-ideal of identities of the infinite dimensional Grassmann algebra (cf. [11]). It is interesting to note that both parts of the above proof use Solomon’s results from [14] on a pseudo-reflection group acting on the tensor product of $K[x_1, \dots, x_n]$ and the Grassmann algebra of an n dimensional vector space.

4. PROOF OF PROPOSITION 2.4

Let G be a pseudo-reflection group, and suppose that $F_n(M)^G$ is generated by n elements (where M is the T-ideal generated by all the proper polynomials of degree greater than 3). As we mentioned in Section 2, then $F_n(M)^G$ is generated by its homogeneous elements f_1, \dots, f_n if and only if their images in $K[x_1, \dots, x_n]$

generate $K[x_1, \dots, x_n]^G$. In the sequel we assume that $f_1, \dots, f_n \in F_n(M)^G$ have this property. By (2.1) the Hilbert series of $F_n(M)$ is

$$H(F_n(M); t_1, \dots, t_n) = \frac{1 + S_{(1,1)}(t_1, \dots, t_n) + S_{(2,1)}(t_1, \dots, t_n)}{\prod_{i=1}^n (1 - t_i)},$$

where $S_\lambda(t_1, \dots, t_n)$ is the Schur function corresponding to the partition $\lambda = (\lambda_1, \dots, \lambda_n)$. On expressing the Schur functions $S_{(1,1)}$ and $S_{(2,1)}$ by the elementary symmetric polynomials we obtain

$$H(F_n(M); t_1, \dots, t_n) = \frac{1 + \sigma_2(t_1, \dots, t_n) - \sigma_3(t_1, \dots, t_n) + \sigma_1\sigma_2(t_1, \dots, t_n)}{\prod_{i=1}^n (1 - t_i^{d_i})}.$$

Since $F_n(M)^G$ is generated by f_1, \dots, f_n , it is an \mathbb{N} -graded homomorphic image of $F_n(M)$, where we give the degrees d_1, \dots, d_n to the generators of $F_n(M)$. Hence in the formal power series

$$D(t) = H(F_n(M); t^{d_1}, \dots, t^{d_n}) - H(F_n(M)^G; t)$$

each coefficient is a non-negative integer. By (2.2) and (3.1) we have

$$D(t) = \frac{F(t)}{\prod_{i=1}^n (1 - t^{d_i})} - H_1(t),$$

where

$$F(t) = (t^{d_1} + \dots + t^{d_n}) \sum_{1 \leq i < j \leq n} t^{d_i + d_j}$$

and

$$(4.1) \quad H_1(t) = \frac{1}{|G|} \sum_{g \in G} \frac{(\omega_1(g)t + \dots + \omega_n(g)t)(\sum_{1 \leq i < j \leq n} \omega_i(g)\omega_j(g)t^2)}{\prod_{i=1}^n (1 - \omega_i(g)t)}.$$

Steinberg proved (see [1, p. 127]) that

$$\sum_{1 \leq i < j \leq n} \omega_i \omega_j : G \rightarrow \mathbb{C}$$

is an irreducible character of G . (Actually, $\sum \omega_i \omega_j$ is the character of $G < Gl(V)$ acting on the second exterior power of V , and we do not need here the irreducibility.) Therefore

$$(\bar{\omega}_1 + \dots + \bar{\omega}_n) \sum_{1 \leq i < j \leq n} \bar{\omega}_i \bar{\omega}_j$$

is a character of degree $n \binom{n}{2}$ of G , and we may decompose it as $\sum_{i=1}^r m_i \chi_i$, where χ_1, \dots, χ_r are pairwise different irreducible characters of G and m_1, \dots, m_r are positive integers. Let I be the ideal of $K[x_1, \dots, x_n]$ generated by the invariants of G of strictly positive degree. Chevalley showed in [2] that the representation of

G on $K[x_1, \dots, x_n]/I$ is equivalent to the regular representation. Following [15] we associate with any irreducible character χ of G the polynomial

$$p_\chi(t) = \sum_{i=0}^{\infty} a_i(\chi)t^i,$$

where $a_i(\chi)$ is the multiplicity of χ in the i th homogeneous component of $K[x_1, \dots, x_n]/I$. (Note that what we call $p_\chi(t)$ is $p_{\bar{\chi}}(t)$ in [15], because there G acts on $K[x_1, \dots, x_n]$ via the adjoint representation.) It turns out that $a_i(\chi) = 0$ if $i > \sum_{j=1}^n (d_j - 1)$ [15, Lemma 2.9]. The coefficient of t^d in the formal power series

$$\frac{1}{|G|} \sum_{g \in G} \frac{(\omega_1(g) + \dots + \omega_n(g)) \sum_{1 \leq i < j \leq n} \omega_i(g)\omega_j(g)}{\prod_{i=1}^n (1 - \omega_i(g)t)}$$

is the scalar product of the character $\sum_{i=1}^r m_i \chi_i$ and the character of G acting on the d th homogeneous component of $K[x_1, \dots, x_n]$. So by [15, 2.6 Proposition and 2.9 Lemma] we have

$$(4.2) \quad H_1(t) = \frac{t^3 \sum_{i=1}^r m_i p_{\chi_i}(t)}{\prod_{i=1}^n (1 - t^{d_i})} = \frac{G(t)}{\prod_{i=1}^n (1 - t^{d_i})},$$

where $G(t)$ is of the form

$$G(t) = \sum_{i=1}^{\binom{n}{2}} t^{e_i}$$

with $0 \leq e_1 \leq \dots \leq e_{\binom{n}{2}} \leq \sum_{i=1}^n (d_i - 1) + 3$.

Now we reduce to the case of irreducible groups as it was done by Guralnick in [9]. Assume that $n \geq 2$ is minimal such that $F_n(M)^G$ is generated by n elements for some pseudo-reflection group G .

1. The same argument as in [9] shows that G is not Abelian.

[9, Lemma 3] remains valid with the same proof for any relatively free algebra instead of the generic matrix algebra. Therefore the minimality of n implies that G is irreducible, and we may assume that G is a complex unitary pseudo-reflection group. We shall use the classification of these groups given in [13].

2. $n > 2$.

Suppose that $n = 2$, and let $\chi = \omega_1 + \omega_2$ denote the character of the given representation of G as a subgroup of $Gl_2(\mathbb{C})$. We showed above that the degree of $G(t)$ (see (4.2)) is at most $d_1 + d_2 + 1$. Though this bound would suffice for our purpose, for sake of completeness we derive some more precise information on the polynomial $G(t)$:

$$\begin{aligned} H_1(t) &= \frac{1}{|G|} \sum_{g \in G} \frac{\chi(g)\omega_1(g)\omega_2(g)t^3}{(1 - \omega_1(g)t)(1 - \omega_2(g)t)} \\ &= \frac{1}{|G|} \frac{t^3}{t^2} \sum_{g \in G} \frac{\chi(g)}{(1 - \omega_1(g^{-1})t^{-1})(1 - \omega_2(g^{-1})t^{-1})} \\ &= \frac{t}{|G|} \sum_{g \in G} \frac{\bar{\chi}(g)}{(1 - \omega_1(g)t^{-1})(1 - \omega_2(g)t^{-1})} \\ &= t \frac{p_\chi(t^{-1})}{(1 - t^{-d_1})(1 - t^{-d_2})} \end{aligned}$$

by a similar argument we used before. The character of the representation of G on the 1st homogeneous component of $K[x_1, \dots, x_n]/I$ is χ , so $p_\chi(t) = t + \sum_{i=2}^{d_1+d_2-2} a_i(\chi)t^i$, hence

$$H_1(t) = t \frac{t^{-1} + \sum_{i=2}^{d_1+d_2-2} a_i(\chi)t^{-i}}{(1-t^{-d_1})(1-t^{-d_2})} = \frac{t^{d_1+d_2} + \sum_{j=3}^{d_1+d_2-1} a_{d_1+d_2+1-j}(\chi)t^j}{(1-t^{d_1})(1-t^{d_2})}.$$

On the other hand, the minimal degree term of $\frac{F(t)}{\prod_{i=1}^n(1-t^{d_i})}$ is $t^{2d_1+d_2}$, thus the coefficient of $t^{d_1+d_2}$ in $D(t) = \frac{F(t)-G(t)}{(1-t^{d_1})(1-t^{d_2})}$ is -1 . This contradicts the assumption that $D(t)$ has non-negative coefficients.

3. G is not one of the groups of no. 23, 28, 30, 35, 36, 37 in the table [13, Table VII]. (The argument in 5 below applies also for these groups. However, first we eliminate them by exhibiting some invariants. This argument shows the role of the polynomial $[x_2, x_1, x_1]$ explicitly.)

Suppose that G is one of these groups. It is well known (see for example [16, 4.2.15. Lemma]) that G can be defined over the reals if and only if G has an invariant of degree 2. So we can suppose that $G < Gl_n(\mathbb{R})$ is an orthogonal group, and then a straightforward calculation shows that

$$x_1^2 + \dots + x_n^2 \in K\langle x_1, \dots, x_n \rangle^G.$$

Thus we can choose f_1, \dots, f_n such that

$$f_1 = y_1^2 + \dots + y_n^2.$$

Consider the invariant

$$f = \sum_{g \in G} g \cdot (y_1(y_1^2 + \dots + y_n^2)y_1).$$

Comparing the condition $K\langle f_1, \dots, f_n \rangle = F_n(M)^G$ and the degrees of the groups under consideration (cf. [13, Table VII]) we conclude that the only invariants of G of degree 4 are the scalar multiples of f_1^2 , therefore $f = af_1^2$ in $F_n(M)$ for some $a \in \mathbb{R}$. Now we have

$$f = \sum_{g \in G} (g_{11}y_1 + \dots + g_{n1}y_n)(y_1^2 + \dots + y_n^2)(g_{11}y_1 + \dots + g_{n1}y_n),$$

and the polynomial

$$h = \sum_{g \in G} (g_{11}x_1 + \dots + g_{n1}x_n)(x_1^2 + \dots + x_n^2)(g_{11}x_1 + \dots + g_{n1}x_n) - a(x_1^2 + \dots + x_n^2)^2$$

is contained in M . The homogeneous component of h of degree 4 in x_i is

$$\left(\sum_{g \in G} g_{i1}^2 - a\right)x_i^4, \quad (i = 1, \dots, n),$$

and it is contained in M , implying that $a = \sum_{g \in G} g_{i1}^2$, $i = 1, \dots, n$. Since for any $g \in G$ there exists an i with $g_{i1} \neq 0$, we have $a \neq 0$. The multihomogeneous component of h of multidegree $(2, 2)$

$$\sum_{g \in G} g_{11}^2 x_1 x_2^2 x_1 + \sum_{g \in G} g_{21}^2 x_2 x_1^2 x_2 - a x_1^2 x_2^2 - a x_2^2 x_1^2 = a(x_1 x_2^2 x_1 + x_2 x_1^2 x_2 - x_1^2 x_2^2 - x_2^2 x_1^2)$$

is contained in M . Make the substitution $x_1 \rightarrow x_1 + 1$, $x_2 \rightarrow x_2 + 1$ in the above polynomial, then we get

$$2ax_1x_2x_1 + \text{ other monomials.}$$

Hence M contains a polynomial of degree 3, which contradicts the definition of M .

4. G is not $G(m, p, n)$ (the groups no. 2 in [13, Table VII]).

Recall the definition of $G(m, p, n)$. Let $n \geq 3$ (the case $n = 2$ was handled in 2.), $m \geq 2$, p be positive integers such that p divides m . Let $A(m, p, n) < Gl_n(\mathbb{C})$ be the group of diagonal matrices whose diagonal entries are m th roots of unity and the determinant is an $\frac{m}{p}$ th root of unity. Consider $Sym(n)$ as the group of permutation matrices in $Gl_n(\mathbb{C})$. Clearly, $Sym(n)$ normalizes $A(m, p, n)$, and $G(m, p, n)$ is defined as the semidirect product

$$G(m, p, n) = A(m, p, n) \rtimes Sym(n).$$

Consider the polynomials

$$h_i = \sum_{\pi \in Sym(n)} \pi \cdot \sigma_i(x_1^m, \dots, x_n^m), \quad i = 1, \dots, n - 1;$$

$$h_n = \sum_{\pi \in Sym(n)} \pi \cdot (x_1 \dots x_n)^{m/p}.$$

The above semidirect decomposition of $G(m, p, n)$ shows that

$$h_1, \dots, h_n \in K\langle x_1, \dots, x_n \rangle^G,$$

and it is well known that their images in $K[x_1, \dots, x_n]$ form a basic set of invariants, with degrees $m, 2m, 3m, \dots, (n - 1)m, \frac{m}{p}n$. So we may suppose that f_1, \dots, f_n are the images of h_1, \dots, h_n under the natural homomorphism $K\langle x_1, \dots, x_n \rangle \rightarrow F_n(M)$. Consider the invariant

$$f = \sum_{g \in G} g \cdot (y_1^{m-1} f_1 y_1).$$

Any $g \in G$ can be written in the form $g = a\pi$ for some $a \in A(m, p, n)$ and $\pi \in Sym(n)$. So we have $g(y_1) = \theta y_{\pi(1)}$ for some permutation π and m th root of unity θ , and

$$f = (n - 1)! m^n / p \sum_{i=1}^n y_i^{m-1} (y_1^m + \dots + y_n^m) y_i.$$

By our hypothesis f is contained in $K\langle f_1, \dots, f_n \rangle$, and since $\deg(f_3), \dots, \deg(f_n) > \deg(f) = 2m$, f can be expressed with f_1, f_2 and f_n . More precisely, there exists a polynomial

$$h = \sum_{i=1}^n x_i^{m-1}(x_1^m + \dots + x_n^m)x_i - (c_1h_1^2 + c_2h_2 + \sum_j a_jh_nb_j) \in M,$$

where $c_1, c_2 \in \mathbb{C}$, and $a_j, b_j \in K\langle x_1, \dots, x_n \rangle$. Consider the multihomogeneous components of h . The coefficient of x_1^{2m} is $1 - c_1$, implying that $c_1 = 1$. Since $n \geq 3$, any monomial of h_n contains the variable x_3 , therefore the sum of the coefficients of the monomials of multidegree (m, m) is $2 - 2c_1 - kc_2$ for some positive integer k , showing that $c_2 = 0$. Thus the polynomial

$$x_1^{m-1}x_2^m x_1 + x_2^{m-1}x_1^m x_2 - x_1^m x_2^m - x_2^m x_1^m$$

is contained in M . After the substitution $x_1 \rightarrow x_1 + 1, x_2 \rightarrow x_2 + 1$ we obtain a polynomial in which the coefficient of $x_1x_2x_1$ is $(m - 1)m$. Again this contradicts the fact that M has no elements of degree 3.

5. G is not one of the groups no. 24, 26, 27, 29, 31, 32, 33, 34 or no. 1 with $n \geq 4$ in [13, Table VII].

Let a, b be functions $\mathbb{N} \rightarrow \mathbb{C}$, and let c be a function $\mathbb{N} \rightarrow \mathbb{N}$. We say that $a(n) = b(n) + O(c(n))$, if $|a(n) - b(n)| \leq Lc(n)$ for some constant L . We need the following lemma (cf. [16, 2.5.9. Lemma]):

Lemma 4.1. *Let $H(t) \in \mathbb{C}[[t]]$ be a formal power series of the form*

$$H(t) = \frac{A(t)}{\prod_{i=1}^n (1 - t^{d_i})} = \sum_{d=0}^{\infty} c_d t^d,$$

where $A(t) \in \mathbb{C}[t]$.

(i)
$$(1 - t)^n H(t) \Big|_{t=1} = \frac{A(1)}{\prod_{i=1}^n d_i}.$$

(ii)
$$(1 - t)^{n-1} H(t) - \frac{A(1)}{\prod_{i=1}^n d_i} \frac{1}{1 - t} \Big|_{t=1} = \frac{\frac{1}{2}A(1) \sum_{i=1}^n (d_i - 1) - A'(1)}{\prod_{i=1}^n d_i}.$$

(iii) *Suppose that $n \geq 3$ and all sets of $n - 1$ of the d_i have greatest common divisor 1. Then we have*

$$c_d = \frac{A(1)}{(n - 1)! \prod_{i=1}^n d_i} d^{n-1} + \frac{\frac{1}{2}A(1) \sum_{i=1}^n d_i - A'(1)}{(n - 2)! \prod_{i=1}^n d_i} d^{n-2} + O(d^{n-3}).$$

(In [16] $n \geq 4$ is required, but the proof works also for $n = 3$.)

Now we investigate the power series $D(t) = \frac{F(t) - G(t)}{\prod_{i=1}^n (1 - t^{d_i})}$. Since $F(1) = G(1) = n \binom{n}{2}$, by Lemma 4.1 (i) $D(t)$ has no pole $\frac{1}{(1 - t)^n}$. By Lemma 4.1 (ii) the coefficient of the pole $\frac{1}{(1 - t)^{n-1}}$ in the Laurent series of $\frac{F(t)}{\prod_{i=1}^n (1 - t^{d_i})}$ is

$$\frac{\frac{1}{2}n \binom{n}{2} \sum_{i=1}^n (d_i - 1) - \frac{3}{2}n(n - 1) \sum_{i=1}^n d_i}{\prod_{i=1}^n d_i} = \frac{n(n - 1)}{4|G|} ((n - 6) \sum_{i=1}^n (d_i - 1) - 6n).$$

By (4.1) the coefficient of the pole $\frac{1}{(1-t)^{n-1}}$ in the Laurent series of $H_1(t)$ is

$$\frac{1}{|G|} \sum_{g \in R} \frac{(1 + \dots + 1 + \omega(g)) \binom{n-1}{2} + (n-1)\omega(g)}{1 - \omega(g)},$$

where $R = \{g \in G \mid g \text{ is a pseudo-reflection}\}$ and $\omega(g)$ is the eigenvalue of g different from 1. It is well known (and one can derive from Lemma 4.1 (i)) that

$$\sum_{g \in R} \frac{1}{1 - \omega(g)} = \frac{1}{2} \sum_{i=1}^n (d_i - 1).$$

Using this formula we get the equalities

$$\sum_{g \in R} \frac{\omega(g)}{1 - \omega(g)} = - \sum_{g \in R} \frac{1}{1 - \bar{\omega}(g)} = -\frac{1}{2} \sum_{i=1}^n (d_i - 1)$$

and

$$\sum_{g \in R} \frac{\omega(g)^2}{1 - \omega(g)} = \sum_{g \in R} \left(-1 - \omega(g) + \frac{1}{1 - \omega(g)}\right) = N - \frac{1}{2} \sum_{i=1}^n (d_i - 1),$$

where N is the number of the reflecting hyperplanes. Thus we have

$$\begin{aligned} \frac{1}{|G|} \sum_{g \in R} \frac{((n-1) + \omega(g)) \binom{n-1}{2} + (n-1)\omega(g)}{1 - \omega(g)} \\ = \frac{n-1}{4|G|} ((n^2 - 6n + 4) \sum_{i=1}^n (d_i - 1) + 4N), \end{aligned}$$

and the coefficient of the pole $\frac{1}{(1-t)^{n-1}}$ in $D(t)$ is

$$\frac{n-1}{2|G|} (-3n^2 - 2 \sum_{i=1}^n (d_i - 1) - 2N) = \frac{G'(1) - F'(1)}{|G|}.$$

(The last equality holds by Lemma 4.1 (ii).)

Denote by δ the greatest common divisor of d_1, \dots, d_n . The assumption $F_n(M)^G = K\langle f_1, \dots, f_n \rangle$ implies that δ divides each of the $e_1, \dots, e_n \binom{n}{2}$, that is, we may write

$$D(t) = \frac{A(s)}{\prod_{i=1}^n (1 - s^{d_i/\delta})} = \sum_{d=0}^{\infty} c_d s^d,$$

where $s = t^\delta$ and $A(s) \in \mathbb{C}[s]$. One can check in [13, Table VII] that for the groups under consideration the numbers $d_1/\delta, \dots, d_n/\delta$ satisfy the condition of Lemma 4.1 (iii), so applying this lemma and the equalities $A(1) = 0$ and $\frac{\partial}{\partial s} A(s) \Big|_{s=1} = \delta(G'(1) - F'(1))$ we obtain

$$\begin{aligned} c_d &= \frac{\delta(G'(1) - F'(1))}{(n-2)! \prod_{i=1}^n d_i} d^{n-2} + O(d^{n-3}) \\ &= \frac{-\delta(n-1)}{2(n-2)!|G|} (3n^2 + 2 \sum_{i=1}^n (d_i - 1) + 2N) d^{n-2} + O(d^{n-3}). \end{aligned}$$

This immediately implies that for sufficiently large d the coefficient of t^d in the power series $D(t)$ is strictly negative, which is a contradiction.

6. G is not the group no. 25 in [13, Table VII].

This group has 24 pseudo-reflections, all of them are of order 3 (see [3, p. 412, Table]), so the number of reflecting hyperplanes is 12. By the calculations in 5 we have

$$F'(1) - G'(1) = 3 \cdot 3^2 + 2 \cdot 24 + 2 \cdot 12.$$

On the other hand

$$F'(1) - G'(1) = 9(d_1 + d_2 + d_3) - (e_1 + \dots + e_9);$$

therefore

$$\frac{e_1 + \dots + e_9}{9} = 16,$$

implying that $e_1 \leq 16$. But we have

$$F(t) = t^{2d_1+d_2} + \text{higher degree terms},$$

and since $2d_1 + d_2 = 21$, the coefficient of t^{e_1} in $D(t) = \frac{F(t)-G(t)}{\prod_{i=1}^9(1-t^{d_i})}$ is strictly negative. This is a contradiction.

7. G is not $Sym(4)$, that is, the group no. 1 with $n = 3$ in [13, Table VII].

One can calculate the power series $D(t)$ directly and conclude that it has a negative coefficient.

We have eliminated all the finite irreducible complex unitary reflection groups, so the proof of Proposition 2.4 is complete.

ACKNOWLEDGEMENTS

I would like to thank the Széchenyi István Scholarship Foundation for the support which made it possible to visit the Penn State University in the first half of 1994. I thank the Department of Mathematics, especially E. Formanek for their hospitality during my visit. I am also grateful to V. Drensky for some earlier conversations on this topic.

REFERENCES

1. N. Bourbaki, *Groupes et Algèbres de Lie*, Chap. IV, V, VI, Hermann, Paris, 1968. MR **39**:1590
2. C. Chevalley, *Invariants of finite groups generated by reflections*, Amer. J. Math. **77** (1955), 778-782. MR **17**:345d
3. A. M. Cohen, *Finite complex reflection groups*, Ann. Sci. École Norm. Sup. **9** (1976), 379-436. MR **54**:10437
4. W. Dicks and E. Formanek, *Poincaré series and a problem of S. Montgomery*, Lin. Mult. Alg. **12** (1982), 21-30. MR **84c**:15032
5. V. Drensky, *Polynomial identities for 2×2 matrices*, Acta Appl. Math. **21** (1990), 137-161. MR **91k**:16024
6. V. Drensky, *Computational techniques for PI-algebras*, Topics in Algebra, Banach Center Publications, Vol. 26, Part 1., PWN-Polish Scientific Publishers, Warsaw, 1990, pp. 17-44. MR **93e**:16036
7. V. Drensky, *Private communication* (1993).
8. E. Formanek, *Noncommutative invariant theory*, Contemporary Mathematics 43, Group Actions on Rings, S. Montgomery ed., Amer. Math. Soc., 1985, pp. 87-119. MR **87d**:16046

9. R. M. Guralnick, *Invariants of finite linear groups acting on relatively free algebras*, Linear Alg. Appl. **72** (1985), 85-92. MR **87d**:16050
10. V. K. Kharchenko, *Noncommutative invariants of finite groups and Noetherian varieties*, J. Pure Appl. Alg. **31** (1984), 83-90. MR **85j**:16052
11. D. Krakowsky and A. Regev, *The polynomial identities of the Grassmann algebra*, Trans. Amer. Math. Soc. **181** (1973), 429-438. MR **48**:4005
12. L. H. Rowen, *Polynomial Identities in Ring Theory*, Academic Press, New York, 1980. MR **82a**:16021
13. G. C. Shephard and J. A. Todd, *Finite unitary reflection groups*, Canad. J. Math. **6** (1954), 274-304. MR **15**:600b
14. L. Solomon, *Invariants of finite reflection groups*, Nagoya Math. J. **22** (1963), 57-64. MR **27**:4872
15. T. A. Springer, *Regular elements of finite reflection groups*, Inv. Math. **25** (1974), 159-198. MR **50**:7371
16. T. A. Springer, *Invariant Theory*, Lecture Notes in Mathematics 585, Springer-Verlag, Berlin, Heidelberg, New York, 1977. MR **56**:5740

MATHEMATICAL INSTITUTE OF THE HUNGARIAN ACADEMY OF SCIENCES, P.O.B. 127, H-1364,
BUDAPEST, HUNGARY

E-mail address: domokos@math-inst.hu