

ON PERFECT ISOMETRIES AND ISOTYPIES IN ALTERNATING GROUPS

PAUL FONG AND MORTON E. HARRIS

ABSTRACT. Perfect isometries and isotypies are constructed for alternating groups between blocks with abelian defect groups and the Brauer correspondents of these blocks. These perfect isometries and isotypies satisfy additional compatibility conditions which imply that an extended Broué conjecture holds for the principal block of an almost simple group with an abelian Sylow p -subgroup and a generalized Fitting subgroup isomorphic to an alternating group.

Let G be a finite group and let \mathcal{O} be a complete discrete valuation ring with field of quotients K of characteristic 0 and residue class field k of characteristic $p > 0$. We suppose that K contains a primitive $|G|$ -th root of unity. In [4, (6.1)] Michel Broué posed the following

Isotypy Conjecture. *Let e be a block of $\mathcal{O}G$ with abelian defect group D and let f be the Brauer correspondent of e in $\mathcal{O}N_G(D)$. Then e and f are isotypic blocks.*

If G has an abelian Sylow p -subgroup, then the conjecture can be posed for the principal block of $\mathcal{O}G$. In this case the authors have shown that the conjecture holds provided an extended conjecture holds for the principal block of almost simple groups with an abelian Sylow p -subgroup (see [10, (5E)]). In this paper the isotypy conjecture for an arbitrary block with abelian defect group is proved for alternating groups. In addition, the extended conjecture is proved for the principal p -block of almost simple groups with abelian Sylow p -subgroups and generalized Fitting subgroup isomorphic to an alternating group.

We recall the basic definitions. Let $\bar{\cdot} : \mathcal{O} \rightarrow k$ be the canonical quotient mapping and let $\bar{\cdot} : \mathcal{O}G \rightarrow kG$ be the induced \mathcal{O} -algebra homomorphism of the group algebras. In particular, $\bar{\cdot} : e \rightarrow \bar{e}$ induces a bijection between central idempotents of $\mathcal{O}G$ and kG . If e is a block idempotent of $\mathcal{O}G$, let ${}_{KGe}\mathbf{Mod}$ be the category of left KGe -modules of finite type and let $\mathcal{R}_K(G, e)$ be the Grothendieck group of ${}_{KGe}\mathbf{Mod}$. Let G^\vee be the set of irreducible characters of G over K . We identify $\mathcal{R}_K(G, e)$ with the free abelian group on

$$(G, e)^\vee = \{\chi \in G^\vee : \chi(ge) = \chi(g) \text{ for all } g \in G\}.$$

Let $\text{CF}(G, K)$ be the K -space of K -valued class functions on G , and let $\text{CF}(G, e, K)$ be the K -subspace of class functions α in $\text{CF}(G, K)$ such that $\alpha(ge) = \alpha(g)$. The

Received by the editors March 26, 1996.

1991 *Mathematics Subject Classification.* Primary 20C15, 20C20; Secondary 20C30.

The first author was supported in part by NSF grant DMS 9100310.

The second author was supported in part by NSA grant MDA 904 92-H-3027.

©1997 American Mathematical Society

K -subspaces of $\text{CF}(G, K)$ and $\text{CF}(G, e, K)$ of class functions vanishing on the p -singular conjugacy classes of G will be denoted by $\text{CF}_{p'}(G, K)$ and $\text{CF}_{p'}(G, e, K)$.

Similar notation will be used for a second finite group H and a block idempotent f of $\mathcal{O}H$. We denote $\mathcal{R}_K(G \times H, e \otimes f)$ also as $\mathcal{R}_K((G, e), (H, f))$. Each μ in $\mathcal{R}_K((G, e), (H, f))$ determines adjoint linear maps

$$R_\mu: \text{CF}(G, e, K) \longrightarrow \text{CF}(H, f, K), \quad I_\mu: \text{CF}(H, f, K) \longrightarrow \text{CF}(G, e, K),$$

where

$$R_\mu(\alpha)(h) = \frac{1}{|G|} \sum_{g \in G} \mu(g, h^{-1})\alpha(g^{-1}), \quad I_\mu(\beta)(g) = \frac{1}{|H|} \sum_{h \in H} \mu(g, h^{-1})\beta(h)$$

for $\alpha \in \text{CF}(G, e, K)$ and $\beta \in \text{CF}(H, f, K)$. Moreover, $R_\mu: \mathcal{R}_K(G, e) \longrightarrow \mathcal{R}_K(H, f)$ and $I_\mu: \mathcal{R}_K(H, f) \longrightarrow \mathcal{R}_K(G, e)$.

Definition. A virtual character μ of $G \times H$ is perfect if the following separation and integrality conditions hold for all $g \in G, h \in H$:

1. $\mu(g, h) = 0$ if $\{g, h\}$ consists of a p -regular element and a p -singular element.
2. $\mu(g, h)$ is divisible in \mathcal{O} by $|C_G(g)|_p$ and by $|C_H(h)|_p$.

If μ is perfect, then R_μ and I_μ induce K -linear mappings

$$\begin{aligned} R_{\mu, p'}: \text{CF}_{p'}(G, e, K) &\longrightarrow \text{CF}_{p'}(H, f, K), \\ I_{\mu, p'}: \text{CF}_{p'}(H, f, K) &\longrightarrow \text{CF}_{p'}(G, e, K) \end{aligned}$$

by restriction. We say μ is an isometry if R_μ is an isometry of $\mathcal{R}_K(G, e)$ onto $\mathcal{R}_K(H, f)$. An isometry μ is thus a pair (J_μ, ϵ_μ) consisting of a bijection and a sign function

$$J_\mu: (G, e)^\vee \longrightarrow (H, f)^\vee, \quad \epsilon_\mu: (G, e)^\vee \longrightarrow \{\pm 1\}$$

such that $R_\mu(\chi) = \epsilon_\mu(\chi) J_\mu(\chi)$ for all $\chi \in (G, e)^\vee$. In particular,

$$\mu(g, h) = \sum_{\chi \in (G, e)^\vee} \epsilon_\mu(\chi) (\chi(g) \times J_\mu(\chi)(h^{-1})).$$

It will be convenient to set $G_P = C_G(P)$ and $G_x = C_G(x)$ when P is a p -subgroup of G and x is a p -element of G . The Brauer morphism $\text{Br}_P^G: \mathcal{O}G \longrightarrow kG_P$ is the \mathcal{O} -linear mapping defined by

$$\text{Br}_P^G(g) = \begin{cases} g & \text{if } g \in G_P, \\ 0 & \text{if } g \notin G_P. \end{cases}$$

Br_P^G preserves the $N_G(P)$ -conjugation action. Let e and \bar{e} be corresponding blocks of $\mathcal{O}G$ and kG . An \bar{e} -subpair is a pair (P, \bar{u}) , where P is a p -subgroup of G , \bar{u} is a block of kG_P , and $(1, \bar{e}) \leq (P, \bar{u})$ is containment in the sense of [1, Definition 3.3]. An e -subpair is a pair (P, u) , where P is a p -subgroup of G , u is a block of $\mathcal{O}G_P$, and (P, \bar{u}) is an \bar{e} -subpair. Containment of such pairs is defined in the obvious way: $(Q, v) \leq (P, u)$ if $(Q, \bar{v}) \leq (P, \bar{u})$. Let $\mathcal{S}_p(G, e)$ denote the set of e -subpairs of G . An \bar{e} -element is a pair (x, \bar{u}) , where x is a p -element of G , \bar{u} is a block of kG_x , and $(\langle x \rangle, \bar{u})$ is an \bar{e} -subpair. An e -element is a pair (x, u) , where x is a p -element of G , u is a block of $\mathcal{O}G_x$, and (x, \bar{u}) is an \bar{e} -element. The decomposition mapping

$$d_G^{(x, u)}: \text{CF}(G, K) \longrightarrow \text{CF}_{p'}(G_x, u, K)$$

is defined by $d_G^{(x,u)}(\alpha)(y) = \alpha(xyu)$ for $\alpha \in \text{CF}(G, K)$, $y \in G_{x,p'}$. If (x, u) is an e -element, then $\alpha(xyu) = \alpha(xyeu)$ by Brauer's Second Main Theorem (see [2, §A2, Théorème A2.1]).

The objects and morphisms of the Brauer category $\mathbf{Br}_e(G)$ are the e -subpairs (P, u) of G and the mappings $(P, u) \rightarrow (Q, v)$ in $\text{Hom}(P, Q)$ induced by conjugation by elements g of G such that $(P, u)^g \leq (Q, v)$. Let (D, e_D) be a maximal e -subpair of G . Each subgroup P of D determines a unique block e_P of $\mathcal{O}G_P$ such that $(P, e_P) \leq (D, e_D)$. If $P = \langle z \rangle$, we also write e_z for e_P . Let $\mathbf{Br}_{e,D}(G)$ be the full subcategory of $\mathbf{Br}_e(G)$ whose objects are such subpairs (P, e_P) . Since any two maximal e -subpairs are conjugate under G , $\mathbf{Br}_{e,D}(G)$ is independent of the choice of (D, e_D) up to a categorical isomorphism. Moreover, the inclusion functor of $\mathbf{Br}_{e,D}(G)$ into $\mathbf{Br}_e(G)$ is a categorical equivalence, since each e -subpair of G is conjugate in G to an e -subpair (P, e_P) contained in (D, e_D) . Similar notation will be used for a block idempotent f of $\mathcal{O}H$ with defect group D .

Definition. ([4, Definition 4.3]) Let e and f be block idempotents of $\mathcal{O}G$ and $\mathcal{O}H$ with a common defect group D . Let (D, e_D) and (D, f_D) be a maximal e -subpair and a maximal f -subpair of G and H . Then e and f are isotypic if the following conditions hold:

- (I1) The inclusion of D in G and H induces an equivalence of the categories $\mathbf{Br}_{e,D}(G)$ and $\mathbf{Br}_{f,D}(H)$.
- (I2) There exists a perfect isometry μ_P in $\mathcal{R}_K((G_P, e_P), (H_P, f_P))$ for each cyclic subgroup P of D such that the diagram

$$(0.1) \quad \begin{array}{ccc} \text{CF}(G, e, K) & \xrightarrow{R_{\mu(1)}} & \text{CF}(H, f, K) \\ \downarrow d_G^{(x, e_P)} & & \downarrow d_H^{(x, f_P)} \\ \text{CF}_{p'}(G_P, e_P, K) & \xrightarrow{R_{\mu_P, p'}} & \text{CF}_{p'}(H_P, f_P, K) \end{array}$$

is commutative for every generator x of P .

We note that the horizontal arrows in the diagram depend only on P , whereas the vertical arrows depend on the generator x of P . The set of perfect isometries $\{\mu_P\}$ is called the local system of the isotopy.

Remark. The actual statement of the Isotypy Conjecture in [4, (6.1)] asserts the following: Let e be a block of $\mathcal{O}G$ with abelian defect group D , let (D, e_D) be a maximal e -subpair of G , and let $T = N_G(D, e_D)$. Then e and e_D are isotypic blocks of $\mathcal{O}G$ and $\mathcal{O}T$. This statement is equivalent to the one given in the introduction of this paper since f and e_D are isotypic blocks of $\mathcal{O}H$ and $\mathcal{O}T$ by the following general fact: Suppose C is a normal subgroup of a finite group H , b is a block of $\mathcal{O}C$, and $T = N_H(b)$. Let u and v be blocks of $\mathcal{O}H$ and $\mathcal{O}T$ covering b such that $u = \text{Tr}_T^H(v)$. Then u and v are isotypic blocks of $\mathcal{O}H$ and $\mathcal{O}T$. Indeed, u and v have a common defect group D . If (P, ϕ) is a v -subpair of T , then $(P, \text{Tr}_{T_P}^{H_P}(\phi))$ is a u -subpair of H . The mapping $\mathcal{S}p(T, v) \rightarrow \mathcal{S}p(H, u)$ defined by $(P, \phi) \mapsto (P, \text{Tr}_{T_P}^{H_P}(\phi))$ implies condition (I1) by [13, Theorem 1] and [4, 4.2, Remark 2]. Let (D, u_D) and (D, v_D) be a maximal u -subpair of H and a maximal v -subpair of T such that $u_D = \text{Tr}_{T_D}^{H_D}(v_D)$. For each cyclic subgroup P of D , the categorical

equivalences

$$\begin{array}{ccc} \mathcal{O}_{T_P v_P} \mathbf{Mod} & \xrightarrow{\text{Ind}_{T_P}^{H_P}} & \mathcal{O}_{H_P u_P} \mathbf{Mod}, \\ \mathcal{O}_{H_P u_P} \mathbf{Mod} & \xrightarrow{v_P \text{Res}_{T_P}^{H_P}} & \mathcal{O}_{T_P v_P} \mathbf{Mod} \end{array}$$

induce perfect isometries μ_P in $\mathcal{R}_K((H_P, u_P), (T_P, v_P))$ satisfying **(I2)**.

The isotypies for the alternating group A_n will be constructed from isotypies for the symmetric group S_n . In §1 we consider a configuration (G, G^+, H, H^+) of finite groups, where G^+ is normal in G , H^+ is normal in H , $G/G^+ \simeq H/H^+$; a configuration (e, e^+, f, f^+) of blocks of $\mathcal{O}G$, $\mathcal{O}G^+$, $\mathcal{O}H$, $\mathcal{O}H^+$, where e covers e^+ and f covers f^+ ; and an isotypy between e and f with local system $\{\mu_P\}$. An isotypy between e^+ and f^+ can be constructed under suitable conditions on $\{\mu_P\}$. In the given context the relevant condition is that of $+$ -equivariance considered in §2. This construction will then be applied to configurations where $G = S_n$, $G^+ = A_n$, e^+ is a block of $\mathcal{O}G^+$ with abelian defect group D^+ , $H = N_G(D^+)$, $H^+ = N_{G^+}(D^+)$, and f^+ is the Brauer correspondent of e^+ . If p is odd, then e and f have D^+ as defect group. In this case Rouquier ([14, Théorème 2.13]) has shown that e and f are isotypic blocks. However, the isotypies in [Ro] are not explicit enough to check the condition of $+$ -equivariance. We therefore give in §3 an explicit $+$ -equivariant isotypy between e and f . §4 contains the relevant character theory of H^+ . The main result there is an analog of Frobenius's theorem on irreducible characters of G^+ which do not extend to characters of G . §5 contains the construction of the isotypy between e^+ and f^+ when $p > 2$, and §6 contains the proof that this isotypy satisfies the extended conjecture of [10, (5E)] when e^+ and f^+ are the principal blocks. §7 treats the case $p = 2$.

1. SOME CONFIGURATIONS OF ISOTYPIES

An isotypy between blocks e and f of $\mathcal{O}G$ and $\mathcal{O}H$ with a common defect group D requires a perfect isometry μ_P for each cyclic subgroup P of D such that (0.1) is commutative for every generator x of P . In practice somewhat less is required. The following reduction steps will be used in our proofs.

Step 1. Suppose e and f are blocks of principal type, that is, $\text{Br}_P^G(e)$ and $\text{Br}_P^H(f)$ are blocks of $\mathcal{O}G_P$ and $\mathcal{O}H_P$ for each subgroup P of D . Then the decomposition mappings $d_G^{(x, e_x)}$ and $d_H^{(x, f_x)}$ in (0.1) are simply d_G^x and d_H^x by Brauer's Second Main Theorem.

Step 2. $\mu_{\langle 1 \rangle}$ need only be an isometry, since $\mu_{\langle 1 \rangle}$ is perfect provided the other conditions hold [4, Remark following 4.6].

Step 3. If **(I1)** holds, the commutativity of (0.1) need only be checked for (P, e_P) with P in a set \mathcal{P} of representatives for the $N_G(D, e_D)$ -conjugacy classes of non-trivial cyclic subgroups of D and for all generators x of such P . This follows by transport of structure. For suppose (0.1) commutes for all generators x of such a P . Fix $g \in N_G(D, e_D)$ and let $Q = P^g$, so that $e_Q = (e_P)^g$. Choose $h \in N_H(D, f_D)$ so that h and g induce the same automorphism of D . Then $Q = P^h$ and $f_Q = (f_P)^h$. Define the perfect isometry μ_Q in $\mathcal{R}_K((G_Q, e_Q), (H_Q, f_Q))$ by $\mu_Q(y, z) = \mu_P(gyg^{-1}, hzh^{-1})$ for $y \in G_Q$, $z \in H_Q$. Then (0.1) with P replaced by Q and x replaced by any generator of Q is commutative.

For the rest of this section σ is a fixed linear character of G . Multiplication by σ gives an action $\chi \mapsto \sigma\chi$ on G^\vee ; an action $e \mapsto \sigma e$ on central idempotents of $\mathcal{O}G$, where $\sigma e = \sum_{g \in G_{p'}} \sigma(g^{-1})c_g g$ if $e = \sum_{g \in G_{p'}} c_g g$; and an action $\bar{e} \mapsto \sigma\bar{e}$ on central idempotents of kG , where $\sigma\bar{e} = \sum_{g \in G_{p'}} \bar{\sigma}(g^{-1})\bar{c}_g g$ if $\bar{e} = \sum_{g \in G_{p'}} \bar{c}_g g$. In particular, $\sigma\bar{e} = \overline{\sigma e}$. If e is a block idempotent of $\mathcal{O}G$, then $\chi \in (G, e)^\vee$ if and only if $\sigma\chi \in (G, \sigma e)^\vee$. Moreover, e and σe have the same defect groups, and $\sigma e = e$ if σ has p -power order. Let e be a fixed block of $\mathcal{O}G$. If P is a p -subgroup of G , set $\sigma_P = \text{Res}_{G_P}^G(\sigma)$ for simplicity of notation. In particular, $\text{Br}_P^G(\sigma e) = \sigma_P \text{Br}_P^G(e)$. If (P, u) is an e -subpair of G , then $(P, \sigma_P u)$ is a σe -subpair of G . More generally, let (P, u) and (Q, v) be subpairs of G . Then $(P, u) \leq (Q, v)$ if and only if $(P, \sigma_P u) \leq (Q, \sigma_Q v)$. Indeed, we need only show this for normal containment, since containment of subpairs is by definition a chain of normal containments. Let $(P, u) \triangleleft (Q, v)$. Then $P \triangleleft Q$, u is stable under Q , and $\text{Br}_Q^{G_P Q}(u)v = v$. So $\sigma_P u$ is stable under Q and $\text{Br}_Q^{G_P Q}(\text{Res}_{G_P Q}^G(\sigma)u) \cdot \sigma_Q v = \sigma_Q v$. Thus $(P, \sigma_P u) \triangleleft (Q, \sigma_Q v)$. A similar argument using σ^{-1} shows that if $(P, \sigma_P u) \triangleleft (Q, \sigma_Q v)$, then $(P, u) \triangleleft (Q, v)$.

Lemma (1A). $\text{Br}_{e,D}(G)$ and $\text{Br}_{\sigma e,D}(G)$ are isomorphic categories under the inclusion of a common defect group D of e and σe in G .

Proof. We fix a maximal e -subpair (D, e_D) of G and the corresponding maximal σe -subpair $(D, \sigma_D e_D)$ of G . If $P \leq D$ and $(P, e_P) \leq (D, e_D)$, then $(P, \sigma_P e_P) \leq (D, \sigma_D e_D)$ and $\sigma_P e_P = (\sigma e)_P$. To complete the argument, we show that

$$\text{Hom}((P, e_P), (Q, e_Q)) = \text{Hom}((P, \sigma_P e_P), (Q, \sigma_Q e_Q))$$

as subsets of $\text{Hom}(P, Q)$. If $(P, e_P)^g = (P^g, (e_P)^g) \leq (Q, e_Q)$ for $g \in G$, then $(P, \sigma_P e_P)^g = (P^g, \sigma_{P^g} e_{P^g}) \leq (Q, \sigma_Q e_Q)$ and $g \in \text{Hom}((P, \sigma_P e_P), (Q, \sigma_Q e_Q))$. A similar argument using the linear character σ^{-1} gives the reverse inclusion. \square

Proposition (1B). Suppose σ is trivial on a defect group D of e and σe . Then e and σe are isotypic blocks with local system $\{\mu_P\}$, where $\mu_P = (J_P, \epsilon_P)$ for each cyclic subgroup P of D is the isometry in $\mathcal{R}_K((G_P, e_P), (G_P, \sigma_P e_P))$ such that $J_P(\zeta) = \sigma_P \zeta$ and $\epsilon_P(\zeta) = 1$ for $\zeta \in (G_P, e_P)^\vee$.

Proof. Condition (I1) for an isotypy holds by (1A). The isometry μ_P is perfect since

$$\mu_P(x, y) = \sum_{\zeta \in (G_P, e_P)^\vee} \zeta(x)\sigma_P(y^{-1})\zeta(y^{-1}) = \sigma_P(y^{-1}) \sum_{\zeta \in (G_P, e_P)^\vee} \zeta(x)\zeta(y^{-1})$$

and the identity isometry of $(G_P, e_P)^\vee$ is perfect. The diagram

$$\begin{CD} \text{CF}(G, e, K) @>R_{\mu_{(1)}}>> \text{CF}(G, \sigma e, K) \\ @Vd_G^{(x, e_P)}VV @VVd_G^{(x, \sigma_P e_P)}V \\ \text{CF}_{p'}(G_P, e_P, K) @>R_{\mu_{P, p'}}>> \text{CF}_{p'}(G_P, \sigma_P e_P, K) \end{CD}$$

is commutative for each cyclic subgroup P of D and for every generator x of P . Indeed, let $\chi \in (G, e)^\vee$ and $x' \in G_{P,p'}$. Then

$$\begin{aligned} d_G^{(x, \sigma_P e_P)}(\mathbf{R}_{\mu(1)}(\chi))(x') &= d_G^{(x, \sigma_P e_P)}(\sigma\chi)(x') \\ &= (\sigma\chi)(xx'\sigma_P e_P), \\ \mathbf{R}_{\mu_P, p'}(d_G^{(x, e_P)}(\chi))(x') &= (\sigma_P \cdot d_G^{(x, e_P)}(\chi))(x') \\ &= \sigma(x')\chi(xx'e_P). \end{aligned}$$

Now

$$xx'\sigma_P e_P = \frac{1}{|G_P|} \sum_{g \in G_P} \sum_{\zeta \in (G_P, e_P)^\vee} \zeta(1)\sigma(g^{-1})\zeta(g^{-1})xx'g.$$

Thus

$$\begin{aligned} (\sigma\chi)(xx'\sigma_P e_P) &= \frac{1}{|G_P|} \sum_{g \in G_P} \sum_{\zeta \in (G_P, e_P)^\vee} \zeta(1)\sigma(g^{-1})\zeta(g^{-1})\sigma(xx'g)\chi(xx'g) \\ &= \frac{1}{|G_P|} \sum_{g \in G_P} \sum_{\zeta \in (G_P, e_P)^\vee} \sigma(xx'g)\zeta(1)\zeta(g^{-1})\chi(xx'g) \\ &= \frac{1}{|G_P|} \sum_{g \in G_P} \sum_{\zeta \in (G_P, e_P)^\vee} \sigma(x'g)\zeta(1)\zeta(g^{-1})\chi(xx'g) \quad \text{since } \sigma(x) = 1 \\ &= \sigma(x')\chi(xx'e_P) \end{aligned}$$

and (1B) holds. □

For the rest of this section we suppose that $\langle \sigma \rangle$ acts regularly on the orbit of e , i.e., $\sigma^i e = e$ if and only if $\sigma^i = 1$. Set $G^+ = \ker \sigma$ and $t = |G/G^+|$. Then G^+ contains the defect group D of e , since t is relatively prime to p . If $\chi \in (G, e)^\vee$, then $\sigma^i \chi \in (G, \sigma^i e)^\vee$ and the t characters $\sigma^i \chi$ are distinct. Thus $\text{Res}_{G^+}^G(\chi)$ is irreducible and the $\sigma^i e$ cover a single block e^+ of $\mathcal{O}G^+$. In other words, $e^+ = \sum_{i=1}^t \sigma^i e$ and e^+ is G -stable. Thus the restriction $\text{Res}_{G^+}^G$ induces a perfect isometry $\rho = (J_\rho, \epsilon_\rho)$ in $\mathcal{R}_K((G, e), (G^+, e^+))$, where $J_\rho(\chi) = \text{Res}_{G^+}^G(\chi)$ and $\epsilon_\rho(\chi) = 1$ for $\chi \in (G, e)^\vee$.

A subpair (P, u) of G covers a subpair (P, u^+) of G^+ if u covers u^+ . Suppose (P, u) is an e -subpair. Then $\text{Br}_P^G(e)u = u$, so that $\text{Br}_P^G(\sigma^i e) \cdot \sigma_P^i u = \sigma_P^i u$. This implies $\langle \sigma_P \rangle$ acts regularly on the orbit of u . Moreover, $G_P^+ = G_P \cap G^+$ is the kernel of σ_P , $G = G^+G_P$, and $G/G^+ \simeq G_P/G_P^+$. In particular, $\text{Res}_{G_P^+}^{G_P}$ induces a perfect isometry $\rho_P = (J_{\rho_P}, \epsilon_{\rho_P})$ in $\mathcal{R}_K((G_P, e_P), (G_P^+, e_P^+))$.

Lemma (1C). *Suppose $\langle \sigma \rangle$ acts regularly on the orbit of e . Then the following hold:*

- (i) *Each e^+ -subpair (P, u^+) of G^+ is covered by a unique e -subpair (P, u) of G .*
- (ii) *Each e -subpair (P, u) of G covers a unique e^+ -subpair (P, u^+) of G^+ .*

Proof. We have $\text{Br}_P^{G^+}(e^+) = \text{Br}_P^G(e^+) = \sum_{i=1}^t \text{Br}_P^G(\sigma^i e)$ since $e^+ = \sum_{i=1}^t \sigma^i e$. If (P, u^+) is an e^+ -subpair of G^+ , then $\sum_i \text{Br}_P^G(\sigma^i e)u^+ = u^+$ and $\text{Br}_P^G(\sigma^i e)u^+ \neq 0$ for some i . So there exists a block u of $\mathcal{O}G_P$ such that $\sigma_P^i u$ covers u^+ and $\text{Br}_P^G(\sigma^i e) \cdot \sigma_P^i u = \sigma_P^i u$. This implies $\text{Br}_P^G(\sigma^j e) \cdot \sigma_P^j u = \sigma_P^j u$ for all j . In particular, $\langle \sigma_P \rangle$

acts regularly on the orbit of u , $u^+ = \sum_{i=1}^t \text{Res}_{G_P^+}^{G_P}(\sigma_P^i)u$, and (P, u) is the unique e -subpair of G covering (P, u^+) . Conversely, suppose (P, u) is an e -subpair of G . Then $\langle \sigma_P \rangle$ acts regularly on the orbit of u , u covers a unique block u^+ of $\mathcal{O}G_P^+$, and $u^+ = \sum_{i=1}^t \sigma_P^i u$. Since $\text{Br}_P^{G^+}(e^+) = \sum_{i=1}^t \text{Br}_P^G(\sigma^i e)$, it follows that $\text{Br}_P^{G^+}(e^+)u^+ = u^+$. Thus (P, u^+) is the unique e^+ -subpair of G^+ covered by (P, u) . \square

Proposition (1D). *Suppose $\langle \sigma \rangle$ acts regularly on the orbit of e and D is a defect group of e . Then e and e^+ are isotypic blocks with local system $\{\rho_P : P \leq D, P \text{ cyclic}\}$.*

Proof. By (1C), u covers u^+ if and only if there is a bijection $\gamma: \mathcal{S}p(G^+, e^+) \rightarrow \mathcal{S}p(G, e)$ such that $\gamma((P, u^+)) = (P, u)$. To show condition (1I) for an isotypy it suffices to show that γ satisfies the following conditions (see [4, 4.2, Remark 2]):

- (C1) γ is injective and $\gamma((P, u^+)^h) = \gamma((P, u^+))^h$ for all $h \in G^+$.
- (C2) Every element in $\mathcal{S}p(G, e)$ is G -conjugate to $\gamma((P, u^+))$ for some element (P, u^+) in $\mathcal{S}p(G^+, e^+)$.
- (C3) If $\gamma((Q, v^+)) = \gamma((P, u^+))^g$ for (P, u^+) and (Q, v^+) in $\mathcal{S}p(G^+, e^+)$ and g in G , then there exist $z \in G_P$ and $h \in G^+$ such that $g = zh$.
- (C4) $\gamma((P, u^+)) \leq \gamma((Q, v^+))$ if and only if $(P, u^+) \leq (Q, v^+)$.

Conditions (C1), (C2), (C3) follow easily from (1C) and the factorization $G = G_P G^+$. Suppose $\gamma((P, u^+)) = (P, u)$, $\gamma((Q, v^+)) = (Q, v)$, and $(P, u^+) \leq (Q, v^+)$. We claim $(P, u) \leq (Q, v)$. It suffices to show this for normal containment. If $(P, u^+) \triangleleft (Q, v^+)$, then $P \triangleleft Q$, u^+ is stable under Q , and $\text{Br}_Q^{G_P^+ Q}(u^+)v^+ = v^+$. Thus u is stable under Q by (1C). Moreover, u covers u^+ viewed as blocks of $\mathcal{O}G_P Q$ and $\mathcal{O}G_P^+ Q$, and $u^+ = \sum_i \text{Res}_{G_P Q}^G(\sigma^i)u$. Since $\text{Br}_Q^{G_P Q}(\text{Res}_{G_P Q}^G(\sigma^i)u) \cdot \sigma_Q^j v = 0$ for $i \neq j$ and since $v^+ = \sum_j \sigma_Q^j v$, it follows that

$$v^+ = \text{Br}_Q^{G_P^+ Q}(u^+)v^+ = \sum_i \text{Br}_Q^{G_P Q}(\text{Res}_{G_P Q}^G(\sigma^i)u)\sigma_Q^i v.$$

Thus $\text{Br}_Q^{G_P Q}(u)v = v$ and $(P, u) \triangleleft (Q, v)$. The argument is reversible, and so (C4) holds.

Let P be a cyclic subgroup of D . The diagram

$$\begin{CD} \text{CF}(G, e, K) @>R_\rho>> \text{CF}(G^+, e^+, K) \\ @Vd_G^{(x, e_P)}VV @VVd_{G^+}^{(x, e_P^+)}V \\ \text{CF}_{P'}(G_P, e_P, K) @>R_{\rho_P \cdot P'}>> \text{CF}_{P'}(G_P^+, e_P^+, K) \end{CD}$$

is commutative for all generators x of P . Indeed, let $\chi \in (G, e)^\vee$ and $x' \in G_{P, P'}^+$. Then

$$\begin{aligned} d_{G^+}^{(x, e_P^+)}(\text{Res}_{G^+}^G(\chi))(x') &= \chi(xx'e_P^+), \\ \text{Res}_{G_P^+}^{G_P}(d_G^{(x, e_P)}(\chi))(x') &= \chi(xx'e_P). \end{aligned}$$

The commutativity now follows since $\chi(xx'e_P^+) = \sum_i \chi(xx'\sigma_P^i e_P) = \chi(xx'e_P)$. \square

Corollary (1E). *Suppose $\langle \sigma \rangle$ acts regularly on the orbit of e and D is a defect group of e . Let $H = N_G(D)$ and $H^+ = N_{G^+}(D)$, and let f and f^+ be the Brauer*

correspondents of e and e^+ in $\mathcal{O}H$ and $\mathcal{O}H^+$. Then e and f are isotypic blocks of $\mathcal{O}G$ and $\mathcal{O}H$ if and only if e^+ and f^+ are isotypic blocks of $\mathcal{O}G^+$ and $\mathcal{O}H^+$.

Proof. We have $\text{Br}_D^G(\sigma^i e) = \sigma_D^i \text{Br}_D^G(e) = \sigma_D^i f$ for $1 \leq i \leq t$. In particular, $\text{Res}_H^G(\sigma)$ acts regularly on the orbit of f , σ_D^i is the Brauer correspondent of $\sigma^i e$, and $f^+ = \sum_{i=1}^t \sigma_D^i f$. Now e and e^+ are isotypic blocks by (1D); f and f^+ are also isotypic blocks by (1D). Thus (1E) holds. \square

2. +EQUIVARIANCE

Let G be a finite group, let G^+ be a subgroup of index 2, and let sgn_G be the sign character of G with kernel G^+ . We write $\chi \mapsto \chi^*$ and $e \mapsto e^*$ for the action of sgn_G on the irreducible characters and blocks of G . Let e and e^+ be blocks of $\mathcal{O}G$ and $\mathcal{O}G^+$. The configuration (G, G^+, e, e^+) is a +configuration if e^+ is the unique block of $\mathcal{O}G^+$ covered by e and if e is the unique block of $\mathcal{O}G$ covering e^+ . This is equivalent to the condition $e = e^+$ or to the conditions e^+ is stable under G and $e^* = e$. Let (H, H^+, f, f^+) be a second +configuration with corresponding sign character sgn_H , and write $\psi \mapsto \psi^*$ and $f \mapsto f^*$ for the action of sgn_H on the irreducible characters and blocks of H . An isometry $\mu = (J, \epsilon)$ in $\mathcal{R}_K((G, e), (H, f))$ is +equivariant with respect to (G^+, H^+) if

$$J(\chi^*) = J(\chi)^*, \quad \epsilon(\chi^*) = \epsilon(\chi)$$

for all $\chi \in (G, e)^\vee$.

Lemma (2A). *Let (G, G^+, e, e^+) and (H, H^+, f, f^+) be +configurations, and let μ be a +equivariant isometry in $\mathcal{R}_K((G, e), (H, f))$.*

- (i) $\chi \in (G, e)^\vee$ is reducible on G^+ if and only if $J(\chi) \in (H, f)^\vee$ is reducible on H^+ .
- (ii) $\mu(g, h) = 0$ if $\text{sgn}_G(g) \neq \text{sgn}_H(h)$.

Proof. (i) holds since $\text{Res}_{G^+}^G(\chi)$ and $\text{Res}_{H^+}^H(J(\chi))$ are reducible if and only if $\chi^* = \chi$ and $J(\chi)^* = J(\chi)$. In (ii) either $g \notin G^+$ or $h \notin H^+$. Now $\mu(g, h)$ is a sum of terms $\epsilon(\chi)\chi(g)J(\chi)(h^{-1})$. If $\text{Res}_{G^+}^G(\chi)$ and $\text{Res}_{H^+}^H(J(\chi))$ are reducible, then one of $\chi(g)$ or $J(\chi)(h^{-1})$ is zero and $\epsilon(\chi)\chi(g)J(\chi)(h^{-1}) = 0$. If $\text{Res}_{G^+}^G(\chi)$ and $\text{Res}_{H^+}^H(J(\chi))$ are irreducible, then $\chi \neq \chi^*$, $J(\chi) \neq J(\chi)^*$, and

$$\begin{aligned} &\epsilon(\chi)\chi(g)J(\chi)(h^{-1}) + \epsilon(\chi^*)\chi^*(g)J(\chi^*)(h^{-1}) \\ &= \epsilon(\chi)\chi(g)J(\chi)(h^{-1})(1 + \text{sgn}_G(g)\text{sgn}_H(h)) \end{aligned}$$

is zero. Thus $\mu(g, h)$ is a sum of zero subsums and $\mu(g, h) = 0$. \square

Let $\mu = (J, \epsilon)$ be a +equivariant isometry in $\mathcal{R}_K((G, e), (H, f))$. An isometry $\mu^+ = (J^+, \epsilon^+)$ in $\mathcal{R}_K((G^+, e^+), (H^+, f^+))$ is μ -induced if one and hence both of the following diagrams are commutative:

$$(2.1) \quad \begin{array}{ccc} \mathcal{R}_K(G, e) & \xrightarrow{R_\mu} & \mathcal{R}_K(H, f) \\ \downarrow \text{Res}_{G^+}^G & & \downarrow \text{Res}_{H^+}^H \\ \mathcal{R}_K(G^+, e^+) & \xrightarrow{R_{\mu^+}} & \mathcal{R}_K(H^+, f^+) \end{array}$$

$$(2.2) \quad \begin{array}{ccc} \mathcal{R}_K(G^+, e^+) & \xrightarrow{R_{\mu^+}} & \mathcal{R}_K(H^+, f^+) \\ \downarrow \text{Ind}_{G^+}^G & & \downarrow \text{Ind}_{H^+}^H \\ \mathcal{R}_K(G, e) & \xrightarrow{R_\mu} & \mathcal{R}_K(H, f) \end{array}$$

All μ -induced isometries $\mu^+ = (J^+, \epsilon^+)$ arise as follows: Let

$$(G^+, e^+)^\vee = (G^+, e^+)_e^\vee \cup (G^+, e^+)_n^\vee,$$

where the characters in $(G^+, e^+)_e^\vee$ extend to G and the characters in $(G^+, e^+)_n^\vee$ do not extend to G . If $\chi^+ \in (G^+, e^+)_e^\vee$ and $\chi^+ = \text{Res}_{G^+}^G(\chi)$ for $\chi \in (G, e)^\vee$, let

$$J^+(\chi^+) = \text{Res}_{H^+}^H(J(\chi)), \quad \epsilon^+(\chi^+) = \epsilon(\chi).$$

$J^+(\chi^+)$ and $\epsilon^+(\chi^+)$ are well-defined since

$$\text{Res}_{H^+}^H(J(\chi^*)) = \text{Res}_{H^+}^H(J(\chi)^*) = \text{Res}_{H^+}^H(J(\chi))$$

and $\epsilon(\chi^*) = \epsilon(\chi)$. If χ_+ and χ_- are characters in $(G^+, e^+)_n^\vee$ such that $\chi_+ + \chi_- = \text{Res}_{G^+}^G(\chi)$ for $\chi \in (G, e)^\vee$, let $\text{Res}_{H^+}^H(J(\chi)) = J(\chi)_+ + J(\chi)_-$ and set

$$J^+(\chi_\pm) = J(\chi)_\pm, \quad \epsilon^+(\chi_\pm) = \epsilon(\chi).$$

$J^+(\chi_\pm)$ depends on the labeling of the constituents χ_\pm and $J(\chi)_\pm$. In all cases χ covers χ^+ if and only if $J(\chi)$ covers $J^+(\chi^+)$. Moreover, $J^+((\chi^+)^g) = J^+(\chi^+)^h$ and $\epsilon^+((\chi^+)^g) = \epsilon^+(\chi^+)$ for $g \in G - G^+$ and $h \in H - H^+$.

Lemma (2B). *Let (G, G^+, e, e^+) and (H, H^+, f, f^+) be $+$ configurations, let μ be a $+$ equivariant isometry in $\mathcal{R}_K((G, e), (H, f))$, and let $\mu^+ = (J^+, \epsilon^+)$ be a μ -induced isometry in $\mathcal{R}_K((G^+, e^+), (H^+, f^+))$. Suppose p is odd and μ is perfect. Then μ^+ is perfect if and only if the generalized character of $G^+ \times H^+$ defined by*

$$\Delta(\mu^+)(g, h) = \sum_{\{\chi_+, \chi_-\} \subseteq (G^+, e^+)_n^\vee} \epsilon^+(\chi_\pm) (\chi_+ - \chi_-)(g) (J^+(\chi_+) - J^+(\chi_-))(h^{-1})$$

is perfect.

Proof. If $(g, h) \in G^+ \times H^+$, then

$$\begin{aligned} \mu(g, h) &= \sum_{\chi \in (G, e)^\vee} \epsilon(\chi) \chi(g) J(\chi)(h^{-1}) \\ &= \sum_{\chi^+ \in (G^+, e^+)_e^\vee} 2\epsilon^+(\chi^+) \chi^+(g) J^+(\chi^+)(h^{-1}) \\ &\quad + \sum_{\{\chi_+, \chi_-\} \subseteq (G^+, e^+)_n^\vee} \epsilon^+(\chi_\pm) (\chi_+ + \chi_-)(g) (J^+(\chi_+) + J^+(\chi_-))(h^{-1}) \\ &= 2\mu^+(g, h) - \sum_{\{\chi_+, \chi_-\} \subseteq (G^+, e^+)_n^\vee} \epsilon^+(\chi_\pm) (\chi_+ - \chi_-)(g) (J^+(\chi_+) - J^+(\chi_-))(h^{-1}). \end{aligned}$$

Thus

$$(2.3) \quad \text{Res}_{G^+ \times H^+}^{G \times H}(\mu) = 2\mu^+ - \Delta(\mu^+),$$

and (2B) follows. □

Remark (2C). If $(\chi_+ - \chi_-)(g)(J^+(\chi_+) - J^+(\chi_-))(h^{-1})$ is perfect for all $\{\chi_+, \chi_-\} \subseteq (G^+, e^+)_n^\vee$, then all μ -induced isometries μ^+ are perfect. We note $\Delta(\mu^+)(g, h) \neq 0$ only if g and h lie in conjugacy classes of G and H which split in G^+ and H^+ .

Suppose G_0 and G_1 are finite groups, G_0^+ and G_1^+ are normal subgroups of G_0 and G_1 of index 2, and sgn_{G_0} and sgn_{G_1} are the corresponding sign characters. Let $G = G_0 \times G_1$, and let $\text{sgn}_G = \text{sgn}_{G_0} \times \text{sgn}_{G_1}$ be the sign character of G with kernel

$$G^+ = (G_0 \times G_1)^+ = \{(g_0, g_1) \in G_0 \times G_1 : \text{sgn}_{G_0}(g_0) = \text{sgn}_{G_1}(g_1)\}.$$

In addition, let $G^{++} = G_0^+ \times G_1^+$ and let sgn_{G^+} be the sign character of G^+ with respect to G^{++} , that is,

$$\text{sgn}_{G^+} = \text{Res}_{G^+}^G(\text{sgn}_{G_0} \times 1_{G_1}) = \text{Res}_{G^+}^G(1_{G_0} \times \text{sgn}_{G_1}).$$

A character χ in G^\vee with components $\chi_i \in G_i^\vee$ is reducible on G^+ if and only if each χ_i is reducible on G_i^+ . In particular, if χ is reducible on G^+ and $\text{Res}_{G^+}^G(\chi) = \chi_+ + \chi_-$, then $\text{Res}_{G^{++}}^{G^+}(\chi_+)$ and $\text{Res}_{G^{++}}^{G^+}(\chi_-)$ are also reducible and all three characters χ, χ_+, χ_- vanish off G^{++} .

Proposition (2D). *Let (G_0, G_0^+, e_0, e_0^+) and (G_1, G_1^+, e_1, e_1^+) be +configurations, let e be the block $e_0 e_1$ of $\mathcal{O}G$, and let e^{++} be the block $e_0^+ e_1^+$ of $\mathcal{O}G^{++}$. Then there exists a unique block e^+ of $\mathcal{O}G^+$ covered by e . Moreover, (G, G^+, e, e^+) and $(G^+, G^{++}, e^+, e^{++})$ are +configurations.*

Proof. Every block of G^+ covered by e covers e^{++} . The hypotheses imply that $\text{sgn}_G e = e$ and e^{++} is stable under G . Suppose first that $\chi_0^+ \chi_1^+$ is a character in $(G^{++}, e^{++})^\vee$ stable under G . Each factor χ_i^+ extends to two characters χ_i and χ_i^* in $(G_i, e_i)^\vee$, and the four characters $\chi_0 \chi_1, \chi_0^* \chi_1^*, \chi_0^* \chi_1, \chi_0 \chi_1^*$ in $(G, e)^\vee$ are distinct. The restrictions of these four characters to G^+ are irreducible and so belong to a unique block e^+ of $\mathcal{O}G^+$. Thus e^+ is the unique block of $\mathcal{O}G^+$ covered by e , and (G, G^+, e, e^+) is a +configuration. The two distinct characters of G^+ extending $\chi_0^+ \chi_1^+$ are

$$\begin{aligned} \chi^+ &= \text{Res}_{G^+}^G(\chi_0 \chi_1) = \text{Res}_{G^+}^G(\chi_0^* \chi_1^*), \\ \chi^- &= \text{Res}_{G^+}^G(\chi_0 \chi_1^*) = \text{Res}_{G^+}^G(\chi_0^* \chi_1), \end{aligned}$$

and $\chi^- = \text{sgn}_{G^+} \chi^+$. Thus $(G^+, G^{++}, e^+, e^{++})$ is a +configuration and $\text{sgn}_{G^+} e^+ = e^+$. Suppose next that $\chi_0^+ \chi_1^+$ is a character in $(G^{++}, e^{++})^\vee$ not stable under G . Then $\chi_0^+ \chi_1^+$ is not stable under G^+ and $\text{Ind}_{G^{++}}^{G^+}(\chi_0^+ \chi_1^+)$ is irreducible. The block e^+ of $\mathcal{O}G^+$ containing $\text{Ind}_{G^{++}}^{G^+}(\chi_0^+ \chi_1^+)$ is the unique block of $\mathcal{O}G^+$ covering e^{++} and $\text{sgn}_{G^+} e^+ = e^+$. Thus (2D) also holds in this case. \square

Let (H_0, H_0^+, f_0, f_0^+) and (H_1, H_1^+, f_1, f_1^+) be a second pair of +configurations, and let $H, H^+, H^{++}, \text{sgn}_H, \text{sgn}_{H^+}, f^+, f^{++}$ have the analogous meanings for this pair. In particular, (H, H^+, f, f^+) and $(H^+, H^{++}, f^+, f^{++})$ are +configurations.

Proposition (2E). *Let (G_i, G_i^+, e_i, e_i^+) and (H_i, H_i^+, f_i, f_i^+) be +configurations and let $\mu_i = (J_i, \epsilon_i)$ be a +equivariant isometry in $\mathcal{R}_K((G_i, e_i), (H_i, f_i))$ for $i = 0, 1$. Then the following hold:*

- (i) $\mu = \mu_0 \times \mu_1$ is a +equivariant isometry in $\mathcal{R}_K((G, e), (H, f))$ with respect to (G^+, H^+) .

(ii) All μ -induced isometries μ^+ in $\mathcal{R}_K((G^+, e^+), (H^+, f^+))$ are +equivariant with respect to (G^{++}, H^{++}) .

Proof. Let $\mu = (J, \epsilon)$, where $J = J_0 \times J_1$ and $\epsilon = \epsilon_0 \times \epsilon_1$. Let $\chi \in (G, e)^\vee$, where $\chi = \chi_0 \chi_1$ and $\chi_i \in (G_i, e_i)^\vee$. Then

$$\begin{aligned} \epsilon(\chi^*) &= \epsilon_0(\chi_0^*)\epsilon_1(\chi_1^*) = \epsilon_0(\chi_0)\epsilon_1(\chi_1) = \epsilon(\chi), \\ J(\chi^*) &= J_0(\chi_0^*)J_1(\chi_1^*) = J_0(\chi_0)^*J_1(\chi_1)^* = J(\chi)^*, \end{aligned}$$

and (i) holds. We set $(\chi^+)^\# = \text{sgn}_{G^+} \chi^+$ and $(\psi^+)^\# = \text{sgn}_{H^+} \psi^+$ for χ^+ in $(G^+)^\vee$ and for ψ^+ in $(H^+)^\vee$. Let $\mu^+ = (J^+, \epsilon^+)$ be a μ -induced isometry in $\mathcal{R}_K((G^+, e^+), (H^+, f^+))$. Let $\chi^+ \in (G^+, e^+)^\vee$ and choose χ in $(G, e)^\vee$ covering χ^+ . Let $\chi = \chi_0 \chi_1$, where $\chi_i \in (G_i, e_i)^\vee$. Then $J(\chi) = J_0(\chi_0)J_1(\chi_1)$ is a character in $(H, f)^\vee$ covering $J^+(\chi^+)$. If $\text{Res}_{G^{++}}^{G^+}(\chi^+)$ is reducible, then $(\chi^+)^\# = \chi^+$ and $\text{Res}_{G_i^+}^{G_i}(\chi_i)$ is reducible for some i . So $\text{Res}_{H_i^+}^{H_i}(J_i(\chi_i))$ is reducible for the same i and $\text{Res}_{H^{++}}^{H^+}(J^+(\chi^+))$ is reducible. Thus $\epsilon^+((\chi^+)^\#) = \epsilon^+(\chi^+)$ and $J^+((\chi^+)^\#) = (J^+(\chi^+))^\#$. If $\text{Res}_{G^{++}}^{G^+}(\chi^+)$ is irreducible, then $\chi^+ = \text{Res}_{G^+}^G(\chi)$ and

$$\begin{aligned} \chi^+ &= \text{Res}_{G^+}^G(\chi_0 \chi_1) = \text{Res}_{G^+}^G(\chi_0^* \chi_1^*), \\ (\chi^+)^\# &= \text{Res}_{G^+}^G(\chi_0 \chi_1^*) = \text{Res}_{G^+}^G(\chi_0^* \chi_1). \end{aligned}$$

Since μ_0 and μ_1 are +equivariant, $\epsilon^+((\chi^+)^\#) = \epsilon^+(\chi^+)$ and

$$\begin{aligned} J^+((\chi^+)^\#) &= \text{Res}_{H^+}^H(J_0(\chi_0)J_1(\chi_1^*)) \\ &= \text{Res}_{H^+}^H(J_0(\chi_0)J_1(\chi_1)^*) \\ &= (\text{Res}_{H^+}^H(J_0(\chi_0)J_1(\chi_1)))^\# \\ &= J^+(\chi^+)^\#. \end{aligned}$$

This completes the proof of (ii). □

We continue with the situation of (2E). Suppose $\mu_i^+ = (J_i^+, \epsilon_i^+)$ is a μ_i -induced isometry in $\mathcal{R}_K((G_i^+, e_i^+), (H_i^+, f_i^+))$ for $i = 0, 1$, and μ^{++} is the isometry $\mu_0^+ \times \mu_1^+$ in $\mathcal{R}_K((G^{++}, e^{++}), (H^{++}, f^{++}))$. We may choose a μ -induced isometry $\mu^+ = (J^+, \epsilon^+)$ such that μ^{++} is μ^+ -induced. Indeed, fix notation so that $J_i^+(\chi_{i,\pm}) = \psi_{i,\pm}$ whenever $J_i(\chi_i) = \psi_i$, $\text{Res}_{G_i^+}^{G_i}(\chi_i) = \chi_{i,+} + \chi_{i,-}$, and $\text{Res}_{H_i^+}^{H_i}(\psi_i) = \psi_{i,+} + \psi_{i,-}$. Suppose $J(\chi) = \psi$, where $\text{Res}_{G^+}^G(\chi) = \chi_+ + \chi_-$ and $\text{Res}_{H^+}^H(\psi) = \psi_+ + \psi_-$ are reducible. Then $\chi = \chi_0 \chi_1$ and $\psi = \psi_0 \psi_1$, where $\text{Res}_{G_i^+}^{G_i}(\chi_i) = \chi_{i,+} + \chi_{i,-}$ and $\text{Res}_{H_i^+}^{H_i}(\psi_i) = \psi_{i,+} + \psi_{i,-}$ are reducible for $i = 0, 1$. We label χ_\pm and ψ_\pm so that

$$\begin{aligned} \text{Res}_{G^{++}}^{G^+}(\chi_+) &= \chi_{0,+} \chi_{1,+} + \chi_{0,-} \chi_{1,-}, \\ \text{Res}_{G^{++}}^{G^+}(\chi_-) &= \chi_{0,+} \chi_{1,-} + \chi_{0,-} \chi_{1,+}, \\ \text{Res}_{H^{++}}^{H^+}(\psi_+) &= \psi_{0,+} \psi_{1,+} + \psi_{0,-} \psi_{1,-}, \\ \text{Res}_{H^{++}}^{H^+}(\psi_-) &= \psi_{0,+} \psi_{1,-} + \psi_{0,-} \psi_{1,+}, \end{aligned} \tag{2.4}$$

and choose J^+ so that $J^+(\chi_\pm) = \psi_\pm$ for all such χ and ψ . Then

$$\begin{CD} \mathcal{R}_K(G^+, e^+) @>R_{\mu^+}>> \mathcal{R}_K(H^+, f^+) \\ @VRes_{G^{++}}VV @VVRes_{H^{++}}V \\ \mathcal{R}_K(G^{++}, e^{++}) @>R_{\mu^{++}}>> \mathcal{R}_K(H^{++}, f^{++}) \end{CD}$$

is a commutative diagram. Thus μ^+ is μ -induced and μ^{++} is μ^+ -induced.

Proposition (2F). *Suppose the hypotheses of (2E) hold and μ_i^+ is a μ_i -induced isometry in $\mathcal{R}_K((G_i^+, e_i^+), (H_i^+, f_i^+))$ for $i = 0, 1$.*

- (i) *There exists a μ -induced isometry μ^+ in $\mathcal{R}_K((G^+, e^+), (H^+, f^+))$ such that $\mu^{++} = \mu_0^+ \times \mu_1^+$ is μ^+ -induced.*
- (ii) *If p is odd, μ_i^+ is perfect for $i = 0, 1$, and μ^+ is as in (i), then μ^+ is perfect.*

Proof. (i) follows from the preceding paragraph. For (ii) it suffices by (2B) to prove that

$$\Delta(\mu^+)(g, h) = \sum_{\{\chi_+, \chi_-\} \subseteq (G^+, e^+)_n^\vee} \epsilon^+(\chi_\pm)(\chi_+ - \chi_-)(g)(\psi_+ - \psi_-)(h^{-1})$$

is perfect on its support $G^{++} \times H^{++}$. We may suppose μ^+ satisfies (2.4). Let $g_0 \in G_0^+, g_1 \in G_1^+, h_0 \in H_0^+, h_1 \in H_1^+$. Then

$$\begin{aligned} \Delta(\mu^+)(g_0g_1, h_0h_1) = & \left[\sum_{\substack{\{\chi_{0,+}, \chi_{0,-}\} \\ \subseteq (G_0, e_0)_n^\vee}} \epsilon_0^+(\chi_{0,\pm})(\chi_{0,+} - \chi_{0,-})(g_0)(\psi_{0,+} - \psi_{0,-})(h_0^{-1}) \right] \\ & \cdot \left[\sum_{\substack{\{\chi_{1,+}, \chi_{1,-}\} \\ \subseteq (G_1, e_1)_n^\vee}} \epsilon_1^+(\chi_{1,\pm})(\chi_{1,+} - \chi_{1,-})(g_1)(\psi_{1,+} - \psi_{1,-})(h_1^{-1}) \right]. \end{aligned}$$

The bracketed sums are perfect generalized characters of $G_0^+ \times H_0^+$ and $G_1^+ \times H_1^+$ since μ_0^+ and μ_1^+ are perfect. Thus $\Delta(\mu^+)$ is perfect. □

3. THE ISOTYPY CONJECTURE IN SYMMETRIC GROUPS

Let G be the symmetric group $S(\Pi)$ on a set Π of cardinality n . By the Nakayama conjecture a p -block e of $\mathcal{O}G$ is parametrized by a p -core λ_0 , and irreducible characters χ_λ in $(G, e)^\vee$ are parametrized by partitions λ of n with p -core λ_0 . In particular, $n = n_0 + pw$, where $n_0 = |\lambda_0|$ and $w \geq 0$ is the so-called weight of e . We suppose that e has an abelian defect group D , or equivalently, that $w < p$. Thus $\Pi = \Pi_0 \cup \Pi_1$ is the disjoint union of sets Π_0 and Π_1 of cardinality n_0 and pw . We may suppose $\Pi_1 = \Gamma \times \Omega$, where $\Gamma = \{1, 2, \dots, p\}$ and Ω is a set of w elements. Let $X = S(\Gamma)$ and $Y = N_X(P)$, where P is a fixed Sylow p -subgroup of X . In particular, Y is a Frobenius group with kernel P of order p and a cyclic complement of order $p - 1$.

We take D as the Sylow p -subgroup P^Ω of $S(\Pi_1)$ and set $H = N_G(D)$, so that $H = H_0 \times H_1$ with $H_0 = S(\Pi_0)$ and $H_1 = Y \wr S(\Omega)$. We view $Y \wr S(\Omega)$ as the product set $Y^\Omega \times S(\Omega)$, where Y^Ω is the set of functions from Ω to Y , $S(\Omega)$ acts

on Y^Ω on the left by $(\sigma f)(i) = f(\sigma^{-1}(i))$ or $(\sigma f)_i = f_{\sigma^{-1}(i)}$, and multiplication in $Y \wr S(\Omega)$ is defined by

$$(f, \sigma)(g, \tau) = (f(\sigma g), \sigma\tau).$$

The Brauer correspondent f of e in $\mathcal{O}H$ has the form $f = f_0 \times f_1$, where f_0 is the block of defect 0 of $\mathcal{O}H_0$ parametrized by λ_0 and f_1 is the principal block of $\mathcal{O}H_1$. We note that f_1 is the unique block of $\mathcal{O}H_1$, since H_1/D is a p' -group and $C_{H_1}(D) = D$.

Let λ be a partition of n with p -core λ_0 and p -quotient $\Lambda = (\lambda_1, \lambda_2, \dots, \lambda_p)$. Then λ_0 can be gotten by removing a sequence $(\nu_1, \nu_2, \dots, \nu_w)$ of p -hooks ν_i from λ . The sign $v(\lambda) = (-1)^{\sum_1^w \ell(\nu_i)}$, where $\ell(\nu_i)$ is the leg length of ν_i , is independent of the choice of $(\nu_1, \nu_2, \dots, \nu_w)$. The residue of the (i, j) -node in the Young diagram of λ is the integer r such that $0 \leq r < p$ and $r \equiv j - i \pmod{p}$. A hook of λ has p -singular length if and only if the hand and foot node residues of the hook are respectively $k, k + 1$ for some k modulo p . We normalize Λ so that λ_i corresponds to hooks of λ of p -singular length and hand node residue $i - 1$. Let λ^* be the dual partition of λ , let λ_0^* be the p -core of λ^* , and let $\Lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_p^*)$ be the p -quotient of λ^* . Then λ_0^* is the dual $(\lambda_0)^*$ of λ_0 and $\lambda_\gamma^* = (\lambda_{p+1-\gamma})^*$ for $1 \leq \gamma \leq p$. In particular, $\lambda = \lambda^*$ if and only if $\lambda_0 = \lambda_0^*$ and $\lambda_\gamma = (\lambda_{p+1-\gamma})^*$ for $1 \leq \gamma \leq p$. Indeed, the (i, j) -hook of λ and the (j, i) -hook of λ^* have the same length. So if this length is p -singular, then the hand node residues of these hooks are of the form k and $-k - 1$ modulo p .

Let $Y^\vee = \{\xi_\gamma : 1 \leq \gamma \leq p\}$. The characters in $(H, f)^\vee$ have the form $\chi_{\lambda_0} \times \psi_\Lambda$, where $(H_0, f_0)^\vee = \{\chi_{\lambda_0}\}$, $\psi_\Lambda \in H_1^\vee$, and Λ is a mapping

$$Y^\vee \longrightarrow \{\text{partitions}\}, \quad \xi_\gamma \mapsto \lambda_\gamma,$$

such that $\sum_\gamma |\lambda_\gamma| = w$. We also represent Λ by the p -tuple $(\lambda_1, \lambda_2, \dots, \lambda_p)$. We recall the definition of ψ_Λ . Let $\Omega = \coprod_{\gamma=1}^p \Omega_\gamma^\Lambda$ be a partition where $|\Omega_\gamma^\Lambda| = |\lambda_\gamma|$, and let $\xi_\Lambda \in (Y^\Omega)^\vee$ be such that $(\xi_\Lambda)_i = \xi_\gamma$ for $i \in \Omega_\gamma^\Lambda$. The stabilizer $H_{1,\Lambda}$ of ξ_Λ in H_1 is $Y^\Omega S(\Omega^\Lambda)$, where

$$S(\Omega^\Lambda) = S(\Omega_1^\Lambda) \times S(\Omega_2^\Lambda) \times \dots \times S(\Omega_p^\Lambda),$$

and ξ_Λ extends to a character $E(\xi_\Lambda)$ of $H_{1,\Lambda}$. Let $\omega_\Lambda = \omega_{\lambda_1} \times \omega_{\lambda_2} \times \dots \times \omega_{\lambda_p}$ be the irreducible character of $S(\Omega^\Lambda)$, where ω_{λ_γ} is the character of $S(\Omega_\gamma^\Lambda)$ parametrized by λ_γ . We may view ω_Λ as a character of $H_{1,\Lambda}$ by inflation. Then

$$\psi_\Lambda = \text{Ind}_{H_{1,\Lambda}}^{H_1} (E(\xi_\Lambda)\omega_\Lambda).$$

We give a formula for $E(\xi_\Lambda)(f, \sigma)$ for $f \in Y^\Omega$ and $\sigma \in S(\Omega^\Lambda)$. Write $\sigma = \sigma_1 \sigma_2 \dots \sigma_d$ as a product of disjoint cycles in $S(\Omega)$ and write

$$(f, \sigma) = (f_1, \sigma_1)(f_2, \sigma_2) \dots (f_d, \sigma_d),$$

where $f_\delta \in Y^{\Omega_\delta^\sigma}$ and Ω_δ^σ is the support of σ_δ . Let $n_\delta = |\Omega_\delta^\sigma|$, let $i_\delta \in \Omega_\delta^\sigma$, and let

$$p_{i_\delta}(f, \sigma) = f(i_\delta)f(\sigma^{-1}(i_\delta)) \dots f(\sigma^{-(n_\delta-1)}(i_\delta)).$$

In particular, $(f_\delta, \sigma_\delta)^{n_\delta} = (p_{i_\delta}(f, \sigma), 1)$. Moreover

$$(3.1) \quad \begin{aligned} p_{i_\delta}({}^h(f, \sigma)) &= h_{i_\delta} p_{i_\delta}(f, \sigma) h_{i_\delta}^{-1}, \\ p_{g(i_\delta)}({}^g(f, \sigma)) &= p_{i_\delta}(f, \sigma) \end{aligned}$$

for $h \in Y^\Omega$ and $g \in S(\Omega)$. Since $(f, \sigma) \in H_{1,\Lambda}$, σ stabilizes each Ω_γ^Λ . Let Δ_γ be index sets such that $\Omega_\gamma^\Lambda = \coprod_{\delta \in \Delta_\gamma} \Omega_\delta^\sigma$ is the orbit decomposition of σ on Ω_γ^Λ . Then

$$(3.2) \quad E(\xi_\Lambda)(f, \sigma) = \prod_\gamma \prod_{\delta \in \Delta_\gamma} \xi_\gamma(p_{i_\delta}(f, \sigma))$$

(see [12, §4.3]). We call $E(\xi_\Lambda)$ the James-Kerber extension of ξ_Λ .

We give a Murnaghan-Nakayama type formula for $\psi_\Lambda(f, \sigma)$ for the preceding element (f, σ) . We fix s with $1 \leq s \leq d$ and consider pairs $(\underline{\zeta}, \underline{\gamma})$ of sequences $\underline{\zeta} = (\zeta_1, \zeta_2, \dots, \zeta_s)$ and $\underline{\gamma} = (\gamma_1, \gamma_2, \dots, \gamma_s)$ with the following properties: ζ_1 is a hook of length $|\sigma_1|$ of the γ_1 -component of Λ ; ζ_2 is a hook of length $|\sigma_2|$ of the γ_2 -component of $(\zeta_1, \gamma_1) * \Lambda$, the p -tuple gotten from Λ by removing ζ_1 from the γ_1 -component of Λ ; ζ_3 is a hook of length $|\sigma_3|$ of the γ_3 -component of $(\zeta_2, \gamma_2) * (\zeta_1, \gamma_1) * \Lambda$, the p -tuple gotten from $(\zeta_1, \gamma_1) * \Lambda$ by removing ζ_2 from the γ_2 -component of $(\zeta_1, \gamma_1) * \Lambda$; and so on. Set

$$(\underline{\zeta}, \underline{\gamma}) * \Lambda = (\zeta_s, \gamma_s) * \dots * (\zeta_2, \gamma_2) * (\zeta_1, \gamma_1) * \Lambda$$

and $v(\underline{\zeta}) = (-1)^{\sum_1^s \ell(\zeta_k)}$.

Proposition (3A). *With the above notation*

$$(3.3) \quad \psi_\Lambda(f, \sigma) = \sum_{(\underline{\zeta}, \underline{\gamma})} v(\underline{\zeta}) \prod_{k=1}^s \xi_{\gamma_k}(p_{i_k}(f, \sigma)) \psi_{(\underline{\zeta}, \underline{\gamma}) * \Lambda}(f', \sigma')$$

where $(f', \sigma') = (f_{s+1}, \sigma_{s+1}) \cdots (f_d, \sigma_d)$ and $(\underline{\zeta}, \underline{\gamma})$ runs over all pairs of the above type.

Proof. We consider first the case $s = 1$. By definition

$$\begin{aligned} \psi_\Lambda(f, \sigma) &= \text{Ind}_{H_{1,\Lambda}}^{H_1} (E(\xi_\Lambda)\omega_\Lambda)(f, \sigma) \\ &= \sum_{g \in S(\Omega^\Lambda) \backslash S(\Omega)} (E(\xi_\Lambda)\omega_\Lambda)^g(f, \sigma), \end{aligned}$$

since $H_1 = Y^\Omega S(\Omega)$ and $H_{1,\Lambda} = Y^\Omega S(\Omega^\Lambda)$. We identify $S(\Omega')$ for $\Omega' \subseteq \Omega$ with the subgroup of $S(\Omega)$ fixing $\Omega - \Omega'$ elementwise. Let $\Omega' = \Omega - \Omega_1^\sigma$ and decompose

$$S(\Omega) = \bigcup_{g \in \Gamma} S(\Omega^\Lambda)gS(\Omega')$$

as a disjoint union of double cosets modulo $(S(\Omega^\Lambda), S(\Omega'))$. We say that a representative g in Γ is good and that the double coset $S(\Omega^\Lambda)gS(\Omega')$ is good if $g(\Omega_1^\sigma) \subseteq \Omega_\gamma^\Lambda$ for some γ . This definition is independent of the choice of coset representative g , since $S(\Omega')$ fixes Ω_1^σ elementwise and $S(\Omega^\Lambda)$ stabilizes each Ω_γ^Λ . Moreover, the integer γ such that $g(\Omega_1^\sigma) \subseteq \Omega_\gamma^\Lambda$ depends only on the double coset. Let Γ' be the subset of good representatives g in Γ . If g and g' are in Γ' and $g(\Omega_1^\sigma)$ and $g'(\Omega_1^\sigma)$ are contained in the same Ω_γ^Λ , then $g = g'$. Indeed, there exists $x \in S(\Omega^\Lambda)$ such that $x(g(i)) = g'(i)$ for all $i \in \Omega_1^\sigma$, so $g^{-1}x^{-1}g' \in S(\Omega')$ and $g' \in S(\Omega^\Lambda)gS(\Omega')$. Thus there is a bijection between good double cosets and components λ_γ of Λ such that $|\lambda_\gamma| \geq |\sigma_1|$.

Given g in Γ , let $S(\Omega^\Lambda)gS(\Omega') = \bigcup_{t \in T_g} S(\Omega^\Lambda)gt$, where T_g is a set of representatives t for the cosets $(g^{-1}S(\Omega^\Lambda)g \cap S(\Omega')) \backslash S(\Omega')$. If $g \in \Gamma'$ and $g(\Omega_1^\sigma) \subseteq \Omega_{\gamma_1}^\Lambda$,

then

$$\begin{aligned}
 (3.4) \quad g^{-1}S(\Omega^\Lambda)g \cap S(\Omega') &= S(g^{-1}(\Omega^\Lambda)) \cap S(\Omega') \\
 &= S(g^{-1}(\Omega_{\gamma_1}^\Lambda) - \Omega_1^\sigma) \times \prod_{\gamma \neq \gamma_1} S(g^{-1}(\Omega_\gamma^\Lambda)).
 \end{aligned}$$

For $g \in \Gamma'$ we say t in T_g and the coset $S(\Omega^\Lambda)gt$ are good if $g^t(f, \sigma) \in H_{1,\Lambda}$. Thus t is good if and only if for each $\delta = 2, 3, \dots, d$, there is a γ such that $gt(\Omega_\delta^\sigma) \subseteq \Omega_\gamma^\Lambda$. This definition is independent of the choice of representative t . Let T'_g be the subset of good representatives t in T_g . It follows that

$$(3.5) \quad \psi_\Lambda(f, \sigma) = \sum_{g \in \Gamma'} \sum_{t \in T'_g} (E(\xi_\Lambda)\omega_\Lambda)^{g^t}(f, \sigma)$$

since $E(\xi_\Lambda)$ and ω_Λ are zero by convention on arguments not in $H_{1,\Lambda}$. To prove (3.3) for the case $s = 1$, it suffices to show that

$$(3.6) \quad \sum_{t \in T'_g} (E(\xi_\Lambda)\omega_\Lambda)^{g^t}(f, \sigma) = \sum_{\zeta_1} v(\zeta_1)\xi_{\gamma_1}(p_{i_1}(f, \sigma))\psi_{(\zeta_1, \gamma_1)*\Lambda}(f', \sigma'),$$

where ζ_1 runs over all hooks of length $|\sigma_1|$ in λ_{γ_1} .

Given $g \in \Gamma'$ such that $g(\Omega_1^\sigma) \subseteq \Omega_{\gamma_1}^\Lambda$ and given t in T'_g , let $\Delta_{gt,\gamma}$ be the index sets such that

$$\Omega_\gamma^\Lambda = \bigcup_{\delta \in \Delta_{gt,\gamma}} gt(\Omega_\delta^\sigma).$$

Let

$$\Delta'_{gt,\gamma} = \begin{cases} \Delta_{gt,\gamma} & \text{if } \gamma \neq \gamma_1, \\ \Delta_{gt,\gamma} - \{1\} & \text{if } \gamma = \gamma_1. \end{cases}$$

Then

$$\begin{aligned}
 E(\xi_\Lambda)^{g^t}(f, \sigma) &= \prod_{\gamma} \prod_{\delta \in \Delta_{gt,\gamma}} \xi_\gamma(p_{gt(i_\delta)}(g^t(f, \sigma))) \\
 &= \xi_{\gamma_1}(p_{gt(i_1)}(g^t(f, \sigma_1))) \left[\prod_{\gamma} \prod_{\delta \in \Delta'_{gt,\gamma}} \xi_\gamma(p_{gt(i_\delta)}(g^t(f', \sigma'))) \right] \\
 &= \xi_{\gamma_1}(p_{i_1}(f_1, \sigma_1)) \left[\prod_{\gamma} \prod_{\delta \in \Delta'_{gt,\gamma}} \xi_\gamma(p_{gt(i_\delta)}(g^t(f', \sigma'))) \right].
 \end{aligned}$$

The usual Murnaghan-Nakayama formula (see [12, 21.1]) gives

$$\begin{aligned}
 \omega_\Lambda(g^t\sigma) &= \prod_{\gamma} \omega_{\lambda_\gamma} \left(\prod_{\delta \in \Delta_{gt,\gamma}} g^t(\sigma_\delta) \right) \\
 &= \omega_{\lambda_{\gamma_1}} \left(\prod_{\delta \in \Delta_{gt,\gamma_1}} g^t(\sigma_\delta) \right) \prod_{\gamma \neq \gamma_1} \omega_{\lambda_\gamma} \left(\prod_{\delta \in \Delta_{gt,\gamma}} g^t(\sigma_\delta) \right) \\
 &= \sum_{\zeta_1} v(\zeta_1)\omega_{\zeta_1*\lambda_{\gamma_1}} \left(\prod_{\delta \in \Delta'_{gt,\gamma_1}} g^t(\sigma_\delta) \right) \prod_{\gamma \neq \gamma_1} \omega_{\lambda_\gamma} \left(\prod_{\delta \in \Delta_{gt,\gamma}} g^t(\sigma_\delta) \right) \\
 &= \sum_{\zeta_1} v(\zeta_1)\omega_{(\zeta_1, \gamma_1)*\Lambda}(g^t\sigma'),
 \end{aligned}$$

where ζ_1 runs over all hooks of length $|\sigma_1|$ in λ_{γ_1} . Thus

$$\begin{aligned} \sum_{t \in T'_g} (\mathbb{E}(\xi_\Lambda)\omega_\Lambda)^{(gt}(f, \sigma)) &= \sum_{\zeta_1} v(\zeta_1)\xi_{\gamma_1}(p_{i_1}(f_1, \sigma_1)) \\ &\times \sum_{t \in T'_g} \left[\prod_{\gamma} \prod_{\delta \in \Delta'_{gt, \gamma}} \xi_\gamma(p_{gt(i_\delta)}(gt(f', \sigma')))\right] \omega_{(\zeta_1, \gamma_1)*\Lambda}^{(gt}\sigma'). \end{aligned}$$

It suffices to show that if ζ_1 is a fixed hook in λ_{γ_1} of length $|\sigma_1|$, then

$$(3.7) \quad \psi_{(\zeta_1, \gamma_1)*\Lambda}(f', \sigma') = \sum_{t \in T'_g} \left[\prod_{\gamma} \prod_{\delta \in \Delta'_{gt, \gamma}} \xi_\gamma(p_{gt(i_\delta)}(gt(f', \sigma')))\right] \omega_{(\zeta_1, \gamma_1)*\Lambda}^{(gt}\sigma').$$

Now $(\zeta_1, \gamma_1) * \Lambda$ corresponds to the p -tuple $\Lambda' = (\lambda'_1, \lambda'_2, \dots, \lambda'_p)$, where

$$\lambda'_\gamma = \begin{cases} \lambda_\gamma & \text{if } \gamma \neq \gamma_1, \\ \zeta_1 * \lambda_\gamma & \text{if } \gamma = \gamma_1. \end{cases}$$

Let $\psi_{\Lambda'}$ be the irreducible character of $Y \wr S(\Omega')$ parametrized by Λ' . We use notation analogous to that for ψ_Λ . Namely, let $\Omega' = \bigcup_{\gamma} \Omega'^{\Lambda'}_\gamma$, where

$$\Omega'^{\Lambda'}_\gamma = \begin{cases} g^{-1}(\Omega^\Lambda_\gamma) & \text{if } \gamma \neq \gamma_1, \\ g^{-1}(\Omega^\Lambda_{\gamma_1}) - \Omega^\sigma_1 & \text{if } \gamma = \gamma_1. \end{cases}$$

So λ'_γ is a partition of $|\Omega'^{\Lambda'}_\gamma|$. Let $\xi_{\Lambda'} \in (Y^{\Omega'})^\vee$ be such that $(\xi_{\Lambda'})_i = \xi_\gamma$ for $i \in \Omega'^{\Lambda'}_\gamma$. Then the stabilizer of $\xi_{\Lambda'}$ in $Y^{\Omega'}S(\Omega')$ is $Y^{\Omega'}S(\Omega'^{\Lambda'})$, where

$$S(\Omega'^{\Lambda'}) = S(\Omega'^{\Lambda'}_1) \times S(\Omega'^{\Lambda'}_2) \times \dots \times S(\Omega'^{\Lambda'}_p).$$

In particular, T'_g is a set of coset representatives of $S(\Omega'^{\Lambda'})$ in $S(\Omega')$ by (3.4). Let $\omega_{\Lambda'} = \omega_{\lambda'_1} \times \omega_{\lambda'_2} \times \dots \times \omega_{\lambda'_p}$ be the irreducible character of $S(\Omega'^{\Lambda'})$, where $\omega_{\lambda'_\gamma}$ is the irreducible character of $S(\Omega'^{\Lambda'}_\gamma)$ parametrized by λ'_γ . Let $\mathbb{E}(\xi_{\Lambda'})$ be the James-Kerber extension of $\xi_{\Lambda'}$ to $Y^{\Omega'}S(\Omega'^{\Lambda'})$. Then

$$\begin{aligned} \psi_{(\zeta_1, \gamma_1)*\Lambda}(f', \sigma') &= \psi_{\Lambda'}(f', \sigma') \\ &= \text{Ind}_{Y^{\Omega'}S(\Omega'^{\Lambda'})}^{Y^{\Omega'}S(\Omega')}(\mathbb{E}(\xi_{\Lambda'})\omega_{\Lambda'})(f', \sigma') \\ &= \sum_{t \in T'_g} (\mathbb{E}(\xi_{\Lambda'})\omega_{\Lambda'})^{(t}(f', \sigma')). \end{aligned}$$

We can now show (3.7). Indeed, ${}^t(f', \sigma') \in S(\Omega'^{\Lambda'})$ if and only if for each $\delta \geq 2$, there exists a γ such that $t(\Omega^\sigma_\delta) \subseteq \Omega'^{\Lambda'}_\gamma$, or equivalently, such that $gt(\Omega^\sigma_\delta) \subseteq \Omega^\Lambda_\gamma$. Moreover, $\Omega'^{\Lambda'}_\gamma = \prod_{\delta \in \Delta'_{gt, \gamma}} t(\Omega^\sigma_\delta)$, since $\Omega^\Lambda_\gamma = \prod_{\delta \in \Delta_{gt, \gamma}} gt(\Omega^\sigma_\delta)$. Thus

$$\begin{aligned} \mathbb{E}(\xi_{\Lambda'})^{(t}(f', \sigma')) &= \prod_{\gamma} \prod_{\delta \in \Delta'_{gt, \gamma}} \xi_\gamma(p_{t(i_\delta)}({}^t(f', \sigma'))) \\ &= \prod_{\gamma} \prod_{\delta \in \Delta_{gt, \gamma}} \xi_\gamma(p_{gt(i_\delta)}(gt(f', \sigma'))) \end{aligned}$$

and

$$\begin{aligned} \omega_{\Lambda'}(t(\sigma')) &= \prod_{\gamma} \omega_{\lambda'_\gamma} \left(\prod_{\delta \in \Delta'_{gt,\gamma}} t(\sigma_\delta) \right) \\ &= \omega_{\lambda'_{\gamma_1}} \left(\prod_{\delta \in \Delta'_{gt,\gamma_1}} gt(\sigma_\delta) \right) \prod_{\gamma \neq \gamma_1} \omega_{\lambda_\gamma} \left(\prod_{\delta \in \Delta_{gt,\gamma}} gt(\sigma_\delta) \right) \\ &= \omega_{\Lambda'}(gt(\sigma')). \end{aligned}$$

This completes the proof of (3.3) for the case $s = 1$. The general case of (3.3) now follows by iteration. \square

Remark. Formula (3.3) does not depend on the structure of Y . Equivalent formulas can also be found in [14, Théorème 2.8] and in [15, Theorem 4.3].

Definition. Let $A(\Gamma)$ be the alternating group on Γ and label ξ_γ in Y^\vee so that ξ_γ and $\xi_{p+1-\gamma}$ have the same restriction to $Y^+ = Y \cap A(\Gamma)$ for $1 \leq \gamma \leq p$. In particular, $\xi_\gamma(1) = 1$ for $\gamma \neq p^*$ and $\xi_\gamma(1) = p - 1$ for $\gamma = p^*$, where $p^* = \frac{1}{2}(p + 1)$. The Rouquier isometry $\mu = (J, \epsilon)$ in $\mathcal{R}_K((G, e), (H, f))$ is defined as follows: For each partition λ of n with p -core λ_0 and p -quotient $\Lambda = (\lambda_1, \lambda_2, \dots, \lambda_p)$, let

$$(3.8) \quad \begin{aligned} J(\chi_\lambda) &= \chi_{\lambda_0} \times \psi_\Lambda, & \epsilon(\chi_\lambda) &= 1 & \text{if } p = 2, \\ J(\chi_\lambda) &= \chi_{\lambda_0} \times \psi_{\hat{\Lambda}}, & \epsilon(\chi_\lambda) &= v(\lambda)(-1)^{|\lambda_{p^*}|} & \text{if } p > 2, \end{aligned}$$

where $\hat{\Lambda}$ is the p -tuple gotten from Λ by replacing λ_{p^*} by $(\lambda_{p^*})^*$.

We define a Rouquier isometry in a more general context. An element $x = (f, 1)$ of D is called standard if the non-identity values of f are all equal. For a standard element x let $\Omega = \Omega_{x,0} \cup \Omega_{x,1}$, where x acts trivially on $\Gamma \times \Omega_{x,0}$ and fixed-point-freely on $\Gamma \times \Omega_{x,1}$. Thus

$$G_x = G_{x,0} \times G_{x,1}, \quad H_x = H_{x,0} \times H_{x,1},$$

where $G_{x,0} = S(\Pi_0 \cup (\Gamma \times \Omega_{x,0}))$, $H_{x,0} = S(\Pi_0) \times (Y \wr S(\Omega_{x,0}))$, and $G_{x,1} = H_{x,1} = P \wr S(\Omega_{x,1})$. Let $H_{x,0} = H_{x,00} \times H_{x,01}$, where $H_{x,00} = S(\Pi_0)$ and $H_{x,01} = Y \wr S(\Omega_{x,0})$. Let

$$e_x = e_{x,0} \times e_{x,1}, \quad f_x = f_{x,00} \times f_{x,01} \times f_{x,1},$$

where $e_{x,0}$ is the block of $\mathcal{O}G_{x,0}$ parametrized by λ_0 , $e_{x,1}$ is the 1-block of $\mathcal{O}G_{x,1}$, $f_{x,00}$ is the block of $\mathcal{O}H_{x,00}$ parametrized by λ_0 , $f_{x,01}$ is the 1-block of $\mathcal{O}H_{x,01}$, and $f_{x,1}$ is the 1-block of $\mathcal{O}H_{x,1}$. In particular, $e_{x,1}$, $f_{x,01}$, and $f_{x,1}$ are the unique blocks of $\mathcal{O}G_{x,1}$, $\mathcal{O}H_{x,01}$, and $\mathcal{O}H_{x,1}$. Let $f_{x,0}$ be the block $f_{x,00} \times f_{x,01}$ of $\mathcal{O}H_{x,0}$. Then $e_{x,0}$ and $f_{x,0}$ have defect group $P^{\Omega_{x,0}}$ and $f_{x,0}$ is the Brauer correspondent of $e_{x,0}$. Let $\mu_{x,0} = (J_{x,0}, \epsilon_{x,0})$ be the Rouquier isometry in $\mathcal{R}_K((G_{x,0}, e_{x,0}), (H_{x,0}, f_{x,0}))$. Let $\mu_{x,1} = (J_{x,1}, \epsilon_{x,1})$ be the identity isometry in $\mathcal{R}_K((G_{x,1}, 1), (H_{x,1}, 1))$. Let $\mu_x = (J_x, \epsilon_x)$ be the isometry $\mu_{x,0} \times \mu_{x,1}$ in $\mathcal{R}_K((G_x, e_x), (H_x, f_x))$. If $\mu_{x,0}$ is perfect, then μ_x is perfect. We call μ_x the Rouquier isometry in $\mathcal{R}_K((G_x, e_x), (H_x, f_x))$. We note that μ_x depends only on $\langle x \rangle$ and not on the generator x .

For each $j = 0, 1, 2, \dots, w$ let x_j be a standard element of D such that $|\Omega_{x_j,1}| = j$. The $w+1$ subgroups $\langle x_j \rangle$ then form a set of representatives for the $N_G(D, e_D)$ -orbits of cyclic subgroups of D , since $N_G(D, e_D) \supseteq (Y \wr S(\Omega))$.

Theorem (3B). *e and f are isotypic blocks of $\mathcal{O}G$ and $\mathcal{O}H$ with local system given by the translates of the Rouquier isometries μ_x for x in $\{x_0, x_1, \dots, x_w\}$.*

Proof. Condition **(I1)** holds by [1, Proposition 4.2] since D is abelian. The blocks e and f are of principal type by a theorem of Marichal and Puig (see [3, (2.2)]). Let $x \in \{x_1, x_2, \dots, x_w\}$. We prove the Rouquier isometries μ_x are perfect by induction on $|D|$. The claim is clear if $|D| = 1$, so we may suppose $D > 1$. By induction $\mu_{x,0}$ and hence μ_x are perfect isometries. We claim the following diagram is commutative:

$$(3.9) \quad \begin{array}{ccc} \mathcal{R}_K(G, e) & \xrightarrow{R_\mu} & \mathcal{R}_K(H, f) \\ \downarrow d_G^x & & \downarrow d_H^x \\ \text{CF}_{p'}(G_x, e_x) & \xrightarrow{R_{\mu_x, p'}} & \text{CF}_{p'}(H_x, f_x) \end{array}$$

Suppose $p = 2$. Then $w = 1$, $G_x = H_x = S(\Pi_0) \times \langle x \rangle$, and λ is a partition with 2-core λ_0 and 2-quotient $(1, 0)$ or $(0, 1)$. The ordinary Murnaghan-Nakayama formula implies the commutativity of (3.9). Suppose $p > 2$. We consider the top-right portion of (3.9). Let $\chi_\lambda \in (G, e)^\vee$, let $y \in H_{x,p'}$, and write $y = y_{00}y_{01}y_1$, where $y_{00} \in H_{x,00}$, $y_{01} \in H_{x,01}$, and $y_1 \in H_{x,1}$. Let $xy_1 = (f_1, \sigma_1)(f_2, \sigma_2) \cdots (f_s, \sigma_s)$, where the σ_i are disjoint cycles of $S(\Omega_{x,1})$. Then

$$\begin{aligned} d_H^x(R_\mu(\chi_\lambda))(y) &= \epsilon(\chi_\lambda)(\chi_{\lambda_0} \times \psi_{\hat{\lambda}})(xy) \\ &= \epsilon(\chi_\lambda)\chi_{\lambda_0}(y_{00})\psi_{\hat{\lambda}}(y_{01}(xy_1)) \\ &= \epsilon(\chi_\lambda)\chi_{\lambda_0}(y_{00}) \sum_{(\zeta, \underline{\gamma})} v(\zeta) \prod_{k=1}^s \xi_{\gamma_k}(p_{i_k}(f_k, \sigma_k)) \psi_{(\zeta, \underline{\gamma}) * \hat{\lambda}}(y_{01}) \end{aligned}$$

by (3A). For each pair $(\zeta, \underline{\gamma})$ in the formula, let $M(\underline{\gamma}) = \{k : \gamma_k = p^*\}$. Then

$$(3.10) \quad d_H^x(R_\mu(\chi_\lambda))(y) = \epsilon(\chi_\lambda)\chi_{\lambda_0}(y_{00}) \sum_{(\zeta, \underline{\gamma})} v(\zeta)(-1)^{|M(\underline{\gamma})|} \psi_{(\zeta, \underline{\gamma}) * \hat{\lambda}}(y_{01})$$

since $p_{i_k}(f_k, \sigma_k) \in P^\#$, so that $\xi_{\gamma_k}(p_{i_k}(f_k, \sigma_k)) = 1$ or -1 according as $\gamma_k \neq p^*$ or $\gamma_k = p^*$.

We consider the left-bottom portion of (3.9). Let $g \in G_{x,p'}$ and write $g = g_0g_1$, where $g_0 \in G_{x,0,p'}$, $g_1 \in G_{x,1,p'}$. Each p' -conjugacy class of $G_{x,1}$ contains a unique conjugacy class of $S(\Omega_{x,1})$, since $S(\Omega_{x,1})$ is a Hall p -complement in $G_{x,1}$. We may suppose without loss of generality that $g_1 = (1, \sigma)$ for some σ in $S(\Omega_{x,1})$. Let $\sigma = \sigma_1\sigma_2 \cdots \sigma_s$ be the cycle decomposition of σ in $S(\Omega_{x,1})$. The Murnaghan-Nakayama formula implies that

$$\begin{aligned} d_G^x(\chi_\lambda)(g) &= \chi_\lambda(xg) \\ &= \chi_\lambda(g_0(xg_1)) \\ &= \sum_{\underline{\nu}} v(\underline{\nu})\chi_{\underline{\nu} * \lambda}(g_0), \end{aligned}$$

where $\underline{\nu}$ runs over sequences $(\nu_1, \nu_2, \dots, \nu_s)$ of hooks ν_k of length $p|\sigma_k|$ of λ . Let τ run over representatives of the p' -conjugacy classes of $S(\Omega_{x,1})$, let η_τ be the characteristic function of the $G_{x,1}$ -conjugacy class of τ , and set

$$\alpha_\lambda = \sum_{\underline{\nu}, \tau} v(\underline{\nu})(\chi_{\underline{\nu} * \lambda} \times \eta_\tau).$$

Then

$$\alpha_\lambda(g_0g_1) = \begin{cases} \sum_{\underline{\nu}} v(\underline{\nu})\chi_{\underline{\nu}*\lambda}(g_0) & \text{if } g_1 \in G_{x,1,p'}, \\ 0 & \text{if } g_1 \notin G_{x,1,p'}. \end{cases}$$

for $g_0 \in G_{x,0}$ and $g_1 \in G_{x,1}$. Thus $d_G^x(\chi_\lambda) = d_{G_x}^1(\alpha_\lambda)$ and

$$\begin{aligned} R_{\mu_x,p'}(d_G^x(\chi_\lambda)) &= R_{\mu_x,p'}(d_{G_x}^1(\alpha_\lambda)) \\ &= d_{H_x}^1(R_{\mu_x}(\alpha_\lambda)), \end{aligned}$$

where the second equality holds since μ_x is perfect by induction. Now

$$\begin{aligned} R_{\mu_x}(\alpha_\lambda) &= \sum_{\underline{\nu}, \tau} v(\underline{\nu})(R_{\mu_{x,0}}(\chi_{\underline{\nu}*\lambda}) \times R_{\mu_{x,1}}(\eta_\tau)) \\ &= \sum_{\underline{\nu}, \tau} v(\underline{\nu})\epsilon_{x,0}(\chi_{\underline{\nu}*\lambda})(\chi_{\lambda_0} \times \psi_{(\underline{\zeta}, \underline{\gamma})*\hat{\Lambda}} \times \eta_\tau), \end{aligned}$$

where $(\underline{\zeta}, \underline{\gamma})$ is obtained from $\underline{\nu}$ as follows: First $\underline{\nu}$ determines a pair $(\tilde{\zeta}, \underline{\gamma})$ of sequences $(\tilde{\zeta}_1, \tilde{\zeta}_2, \dots, \tilde{\zeta}_s)$ and $(\gamma_1, \gamma_2, \dots, \gamma_s)$, where $\tilde{\zeta}_k$ is a hook of length $|\sigma_k|$ of the γ_k -component of Λ . So $\underline{\nu}$ determines a pair $(\underline{\zeta}, \underline{\gamma})$ of sequences $(\zeta_1, \zeta_2, \dots, \zeta_s)$ and $(\gamma_1, \gamma_2, \dots, \gamma_s)$, where ζ_k is a hook of length $|\sigma_k|$ of the γ_k -component of $\hat{\Lambda}$, and $\zeta_k = \tilde{\zeta}_k$ or $(\tilde{\zeta}_k)^*$ according as $\gamma_k \neq p^*$ or $\gamma_k = p^*$. Thus

$$R_{\mu_x,p'}(d_G^x(\chi_\lambda)) = \sum_{\underline{\nu}, \tau} v(\underline{\nu})\epsilon_{x,0}(\chi_{\underline{\nu}*\lambda}) d_{H_x}^1(\chi_{\lambda_0} \times \psi_{(\underline{\zeta}, \underline{\gamma})*\hat{\Lambda}} \times \eta_\tau).$$

If $y \in H_{x,p'}$ and $y = y_{00}y_{01}y_1$, where $y_{00} \in H_{x,00}$, $y_{01} \in H_{x,01}$, and $y_1 \in H_{x,1}$, then

$$(3.11) \quad R_{\mu_x,p'}(d_G^x(\chi_\lambda))(y) = \sum_{\underline{\nu}} v(\underline{\nu})\epsilon_{x,0}(\chi_{\underline{\nu}*\lambda})\chi_{\lambda_0}(y_{00})\psi_{(\underline{\zeta}, \underline{\gamma})*\hat{\Lambda}}(y_{01}).$$

The commutativity of the diagram (3.9) holds by (3.10) and (3.11) if

$$(3.12) \quad \epsilon(\chi_\lambda)v(\underline{\zeta})(-1)^{|M(\underline{\gamma})|} = v(\underline{\nu})\epsilon_{x,0}(\chi_{\underline{\nu}*\lambda})$$

for corresponding $(\underline{\zeta}, \underline{\gamma})$ and $\underline{\nu}$. Now

$$\begin{aligned} \epsilon(\chi_\lambda) &= v(\lambda)(-1)^{|\lambda_{p^*}|}, \\ \epsilon_{x,0}(\chi_{\underline{\nu}*\lambda}) &= v(\underline{\nu}*\lambda)(-1)^{|(\underline{\nu}*\lambda)_{p^*}|}. \end{aligned}$$

So (3.12) becomes

$$v(\lambda)(-1)^{|\lambda_{p^*}|}v(\underline{\zeta})(-1)^{|M(\underline{\gamma})|} = v(\underline{\nu})v(\underline{\nu}*\lambda)(-1)^{|(\underline{\nu}*\lambda)_{p^*}|}.$$

Since $|(\underline{\nu}*\lambda)_{p^*}| = |\lambda_{p^*}| - \sum_{k \in M(\underline{\gamma})} |\zeta_k|$, (3.12) is equivalent to

$$(3.13) \quad \begin{aligned} v(\lambda)v(\underline{\nu}*\lambda) &= v(\underline{\nu})v(\underline{\zeta})(-1)^{|\lambda_{p^*}|}(-1)^{|M(\underline{\gamma})|}(-1)^{|(\underline{\nu}*\lambda)_{p^*}|} \\ &= v(\underline{\nu})v(\underline{\zeta}) \prod_{k \in M(\underline{\gamma})} (-1)^{(|\zeta_k|-1)}. \end{aligned}$$

But $v(\lambda)v(\underline{\nu}*\lambda) = v(\tilde{\zeta})v(\underline{\nu})$ by [8, (2.15)] or [9, Théorème 1.4, b)]. Since

$$v(\underline{\zeta})v(\tilde{\zeta}) = \prod_{k \in M(\underline{\gamma})} (-1)^{(|\zeta_k|-1)},$$

(3.13) holds and (3.9) is commutative. This argument for the commutativity of (3.9) still applies if x is replaced by any generator of $\langle x \rangle$, since the lower horizontal arrow in (3.9) depends only on $\langle x \rangle$. (3B) now follows by transport of structure, as remarked in §1. □

4. AN ANALOG OF FROBENIUS’S THEOREM

Let G and G^+ be the symmetric group $S(\Pi)$ and the alternating group $A(\Pi)$ on Π . An irreducible character χ_λ of G is reducible on G^+ if and only if λ is self-dual. For such λ let $\eta(\lambda)$ be the partition of n with parts $\eta_{11}(\lambda), \eta_{22}(\lambda), \dots, \eta_{ss}(\lambda)$, where $\eta_{ii}(\lambda)$ is the hook length of the (i, i) -node of λ and s is the number of diagonal nodes of λ . Since the $\eta_{ii}(\lambda)$ are distinct odd integers, the conjugacy class $\mathcal{C}_{\eta(\lambda)}$ of G parametrized by $\eta(\lambda)$ lies in G^+ and splits in G^+ . The mapping $\lambda \mapsto \eta(\lambda)$ thus induces a bijection $\chi_\lambda \mapsto \mathcal{C}_{\eta(\lambda)}$ from irreducible characters χ_λ of G which split on G^+ to conjugacy classes $\mathcal{C}_{\eta(\lambda)}$ of G which lie in G^+ and split in G^+ . Let $\text{Res}_{G^+}^G(\chi_\lambda) = \chi_{\lambda,+} + \chi_{\lambda,-}$ and $\mathcal{C}_{\eta(\lambda)} = \mathcal{C}_{\eta(\lambda),+} \cup \mathcal{C}_{\eta(\lambda),-}$ be the corresponding decompositions. We use the convention that $\sqrt{a} = \sqrt{-a}i$ for negative real a , where i is a fixed primitive 4-th root of unity in K . In addition, we write $\sqrt{\pm\eta(\lambda)}$ for $\sqrt{\pm\eta_{11}(\lambda)\eta_{22}(\lambda)\cdots\eta_{ss}(\lambda)}$. The following is Frobenius’s Theorem [11, §3].

Theorem (4A). *Suppose χ_λ is an irreducible character of G which splits on G^+ . Let $g \in G^+$. Then $(\chi_{\lambda,+} - \chi_{\lambda,-})(g) \neq 0$ if and only if g is in $\mathcal{C}_{\eta(\lambda)}$. Moreover, $\chi_{\lambda,\pm}$ and $\mathcal{C}_{\eta(\lambda),\pm}$ may be labeled so that*

$$\begin{aligned} \chi_{\lambda,\pm}(g) &= \frac{1}{2}(\epsilon_\lambda + \sqrt{\epsilon_\lambda\eta(\lambda)}) \quad \text{if } g \in \mathcal{C}_{\eta(\lambda),\pm}, \\ \chi_{\lambda,\pm}(g) &= \frac{1}{2}(\epsilon_\lambda - \sqrt{\epsilon_\lambda\eta(\lambda)}) \quad \text{if } g \in \mathcal{C}_{\eta(\lambda),\mp}, \end{aligned}$$

where $\epsilon_\lambda = (-1)^{\frac{1}{2}(n-s)}$.

We suppose for the rest of this section that $\Pi = \Gamma \times \Omega$, where $\Gamma = \{1, 2, \dots, p\}$ and $|\Omega| = w < p$. Let X and X^+ be the symmetric group $S(\Gamma)$ and the alternating group $A(\Gamma)$ on Γ ; let $Y = N_X(P)$ and $Y^+ = N_{X^+}(P)$, where P is a fixed Sylow p -subgroup of X ; and let $H = N_G(D)$ and $H^+ = N_{G^+}(D)$, where D is the abelian Sylow p -subgroup P^Ω of G . We denote the sign characters of $G, H, X, Y, S(\Omega)$ with respect to $G^+, H^+, X^+, Y^+, A(\Omega)$ by $\text{sgn}_G, \text{sgn}_H, \text{sgn}_X, \text{sgn}_Y, \text{sgn}_{S(\Omega)}$. In particular, $\text{sgn}_H = \text{Res}_H^G(\text{sgn}_G)$ and $\text{sgn}_Y = \text{Res}_Y^X(\text{sgn}_X)$. If (f, σ) is an element of $H = Y \wr S(\Omega)$ with $f \in Y^\Omega$ and $\sigma \in S(\Omega)$, then

$$(4.1) \quad \text{sgn}_H(f, \sigma) = \text{sgn}_{S(\Omega)}(\sigma) \prod_{i \in \Omega} \text{sgn}_Y(f(i)).$$

Thus (f, σ) is in H^+ if and only if $\text{sgn}_{S(\Omega)}(\sigma) = \prod_{i \in \Omega} \text{sgn}_Y(f(i))$. The next two results hold for any finite group Y and a normal subgroup Y^+ of Y .

Lemma (4B). *Let $(f, \sigma) \in Y \wr S(\Omega)$ and let $\Omega = \coprod_{\delta=1}^d \Omega_\delta^\sigma$ be the orbit decomposition of σ on Ω . Then the following hold:*

- (i) $(f, \sigma)^{Y \wr S(\Omega)} \cap (Y^+ \wr S(\Omega)) \neq \emptyset$ if and only if $p_{i_\delta}(f, \sigma) \in Y^+$ for all δ .
- (ii) Let $(f, \sigma) \in Y^+ \wr S(\Omega)$ and $g \in Y^\Omega$. Then ${}^{(g,1)}(f, \sigma) \in (Y^+ \wr S(\Omega))$ if and only if g is constant modulo Y^+ on Ω_δ^σ for $1 \leq \delta \leq d$.

Proof. Part (i) follows from (3.1). For (ii) we note that $(g,1)(f, \sigma) \in (Y^+ \wr S(\Omega))$ if and only if $g(i)f(i)(g(\sigma^{-1}(i)))^{-1} \in Y^+$ for all $i \in \Omega$. Since $f \in (Y^+)^{\Omega}$, this is equivalent to saying $g(i)(g(\sigma^{-1}(i)))^{-1} \in Y^+$ for all $i \in \Omega$. \square

Lemma (4C). *Let ψ^+ be a character of Y^+ , let $\psi = \text{Ind}_{Y^+}^Y(\psi^+)$, and let $E((\psi^+)^{\Omega})$ and $E(\psi^{\Omega})$ be the James-Kerber extensions of $(\psi^+)^{\Omega}$ and ψ^{Ω} to $Y^+ \wr S(\Omega)$ and $Y \wr S(\Omega)$. Then*

$$E(\psi^{\Omega}) = \text{Ind}_{Y^+ \wr S(\Omega)}^{Y \wr S(\Omega)}(E((\psi^+)^{\Omega})).$$

Proof. Let $(f, \sigma) \in (Y \wr S(\Omega))$ and let $\Omega = \coprod_{\delta=1}^d \Omega_{\delta}^{\sigma}$ be the orbit decomposition of σ on Ω . If $(f, \sigma)^{Y \wr S(\Omega)} \cap (Y^+ \wr S(\Omega)) = \emptyset$, then $\text{Ind}_{Y^+ \wr S(\Omega)}^{Y \wr S(\Omega)}(E((\psi^+)^{\Omega}))(f, \sigma) = 0$. On the other hand, by (4B), (i) there exists δ such that $p_{i_{\delta}}(f, \sigma) \notin Y^+$, so that

$$E(\psi^{\Omega})(f, \sigma) = \prod_{\delta} \psi(p_{i_{\delta}}(f, \sigma)) = 0.$$

Thus (4C) holds in this case. We may then suppose that $(f, \sigma) \in (Y^+ \wr S(\Omega))$ and that $p_{i_{\delta}}(f, \sigma) \in Y^+$ for all δ . Let T be a transversal for Y^+ in Y . Then

$$\begin{aligned} E(\psi^{\Omega})(f, \sigma) &= \prod_{\delta} \psi(p_{i_{\delta}}(f, \sigma)) \\ &= \prod_{\delta} \sum_{t \in T} \psi^+(tp_{i_{\delta}}(f, \sigma)t^{-1}) \\ &= \sum_{(t_{\delta}) \in T^d} \prod_{\delta} \psi^+(t_{\delta}p_{i_{\delta}}(f, \sigma)t_{\delta}^{-1}). \end{aligned}$$

T^{Ω} is a transversal for $(Y^+)^{\Omega}$ in Y^{Ω} . Let T_{σ}^{Ω} be the subset of all $g \in T^{\Omega}$ such that g is constant on Ω_{δ}^{σ} for all δ . We may identify such a g with the d -tuple (t_{δ}) , where $t_{\delta} = g(i_{\delta})$. This gives a bijection of T_{σ}^{Ω} with T^d . Then (4B), (ii) and (3.1) imply that

$$\begin{aligned} \text{Ind}_{Y^+ \wr S(\Omega)}^{Y \wr S(\Omega)}(E((\psi^+)^{\Omega}))(f, \sigma) &= \sum_{g \in T_{\sigma}^{\Omega}} E((\psi^+)^{\Omega})(g(f, \sigma)g^{-1}) \\ &= \sum_{(t_{\delta}) \in T^d} \prod_{\delta} \psi^+(t_{\delta}p_{i_{\delta}}(f, \sigma)t_{\delta}^{-1}), \end{aligned}$$

and (4C) holds. \square

Y is a Frobenius group with kernel $P = \langle a \rangle$ of order p and cyclic complement $\langle b \rangle$ of order $p - 1$. The set $\mathcal{R} = \{a, b^i : 1 \leq i \leq p - 1\}$ is a transversal for the conjugacy classes of Y ; the classes contained in Y^+ are represented by a and b^{2i} for $1 \leq i \leq \frac{1}{2}(p - 1)$. Of these only a^Y splits in Y^+ , say $a^Y = a_+^{Y^+} \cup a_-^{Y^+}$. We choose $\xi_1 = \text{sgn}_Y$, $\xi_p = 1_Y$, and label the characters ξ_{γ} in Y^{\vee} so that $\xi_{p+1-\gamma} = \xi_{\gamma} \text{sgn}_Y$ for $1 \leq \gamma \leq p$. We also write $\xi_{p+1-\gamma}$ as ξ_{γ}^* . In particular, $\text{Res}_{Y^+}^Y(\xi_{p^*}) = \xi_{p^*,+} + \xi_{p^*,-}$, where $p^* = \frac{1}{2}(p + 1)$. We label $\xi_{p^*,\pm}$ and a_{\pm} so that

$$\xi_{p^*,+}(x) - \xi_{p^*,-}(x) = \pm\sqrt{\epsilon_p p} \quad \text{for } x \in (a_{\pm})^{Y^+},$$

where $\epsilon_p = (-1)^{\frac{p-1}{2}}$. As in §3 we parametrize the irreducible characters ψ_{Λ} of H by mappings

$$\Lambda: Y^{\vee} \longrightarrow \{\text{partitions}\}, \quad \xi_{\gamma} \mapsto \lambda_{\gamma},$$

where $\sum_{\gamma} |\lambda_{\gamma}| = w$. We also represent Λ by the p -tuple $(\lambda_1, \lambda_2, \dots, \lambda_p)$. In particular, $\text{sgn}_H = \psi_{\Lambda_{\text{sgn}}}$, where $\Lambda_{\text{sgn}} = (\{1^w\}, \emptyset, \emptyset, \dots, \emptyset)$. Indeed, the stabilizer $H_{\Lambda_{\text{sgn}}}$ of Λ_{sgn} is H , and

$$\begin{aligned} E(\xi_{\Lambda_{\text{sgn}}})(f, \sigma) &= \prod_{\delta \in \Delta_1} \xi_1(p_{i_{\delta}}(f, \sigma)) \\ &= \prod_{\delta \in \Delta_1} \prod_{i \in \Omega_{\delta}^{\sigma}} \text{sgn}_Y(f(i)) \\ &= \prod_{i \in \Omega} \text{sgn}_Y(f(i)) \end{aligned}$$

by (3.2). Thus $\psi_{\Lambda_{\text{sgn}}} = \text{sgn}_H$. Let $*$ be the duality $\Lambda \mapsto \Lambda^*$, where $\Lambda^*: \xi_{\gamma} \mapsto (\lambda_{p+1-\gamma})^*$. We say Λ is a splitting mapping of H if ψ_{Λ} is a splitting character of H , i.e., $\text{Res}_{H^+}^H(\psi_{\Lambda}) = \psi_{\Lambda,+} + \psi_{\Lambda,-}$ is reducible on H^+ . The main result of this section is an analog of Frobenius’s theorem for splitting characters ψ_{Λ} .

Proposition (4D). *Let $\psi_{\Lambda} \in H^{\vee}$. Then $\text{sgn}_H \psi_{\Lambda} = \psi_{\Lambda^*}$. In particular, ψ_{Λ} is a splitting character if and only if $\Lambda = \Lambda^*$.*

Proof. Clearly

$$\begin{aligned} \psi_{\Lambda} \text{sgn}_H &= (\text{Ind}_{H_{\Lambda}}^H(E(\xi_{\Lambda})\omega_{\Lambda})) \text{sgn}_H \\ &= \text{Ind}_{H_{\Lambda}}^H(E(\xi_{\Lambda})\omega_{\Lambda} \text{sgn}_{H_{\Lambda}}) \end{aligned}$$

where $\text{sgn}_{H_{\Lambda}} = \text{Res}_{H_{\Lambda}}^H(\text{sgn}_H)$. Let $(f, \sigma) \in H_{\Lambda}$, and write $(f, \sigma) = \prod_{\gamma} (f_{\gamma}, \sigma_{\gamma})$, where $(f_{\gamma}, \sigma_{\gamma})$ in $Y \wr S(\Omega_{\gamma}^{\Lambda})$. Let $\Omega_{\gamma}^{\Lambda} = \prod_{\delta \in \Delta_{\gamma}} \Omega_{\delta}^{\sigma}$ be the orbit decomposition of σ_{γ} on $\Omega_{\gamma}^{\Lambda}$. Since

$$\text{sgn}_{H_{\Lambda}}(f, \sigma) = \prod_{\gamma} \left[\prod_{\delta \in \Delta_{\gamma}} \text{sgn}_Y(p_{i_{\delta}}(f_{\gamma}, \sigma_{\gamma})) \right] \text{sgn}_{S(\Omega_{\gamma}^{\Lambda})}(\sigma_{\gamma}),$$

we have that

$$\begin{aligned} & (E(\xi_{\Lambda})\omega_{\Lambda} \text{sgn}_{H_{\Lambda}})(f, \sigma) \\ &= \prod_{\gamma} \left[\prod_{\delta \in \Delta_{\gamma}} \xi_{\gamma}(p_{i_{\delta}}(f_{\gamma}, \sigma_{\gamma})) \text{sgn}_Y(p_{i_{\delta}}(f_{\gamma}, \sigma_{\gamma})) \right] \omega_{\lambda_{\gamma}}(\sigma_{\gamma}) \text{sgn}_{S(\Omega_{\gamma}^{\Lambda})}(\sigma_{\gamma}) \\ &= \prod_{\gamma} \left[\prod_{\delta \in \Delta_{\gamma}} \xi_{p+1-\gamma}(p_{i_{\delta}}(f_{\gamma}, \sigma_{\gamma})) \right] \omega_{\lambda_{\gamma}^*}(\sigma_{\gamma}) \\ &= (E(\xi_{\Lambda^*})\omega_{\Lambda^*})(f, \sigma). \end{aligned}$$

Thus $\psi_{\Lambda} \text{sgn}_H = \psi_{\Lambda^*}$, and $\text{Res}_{H^+}^H(\psi_{\Lambda})$ is reducible if and only if $\Lambda = \Lambda^*$. □

Each element of H is Y^{Ω} -conjugate to an element (f, σ) of the following type: Let $\Omega = \prod_{\delta}^d \Omega_{\delta}^{\sigma}$ be the orbit decomposition of σ , let $(f, \sigma) = \prod_{\delta} (f_{\delta}, \sigma_{\delta})$, where $f_{\delta} \in Y^{\Omega_{\delta}^{\sigma}}$ and σ_{δ} is the restriction of σ to Ω_{δ}^{σ} , and let $n_{\delta} = |\Omega_{\delta}^{\sigma}|$. Then f_{δ} takes a value r_{δ} on \mathcal{R} on a chosen element i_{δ} in Ω_{δ}^{σ} and value 1 on $\Omega_{\delta}^{\sigma} - \{i_{\delta}\}$. In particular, $p_{i_{\delta}}(f, \sigma) = r_{\delta}$. The data $\prod_{\delta} [r_{\delta}, n_{\delta}]$ is called the cycle type of (f, σ) and characterizes the conjugacy class \mathcal{C} of (f, σ) in H . The cycle type of (f, σ) in H and the cycle type of (f, σ) in $S(\Gamma \times \Omega)$ are related as follows: Let $\Gamma = \prod_{\kappa=1}^{k_{\delta}} \Gamma_{\delta, \kappa}$ be the orbit decomposition of $\langle p_{i_{\delta}}(f, \sigma) \rangle$. The subsets $\Gamma_{\delta, \kappa} \times \Omega_{\delta}$ for $1 \leq \kappa \leq k_{\delta}$ are the orbits of $(f_{\delta}, \sigma_{\delta})$ on $\Gamma \times \Omega_{\delta}^{\sigma}$. Thus the cycle type of (f, σ) in $S(\Gamma \times \Omega)$ is the partition

with parts $|\Gamma_{\delta,\kappa}||\Omega_\delta|$ for $1 \leq \delta \leq d$ and $1 \leq \kappa \leq k_\delta$. A conjugacy class \mathcal{C} of H is a splitting class if $\mathcal{C} \subseteq H^+$ and $\mathcal{C} = \mathcal{C}_+ \cup \mathcal{C}_-$ is the union of two conjugacy classes of H^+ . The following result is [6, §4.2].

Lemma (4E). *A conjugacy class \mathcal{C} of H is a splitting class if and only if its cycle type satisfies the following conditions:*

- (i) *Cycles $[a, k]$ occur only for odd k , and such cycles have distinct lengths k .*
- (ii) *Cycles $[b^i, j]$ occur only for odd i and even j .*

A splitting class \mathcal{C} thus has cycle type

$$(4.2) \quad \prod_{\delta=1}^d [a, k_\delta] \prod_{i=1}^{p^*-1} \prod_j [b^{2i-1}, 2j]^{w_{ij}},$$

where the k_δ are distinct odd integers. The cycle type of a splitting \mathcal{C} is p -regular if no cycle of type $[a, k]$ occurs, p -singular if no cycle of type $[b^{2i-1}, 2j]$ occurs. A splitting class \mathcal{C} consists of p -singular elements if and only if $d > 0$. If $(f, \sigma) \in \mathcal{C}$, then

$$C_{Y\Omega}(f, \sigma) \simeq \langle a \rangle^d \times \prod_{i,j} \langle b \rangle^{w_{ij}}.$$

In particular,

$$(4.3) \quad |C_H(f, \sigma)|_p = p^d.$$

We define a bijection between splitting mappings Λ and splitting classes \mathcal{C} of H . If Λ is a splitting mapping, let \mathcal{C}_Λ be the splitting class of the following cycle type (4.2): d is the number of diagonal nodes of λ_{p^*} , k_δ is the hook length $\eta_{\delta\delta}(\lambda_{p^*})$, and w_{ij} is the number of parts of length j in λ_i . Conversely, if \mathcal{C} is a splitting class of cycle type (4.2), let $\Lambda_{\mathcal{C}}$ be the following p -tuple $(\lambda_1, \lambda_2, \dots, \lambda_p)$: λ_i for $i < p^*$ is the partition having a part j for each cycle of type $[b^{2i-1}, 2j]$, λ_{p^*} is the unique self-dual partition such that $\eta_{\delta\delta}(\lambda_{p^*}) = k_\delta$ for $1 \leq \delta \leq d$, and λ_i for $i > p^*$ is the partition λ_{p+1-i}^* . Then $(\Lambda_{\mathcal{C}})^* = \Lambda_{\mathcal{C}}$ and $\Lambda_{\mathcal{C}}$ is a splitting mapping.

If Λ is a splitting mapping of H , let Λ' and Λ'' be the p -tuples gotten from Λ by replacing λ_{p^*} by \emptyset in the one case and by replacing λ_γ for $\gamma \neq p^*$ by \emptyset in the other case. We call Λ' and Λ'' the p -regular and the p -singular parts of Λ . The p -tuple with all components \emptyset will be called the trivial p -tuple. Let $\Omega = \Omega' \cup \Omega''$, where

$$\Omega' = \prod_{\gamma \neq p^*} \Omega_\gamma^\Lambda, \quad \Omega'' = \Omega_{p^*}^\Lambda,$$

and $|\Omega_\gamma^\Lambda| = |\lambda_\gamma|$ for all γ . Let $w' = |\Omega'|$, $w'' = |\Omega''|$, $H' = Y \wr S(\Omega')$, $H'' = Y \wr S(\Omega'')$. We may regard a non-trivial Λ' or Λ'' as a splitting mapping parametrizing a splitting character $\psi_{\Lambda'}$ or $\psi_{\Lambda''}$ of H' or H'' , since $(\Lambda')^* = \Lambda'$ and $(\Lambda'')^* = \Lambda''$. In that case let $(H')^+ = H' \cap A(\Gamma \times \Omega')$ and $(H'')^+ = H'' \cap A(\Gamma \times \Omega'')$, and let

$$\text{Res}_{(H')^+}^{H'}(\psi_{\Lambda'}) = \psi_{\Lambda',+} + \psi_{\Lambda',-} \quad \text{and} \quad \text{Res}_{(H'')^+}^{H''}(\psi_{\Lambda''}) = \psi_{\Lambda'',+} + \psi_{\Lambda'',-}.$$

Suppose \mathcal{C}_Λ has cycle type (4.2). If Λ' is trivial, then $H' = (H')^+ = 1$ and we define $\mathcal{C}_{\Lambda'}$ as the identity class of H' . If Λ' is non-trivial, then $\mathcal{C}_{\Lambda'} = \mathcal{C}_{\Lambda',+} \cup \mathcal{C}_{\Lambda',-}$ is a splitting class of H' of cycle type $\prod_{i,j} [b^{2i-1}, 2j]^{w_{ij}}$. If Λ'' is trivial, then $H'' = (H'')^+ = 1$ and we define $\mathcal{C}_{\Lambda''}$ as the identity class of H'' . If Λ'' is non-trivial, then $\mathcal{C}_{\Lambda''} = \mathcal{C}_{\Lambda'',+} \cup \mathcal{C}_{\Lambda'',-}$ is a splitting class of H'' of cycle type $\prod_\delta [a, k_\delta]$. Moreover, the irreducible character $\omega_{\lambda_{p^*}}$ of $S(\Omega'')$ splits as $\omega_{\lambda_{p^*,+}} + \omega_{\lambda_{p^*,-}}$ on $A(\Omega'')$.

Proposition (4F). *Suppose Λ is a splitting mapping of H such that Λ equals its p -singular part Λ'' . Let $(f, \sigma) \in H^+$. Then $(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \neq 0$ if and only if $(f, \sigma) \in \mathcal{C}_\Lambda$. Moreover, $\psi_{\Lambda,\pm}$ and $\mathcal{C}_{\Lambda,\pm}$ may be labeled so that*

$$(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) = \pm(\sqrt{\epsilon_p p})^d \sqrt{\epsilon_{\lambda_{p^*}} \eta(\lambda_{p^*})} \quad \text{for } (f, \sigma) \in \mathcal{C}_{\Lambda,\pm},$$

where d is the number of diagonal nodes in λ_{p^*} .

Proof. For simplicity of notation we write ξ for ξ_{p^*} and λ for λ_{p^*} . By hypothesis $H_\Lambda = H$ and $\psi_\Lambda = E(\xi_\Lambda)\omega_\Lambda$. Suppose $w = 1$. Then $H = Y$, $H^+ = Y^+$, $\psi_\Lambda = \xi$, and $\mathcal{C}_\Lambda = a^Y$. So (4F) holds if we choose $\psi_{\Lambda,\pm} = \xi_\pm$ and $\mathcal{C}_{\Lambda,\pm} = (a_\pm)^{Y^+}$. We may suppose $w > 1$. The irreducible character $\theta_\Lambda = (\xi_+)^{\Omega}$ of $(Y^+)^{\Omega}$ is stable under $Y^+ \wr S(\Omega)$. Let $E(\theta_\Lambda)$ be its James-Kerber extension to $Y^+ \wr S(\Omega)$. Since $\xi = \text{Ind}_{Y^+}^Y(\xi_+)$, it follows by (4C) that

$$E(\xi_\Lambda) = \text{Ind}_{Y^+ \wr S(\Omega)}^H(E(\theta_\Lambda)).$$

Frobenius reciprocity and Mackey decomposition give

$$\begin{aligned} \text{Res}_{H^+}^H(\psi_\Lambda) &= \text{Res}_{H^+}^H(E(\xi_\Lambda)\omega_\Lambda) \\ &= \text{Res}_{H^+}^H(\text{Ind}_{Y^+ \wr S(\Omega)}^H(E(\theta_\Lambda))) \text{Res}_{H^+}^H(\omega_\Lambda) \\ &= \text{Ind}_{Y^+ \wr A(\Omega)}^{H^+}(\text{Res}_{Y^+ \wr A(\Omega)}^{Y^+ \wr S(\Omega)}(E(\theta_\Lambda))) \text{Res}_{H^+}^H(\omega_\Lambda). \end{aligned}$$

Let $(Y^\Omega)^+ = Y^\Omega \cap G^+$ and let $\Theta_{\Lambda,+} = \text{Ind}_{Y^+ \wr A(\Omega)}^{(Y^\Omega)^+ \wr A(\Omega)}(\text{Res}_{Y^+ \wr A(\Omega)}^{Y^+ \wr S(\Omega)}(E(\theta_\Lambda)))$. Then

$$\begin{aligned} \text{Res}_{H^+}^H(\psi_\Lambda) &= \text{Ind}_{(Y^\Omega)^+ \wr A(\Omega)}^{H^+}(\Theta_{\Lambda,+}) \text{Res}_{H^+}^H(\omega_\Lambda) \\ &= \text{Ind}_{(Y^\Omega)^+ \wr A(\Omega)}^{H^+}(\Theta_{\Lambda,+} \text{Res}_{(Y^\Omega)^+ \wr A(\Omega)}^H(\omega_\Lambda)) \\ &= \text{Ind}_{(Y^\Omega)^+ \wr A(\Omega)}^{H^+}(\Theta_{\Lambda,+}(\omega_{\Lambda,+} + \omega_{\Lambda,-})). \end{aligned}$$

Since $\text{Res}_{H^+}^H(\psi_\Lambda) = \psi_{\Lambda,+} + \psi_{\Lambda,-}$, it follows that $\Theta_{\Lambda,+}$ is an irreducible character of $(Y^\Omega)^+ \wr A(\Omega)$. We label $\psi_{\Lambda,\pm}$ so that

$$(4.5) \quad \begin{aligned} \psi_{\Lambda,+} &= \text{Ind}_{(Y^\Omega)^+ \wr A(\Omega)}^{H^+}(\Theta_{\Lambda,+}\omega_{\Lambda,+}), \\ \psi_{\Lambda,-} &= \text{Ind}_{(Y^\Omega)^+ \wr A(\Omega)}^{H^+}(\Theta_{\Lambda,+}\omega_{\Lambda,-}). \end{aligned}$$

Let $H^+ = \langle (Y^\Omega)^+ \wr A(\Omega), s \rangle$, where $s = (g, \tau)$ with $g \notin (Y^\Omega)^+$ and $\tau \notin A(\Omega)$, and set $\Theta_{\Lambda,-} = (\Theta_{\Lambda,+})^s$. Then

$$\begin{aligned} \text{Res}_{(Y^+)^{\Omega}}^{(Y^\Omega)^+ \wr A(\Omega)}(\Theta_{\Lambda,+}) &= \sum_{\substack{(\epsilon_1, \epsilon_2, \dots, \epsilon_w) \\ \epsilon_1 \epsilon_2 \dots \epsilon_w = +}} \xi_{\epsilon_1} \times \xi_{\epsilon_2} \times \dots \times \xi_{\epsilon_w}, \\ \text{Res}_{(Y^+)^{\Omega}}^{(Y^\Omega)^+ \wr A(\Omega)}(\Theta_{\Lambda,-}) &= \sum_{\substack{(\epsilon_1, \epsilon_2, \dots, \epsilon_w) \\ \epsilon_1 \epsilon_2 \dots \epsilon_w = -}} \xi_{\epsilon_1} \times \xi_{\epsilon_2} \times \dots \times \xi_{\epsilon_w}. \end{aligned}$$

In particular, $\Theta_{\Lambda,-} \neq \Theta_{\Lambda,+}$. By (4.5)

$$(4.6) \quad \psi_{\Lambda,+} - \psi_{\Lambda,-} = \begin{cases} (\Theta_{\Lambda,+} - \Theta_{\Lambda,-})(\omega_{\Lambda,+} - \omega_{\Lambda,-}) & \text{on } (Y^\Omega)^+ \wr A(\Omega), \\ 0 & \text{off } (Y^\Omega)^+ \wr A(\Omega). \end{cases}$$

Suppose $(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \neq 0$. Then $(f, \sigma) \in (Y^\Omega)^+ A(\Omega)$ by (4.6), and

$$(\Theta_{\Lambda,+} - \Theta_{\Lambda,-})(f, \sigma) \neq 0, \quad (\omega_{\Lambda,+} - \omega_{\Lambda,-})(\sigma) \neq 0.$$

(4A) implies that σ is in the class $\mathcal{C}_{\eta(\lambda)}$ of $S(\Omega)$ and

$$(\omega_{\Lambda,+} - \omega_{\Lambda,-})(\sigma) = \pm \sqrt{\epsilon_\lambda \eta(\lambda)}.$$

Since (f, σ) is in a splitting class of H , (4E) then implies that $(f, \sigma) \in \mathcal{C}_\Lambda$. We may suppose $p_{i_\delta}((f, \sigma)) = a$ for all δ . The proof of (4C) then gives

$$\Theta_{\Lambda,+}((f, \sigma)) = \sum_{h \in (Y^\Omega)^+ / (Y^+)^{\Omega}} E(\theta_\Lambda)(h(f, \sigma)) = \sum_{\substack{(\epsilon_1, \epsilon_2, \dots, \epsilon_d) \\ \epsilon_1 \epsilon_2 \dots \epsilon_d = +}} \prod_{\delta=1}^d \xi_{\epsilon_\delta}(a).$$

Since $\Theta_{\Lambda,+}$ is $(Y^\Omega)^+ S(\Omega)$ -invariant and $\Theta_{\Lambda,-} = (\Theta_{\Lambda,+})^s$,

$$\Theta_{\Lambda,-}((f, \sigma)) = \sum_{\substack{(\epsilon_1, \epsilon_2, \dots, \epsilon_d) \\ \epsilon_1 \epsilon_2 \dots \epsilon_d = -}} \prod_{\delta=1}^d \xi_{\epsilon_\delta}(a).$$


Thus

$$(\Theta_{\Lambda,+} - \Theta_{\Lambda,-})((f, \sigma)) = (\xi_+(a) - \xi_-(a))^d = (\sqrt{\epsilon_p p})^d.$$

We then label $\psi_{\Lambda,\pm}$ and $\mathcal{C}_{\Lambda,\pm}$ so that (4F) holds. □

Proposition (4G). *Suppose Λ is a splitting mapping of H such that Λ equals its p -regular part Λ' . Let $(f, \sigma) \in H^+$. Then $(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \neq 0$ only if (f, σ) is in a p -regular splitting class of H .*

Proof. Let $H_\Lambda^+ = H_\Lambda \cap H^+$ and $\Theta_\Lambda = \text{Res}_{H_\Lambda^+}^{H_\Lambda} (E(\xi_\Lambda)\omega_\Lambda)$. Then Θ_Λ is irreducible since $(E(\xi_\Lambda)\omega_\Lambda)(h, 1) \neq 0$ for any element h in $Y^\Omega - (Y^\Omega)^+$ taking value b on some $i \in \Omega$ and value 1 on $\Omega - \{i\}$. We fix such an h . Let τ be an involution in $S(\Omega)$ interchanging Ω_γ^Λ and $\Omega_{p+1-\gamma}^\Lambda$ for $1 \leq \gamma < p^*$. Let $t = (g, \tau)$, where $g = 1$ if $\tau \in A(\Omega)$ and $g = h$ if $\tau \notin A(\Omega)$. Then $t \in N_{H^+}(H_\Lambda^+)$ and $T_\Lambda = \langle H_\Lambda^+, t \rangle$ is an extension of H_Λ^+ of order 2.



We claim T_Λ stabilizes Θ_Λ . Indeed, let $(f, \sigma) \in H_\Lambda^+$, and write $(f, \sigma) = \prod_{\gamma=1}^p (f_\gamma, \sigma_\gamma)$, where $f_\gamma \in Y^{\Omega_\gamma^\Lambda}$, $\sigma_\gamma \in S(\Omega_\gamma^\Lambda)$. Then

$$\begin{aligned}
 \omega_\Lambda^t(\sigma) &= \prod_{\gamma=1}^p \omega_{\lambda_\gamma}(t\sigma_\gamma) \\
 &= \prod_{\gamma=1}^p \omega_{\lambda_\gamma^*}(\sigma_\gamma) \\
 &= \prod_{\gamma=1}^p \omega_{\lambda_\gamma}(\sigma_\gamma) \operatorname{sgn}_{S(\Omega_\gamma^\Lambda)}(\sigma_\gamma) \\
 &= (\omega_\Lambda \operatorname{sgn}_{S(\Omega)})(\sigma).
 \end{aligned}
 \tag{4.7}$$

On the other hand, if $(f_\gamma, \sigma_\gamma)$ decomposes into cycles $\prod_{\delta \in \Delta_\gamma} (f_\delta, \sigma_\delta)$, then

$$\begin{aligned}
 E(\xi_\Lambda)^t(f, \sigma) &= E(\xi_\Lambda)^{(g,1)}((1,\tau)(f, \sigma)) \\
 &= E(\xi_\Lambda)^{(1,\tau)}(f, \sigma) \\
 &= \prod_{\gamma} \prod_{\delta \in \Delta_\gamma} \xi_{p+1-\gamma}(p_{\tau(i_\delta)}((1,\tau)(f, \sigma))) \\
 &= \prod_{\gamma} \prod_{\delta \in \Delta_\gamma} \xi_\gamma^*(p_{i_\delta}(f, \sigma)) \\
 &= \prod_{\gamma} \prod_{\delta \in \Delta_\gamma} (\xi_\gamma(p_{i_\delta}(f, \sigma)) \operatorname{sgn}_Y(p_{i_\delta}(f, \sigma))) \\
 &= E(\xi_\Lambda)(f, \sigma) \operatorname{sgn}_{S(\Omega)}(\sigma),
 \end{aligned}
 \tag{4.8}$$

since $\prod_{\gamma} \prod_{i_\delta \in \Delta_\gamma} \operatorname{sgn}_Y(p_{i_\delta}(f, \sigma)) = \prod_{i \in \Omega} \operatorname{sgn}_Y(f(i)) = \operatorname{sgn}_{S(\Omega)}(\sigma)$. So T_Λ stabilizes Θ_Λ by (4.7) and (4.8).

Let Θ_Λ^+ and Θ_Λ^- be the two extensions of Θ_Λ to T_Λ . By Mackey decomposition we may label $\psi_{\Lambda, \pm}$ so that

$$\psi_{\Lambda,+} = \operatorname{Ind}_{T_\Lambda}^{H^+}(\Theta_\Lambda^+), \quad \psi_{\Lambda,-} = \operatorname{Ind}_{T_\Lambda}^{H^+}(\Theta_\Lambda^-).
 \tag{4.9}$$

We note for later use that any odd element s of Y^Ω normalizes T_Λ and interchanges Θ_Λ^+ and Θ_Λ^- , since $\psi_{\Lambda,+}^s = \psi_{\Lambda,-}$.

Suppose $(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \neq 0$. Replacing (f, σ) by a conjugate, we may suppose by (4.9) that $(f, \sigma) \in T_\Lambda$ and $(\Theta_\Lambda^+ - \Theta_\Lambda^-)(f, \sigma) \neq 0$. Thus $(f, \sigma) \notin H_\Lambda^+$. Elements of $T_\Lambda - H_\Lambda^+$ have the form (h, ρ) , where ρ interchanges $\prod_{\gamma < p^*} \Omega_\gamma^\Lambda$ and $\prod_{\gamma > p^*} \Omega_\gamma^\Lambda$. In particular, ρ contains no cycles of odd length. Thus (f, σ) is contained in a p -regular splitting class of H . This completes the proof of (4G). \square

Proposition (4H). *Suppose Λ is a splitting mapping of H , Λ' and Λ'' are the p -regular and p -singular parts of Λ , and H' and H'' are the subgroups of H associated to Λ' and Λ'' .*

- (i) *If $(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \neq 0$, then (f, σ) is H^+ -conjugate to $(f', \sigma')(f'', \sigma'')$, where (f', σ') is in a p -regular splitting class of H' and (f'', σ'') is in the p -singular splitting class $\mathcal{C}_{\Lambda''}$ of H'' .*

(ii) Given labelings of $\psi_{\Lambda',\pm}$ and $\psi_{\Lambda'',\pm}$, there exists a labeling of $\psi_{\Lambda,\pm}$ such that

$$(4.10) \quad \begin{aligned} (\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) &= (\psi_{\Lambda',+} - \psi_{\Lambda',-})(f', \sigma') \\ &\quad \times (\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(f'', \sigma'') \end{aligned}$$

for all $(f, \sigma) = (f', \sigma')(f'', \sigma'')$ with (f', σ') a p -regular element of $(H')^+$ and (f'', σ'') an arbitrary element of $(H'')^+$.

Proof. (4H) follows trivially from (4F) or (4G) if Λ' or Λ'' is trivial. So we may suppose Λ' and Λ'' are non-trivial. Let ξ_Λ be the irreducible character $\prod_\gamma (\xi_\gamma)^{\Omega_\Lambda}$ of Y^Ω . Let $\xi_{\Lambda'}$ and $\xi_{\Lambda''}$ be the irreducible characters of $Y^{\Omega'}$ and $Y^{\Omega''}$ defined by

$$\xi_{\Lambda'} = \prod_{\gamma \neq p^*} (\xi_\gamma)^{\Omega_\Lambda}, \quad \xi_{\Lambda''} = (\xi_{p^*})^{\Omega_\Lambda}.$$

The stabilizers of $\xi_{\Lambda'}$ in H' and $\xi_{\Lambda''}$ in H'' are $H_{\Lambda'} = Y^{\Omega'} S(\Omega^{\Lambda'})$ and $H_{\Lambda''} = Y^{\Omega''} S(\Omega^{\Lambda''})$, where $S(\Omega^{\Lambda'})$ and $S(\Omega^{\Lambda''})$ are the components of $S(\Omega^\Lambda)$ on Ω' and Ω'' . Let $E(\xi_{\Lambda'})$ and $E(\xi_{\Lambda''})$ be the James-Kerber extensions of $\xi_{\Lambda'}$ and $\xi_{\Lambda''}$ to $H_{\Lambda'}$ and $H_{\Lambda''}$, and let $\omega_{\Lambda'}$ and $\omega_{\Lambda''}$ be the irreducible characters of $S(\Omega^{\Lambda'})$ and $S(\Omega^{\Lambda''})$ parametrized by Λ' and Λ'' . Then

$$\psi_{\Lambda'} = \text{Ind}_{H_{\Lambda'}}^{H'}(E(\xi_{\Lambda'})\omega_{\Lambda'}), \quad \psi_{\Lambda''} = E(\xi_{\Lambda''})\omega_{\Lambda''},$$

and

$$E(\xi_\Lambda)\omega_\Lambda = E(\xi_{\Lambda'})\omega_{\Lambda'} \times E(\xi_{\Lambda''})\omega_{\Lambda''}.$$

Let

$$\Theta_\Lambda = \text{Res}_{H_\Lambda^+}^{H_\Lambda} (E(\xi_\Lambda)\omega_\Lambda), \quad \Theta_{\Lambda'} = \text{Res}_{H_{\Lambda'}^+}^{H_{\Lambda'}} (E(\xi_{\Lambda'})\omega_{\Lambda'}).$$


Then Θ_Λ and $\Theta_{\Lambda'}$ are irreducible since $(E(\xi_\Lambda)\omega_\Lambda)(f', 1)$ and $(E(\xi_{\Lambda'})\omega_{\Lambda'})(f', 1)$ are non-zero for any element f' in $Y^{\Omega'} - (Y^{\Omega'})^+$ taking value b on some $i \in \Omega'$ and value 1 on $\Omega' - \{i\}$. We fix such an f' . On the other hand,

$$\text{Res}_{(H'')^+}^{H''}(\psi_{\Lambda''}) = \psi_{\Lambda'',+} + \psi_{\Lambda'',-}, \quad \text{Res}_{H_{\Lambda'}^+ \times H_{\Lambda''}^+}^{H_\Lambda^+}(\Theta_\Lambda) = \Theta_{\Lambda,+} + \Theta_{\Lambda,-}$$

are reducible. Given a labeling of $\psi_{\Lambda'',\pm}$, we label $\Theta_{\Lambda,\pm}$ so that $\Theta_{\Lambda,+} = \Theta_{\Lambda'} \times \psi_{\Lambda'',+}$ and $\Theta_{\Lambda,-} = \Theta_{\Lambda'} \times \psi_{\Lambda'',-}$.

Let $s = (h, 1)$ in Y^Ω be such that h takes the value b once on Ω' and once on Ω'' , and the value 1 everywhere else. Then $H_\Lambda^+ = \langle H_{\Lambda'}^+ \times H_{\Lambda''}^+, s \rangle$. Let $\tau \in S(\Omega')$ be an involution interchanging Ω_γ^Λ and $\Omega_{p+1-\gamma}^\Lambda$ for $1 \leq \gamma < p^*$. Let $t = (g, \tau)$, where g is the identity of $Y^{\Omega'}$ if $\tau \in A(\Omega')$, and $g = f'$ if $\tau \notin A(\Omega')$. Then $t \in N_{(H')^+}(H_{\Lambda'}^+)$, $T_{\Lambda'} = \langle H_{\Lambda'}^+, t \rangle$ is an extension of $H_{\Lambda'}^+$ of order 2, and $[s, t] \in H_{\Lambda'}^+ \times H_{\Lambda''}^+$. In

particular, if $T_\Lambda = \langle H_\Lambda^+, t \rangle$, then $T_\Lambda = \langle H_{\Lambda'}^+ \times H_{\Lambda''}^+, s, t \rangle = \langle T_{\Lambda'} \times H_{\Lambda''}^+, s \rangle$.



$T_{\Lambda'}$ stabilizes $\Theta_{\Lambda'}$ as in the proof of (4G). In particular, $\Theta_{\Lambda'}$ has two extensions $\Theta_{\Lambda'}^+$ and $\Theta_{\Lambda'}^-$ to $T_{\Lambda'}$, $\text{Ind}_{H_{\Lambda'}^+}^{T_{\Lambda'}}(\Theta_{\Lambda'}) = \Theta_{\Lambda'}^+ + \Theta_{\Lambda'}^-$, and $\text{Ind}_{T_{\Lambda'}}^{(H')^+}(\Theta_{\Lambda'}^+ + \Theta_{\Lambda'}^-) = \psi_{\Lambda',+} + \psi_{\Lambda',-}$. Given a labeling of $\psi_{\Lambda',\pm}$, we label $\Theta_{\Lambda'}^\pm$ so that

$$\psi_{\Lambda',+} = \text{Ind}_{T_{\Lambda'}}^{(H')^+}(\Theta_{\Lambda'}^+), \quad \psi_{\Lambda',-} = \text{Ind}_{T_{\Lambda'}}^{(H')^+}(\Theta_{\Lambda'}^-).$$

Then $T_{\Lambda'} \times H_{\Lambda''}^+$ stabilizes Θ_Λ since $\Theta_\Lambda = \text{Ind}_{H_{\Lambda'}^+ \times H_{\Lambda''}^+}^{H_\Lambda^+}(\Theta_{\Lambda',+})$, and T_Λ stabilizes Θ_Λ since $s \in H_\Lambda^+$. Now

$$\begin{aligned} (\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',+})^s &= \Theta_{\Lambda'}^- \times \psi_{\Lambda'',-}, \\ (\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',-})^s &= \Theta_{\Lambda'}^- \times \psi_{\Lambda'',+}, \end{aligned}$$

since any odd element of $Y^{\Omega'}$ normalizes $T_{\Lambda'}$ and interchanges $\Theta_{\Lambda'}^+$ and $\Theta_{\Lambda'}^-$, and any odd element of $Y^{\Omega''}$ interchanges $\psi_{\Lambda'',+}$ and $\psi_{\Lambda'',-}$. Thus

$$\begin{aligned} \Xi_{\Lambda,+} &= \text{Ind}_{T_{\Lambda'} \times H_{\Lambda''}^+}^{T_\Lambda}(\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',+}), \\ \Xi_{\Lambda,-} &= \text{Ind}_{T_{\Lambda'} \times H_{\Lambda''}^+}^{T_\Lambda}(\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',-}) \end{aligned}$$

are distinct irreducible characters, and

$$(4.11) \quad \Xi_{\Lambda,+} - \Xi_{\Lambda,-} = \begin{cases} (\Theta_{\Lambda'}^+ - \Theta_{\Lambda'}^-) \times (\psi_{\Lambda'',+} - \psi_{\Lambda'',-}) & \text{on } T_{\Lambda'} \times H_{\Lambda''}^+, \\ 0 & \text{on } T_\Lambda - (T_{\Lambda'} \times H_{\Lambda''}^+). \end{cases}$$

We may label $\psi_{\Lambda,\pm}$ so that

$$(4.12) \quad \begin{aligned} \psi_{\Lambda,+} &= \text{Ind}_{T_\Lambda}^{H^+}(\Xi_{\Lambda,+}) = \text{Ind}_{T_{\Lambda'} \times H_{\Lambda''}^+}^{H^+}(\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',+}), \\ \psi_{\Lambda,-} &= \text{Ind}_{T_\Lambda}^{H^+}(\Xi_{\Lambda,-}) = \text{Ind}_{T_{\Lambda'} \times H_{\Lambda''}^+}^{H^+}(\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',-}). \end{aligned}$$

Indeed, Mackey decomposition implies that

$$\text{Res}_{H^+}^H(\psi_\Lambda) = \text{Ind}_{H_\Lambda^+}^{H^+}(\Theta_\Lambda) = \text{Ind}_{T_\Lambda}^{H^+}(\text{Ind}_{H_\Lambda^+}^{T_\Lambda}(\Theta_\Lambda)),$$

since $\psi_\Lambda = \text{Ind}_{H_\Lambda}^H (E(\xi_\Lambda)\omega_\Lambda)$. But

$$\begin{aligned} \text{Ind}_{H_\Lambda}^{T_\Lambda}(\Theta_\Lambda) &= \text{Ind}_{H_\Lambda}^{T_\Lambda}(\text{Ind}_{H_{\Lambda'}^+ \times H_{\Lambda''}^+}^{H_\Lambda^+}(\Theta_{\Lambda'} \times \psi_{\Lambda'',+})) \\ &= \text{Ind}_{T_{\Lambda'} \times H_{\Lambda''}^+}^{T_\Lambda}(\text{Ind}_{H_{\Lambda'}^+ \times H_{\Lambda''}^+}^{T_{\Lambda'} \times H_{\Lambda''}^+}(\Theta_{\Lambda'} \times \psi_{\Lambda'',+})) \\ &= \text{Ind}_{T_{\Lambda'} \times H_{\Lambda''}^+}^{T_\Lambda}((\Theta_{\Lambda'}^+ + \Theta_{\Lambda'}^-) \times \psi_{\Lambda'',+}) \\ &= \bar{\Xi}_{\Lambda,+} + \bar{\Xi}_{\Lambda,-}. \end{aligned}$$

Thus (4.12) holds since $\text{Res}_{H^+}^H(\psi_\Lambda) = \psi_{\Lambda,+} + \psi_{\Lambda,-}$.

Suppose $(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \neq 0$. By (4.11) and (4.12) there exist (f', σ') in $T_{\Lambda'} - H_{\Lambda'}^+$ and (f'', σ'') in $H_{\Lambda''}^+$ such that (f, σ) is H^+ -conjugate to $(f', \sigma')(f'', \sigma'')$ and

$$(4.13) \quad (\Theta_{\Lambda'}^+ - \Theta_{\Lambda'}^-)(f', \sigma') \neq 0, \quad (\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(f'', \sigma'') \neq 0.$$

(4H),(i) now follows by (4F) and the last paragraph of the proof of (4G).

Let $(f, \sigma) = (f', \sigma')(f'', \sigma'') \in H$ with $(f', \sigma') \in (H')_{p'}^+$ and $(f'', \sigma'') \in (H'')^+$. If (f', σ') or (f'', σ'') is not in a splitting class of H' or H'' , then (f, σ) is not in a splitting class of H . Thus both sides of (4.10) are zero and (4.10) holds. So we may suppose (f', σ') and (f'', σ'') are in splitting classes of H' and H'' . Suppose (f'', σ'') is not in $\mathcal{C}_{\Lambda''}$. Then the right-hand side of (4.10) is zero by (4F). Suppose the left-hand side of (4.10) is non-zero. By (4H) (i) there exist u' in a p -regular splitting class of H' , u'' in a p -singular splitting class of H'' , and $h \in H^+$ such that $((f', \sigma')(f'', \sigma''))^h = u'u''$. Moreover, $u'' \in \mathcal{C}_{\Lambda''}$. In particular, u'' is a product of cycles of p -singular length as an element of $S(\Gamma \times \Omega'')$. A comparison of the p -regular and p -singular cycle types of these elements as elements of $S(\Gamma \times \Omega)$ shows that $(f', \sigma')^h = u'$, $(f'', \sigma'')^h = u''$. Since u'' has no fixed-points on Ω'' , neither does (f'', σ'') . So h stabilizes Ω' and Ω'' , and $h \in H'H''$. Thus (f', σ') is conjugate to u' in H' , (f'', σ'') is conjugate to u'' in H'' , and $(f'', \sigma'') \in \mathcal{C}_{\Lambda''}$. This is a contradiction. So we may suppose (f', σ') is in a p -regular splitting class of H' and (f'', σ'') is in $\mathcal{C}_{\Lambda''}$.

Suppose there exists (g, τ) in H^+ such that $(g, \tau)(f, \sigma) = (h', \rho')(h'', \rho'')$, where $(h', \rho') \in T_{\Lambda'}$, $(h'', \rho'') \in H_{\Lambda''}^+$, and

$$(4.14) \quad (\Theta_{\Lambda'}^+ - \Theta_{\Lambda'}^-)(h', \rho') \neq 0, \quad (\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(h'', \rho'') \neq 0.$$

If $\tau(i_\delta) \in \Omega''$ for some $i_\delta \in \Omega'$, then $p_{\tau(i_\delta)}(h'', \rho'') \in a^Y$ by (4F). But

$$\begin{aligned} p_{\tau(i_\delta)}(h'', \rho'') &= p_{\tau(i_\delta)}((g, \tau)(f, \sigma)) \\ &= g(\tau(i_\delta))p_{i_\delta}(f, \sigma)g(\tau(i_\delta))^{-1} \\ &= g(\tau(i_\delta))p_{i_\delta}(f', \sigma')g(\tau(i_\delta))^{-1} \end{aligned}$$

and $p_{i_\delta}(f', \sigma') \in (b^i)^Y$ for odd i by hypothesis. This is impossible. Thus τ stabilizes Ω' and Ω'' and $(g, \tau) = (g', \tau')(g'', \tau'')$, where $(g', \tau') \in H'$ and $(g'', \tau'') \in H''$. Then

$$\begin{aligned} &(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \\ &= \text{Ind}_{T_{\Lambda'} \times H_{\Lambda''}^+}^{H^+}((\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',+}) - (\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',-}))(f, \sigma) \\ &= \text{Ind}_{T_{\Lambda'} \times H_{\Lambda''}^+}^{(H' \times H'')^+}((\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',+}) - (\Theta_{\Lambda'}^+ \times \psi_{\Lambda'',-}))(f, \sigma). \end{aligned}$$

Let \mathcal{D} be a transversal for $T_{\Lambda'}$ in $(H')^+$. Then $\mathcal{D} \cup \mathcal{D}s$ is a transversal for $T_{\Lambda'} \times H_{\Lambda''}^+$ in $(H' \times H'')^+$, where $s = (h, 1)$ is the element introduced earlier in the proof. It follows that

$$\begin{aligned} &(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \\ &= \text{Ind}_{T_{\Lambda'}}^{(H')^+} (\Theta_{\Lambda'}^+)(f', \sigma') \times (\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(f'', \sigma'') \\ &\quad + \text{Ind}_{T_{\Lambda'}}^{(H')^+} (\Theta_{\Lambda'}^-)(f', \sigma') \times (\psi_{\Lambda'',-} - \psi_{\Lambda'',+})(f'', \sigma'') \\ &= (\psi_{\Lambda',+} - \psi_{\Lambda',-})(f', \sigma') \times (\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(f'', \sigma''), \end{aligned}$$

and (4.10) holds. Suppose no (g, τ) in H^+ satisfies (4.14). Then $(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma)$ is 0 by (4.12). Thus (4.10) holds unless

$$(\psi_{\Lambda',+} - \psi_{\Lambda',-})(f', \sigma') \neq 0, \quad (\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(f'', \sigma'') \neq 0.$$

In this case there exists (g', τ') in $(H')^+$ such that $(g', \tau')(f', \sigma') = (h', \rho')$ is in $T_{\Lambda'}$ and $(\Theta_{\Lambda'}^+ - \Theta_{\Lambda'}^-)(h', \rho') \neq 0$. This is impossible since $(g', \tau')(f'', \sigma'') = (f'', \sigma'')$, so that (4.14) holds. This completes the proof of (4H), (ii). \square

Corollary (4I). *Let Λ be a splitting mapping of H . If $(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \neq 0$, then*

$$(\psi_{\Lambda,+} - \psi_{\Lambda,-})(f, \sigma) \equiv 0 \pmod{\sqrt{|C_H(f, \sigma)|_p}}$$

and $|C_H(f, \sigma)|_p = p^d$, where d is the number of diagonal nodes in λ_{p^*} .

Proof. This follows from (4H), (4F), and (4.3). \square

5. THE ISOTYPY CONJECTURE FOR ALTERNATING GROUPS FOR $p > 2$

We suppose p is odd in this section. Let G and G^+ be the symmetric and alternating groups on the set Π of n elements. Let e and e^+ be blocks of $\mathcal{O}G$ and $\mathcal{O}G^+$ such that e covers e^+ . In particular, e and e^+ have the same defect group D . If D is non-trivial, then there exists χ_λ in $(G, e)^\vee$ such that $\text{Res}_{G^+}^G(\chi_\lambda)$ is irreducible. Indeed, let λ_0 be the p -core parametrizing e , and choose a partition λ of n with p -core λ_0 and a p -quotient Λ such that $\Lambda^* \neq \Lambda$. In particular, e^+ is the unique block of $\mathcal{O}G^+$ covered by e .

Let $H = N_G(D)$ and $H^+ = N_{G^+}(D)$. Let $f = \text{Br}_D^G(e)$ and $f^+ = \text{Br}_D^{G^+}(e^+)$ be the Brauer correspondents of e and e^+ in $\mathcal{O}H$ and $\mathcal{O}H^+$. In particular, f^+ is H -stable since e^+ is G -stable. Moreover, $ff^+ = f$, since $ee^+ = e$ and $\text{Br}_D^G(e^+) = \text{Br}_D^{G^+}(e^+)$. Thus f covers f^+ , and f^+ is the unique block of $\mathcal{O}H^+$ covered by f . We suppose from now on that D is abelian and that $|D| = p^w$ with $1 \leq w < p$. Then e and f are isotypic blocks by (3B). If $\lambda_0^* \neq \lambda_0$, then e^+ and f^+ are isotypic blocks by (1E). So we also suppose from now on that $\lambda_0^* = \lambda_0$. Thus $e^* = e = e^+$, $f^* = f = f^+$, and (G, G^+, e, e^+) and (H, H^+, f, f^+) are +configurations.

We use the notation of §3: $\Pi = \Pi_0 \cup \Pi_1$, where Π_0 and Π_1 are disjoint sets of n_0 and pw elements; $\Pi_1 = \Gamma \times \Omega$, where $\Gamma = \{1, 2, \dots, p\}$ and $|\Omega| = w$; $X = S(\Gamma)$ and $Y = N_X(P)$, where P is a Sylow p -subgroup of X ; and D is the Sylow p -subgroup P^Ω of $S(\Pi_1)$. Let

$$(5.1) \quad \begin{aligned} H_0 &= S(\Pi_0), & H_1 &= Y \wr S(\Omega), \\ H_0^+ &= A(\Pi_0), & H_1^+ &= (Y \wr S(\Omega))^+, \end{aligned}$$

and set $H^{++} = H_0^+ \times H_1^+$. Then $H = H_0 \times H_1$, $H > H^+ \geq H^{++}$, and H^+/H^{++} has order 1 or 2 according as $|\Pi_0| \leq 1$ or $|\Pi_0| > 1$. We have

$$(5.2) \quad f^+ = f = e_{\lambda_0} \times 1_{H_1},$$

where e_{λ_0} is the block of defect 0 of $\mathcal{O}H_0$ parametrized by λ_0 . Let χ_{λ_0} denote the character in $(H_0, e_{\lambda_0})^\vee$. Since λ_0 is self-dual, either $|\lambda_0| \leq 1$ or $|\lambda_0| \geq 3$. It will be convenient to interpret H_0 and H_0^+ as the identity group when $|\lambda_0| \leq 1$ and to omit identity factors in expressions such as (5.2). We fix a maximal e^+ -subpair (D, e_D^+) and a maximal f^+ -subpair (D, f_D^+) as follows: $C_{G^+}(D) = C_{H^+}(D) = H_0^+ \times D$. If $|\Pi_0| \leq 1$, let $e_D^+ = f_D^+$ be the 1-block; if $|\Pi_0| \geq 3$, let $e_D^+ = f_D^+ = e_{\lambda_0,+}$, where $e_{\lambda_0,+}$ is the block of $\mathcal{O}H_0^+$ containing an irreducible constituent $\chi_{\lambda_0,+}$ of $\text{Res}_{H_0^+}^{H_0}(\chi_{\lambda_0})$. In both cases

$$(5.3) \quad N_{G^+}(D, e_D^+) = N_{H^+}(D, f_D^+) = H^{++}.$$

We describe certain subpairs in $\mathbf{Br}_{e^+}(G^+)$ and $\mathbf{Br}_{f^+}(H^+)$. Every e^+ -subpair (Q, u^+) is covered by an e -subpair (Q, u) in the sense that u covers u^+ . Conversely, any subpair (Q, u^+) of G^+ covered by an e -subpair (Q, u) is an e^+ -subpair. Indeed, these claims follow from the relation $\text{Br}_Q^G(e) = \text{Br}_Q^{G^+}(e^+)$. If $|\Pi_0| \leq 1$, then e^+ and f^+ are the 1-blocks of $\mathcal{O}G^+$ and $\mathcal{O}H^+$, and Brauer’s Third Main Theorem implies the subpairs in $\mathbf{Br}_{e^+}(G^+)$ and $\mathbf{Br}_{f^+}(H^+)$ are (Q, e_Q^+) and (Q, f_Q^+) , where e_Q^+ and f_Q^+ are the 1-blocks of $\mathcal{O}G_Q^+$ and $\mathcal{O}H_Q^+$. So we suppose $|\Pi_0| \geq 3$. Let x be a standard non-identity element of D in the sense of §3, and recall that

$$G_x = G_{x,0} \times G_{x,1}, \quad H_x = H_{x,0} \times H_{x,1},$$

where $G_{x,0} = S(\Pi_0 \cup (\Gamma \times \Omega_{x,0}))$, $H_{x,0} = S(\Pi_0) \times (Y \wr S(\Omega_{x,0}))$, and $G_{x,1} = H_{x,1} = P \wr S(\Omega_{x,1})$. Let $H_{x,0} = H_{x,00} \times H_{x,01}$, where $H_{x,00} = S(\Pi_0)$ and $H_{x,01} = Y \wr S(\Omega_{x,0})$. Let

$$e_x = e_{x,0} \times e_{x,1}, \quad f_x = f_{x,00} \times f_{x,01} \times f_{x,1},$$

where $e_{x,0}$ is the block of $\mathcal{O}G_{x,0}$ parametrized by λ_0 , $e_{x,1}$ is the 1-block of $\mathcal{O}G_{x,1}$, $f_{x,00}$ is the block of $\mathcal{O}H_{x,00}$ parametrized by λ_0 , $f_{x,01}$ is the 1-block of $\mathcal{O}H_{x,01}$, and $f_{x,1}$ is the 1-block of $\mathcal{O}H_{x,1}$. Let $f_{x,0}$ be the block $f_{x,00} \times f_{x,01}$ of $\mathcal{O}H_{x,0}$. Let G_x^+ and H_x^+ be the intersections of G_x and H_x with $A(\Pi)$. Similarly, let $G_{x,0}^+$ and $H_{x,0}^+$ be the intersections of $G_{x,0}$ and $H_{x,0}$ with $A(\Pi_0 \cup (\Gamma \times \Omega_{x,0}))$, and let $G_{x,1}^+$ and $H_{x,1}^+$ be the intersections of $G_{x,1}$ and $H_{x,1}$ with $A(\Gamma \times \Omega_{x,1})$. In particular, $G_x > G_x^+$ if $|\Pi_0| > 1$ or $w > 1$. Let $G_x^{++} = G_{x,0}^+ \times G_{x,1}^+$ and $H_x^{++} = H_{x,0}^+ \times H_{x,1}^+$. If $w > 1$, then e_x and f_x cover unique blocks e_x^+ and f_x^+ of $\mathcal{O}G_x^+$ and $\mathcal{O}H_x^+$. Indeed, let $\chi_x = \chi_{x,0} \times \chi_{x,1}$, where $\chi_{x,0} \in (G_{x,0}, e_{x,0})^\vee$, $\chi_{x,1} \in (G_{x,1}, e_{x,1})^\vee$, and $\chi_{x,0}$ or $\chi_{x,1}$ is irreducible on $G_{x,0}^+$ or $G_{x,1}^+$. Such a $\chi_{x,0}$ or $\chi_{x,1}$ exists since $|\Omega_{x,0}| > 0$ or $|\Omega_{x,1}| \geq 2$. So $\chi_x \in (G_x, e_x)^\vee$, χ_x is irreducible on G_x^+ , and e_x covers a unique block e_x^+ of $\mathcal{O}G_x^+$. A similar argument shows that f_x covers a unique block f_x^+ of $\mathcal{O}H_x^+$.

Proposition (5A). *Let x be a non-identity standard element in D . If $|\Pi_0| \leq 1$ or $w > 1$, then $\text{Br}_{\langle x \rangle}^{G^+}(e^+)$ and $\text{Br}_{\langle x \rangle}^{H^+}(f^+)$ are blocks of $\mathcal{O}G_x^+$ and $\mathcal{O}H_x^+$. If $|\Pi_0| \geq 3$ and $w = 1$, then $D = \langle x \rangle$, $G_x^+ = H_x^+ = H_0^+ \times D$, and*

$$\text{Br}_{\langle x \rangle}^{G^+}(e^+) = \text{Br}_{\langle x \rangle}^{H^+}(f^+) = e_{\lambda_0,+} + e_{\lambda_0,-}.$$

Proof. If $|\Pi_0| \leq 1$, then e^+ and f^+ are the principal blocks and (5A) is clear. If $|\Pi_0| \geq 3$ and $w > 1$, then (5A) follows from the preceding paragraph and the fact that e and f are blocks of principal type. If $|\Pi_0| \geq 3$ and $w = 1$, then the result follows from our discussion of the maximal e^+ -subpair (D, e_D^+) and the maximal f^+ -subpair (D, f_D^+) . \square

Proposition (5B). *Let $\mu = (J, \epsilon)$ be the Rouquier isometry in $\mathcal{R}_K((G, e), (H, f))$. Then the following hold:*

- (i) μ is +equivariant with respect to (G^+, H^+) .
- (ii) Every μ -induced isometry $\mu^+ = (J^+, \epsilon^+)$ in $\mathcal{R}_K((G^+, e^+), (H^+, f^+))$ is perfect.

Proof. By definition (3.8) of the Rouquier isometry

$$J(\text{sgn}_G \chi_\lambda) = J(\chi_{\lambda^*}) = \chi_{\lambda_0^*} \times \psi_{\widehat{\Lambda}^*} = \text{sgn}_{H_0} \chi_{\lambda_0} \times \text{sgn}_{H_1} \psi_{\widehat{\Lambda}} = \text{sgn}_H J(\chi_\lambda).$$

Moreover, $\epsilon(\text{sgn}_G \chi_\lambda) = \epsilon(\chi_\lambda)$, since $|\lambda_{\frac{1}{2}(p+1)}| = |\lambda_{\frac{1}{2}(p+1)}^*|$ and $v(\lambda) = v(\lambda^*)$. Here the last equality holds since the dual of a sequence of p -hooks leading from λ to λ_0 is a sequence of p -hooks from λ^* to λ_0^* , and the leg and arm lengths of a p -hook have the same parity.

For (ii) it suffices by (2B) to show that

$$\Phi_\lambda(g, h) = (\chi_{\lambda,+} - \chi_{\lambda,-})(g)(J^+(\chi_{\lambda,+}) - J^+(\chi_{\lambda,-}))(h^{-1})$$

is perfect on $G^+ \times H^+$ for all self-dual partitions λ with p -core λ_0 . Suppose $\Phi_\lambda(g, h) \neq 0$ for $g \in G^+, h \in H^+$. Then g and h are in splitting classes of G and H , and

$$(5.4) \quad (\chi_{\lambda,+} - \chi_{\lambda,-})(g) \neq 0, \quad (J^+(\chi_{\lambda,+}) - J^+(\chi_{\lambda,-}))(h^{-1}) \neq 0.$$

The first inequality and (4A) imply that g has cycle type $\eta(\lambda)$. If d is the number of $\eta_{ii}(\lambda)$ divisible by p , then

$$(\chi_{\lambda,+} - \chi_{\lambda,-})(g) \equiv 0 \pmod{\sqrt{\epsilon_\lambda p^d}}$$

and $|C_G(g)|_p = p^d$. Moreover, g is p -singular if and only if $d > 0$. The second inequality implies that $h \in H^{++}$. Indeed, if $|\Pi_0| \leq 1$, then $h \in H^{++}$ since $H^+ = H^{++}$. If $|\Pi_0| \geq 3$, then χ_{λ_0} and $\psi_{\widehat{\Lambda}}$ are reducible on H_0^+ and H_1^+ , and $J^+(\chi_{\lambda,\pm})$ has support on H^{++} . Thus $h = h_0(f_1, \sigma_1)$, where $h_0 \in H_0^+, (f_1, \sigma_1) \in H_1^+$, and

$$(\chi_{\lambda_0,+} - \chi_{\lambda_0,-})(h_0^{-1}) \neq 0, \quad (\psi_{\widehat{\Lambda},+} - \psi_{\widehat{\Lambda},-})((f_1, \sigma_1)^{-1}) \neq 0.$$

Here $(\chi_{\lambda_0,+} - \chi_{\lambda_0,-})(h_0^{-1})$ is interpreted as 1 if $|\Pi_0| \leq 1$. Since $\chi_{\lambda_0,\pm}$ has defect 0, it follows that h_0 is p -regular and $(\chi_{\lambda_0,+} - \chi_{\lambda_0,-})(h_0)$ is divisible by $|C_{H_0}(h_0)|_p$. On the other hand, (4I) implies that $|C_{H_1}(f_1, \sigma_1)|_p = p^d$ and

$$(\psi_{\widehat{\Lambda},+} - \psi_{\widehat{\Lambda},-})(f_1, \sigma_1) \equiv 0 \pmod{\sqrt{|C_{H_1}(f_1, \sigma_1)|_p}}.$$

Moreover, h is p -singular if and only if $d > 0$. It now follows that Φ_λ is perfect.

For each $j = 1, 2, \dots, w$, let x_j be a standard element of D such that $|\Omega_{x_j,1}| = j$. We claim the w subgroups $\langle x_j \rangle$ form a set of representatives for the $N_{G^+}(D, e_D^+)$ -orbits of non-trivial cyclic subgroups of D . This is trivially true when $w = 1$, so we may suppose $w > 1$. The w subgroups $\langle x_j \rangle$ form a set of representatives for the

$(Y \wr S(\Omega))$ -orbits of non-trivial cyclic subgroups of D . But the $(Y \wr S(\Omega))$ -orbit of $\langle x_j \rangle$ is the $(Y \wr S(\Omega))^+$ -orbit of $\langle x_j \rangle$, since

$$C_{Y \wr S(\Omega)}(x_j) = (Y \wr S(\Omega_{x_j,0})) \times (P \wr S(\Omega_{x_j,1}))$$

contains odd permutations when $w > 1$. Since $(Y \wr S(\Omega))^+ \leq N_{G^+}(D, e_D^+)$, the claim holds. \square

We will use the isotypy (3B) between e and f to define an isometry μ^+ and perfect isometries μ_x^+ for x in $\{x_1, x_2, \dots, x_w\}$ so that the diagrams

$$(5.5) \quad \begin{array}{ccc} \mathcal{R}_K(G^+, e^+) & \xrightarrow{R_{\mu^+}} & \mathcal{R}_K(H^+, f^+) \\ \downarrow d_{G^+}^{x, e_x^+} & & \downarrow d_{H^+}^{x, f_x^+} \\ \text{CF}_{p'}(G_x^+, e_x^+) & \xrightarrow{R_{\mu_x^+, p'}} & \text{CF}_{p'}(H_x^+, f_x^+) \end{array}$$

are commutative. The definitions require a consistent labeling of splitting characters and splitting classes.

The following three configurations (5.6), (5.7), (5.8) are basic for the labeling:

$$(5.6) \quad \begin{array}{ccc} G_{x,0} & \xrightarrow{\mu_{x,0}} & H_{x,0} = H_{x,00} \times H_{x,01} \\ \uparrow & & \uparrow \\ G_{x,0}^+ & \xrightarrow{\mu_{x,0}^+} & H_{x,0}^+ \\ & & \uparrow \\ & & H_{x,00}^+ \times H_{x,01}^+ \end{array}$$

where $\mu_{x,0}$ is the Rouquier isometry. If $G_{x,0} = G_{x,0}^+$ or $G_{x,0} = H_{x,0}$, then $\mu_{x,0}$ is the identity isometry. Here $G_{x,0} = G_{x,0}^+$ if $|\Pi_0| \leq 1$ and $j = w$; $G_{x,0} = H_{x,0}$ if $j = w$. In these cases, we fix $\mu_{x,0}^+$ as the identity isometry. If $G_{x,0} > G_{x,0}^+$, then $(G_{x,0}, G_{x,0}^+, e_{x,0}, e_{x,0}^+)$ and $(H_{x,0}, H_{x,0}^+, f_{x,0}, f_{x,0}^+)$ are +configurations, and $\mu_{x,0}$ is +equivariant with respect to $(G_{x,0}^+, H_{x,0}^+)$ by (5B), (i). We will fix $\mu_{x,0}^+$ in what follows so that $\mu_{x,0}^+$ is $\mu_{x,0}$ -induced and hence perfect by (5B), (ii).

$$(5.7) \quad \begin{array}{ccc} G_{x,1} & \xrightarrow{\mu_{x,1}} & H_{x,1} \\ \uparrow & & \uparrow \\ G_{x,1}^+ & \xrightarrow{\mu_{x,1}^+} & H_{x,1}^+ \end{array}$$

where $\mu_{x,1}$ is the identity isometry. We fix $\mu_{x,1}^+$ as the identity isometry. If $G_{x,1} > G_{x,1}^+$, then $(G_{x,1}, G_{x,1}^+, e_{x,1}, e_{x,1}^+)$ and $(H_{x,1}, H_{x,1}^+, f_{x,1}, f_{x,1}^+)$ are +configurations, $\mu_{x,1}$ is trivially +equivariant with respect to $(G_{x,1}^+, H_{x,1}^+)$, and $\mu_{x,1}^+$ is trivially $\mu_{x,1}$ -induced. Here $G_{x,1} > G_{x,1}^+$ if and only if $j > 1$.

$$\begin{array}{ccc}
 G_x = G_{x,0} \times G_{x,1} & \xrightarrow{\mu_x = \mu_{x,0} \times \mu_{x,1}} & H_x = H_{x,0} \times H_{x,1} \\
 \uparrow & & \uparrow \\
 G_x^+ & \xrightarrow{\mu_x^+} & H_x^+ \\
 \uparrow & & \uparrow \\
 G_x^{++} = G_{x,0}^+ \times G_{x,1}^+ & \xrightarrow{\mu_{x,0}^+ \times \mu_{x,1}^+} & H_x^{++} = H_{x,0}^+ \times H_{x,1}^+ \\
 & & \uparrow \\
 & & H_{x,00}^+ \times H_{x,01}^+ \times H_{x,1}^+
 \end{array}
 \tag{5.8}$$

where $\mu_{x,0}, \mu_{x,1}, \mu_{x,0}^+, \mu_{x,1}^+$ are the perfect isometries in (5.6) and (5.7). If $G_x = G_x^+$, then $G_{x,0} = G_{x,0}^+$ and $G_{x,1} = G_{x,1}^+$, that is, $|\Pi_0| \leq 1$ and $w = 1$. Thus $G_x = G_x^+ = G_x^{++}$ and μ_x is the identity isometry. In this case we fix μ_x^+ as the identity isometry. If $G_x > G_x^+$ and $G_x^+ = G_x^{++}$, then either $G_{x,0} = G_{x,0}^+$ or $G_{x,1} = G_{x,1}^+$, that is, either $|\Pi_0| \leq 1$ and $1 < j = w$, or $|\Pi_0 \cup (\Gamma \times \Omega_{x,0})| \geq 3$ and $j = 1$. In this case we fix $\mu_x^+ = \mu_{x,0}^+ \times \mu_{x,1}^+$. If $G_x > G_x^+ > G_x^{++}$, then

$$\begin{array}{ll}
 (G_x, G_x^+, e_x, e_x^+), & (H_x, H_x^+, f_x, f_x^+), \\
 (G_x^+, G_x^{++}, e_x^+, e_x^{++}), & (H_x^+, H_x^{++}, f_x^+, f_x^{++})
 \end{array}$$

are $+$ -configurations. Moreover, μ_x is perfect and $+$ -equivariant with respect to (G_x^+, H_x^+) by (2E), (i). We will fix μ_x^+ in what follows so that μ_x^+ is μ_x -induced and so that $\mu_{x,0}^+ \times \mu_{x,1}^+$ is μ_x^+ -induced. This is possible by (2F), (i). In particular, μ_x^+ is perfect by (5B), (ii).

We fix μ_x^+ for $x = x_j$ in the remaining cases as follows:

Step 1. If $|\lambda_0| \geq 3$, then the character χ_{λ_0} of $H_{x,00}$ splits on $H_{x,00}^+$. We label the irreducible characters $\chi_{\lambda_0, \pm}$ and the conjugacy classes $\mathcal{C}_{\eta(\lambda_0), \pm}$ of $H_{x,00}^+$ so that (4A) holds. This labeling is fixed for all j . If $|\lambda_0| \leq 1$, then $H_{x,00} = H_{x,00}^+ = 1$ and we omit this step.

Step 2. If $|\Omega_{x,0}| \geq 1$, let Λ' run over the p -regular splitting mappings of $H_{x,01}$. The splitting characters $\psi_{\Lambda'}$ of $H_{x,01}$ satisfy the hypotheses of (4G); the splitting classes $\mathcal{C}_{\Lambda'}$ of $H_{x,01}$ have p -regular cycle type. We label arbitrarily the irreducible characters $\psi_{\Lambda', \pm}$ of $H_{x,01}^+$ and the conjugacy classes $\mathcal{C}_{\Lambda', \pm}$ of $H_{x,01}^+$. If $|\Omega_{x,0}| = 0$, then $H_{x,01} = H_{x,01}^+ = 1$ and we omit this step.

Step 3. Let Λ'' run over the p -singular splitting mappings of $Y \wr S(\Omega_{x,1})$. Such mappings exist since $|\Omega_{x,1}| \geq 1$. The splitting characters $\psi_{\Lambda''}$ of $Y \wr S(\Omega_{x,1})$ satisfy the hypotheses of (4F); the splitting classes $\mathcal{C}_{\Lambda''}$ of $Y \wr S(\Omega_{x,1})$ have p -singular cycle type. We label the irreducible characters $\psi_{\Lambda'', \pm}$ of $Y \wr S(\Omega_{x,1})$ and the conjugacy classes $\mathcal{C}_{\Lambda'', \pm}$ of $Y \wr S(\Omega_{x,1})$ so that (4F) holds.

Step 4. Let λ' run over all self-dual partitions of $|\Pi_0 \cup (\Gamma \times \Omega_{x,0})|$ with p -core λ_0 and p -regular p -quotient λ' . Suppose $|\Omega_{x,0}| = 0$. Then λ' is trivial, $\lambda' = \lambda_0$, $G_{x,0} = H_{x,0} = H_{x,00}$, and $\chi_{\lambda', \pm}$ and $\mathcal{C}_{\eta(\lambda'), \pm}$ have already been labeled in step 1. Suppose $|\Omega_{x,0}| \geq 1$. Then λ' is a p -regular splitting mapping of $H_{x,01}$ of the type

considered in step 2. The conjugacy class $\mathcal{C}_{\eta(\lambda')}$ of $G_{x,0}$ contains the conjugacy class $\mathcal{C}_{\eta(\lambda_0)}\mathcal{C}_{\Lambda'}$ of $H_{x,0}$. If $|\lambda_0| \geq 3$, we label $\mathcal{C}_{\eta(\lambda'),\pm}$ so that

$$\begin{aligned} \mathcal{C}_{\eta(\lambda'),+} &= (\mathcal{C}_{\eta(\lambda_0),+}\mathcal{C}_{\Lambda',+})^{G_{x,0}^+} = (\mathcal{C}_{\eta(\lambda_0),-}\mathcal{C}_{\Lambda',-})^{G_{x,0}^+}, \\ \mathcal{C}_{\eta(\lambda'),-} &= (\mathcal{C}_{\eta(\lambda_0),+}\mathcal{C}_{\Lambda',-})^{G_{x,0}^+} = (\mathcal{C}_{\eta(\lambda_0),-}\mathcal{C}_{\Lambda',+})^{G_{x,0}^+}. \end{aligned}$$

If $|\lambda_0| \leq 1$, then $H_{x,00} = H_{x,00}^+ = 1$ and we label $\mathcal{C}_{\eta(\lambda'),\pm}$ so that $\mathcal{C}_{\eta(\lambda'),+} = (\mathcal{C}_{\Lambda',+})^{G_{x,0}^+}$ and $\mathcal{C}_{\eta(\lambda'),-} = (\mathcal{C}_{\Lambda',-})^{G_{x,0}^+}$. We then label the irreducible characters $\chi_{\lambda',\pm}$ of $G_{x,0}^+$ so that (4A) holds. We label the irreducible characters $(\chi_{\lambda_0}\psi_{\Lambda'})_{\pm}$ of $H_{x,0}^+$ so that

$$\begin{aligned} \text{Res}_{H_{x,00}^+ H_{x,01}^+}^{H_{x,0}^+} ((\chi_{\lambda_0}\psi_{\Lambda'})_+) &= \chi_{\lambda_0,+}\psi_{\Lambda',+} + \chi_{\lambda_0,-}\psi_{\Lambda',-}, \\ \text{Res}_{H_{x,00}^+ H_{x,01}^+}^{H_{x,0}^+} ((\chi_{\lambda_0}\psi_{\Lambda'})_-) &= \chi_{\lambda_0,+}\psi_{\Lambda',-} + \chi_{\lambda_0,-}\psi_{\Lambda',+} \end{aligned}$$

for $|\lambda_0| \geq 3$, and so that

$$(\chi_{\lambda_0}\psi_{\Lambda'})_+ = 1_{S(\Pi_0)} \times \psi_{\Lambda',+}, \quad (\chi_{\lambda_0}\psi_{\Lambda'})_- = 1_{S(\Pi_0)} \times \psi_{\Lambda',-}$$

for $|\lambda_0| \leq 1$.

Step 5. Let $\mu_{x,0} = (J_{x,0}, \epsilon_{x,0})$. We fix a $\mu_{x,0}$ -induced isometry $\mu_{x,0}^+ = (J_{x,0}^+, \epsilon_{x,0}^+)$ so that

$$J_{x,0}^+(\chi_{\lambda',\pm}) = J_{x,0}(\chi_{\lambda'})_{\pm} = (\chi_{\lambda_0}\psi_{\Lambda'})_{\pm}$$

for the λ' in step 4. Here $\widehat{\Lambda}' = \Lambda'$, since $\Lambda^* = \Lambda$.

Step 6. We label arbitrarily the irreducible characters and the conjugacy classes of $G_{x,1}^+ = H_{x,1}^+$ occurring in splitting irreducible characters and conjugacy classes of $G_{x,1} = H_{x,1}$. Here $\mu_{x,1}^+$ was already chosen as the identity isometry. If $G_x > G_x^+ > G_x^{++}$, we choose a μ_x -induced isometry μ_x^+ such $\mu_{x,0}^+ \times \mu_{x,1}^+$ is μ_x^+ -induced. In the remaining cases, we choose μ_x^+ as in the comments following (5.8). In all cases, μ_x^+ is μ_x -induced, $\mu_{x,0}^+ \times \mu_{x,1}^+$ is μ_x^+ -induced, and μ_x^+ is perfect.

Step 7. Let Λ run over all splitting mappings of H_1 such that the p -regular and p' -regular parts Λ' and Λ'' of Λ are splitting mappings of $H_{x,01}$ and $Y \wr S(\Omega_{x,1})$. We label the irreducible characters $\psi_{\Lambda,\pm}$ of H_1^+ so that (4H) (ii) holds. The conjugacy class \mathcal{C}_{Λ} of H_1 contains the conjugacy class $\mathcal{C}_{\Lambda'}\mathcal{C}_{\Lambda''}$ of $H_{x,01}H_{x,1}$. If $|\Omega_{x,0}| \geq 1$, we label $\mathcal{C}_{\Lambda,\pm}$ so that

$$\begin{aligned} \mathcal{C}_{\Lambda,+} &= (\mathcal{C}_{\Lambda',+}\mathcal{C}_{\Lambda'',+})^{H_1^+} = (\mathcal{C}_{\Lambda',-}\mathcal{C}_{\Lambda'',-})^{H_1^+}, \\ \mathcal{C}_{\Lambda,-} &= (\mathcal{C}_{\Lambda',+}\mathcal{C}_{\Lambda'',-})^{H_1^+} = (\mathcal{C}_{\Lambda',-}\mathcal{C}_{\Lambda'',+})^{H_1^+}. \end{aligned}$$

If $|\Omega_{x,0}| = 0$, then $H_{x,01} = 1$, and we label $\mathcal{C}_{\Lambda,\pm}$ so that $\mathcal{C}_{\Lambda,+} = \mathcal{C}_{\Lambda'',+}$ and $\mathcal{C}_{\Lambda,-} = \mathcal{C}_{\Lambda'',-}$.

Step 8. Let λ run over all self-dual partitions of $|\Pi|$ with p -core λ_0 and p -quotient Λ as in step 7. Let λ' and λ'' be the self-dual partitions gotten from λ by removing respectively all (i, i) -hooks of p -singular length and all (i, i) -hooks of p -regular length. We call λ' and λ'' the p -regular and p' -regular parts of λ . In particular, λ'

has p -core λ_0 and p -quotient Λ' , and λ'' has p -core \emptyset and p -quotient Λ'' . If $|\lambda'| \geq 3$, we label the conjugacy classes $\mathcal{C}_{\eta(\lambda),\pm}$ of G^+ so that

$$\begin{aligned} \mathcal{C}_{\eta(\lambda),+} &= (\mathcal{C}_{\eta(\lambda'),+}\mathcal{C}_{\Lambda'',+})^{G^+} = (\mathcal{C}_{\eta(\lambda'),-}\mathcal{C}_{\Lambda'',-})^{G^+}, \\ \mathcal{C}_{\eta(\lambda),-} &= (\mathcal{C}_{\eta(\lambda'),+}\mathcal{C}_{\Lambda'',-})^{G^+} = (\mathcal{C}_{\eta(\lambda'),-}\mathcal{C}_{\Lambda'',+})^{G^+}. \end{aligned}$$

If $|\lambda'| \leq 1$, then $G_{x,0} = 1$, and we label the conjugacy classes $\mathcal{C}_{\eta(\lambda),\pm}$ of G^+ so that $\mathcal{C}_{\eta(\lambda),+} = \mathcal{C}_{\Lambda'',+}$ and $\mathcal{C}_{\eta(\lambda),-} = \mathcal{C}_{\Lambda'',-}$. We then label the irreducible characters $\chi_{\lambda,\pm}$ of G^+ so that (4A) holds.

Step 9. The λ in step 8 determine splitting classes $\mathcal{C}_{\eta(\lambda_0),\Lambda} = \mathcal{C}_{\eta(\lambda_0)}\mathcal{C}_{\Lambda}$ of H . If $|\lambda_0| \geq 3$, we label the classes $\mathcal{C}_{\eta(\lambda_0),\Lambda,\pm}$ of H^+ so that

$$\begin{aligned} \mathcal{C}_{\eta(\lambda_0),\Lambda,+} &= (\mathcal{C}_{\eta(\lambda_0),+}\mathcal{C}_{\Lambda,+})^{H^+} = (\mathcal{C}_{\eta(\lambda_0),-}\mathcal{C}_{\Lambda,-})^{H^+}, \\ \mathcal{C}_{\eta(\lambda_0),\Lambda,-} &= (\mathcal{C}_{\eta(\lambda_0),+}\mathcal{C}_{\Lambda,-})^{H^+} = (\mathcal{C}_{\eta(\lambda_0),-}\mathcal{C}_{\Lambda,+})^{H^+}. \end{aligned}$$

We label the irreducible constituents $(\chi_{\lambda_0}\psi_{\Lambda})_{\pm}$ of $\text{Res}_{H^+}^H(\chi_{\lambda_0}\psi_{\Lambda})$ so that

$$\begin{aligned} \text{Res}_{H_0^+ H_1^+}^{H^+}((\chi_{\lambda_0}\psi_{\Lambda})_+) &= \chi_{\lambda_0,+}\psi_{\Lambda,+} + \chi_{\lambda_0,-}\psi_{\Lambda,-}, \\ \text{Res}_{H_0^+ H_1^+}^{H^+}((\chi_{\lambda_0}\psi_{\Lambda})_-) &= \chi_{\lambda_0,+}\psi_{\Lambda,-} + \chi_{\lambda_0,-}\psi_{\Lambda,+}. \end{aligned}$$

If $|\lambda_0| \leq 1$, then $H_{x,0} = 1$, and we label the classes $\mathcal{C}_{\eta(\lambda_0),\Lambda,\pm}$ of H^+ so that $\mathcal{C}_{\eta(\lambda_0),\Lambda,+} = \mathcal{C}_{\Lambda,+}$ and $\mathcal{C}_{\eta(\lambda_0),\Lambda,-} = \mathcal{C}_{\Lambda,-}$. We label the irreducible constituents $(\chi_{\lambda_0}\psi_{\Lambda})_{\pm}$ of $\text{Res}_{H^+}^H(\chi_{\lambda_0}\psi_{\Lambda})$ so that

$$(\chi_{\lambda_0}\psi_{\Lambda})_+ = 1_{S(\Pi_0)} \times \psi_{\Lambda,+}, \quad (\chi_{\lambda_0}\psi_{\Lambda})_- = 1_{S(\Pi_0)} \times \psi_{\Lambda,-}.$$

Step 10. Let λ run over all self-dual partitions of $|\Pi|$ with p -core λ_0 and a p -regular p -quotient Λ . We label arbitrarily the irreducible characters $\chi_{\lambda,\pm}$ and the conjugacy classes $\mathcal{C}_{\Lambda,\pm}$ of G^+ . Similarly, we label arbitrarily the irreducible constituents $(\chi_{\lambda_0}\psi_{\Lambda})_{\pm}$ of $\text{Res}_{H^+}^H(\chi_{\lambda_0}\psi_{\Lambda})$ and the conjugacy classes $\mathcal{C}_{\eta(\lambda_0),\Lambda,\pm} = (\mathcal{C}_{\eta(\lambda_0)}\mathcal{C}_{\Lambda})_{\pm}$ of H^+ .

Step 11. Let λ run over all self-dual partitions of $|\Pi|$ with p -core λ_0 . Let $\mu = (J, \epsilon)$. We choose the μ -induced isometry $\mu^+ = (J^+, \epsilon^+)$ so that

$$J^+(\chi_{\lambda,\pm}) = \begin{cases} J(\chi_{\lambda})_{\pm} = (\chi_{\lambda_0}\psi_{\Lambda})_{\pm} & \text{if } (-\epsilon_p)^{d_{\lambda}}\sigma_{\lambda} = 1, \\ J(\chi_{\lambda})_{\pm} = (\chi_{\lambda_0}\psi_{\Lambda})_{\mp} & \text{if } (-\epsilon_p)^{d_{\lambda}}\sigma_{\lambda} = -1, \end{cases}$$

where d_{λ} is the number of diagonal nodes of λ'' and

$$\sigma_{\lambda} = \frac{\sqrt{\epsilon_{\lambda'}}(\sqrt{\epsilon_p})^{d_{\lambda}}\sqrt{\epsilon_{\lambda_{p^*}}}}{\sqrt{\epsilon_{\lambda}}}.$$

Note that $\sigma_{\lambda} = \pm 1$. Indeed, if $\eta_{ii}(\lambda_{p^*}) = 2u_i + 1$, then

$$\begin{aligned} \epsilon_{\lambda_{p^*}} &= (-1)^{\frac{1}{2}[(\sum(2u_i+1))-d_{\lambda}]} \\ &= (-1)^{\sum u_i}, \end{aligned}$$

and

$$\begin{aligned} \epsilon_{\lambda''} &= (-1)^{\frac{1}{2}[(\sum(2u_i+p))-d_{\lambda}]} \\ &= (-1)^{(\sum u_i p) + d_{\lambda}(\frac{p-1}{2})}. \end{aligned}$$

Since $\sum u_i p \equiv \sum u_i \pmod{2}$ and $\epsilon_p = (-1)^{\binom{p-1}{2}}$, it follows that $\epsilon_{\lambda'} = (\epsilon_p)^{d_\lambda} \epsilon_{\lambda_{p^*}}$ and $\epsilon_\lambda = \epsilon_{\lambda'} (\epsilon_p)^{d_\lambda} \epsilon_{\lambda_{p^*}}$. Thus $\sigma_\lambda = \pm 1$.

Proposition (5C). *Let λ be a self-dual partition of n with p -core λ_0 and p -quotient Λ , and let $x \in \{x_1, x_2, \dots, x_w\}$. Then*

$$R_{\mu_x^+, p'}(d_{G^+}^x(\chi_{\lambda,+} - \chi_{\lambda,-}))(y) = 0$$

unless $|\lambda_{p^*}| = |\Omega_{x,1}|$ and $y \in (H_{x,0}^+ H_{x,01}^+ H_{x,1}^+)_{p'}$. Moreover, if $|\lambda_{p^*}| = |\Omega_{x,1}|$ and $y = y_{00} y_{01} y_1$ with $y_{00} \in (H_{x,00}^+)_{p'}$, $y_{01} \in (H_{x,01}^+)_{p'}$, $y_1 \in (H_{x,1}^+)_{p'}$, then

$$R_{\mu_x^+, p'}(d_{G^+}^x(\chi_{\lambda,+} - \chi_{\lambda,-}))(y) = \sigma_\lambda \epsilon_{x,0}(\chi_{\lambda'}) (\chi_{\lambda_0,+} - \chi_{\lambda_0,-})(y_{00}) \\ \times (\psi_{\Lambda',+} - \psi_{\Lambda',-})(y_{01}) (\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(xy_1).$$

Here λ' is the p -regular part of λ ; $\psi_{\Lambda'}$ and $\psi_{\Lambda''}$ are the irreducible characters of $Y \wr S(\Omega_{x,0})$ and $Y \wr S(\Omega_{x,1})$ parametrized by the p -regular and p -singular parts Λ' and Λ'' of Λ ; and the factor $(\chi_{\lambda_0,+} - \chi_{\lambda_0,-})$ or $(\psi_{\Lambda',+} - \psi_{\Lambda',-})$ is interpreted as 1 when $|\lambda_0| \leq 1$ or $|\Omega_{x,0}| = 0$.

Proof. Let $\rho_{\Lambda'', \pm} = \text{Res}_{(P \wr S(\Omega_{x,1}))^+}^{(Y \wr S(\Omega_{x,1}))^+}(\psi_{\Lambda'', \pm})$ if $|\lambda_{p^*}| = |\Omega_{x,1}|$, and let $\rho_{\Lambda'', \pm}$ be the zero function if $|\lambda_{p^*}| \neq |\Omega_{x,1}|$. We show first that

$$(5.9) \quad d_{G^+}^x(\chi_{\lambda,+} - \chi_{\lambda,-}) = \sigma_\lambda \text{Ind}_{G_{x,0}^+}^{G_x^+} (d_{G_{x,0}^+}^1(\chi_{\lambda',+}) \times d_{G_{x,1}^+}^x(\rho_{\Lambda'',+} - \rho_{\Lambda'',-})),$$

where $d_{G_{x,0}^+}^1(\chi_{\lambda',+})$ is interpreted as 1 when $|\lambda'| \leq 1$. Indeed, let $z_0 z_1 \in G_{x,p'}^+$, where $z_0 \in G_{x,0}$ and $z_1 \in G_{x,1}$. By (4A), $(\chi_{\lambda,+} - \chi_{\lambda,-})(xz_0 z_1) \neq 0$ only if $xz_0 z_1 \in \mathcal{C}_{\eta(\lambda)}$, or equivalently, only if $|\lambda_{p^*}| = |\Omega_{x,1}|$, z_0 is in the class $\mathcal{C}_{\eta(\lambda')}$ of $G_{x,0}$, and xz_1 is in the class $\mathcal{C}_{\Lambda''}$ of $G_{x,1}$. Thus (5.9) holds if $|\lambda_{p^*}| \neq |\Omega_{x,1}|$, since both sides of (5.9) are zero. Suppose $|\lambda_{p^*}| = |\Omega_{x,1}|$. Then

$$d_{G^+}^x(\chi_{\lambda,+} - \chi_{\lambda,-})(z_0 z_1) = \begin{cases} \pm \sqrt{\epsilon_\lambda \eta(\lambda)} & \text{if } xz_0 z_1 \in \mathcal{C}_{\eta(\lambda), \pm}, \\ 0 & \text{if } xz_0 z_1 \notin \mathcal{C}_{\eta(\lambda), \pm}. \end{cases}$$

On the other hand, $\text{Ind}_{G_{x,0}^+}^{G_x^+} (d_{G_{x,0}^+}^1(\chi_{\lambda',+}) \times d_{G_{x,1}^+}^x(\rho_{\Lambda'',+} - \rho_{\Lambda'',-}))$ has support on $G_{x,p'}^+$, where it is equal to

$$(5.10) \quad d_{G_{x,0}^+}^1(\chi_{\lambda',+} - \chi_{\lambda',-}) \times d_{G_{x,1}^+}^x(\rho_{\Lambda'',+} - \rho_{\Lambda'',-}).$$

Moreover, (4A) and (4F) imply that $(\chi_{\lambda',+} - \chi_{\lambda',-})(z_0)$ and $(\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(xz_1)$ are zero except for

$$(\chi_{\lambda',+} - \chi_{\lambda',-})(z_0) = \pm \sqrt{\epsilon_{\lambda'} \eta(\lambda')} \quad \text{if } z_0 \in \mathcal{C}_{\eta(\lambda'), \pm}, \\ (\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(xz_1) = \pm (\sqrt{\epsilon_p p})^{d_\lambda} \sqrt{\epsilon_{\lambda_{p^*}} \eta(\lambda_{p^*})} \quad \text{if } xz_1 \in \mathcal{C}_{\Lambda'', \pm}.$$

Here $(\chi_{\lambda',+} - \chi_{\lambda',-})(z_0)$ and $\pm \sqrt{\epsilon_{\lambda'} \eta(\lambda')}$ are interpreted as 1 when $|\lambda'| \leq 1$. Since

$$\epsilon_\lambda = \epsilon_{\lambda'} (\epsilon_p)^{d_\lambda} \epsilon_{\lambda_{p^*}}, \quad \sqrt{\eta(\lambda)} = \sqrt{\eta(\lambda') p^{d_\lambda} \eta(\lambda_{p^*})},$$

(5.9) follows from the labeling in step 8.

Suppose $|\lambda'| \geq 3$. Then

$$\begin{aligned}
 & R_{\mu_x^+, p'}(d_{G^+}^x(\chi_{\lambda, +} - \chi_{\lambda, -})) \\
 &= R_{\mu_x^+, p'}(\sigma_\lambda \text{Ind}_{G_x^{++}}^{G_x^+}(d_{G_x^+, 0}^1(\chi_{\lambda', +}) \times d_{G_x^+, 1}^x(\rho_{\Lambda'', +} - \rho_{\Lambda'', -}))) \\
 &= \sigma_\lambda \text{Ind}_{H_x^{++}}^{H_x^+}(R_{\mu_{x,0}^+ \times \mu_{x,1}^+, p'}(d_{G_x^+, 0}^1(\chi_{\lambda', +}) \times d_{G_x^+, 1}^x(\rho_{\Lambda'', +} - \rho_{\Lambda'', -}))) \\
 &\quad (\text{since } \mu_x^+ \text{ induces } \mu_{x,0}^+ \times \mu_{x,1}^+) \\
 (5.11) \quad &= \sigma_\lambda \text{Ind}_{H_x^{++}}^{H_x^+}(R_{\mu_{x,0}^+, p'}(d_{G_x^+, 0}^1(\chi_{\lambda', +})) \times R_{\mu_{x,1}^+, p'}(d_{G_x^+, 1}^x(\rho_{\Lambda'', +} - \rho_{\Lambda'', -}))) \\
 &= \sigma_\lambda \text{Ind}_{H_x^{++}}^{H_x^+}(d_{H_x^+, 0}^1(R_{\mu_{x,0}^+}(\chi_{\lambda', +})) \times d_{H_x^+, 1}^x(R_{\mu_{x,1}^+}(\rho_{\Lambda'', +} - \rho_{\Lambda'', -}))) \\
 &\quad (\text{since } \mu_{x,0}^+ \text{ is perfect by (5B) and } \mu_{x,1}^+ \text{ is the identity isometry}) \\
 &= \sigma_\lambda d_{H_x^+}^x(\text{Ind}_{H_x^{++}}^{H_x^+}(R_{\mu_{x,0}^+ \times \mu_{x,1}^+}(\chi_{\lambda', +} \times (\rho_{\Lambda'', +} - \rho_{\Lambda'', -}))).
 \end{aligned}$$

Thus $R_{\mu_x^+, p'}(d_{G^+}^x(\chi_{\lambda, +} - \chi_{\lambda, -}))$ has support on $(H_x^{++})_{p'}$, where it is equal to

$$\sigma_\lambda d_{H_x^+}^x(R_{\mu_{x,0}^+}(\chi_{\lambda', +} - \chi_{\lambda', -}) \times R_{\mu_{x,1}^+}(\rho_{\Lambda'', +} - \rho_{\Lambda'', -})).$$

In turn, $R_{\mu_{x,0}^+}(\chi_{\lambda', +} - \chi_{\lambda', -})$ has support on $H_{x,00}^+ H_{x,01}^+$, where it is equal to

$$\epsilon_{x,0}(\chi_{\lambda'}) (\chi_{\lambda_{0,+}} - \chi_{\lambda_{0,-}}) \times (\psi_{\Lambda', +} - \psi_{\Lambda', -}).$$

Thus $R_{\mu_x^+, p'}(d_{G^+}^x(\chi_{\lambda, +} - \chi_{\lambda, -}))$ has support on $(H_{x,00}^+ H_{x,01}^+ H_{x,1}^+)_{p'}$, and the first half of (5C) holds. The argument shows that

$$\begin{aligned}
 R_{\mu_x^+, p'}(d_{G^+}^x(\chi_{\lambda, +} - \chi_{\lambda, -}))(y) &= \sigma_\lambda \epsilon_{x,0}(\chi_{\lambda'}) (\chi_{\lambda_{0,+}} - \chi_{\lambda_{0,-}})(y_{00}) \\
 &\quad \times (\psi_{\Lambda', +} - \psi_{\Lambda', -})(y_{01}) (\psi_{\Lambda'', +} - \psi_{\Lambda'', -})(xy_1)
 \end{aligned}$$

if $|\lambda_{p^*}| = |\Omega_{x,1}|$ and $y = y_{00}y_{01}y_1$ with $y_{00} \in (H_{x,00}^+)_{p'}$, $y_{01} \in (H_{x,01}^+)_{p'}$, and $y_1 \in (H_{x,1}^+)_{p'}$. Thus (5C) holds when $|\lambda'| \geq 3$.

Suppose $|\lambda'| \leq 1$. Then $G_x^+ = G_x^{++}$, and (5.11) reduces to

$$\begin{aligned}
 R_{\mu_x^+, p'}(d_{G^+}^x(\chi_{\lambda, +} - \chi_{\lambda, -})) &= R_{\mu_x^+, p'}(\sigma_\lambda d_{G_x^+, 1}^x(\rho_{\Lambda'', +} - \rho_{\Lambda'', -})) \\
 &= \sigma_\lambda R_{\mu_{x,1}^+, p'}(d_{G_x^+, 1}^x(\rho_{\Lambda'', +} - \rho_{\Lambda'', -})) \\
 &= \sigma_\lambda d_{H_{x,1}^+}^x(R_{\mu_{x,1}^+}(\rho_{\Lambda'', +} - \rho_{\Lambda'', -})).
 \end{aligned}$$

Then (5C) holds trivially in this case, since $(H_{x,00}^+ H_{x,01}^+ H_{x,1}^+)_{p'} = (H_{x,1}^+)_{p'}$. □

Proposition (5D). *Let λ be a self-dual partition of n with p -core λ_0 and p -quotient Λ , and let $x \in \{x_1, x_2, \dots, x_w\}$. Then*

$$d_{H^+}^x(R_{\mu^+}((\chi_{\lambda, +} - \chi_{\lambda, -}))(y) = 0$$

unless $|\lambda_{p^*}| = |\Omega_{x,1}|$ and $y \in (H_{x,00}^+ H_{x,01}^+ H_{x,1}^+)_{p'}$. Moreover, if $|\lambda_{p^*}| = |\Omega_{x,1}|$ and $y = y_{00}y_{01}y_1$ with $y_{00} \in (H_{x,00}^+)_{p'}$, $y_{01} \in (H_{x,01}^+)_{p'}$, $y_1 \in (H_{x,1}^+)_{p'}$, then

$$\begin{aligned}
 d_{H^+}^x(R_{\mu^+}(\chi_{\lambda, +} - \chi_{\lambda, -}))(y) &= \sigma_\lambda (-\epsilon_p)^{d_\lambda} \epsilon(\chi_\lambda) (\chi_{\lambda_{0,+}} - \chi_{\lambda_{0,-}})(y_{00}) \\
 &\quad \times (\psi_{\Lambda', +} - \psi_{\Lambda', -})(y_{01}) (\psi_{\Lambda'', +} - \psi_{\Lambda'', -})(xy_1).
 \end{aligned}$$

Here $\psi_{\Lambda'}$ and $\psi_{\Lambda''}$ are the irreducible characters of $Y \wr S(\Omega_{x,0})$ and $Y \wr S(\Omega_{x,0})$ parametrized by the p -regular and p -singular parts Λ' and Λ'' of Λ , and the factor $(\chi_{\lambda_0,+} - \chi_{\lambda_0,-})$ or $(\psi_{\Lambda',+} - \psi_{\Lambda',-})$ is interpreted as 1 when $|\lambda_0| \leq 1$ or $|\Omega_{x,0}| = 0$.

Proof. For $y \in (H_x^+)_{p'}$ we have

$$\begin{aligned} d_{H^+}^x(\mathbf{R}_{\mu^+}(\chi_{\lambda,+} - \chi_{\lambda,-}))(y) &= \mathbf{R}_{\mu^+}(\chi_{\lambda,+} - \chi_{\lambda,-})(xy) \\ &= (-\epsilon_p)^{d_\lambda} \sigma_{\lambda \in (\chi_\lambda)}(\mathbf{J}^+(\chi_{\lambda,+}) - \mathbf{J}^+(\chi_{\lambda,-}))(xy). \end{aligned}$$

Let $y = y_{00}y_{01}y_1$, where $y_{00} \in (H_{x,00})_{p'}$, $y_{01} \in (H_{x,01})_{p'}$, $y_1 \in (H_{x,1})_{p'}$. Thus $y_{00} \in (H_0)_{p'}$, $y_{01}y_1 \in (H_1)_{p'}$, and

$$\begin{aligned} &(\mathbf{J}^+(\chi_{\lambda,+}) - \mathbf{J}^+(\chi_{\lambda,-}))(xy) \\ &= \begin{cases} (\chi_{\lambda_0,+} - \chi_{\lambda_0,-})(y_{00})(\psi_{\Lambda,+} - \psi_{\Lambda,-})(xy_{01}y_1) & \text{if } xy \in H_0^+ H_1^+, \\ 0 & \text{if } xy \notin H_0^+ H_1^+ \end{cases} \end{aligned}$$

by the labeling of $\mathbf{J}^+(\chi_{\lambda,\pm})$ in steps 8–10. Suppose $d_{H^+}^x(\mathbf{R}_{\mu^+}(\chi_{\lambda,+} - \chi_{\lambda,-}))(y) \neq 0$. Then $y_{00} \in (H_{x,00}^+)_{p'}$, $y_{01}y_1 \in (H_1^+)_{p'}$, and $(\psi_{\Lambda,+} - \psi_{\Lambda,-})(xy_{01}y_1) \neq 0$. Let H_1' and H_1'' be the subgroups of H_1 associated to Λ' and Λ'' . By (4H) (i) there exist elements u' in a p -regular splitting class of H_1' , u'' in a p -singular splitting class of H_1'' , and $h \in H_1^+$ such that $(xy_{01}y_1)^h = u'u''$. A comparison of the p -regular and p -singular cycle types of these elements as elements of $S(\Gamma \times \Omega)$ shows that $(y_{01})^h = u'$, $(xy_1)^h = u''$. Thus $|\lambda_{p^*}| = |\Omega_{x,1}|$, $y_{01} \in H_{x,01}^+$, $y_1 \in H_{x,1}^+$, and the first part of (5D) holds. Now suppose $|\lambda_{p^*}| = |\Omega_{x,1}|$, $y_{00} \in (H_{x,00}^+)_{p'}$, $y_{01} \in (H_{x,01}^+)_{p'}$, and $y_1 \in (H_{x,1}^+)_{p'}$. We can then choose $Y \wr S(\Omega_{x,0})$ and $Y \wr S(\Omega_{x,1})$ as the subgroups H_1' and H_1'' of H_1 associated to Λ' and Λ'' . The second part of (5D) now follows from (4H) (ii), the labelings in steps 8–9, and the definition of $\mathbf{J}^+(\chi_{\lambda,\pm})$. \square

Proposition (5E). *The diagram*

$$(5.12) \quad \begin{array}{ccc} \mathcal{R}_K(G^+, e^+) & \xrightarrow{\mathbf{R}_{\mu^+}} & \mathcal{R}_K(H^+, f^+) \\ \downarrow d_{G^+}^x & & \downarrow d_{H^+}^x \\ CF_{p'}(G_x^+, e_x^+) & \xrightarrow{\mathbf{R}_{\mu_x^+, p'}} & CF_{p'}(H_x^+, f_x^+) \end{array}$$

is commutative for all x in $\{x_1, x_2, \dots, x_w\}$.

Proof. Suppose $\chi^+ = \text{Res}_{G^+}^G(\chi)$ for $\chi \in (G, e)^\vee$. Then

$$d_{H^+}^x(\mathbf{R}_{\mu^+}(\chi^+)) = \mathbf{R}_{\mu_x^+, p'}(d_{G^+}^x(\chi^+)),$$

that is, (5.12) is commutative for such χ^+ . Indeed, $d_{H^+}^x \circ \mathbf{R}_{\mu^+} = \mathbf{R}_{\mu_x, p'} \circ d_{G^+}^x$ by (3B), and

$$\mathbf{R}_{\mu^+} \circ \text{Res}_{G^+}^G = \text{Res}_{H^+}^H \circ \mathbf{R}_{\mu^+}, \quad \mathbf{R}_{\mu_x^+, p'} \circ \text{Res}_{G_x^+}^{G_x} = \text{Res}_{H_x^+}^{H_x} \circ \mathbf{R}_{\mu_x},$$

since μ^+ is μ -induced and μ_x^+ is μ_x -induced. Thus

$$\begin{aligned} d_{H^+}^x \circ R_{\mu^+} \circ \text{Res}_{G^+}^G &= d_{H^+}^x \circ \text{Res}_{H^+}^H \circ R_{\mu} \\ &= \text{Res}_{H_{x,p'}^+}^{H_{x,p'}} \circ d_H^x \circ R_{\mu} \\ &= \text{Res}_{H_{x,p'}^+}^{H_{x,p'}} \circ R_{\mu_{x,p'}} \circ d_G^x \\ &= R_{\mu_{x,p'}^+} \circ \text{Res}_{G_{x,p'}^+}^{G_{x,p'}} \circ d_G^x \\ &= R_{\mu_{x,p'}^+} \circ d_{G^+}^x \circ \text{Res}_{G^+}^G, \end{aligned}$$

since μ_x and μ_x^+ are perfect isometries.

Suppose $\chi^+ = \chi_{\lambda,+} - \chi_{\lambda,-}$, where $\chi_{\lambda} \in (G, e)^\vee$, λ is self-dual, and $G_x^+ > G_x^{++}$. The commutativity of (5.12) for χ^+ follows from the expressions for

$$R_{\mu_{x,p'}^+}(d_{G^+}^x(\chi_{\lambda,+} - \chi_{\lambda,-}))(y), \quad d_{H^+}^x(R_{\mu^+}(\chi_{\lambda,+} - \chi_{\lambda,-}))(y)$$

given in (5C) and (5D). Indeed, we may suppose that Λ' is a p -regular splitting mapping of $Y \wr S(\Omega_{x,0})$ and Λ'' is a p -singular splitting mapping of $Y \wr S(\Omega_{x,1})$. It remains to show that

$$(5.13) \quad \sigma_{\lambda} \epsilon_{x,0}(\chi_{\lambda'}) = (-\epsilon_p)^{d_{\lambda}} \sigma_{\lambda} \epsilon(\chi_{\lambda}).$$

But $\epsilon(\chi_{\lambda}) = v(\lambda)(-1)^{|\lambda_{p^*}|} = (-1)^{d_{\lambda}}$ since λ is self-dual, $\epsilon_{x,0}(\chi_{\lambda'}) = v(\lambda')$ since λ' is p -regular, and $v(\lambda)v(\lambda') = (\epsilon_p)^{d_{\lambda}}$. Thus (5.13) holds.

Suppose $\chi^+ = \chi_{\lambda,+} - \chi_{\lambda,-}$, where $\chi_{\lambda} \in (G, e)^\vee$, λ is self-dual, and $G_x^+ = G_x^{++}$. Then $|\Pi_0 \cup (\Gamma \times \Omega_{x,0})| \leq 1$ or $|\Omega_{x,1}| = 1$. If $|\Pi_0 \cup (\Gamma \times \Omega_{x,0})| \leq 1$, then $G_x^+ = H_x^+$, μ_x^+ is the identity isometry, and (5.12) holds trivially. If $|\Omega_{x,1}| = 1$, then

$$G_x^+ = A(\Pi_0 \cup (\Gamma \times \Omega_{x,0}) \times P, \quad H_x^+ = (S(\Pi_0) \times (Y \wr S(\Omega_{x,0})))^+ \times P$$

The previous arguments apply if $(\psi_{\Lambda'',+} - \psi_{\Lambda'',-})(y_1)$ and $(\rho_{\Lambda'',+} - \rho_{\Lambda'',-})(y_1)$ are interpreted as 1 when $|\lambda_0| \leq 1$ or $|\Omega_{x,0}| = 0$. This completes the proof of (5E). \square

Theorem (5F). *e^+ and f^+ are isotypic blocks of $\mathcal{O}G^+$ and $\mathcal{O}H^+$ with local system given by the translates of the μ_x^+ for x in $\{x_1, x_2, \dots, x_w\}$.*

Proof. The inclusion of D in G^+ and H^+ induces an equivalence of $\mathbf{Br}_{e^+,D}(G^+)$ and $\mathbf{Br}_{f^+,D}(H^+)$ by [1, (4.1)]. We claim the diagram (5.12) with x replaced by any generator of $\langle x \rangle$ is a commutative diagram. Indeed, the horizontal arrows in (5.12) depend only on $\langle x \rangle$ and not on x . The proofs remain valid when x is replaced by any generator of $\langle x \rangle$. Since $x \in \{x_1, x_2, \dots, x_w\}$ and the w subgroups $\langle x_j \rangle$ form a set of representatives for the $N_{G^+}(D, e_D^+)$ -orbits of non-trivial cyclic subgroups of D , (5F) holds by transport of structure, as remarked in §1. \square

6. COMPATIBILITY

The Broué conjecture for the principal block holds for all finite groups if it holds for all finite simple non-abelian groups X and if the isotypies for X satisfy compatibility conditions with regard to extensions of X by p' -groups of automorphisms (see [10, (5E)]). We show that the isotypies constructed in §5 for alternating groups satisfy these compatibility conditions. We suppose $p > 2$ throughout this section.

Let G and G^+ be the symmetric and the alternating groups of degree n , let e and e^+ be the 1-blocks of $\mathcal{O}G$ and $\mathcal{O}G^+$, and let e and e^+ have the common non-trivial abelian defect group D . Moreover, let $H = N_G(D)$ and $H^+ = N_{G^+}(D)$, and let f

and f^+ be the Brauer correspondents of e and e^+ . In particular, e^+ and f^+ are stable under G and H . On the other hand, $e = e^*$ and $f = f^*$ hold if and only if $|\Pi_0| \leq 1$. The maximal subpairs $(D, e_D), (D, f_D), (D, e_D^+), (D, f_D^+)$ for e, f, e^+, f^+ are unique since e, f, e^+, f^+ are the 1-blocks. Let $\{\mu_P\}$ and $\{\mu_P^+\}$ be the local systems for the isotopies between e and f and between e^+ and f^+ constructed in §3 and §5.

Let $\tilde{G} = \text{Aut}(G^+)$, so that $\tilde{G} \geq G$ and $|\tilde{G}/G| = 1$ or 2 according as $n \neq 6$ or $n = 6$. Let $\tilde{H} = N_{\tilde{G}}(D)$. The inclusion of \tilde{H} in \tilde{G} induces an isomorphism $\alpha: \tilde{G}/G^+ \rightarrow \tilde{H}/H^+$, since $\tilde{G} = \tilde{H}G^+$. The compatibility conditions from [10, §3] are of three types. The first type $(\tilde{\mathbf{F}}1), (\tilde{\mathbf{F}}2'), (\tilde{\mathbf{F}}3)$ relate to fusion:

- $(\tilde{\mathbf{F}}1)$ \tilde{G}/G^+ and \tilde{H}/H^+ are p' -groups stabilizing e^+ and f^+ .
- $(\tilde{\mathbf{F}}2')$ Given x in $N_{\tilde{G}}(D)$, there exists y in $N_{\tilde{H}}(D)$ with $\alpha(xG^+) = yH^+$ such that x and y induce the same mapping on D . Similarly, given y in $N_{\tilde{H}}(D)$, there exists x in $N_{\tilde{G}}(D)$ with $\alpha(xG^+) = yH^+$ such that x and y induce the same mapping on D .
- $(\tilde{\mathbf{F}}3)$ $\alpha(\tilde{G}_P G^+/G^+) = \tilde{H}_P H^+/H^+$ for all cyclic subgroups P of D .

$(\tilde{\mathbf{F}}1)$ holds since e^+ and f^+ are the 1-blocks. $(\tilde{\mathbf{F}}2')$ and $(\tilde{\mathbf{F}}3)$ follow easily from $\tilde{G} = \tilde{H}G^+$ and the structure of \tilde{G}_P and \tilde{H}_P .

The second type $(\tilde{\mathbf{I}}1)$ imposes extendibility conditions on the isometries $\mu_P^+ = (J_P^+, \epsilon_P^+)$ for all cyclic $P \leq D$. Namely, let $\alpha_P: \tilde{G}_P/G_P^+ \rightarrow \tilde{H}_P/H_P^+$ be the isomorphism defined by $\alpha_P(xG_P^+) = \alpha(xG^+) \cap \tilde{H}_P$ for $x \in \tilde{G}_P$, and set

$$\Delta(\alpha_P) = \{(x, y) : x \in \tilde{G}_P, y \in \tilde{H}_P, \alpha_P(xG_P^+) = yH_P^+\},$$

$$\Delta(\mu_P^+) = \{\chi_P^+ \times \bar{\psi}_P^+ : \chi_P^+ \in (G_P^+, e_P^+)^\vee, \psi_P^+ \in (H_P^+, f_P^+)^\vee, J_P^+(\chi_P^+) = \psi_P^+\}.$$

In particular, $\Delta(\alpha_P) = \Delta(\alpha) \cap (\tilde{G}_P \times \tilde{H}_P)$. Then the following hold:

- $(\tilde{\mathbf{I}}1.1)$ μ_P^+ is stable under $\Delta(\alpha_P)$.
- $(\tilde{\mathbf{I}}1.2)$ Each $\chi_P^+ \times \bar{\psi}_P^+$ in $\Delta(\mu_P^+)$ has a $\Delta(\alpha_P)$ -equivariant extension to a character $E_P(\chi_P^+ \times \bar{\psi}_P^+)$ of its stabilizer $\Delta(\alpha_P)_{\chi_P^+ \times \bar{\psi}_P^+}$ in $\Delta(\alpha_P)$.
- $(\tilde{\mathbf{I}}1.3)$ The extension of μ_P^+ to $\Delta(\alpha_P)$ given by

$$E_P(\mu_P^+) = \sum_{\chi_P^+ \times \bar{\psi}_P^+ \in \Delta(\mu_P^+)} \epsilon_{\mu_P^+}(\chi_P^+) E_P(\chi_P^+ \times \bar{\psi}_P^+),$$

where $E_P(\chi_P^+ \times \bar{\psi}_P^+)(x, y) = 0$ for $(x, y) \notin \Delta(\alpha_P)_{\chi_P^+ \times \bar{\psi}_P^+}$, is perfect in the sense that

- $(\tilde{\mathbf{I}}1.3a)$ $E_P(\mu_P^+)$ separates p -singular and p -regular elements of $\Delta(\alpha_P)$, and
- $(\tilde{\mathbf{I}}1.3b)$ $E_P(\mu_P^+)(x, y)$ is divisible in \mathcal{O} by $|C_{G_P^+}(x)|$ and $|C_{H_P^+}(y)|$ for all (x, y) in $\Delta(\alpha_P)$.

We verify $(\tilde{\mathbf{I}}1)$ when $n \neq 6$. If $e = e^*$, then μ_P^+ is stable under $\Delta(\alpha_P)$ by the construction in §5. If $e \neq e^*$, then each term $\chi_P^+ \times \bar{\psi}_P^+$ in μ_P^+ is even stable under $\Delta(\alpha_P)$. Let χ_P^+ and ψ_P^+ be characters corresponding under J_P^+ . Then $\Delta(\alpha_P)_{\chi_P^+ \times \bar{\psi}_P^+}$ is $G_P^+ \times H_P^+$ or $\Delta(\alpha_P)$. In the first case we set $E_P(\chi_P^+ \times \bar{\psi}_P^+) = \chi_P^+ \times \bar{\psi}_P^+$. In the second case χ_P^+ and ψ_P^+ extend to irreducible characters χ_P and ψ_P in $(G_P, e_P)^\vee$ and $(H_P, f_P)^\vee$ corresponding under μ_P , and we set

$$E_P(\chi_P^+ \times \bar{\psi}_P^+) = \text{Res}_{\Delta(\alpha_P)}^{G_P \times H_P}(\chi_P \times \bar{\psi}_P).$$

By (1D) and (2.3)

$$(6.1) \quad E_P(\mu_P^+)(x, y) = \begin{cases} \mu_P(x, y) & \text{if } e \neq e^*, \\ \frac{1}{2}\mu_P(x, y) + \frac{1}{2}\Delta(\mu_P^+)(x, y) & \text{if } e = e^*. \end{cases}$$

where $\Delta(\mu_P^+)$ is the generalized character of $G_P^+ \times H_P^+$ of (2B) extended by zero to $\Delta(\alpha_P)$. Since μ_P and $\Delta(\mu_P^+)$ are perfect, it follows that all conditions of $(\tilde{\mathbf{I1}})$ hold.

The third type $(\tilde{\mathbf{G1}})$ and $(\tilde{\mathbf{G2}})$ are glueing conditions on the extensions in $(\tilde{\mathbf{I1}})$.

$$(\tilde{\mathbf{G1}}) \quad E_{\langle 1 \rangle}(\mu_{\langle 1 \rangle}^+)(xx', yy') = 0 \text{ for all } (xx', yy') \in \Delta(\alpha) \text{ such that } x \in G_p^+, y \in H_p^+, x' \in \tilde{G}_{x,p'}, y' \in \tilde{H}_{y,p'}, \text{ and } x \not\sim y \text{ in } G^+.$$

$$(\tilde{\mathbf{G2}}) \quad E_{\langle 1 \rangle}(\mu_{\langle 1 \rangle}^+)(zx', zy') = E_{\langle z \rangle}(\mu_{\langle z \rangle}^+)(x', y') \text{ for all } (zx', zy') \in \Delta(\alpha) \text{ such that } z \in D, x' \in C_{\tilde{X}}(z)_{p'}, y' \in C_{\tilde{Y}}(z)_{p'}.$$

We verify $(\tilde{\mathbf{G1}})$ and $(\tilde{\mathbf{G2}})$ when $n \neq 6$. Suppose $e \neq e^*$. Then $E_{\langle 1 \rangle}(\mu_{\langle 1 \rangle}^+) = \text{Res}_{\Delta(\alpha)}^{G \times H}(\mu)$ and $E_P(\mu_P^+) = \text{Res}_{\Delta(\alpha_P)}^{G_P \times H_P}(\mu_P)$, so $(\tilde{\mathbf{G2}})$ holds. Let $(xx', yy') \in \Delta(\alpha)$ be as in $(\tilde{\mathbf{G1}})$. Since $G_x \not\leq G^+$ when $|\Pi_0| \geq 3$, it follows that $x \not\sim y$ in G , and $(\tilde{\mathbf{G1}})$ holds. Suppose $e = e^*$. Let $(xx', yy') \in \Delta(\alpha)$ be as in $(\tilde{\mathbf{G1}})$. If $x \not\sim y$ in G , then $\mu_{\langle 1 \rangle}(xx', yy') = 0$. Moreover, $\Delta(\mu_{\langle 1 \rangle}^+)(xx', yy') = 0$, since each summand

$$(\chi_+ - \chi_-)(xx') (\bar{\psi}_+ - \bar{\psi}_-)(yy')$$

in $\Delta(\mu_{\langle 1 \rangle}^+)(xx', yy')$ is zero by (4A), (4F), (4G), or (4H). Thus $E_{\langle 1 \rangle}(\mu_{\langle 1 \rangle}^+)(xx', yy') = 0$ by (6.1), and $(\tilde{\mathbf{G1}})$ holds. If $x \sim y$ in G , then $n = p + 1$ or p ; otherwise $G_x > G_x^+$ and $x \sim y$ in G^+ . In this case $(G, e)^\vee = \{\chi_1, \chi_2, \dots, \chi_p\}$, where χ_i is parametrized by the partition λ_i of n with a p -hook of leg length $i - 1$. So $\text{Res}_{G^+}(\chi_i) = \chi_i^+$ is irreducible for $1 \leq i \leq p^* - 1$, $\text{Res}_{G^+}(\chi_{p^*}) = \chi_{p^*,+} + \chi_{p^*,-}$ is reducible, and

$$(G^+, e^+)^\vee = \{\chi_1^+, \chi_2^+, \dots, \chi_{p^*-1}^+, \chi_{p^*,+}, \chi_{p^*,-}\}.$$

$(H, f)^\vee$ may be identified with $Y^\vee = \{\xi_1, \xi_2, \dots, \xi_p\}$. So $\text{Res}_{H^+}(\xi_i) = \xi_i^+$ is irreducible for $1 \leq i \leq p^* - 1$, $\text{Res}_{H^+}(\xi_{p^*}) = \xi_{p^*,+} + \xi_{p^*,-}$ is reducible, and

$$(H^+, f^+)^\vee = \{\xi_1^+, \xi_2^+, \dots, \xi_{p^*-1}^+, \xi_{p^*,+}, \xi_{p^*,-}\}.$$

The isometries μ and μ^+ are

$$\begin{aligned} \chi_i &\mapsto (-1)^{i+1} \xi_i \quad \text{for } i \neq p^*, & \chi_{p^*} &\mapsto (-1)^{p^*} \xi_{p^*} \\ \chi_i^+ &\mapsto (-1)^{i+1} \xi_i^+ \quad \text{for } i \neq p^* & \chi_{p^*,\pm} &\mapsto -\epsilon_p \xi_{p^*,\pm}, \end{aligned}$$

A direct calculation gives $(\tilde{\mathbf{G1}})$.

Suppose $(zx', zy') \in \Delta(\alpha)$, where $z \in D, x' \in G_{z,p'}, y' \in H_{z,p'}$. If $x' \notin G^+$ and $y' \notin H^+$, then

$$(6.2) \quad E_{\langle 1 \rangle}(\mu_{\langle 1 \rangle}^+)(zx', zy') = E_{\langle z \rangle}(\mu_{\langle z \rangle}^+)(x', y')$$

by (6.1), since $\mu_{\langle 1 \rangle}(zx', zy') = \mu_{\langle z \rangle}(x', y')$. If $x' \in G^+$ and $y' \in H^+$, then (6.2) is just $\mu_{\langle 1 \rangle}^+(zx', zy') = \mu_{\langle z \rangle}^+(x', y')$. Thus $(\tilde{\mathbf{G}}2)$ holds.

We now suppose $n = 6$. Then $G^+ \simeq A(6) \simeq \text{PSL}(2, 9)$, $\tilde{G} = \text{Aut } G^+ \simeq \text{P}\Gamma\text{L}(2, 9)$, and $p = 3$ or 5 . Let G_1, G_2, G_3 be the maximal subgroups of \tilde{G} containing G^+ , labeled so that

$$G_1 \simeq S(6), \quad G_2 \simeq \text{PGL}(2, 9), \quad G_3 \simeq M_{10}.$$

We follow Atlas notation [7, page 5] for the character table of \tilde{G} :

TABLE (6.3). \tilde{G}

360	8	9	9	4	5	5	24	24	4	3	3	10	4	4	5	5	2	4	4			
1A	2A	3A	3B	4A	5A	B*	2B	2C	4B	6A	6B	2D	8A	B*	10A	B*	4C	8C	D**			
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
5	1	2	-1	-1	0	0	3	-1	1	0	-1	•	0	0	0	0	0	0	•	0	0	0
5	1	-1	2	-1	0	0	-1	3	1	-1	0	•	0	0	0	0	0	0	•	0	0	0
8	0	-1	-1	0	-b5	*	•	0	0	0	0	0	2	0	0	b5	*	•	0	0	0	0
8	0	-1	-1	0	*-b5	•	0	0	0	0	0	0	2	0	0	*	b5	•	0	0	0	0
9	1	0	0	1	-1	-1	3	3	-1	0	0	0	-1	1	1	-1	-1	1	-1	-1	-1	-1
10	-2	1	1	0	0	0	2	-2	0	-1	1	0	r2	-r2	0	0	0	0	i2	-i2	-i2	-i2

The columns correspond to conjugacy classes C of \tilde{G} , the first two entries in each column being $|C_{G^+}(g)|$ for g in C and the Atlas label of the class. Also $b5 = \frac{1}{2}(-1 + \sqrt{5})$, $r2 = \sqrt{2}$, $i2 = i\sqrt{2}$.

The natural inclusion of $\tilde{H} = N_{\tilde{G}}(D)$ in \tilde{G} induces an isomorphism $\alpha: \tilde{G}/G^+ \rightarrow \tilde{H}/H^+$ since $\tilde{G} = \tilde{H}G^+$. Also $H^+ \simeq 3^2: 4$, $\tilde{H} \simeq 3^2: [2^4]$ if $p = 3$, and $H^+ \simeq D_{10}$, $\tilde{H} \simeq 10: 4$ if $p = 5$. Let $H_i = N_{G_i}(D)$, so that

$$\begin{aligned} H_1 \simeq 3^2: D_8, \quad H_2 \simeq 3^2: 8, \quad H_3 \simeq 3^2: Q_8 & \text{ if } p = 3, \\ H_1 \simeq 5: 4, \quad H_2 \simeq D_{20}, \quad H_3 \simeq 5: 4 & \text{ if } p = 5. \end{aligned}$$

Let α_i be the restriction of α to G_i/G^+ for $i = 1, 2, 3$, and set

$$\Delta(\alpha_i) = \{(x, y) : x \in G_i, y \in H_i, \alpha_i(xG^+) = yH^+\}.$$

We verify the compatibility conditions $(\tilde{\mathbf{I}}1)$, $(\tilde{\mathbf{G}}1)$, and $(\tilde{\mathbf{G}}2)$ for the two cases $p = 3$ and $p = 5$.

Suppose $p = 3$. Then $D \simeq 3^2$, $H^+ \simeq 3^2: 4$, and $\tilde{H} \simeq 3^2: [2^4]$. The characters 1, $5_1, 5_2, 8_1, 8_2, 10$ in $(G^+, e^+)^\vee$ have stabilizers $\tilde{G}, G_1, G_1, G_2, \tilde{G}$ in \tilde{G} . Only the characters 1, $5_1, 5_2, 8_1, 8_2$ extend to their stabilizers. Let $E(1), E(5_1), E(5_2), E(8_1), E(8_2)$ be the extensions given in (6.3). The following is the character table of \tilde{H} :

TABLE (6.4). \tilde{H} , $p = 3$

36	4	4	4	9	9	6	6	3	3	4	4	4	4	2	2
1A	2A	4A	4A	3A	3B	2B	2C	6A	6B	8A	8A	8B	8B	4C	4C
1	1	1	1	1	1	:	1	1	1	1	:	1	1	1	1
1	1	-1	-1	1	1	:	1	-1	1	-1	:	i	$-i$	i	$-i$
1	-1	i	$-i$	1	1	•	0	0	0	0	:	z_8	z_8^7	z_8^5	z_8^3
1	-1	$-i$	i	1	1	•	0	0	0	0	:	z_8^7	z_8	z_8^3	z_8^5
4	0	0	0	1	-2	:	0	2	-1	0	•	0	0	0	0
4	0	0	0	-2	1	:	2	0	0	-1	•	0	0	0	0

The columns correspond to conjugacy classes C of \tilde{H} , the first two entries being $|C_{H^+}(h)|$ for h in C and the Atlas name of the \tilde{G} -class containing C . The characters $1_1, 1_2, 1_3, 1_4, 4_1, 4_2$ in $(H^+, f^+)^\vee$ have stabilizers $\tilde{H}, \tilde{H}, H_2, H_2, H_1, H_1$ in \tilde{H} . Only the characters $1_1, 1_3, 1_4, 4_1, 4_2$ extend to their stabilizers. Let $E(1_1), E(1_3), E(1_4), E(4_1), E(4_2)$ be the extensions given in (6.4). The isometry $\mu^+ = (J^+, \epsilon^+)$

$$1 \mapsto 1_1, \quad 10 \mapsto 1_2, \quad 5_1 \mapsto -4_2, \quad 5_2 \mapsto -4_1, \quad 8_1 \mapsto -1_3, \quad 8_2 \mapsto -1_4$$

is clearly stable under $\Delta(\alpha)$.

Let χ^+ and ψ^+ be characters in $(G^+, e^+)^\vee$ and $(H^+, f^+)^\vee$ corresponding under J^+ , and set

$$E(\chi^+ \times \overline{\psi^+}) = \begin{cases} E(\chi^+) \times \overline{E(\psi^+)}, \\ \text{sgn}_{\Delta(\alpha_1)}(E(\chi^+) \times \overline{E(\psi^+)}) \end{cases}$$

according as $(\chi^+, \psi^+) = (1, 1_1), (8_1, 1_3), (8_2, 1_4)$ or $(\chi^+, \psi^+) = (5_1, 4_2), (5_2, 4_1)$. Here the characters on the right-hand side are to be restricted to the stabilizer of $\chi^+ \times \overline{\psi^+}$ in $\Delta(\alpha)$. The character $10 \times 1_2$ extends to a character $E(10 \times 1_2)$ of $\Delta(\alpha)$. Indeed, 10 and 1_2 extend to characters $E_1(10)$ and $E_1(1_2)$ of G_1 and H_1 , where $E_1(10)$ and $E_1(1_2)$ are the extensions given in Tables (6.3) and (6.4). Then

$$E_1(10 \times 1_2) = \text{Res}_{\Delta(\alpha_1)}^{G_1 \times H_1}(E_1(10) \times E_1(1_2))$$

is $\Delta(\alpha)$ -stable, so that $E_1(10 \times 1_2)$ extends to a character $E(10 \times 1_2)$ of $\Delta(\alpha)$. The assignment $\chi^+ \times \overline{\psi^+} \mapsto E(\chi^+ \times \overline{\psi^+})$ is $\Delta(\alpha)$ -equivariant, and we may define

$$E(\mu^+) = \sum_{\chi^+ \in (G^+, e^+)^\vee} \epsilon^+(\chi^+) E(\chi^+ \times \overline{\psi^+}).$$

$E(\mu^+)(x, y)$ satisfies conditions **(I1.3a)** and **(I1.3b)** for $(x, y) \in \Delta(\alpha) - (G^+ \times H^+)$. Indeed, it suffices to take $(x, y) \in \Delta(\alpha_1) - (G^+ \times H^+)$, since x in $\tilde{G} - G^+$ commutes with a non-trivial 3-element only if $x \in G_1$.

Let $Q = \langle z \rangle$ be a non-trivial cyclic subgroup of D . Then

$$G_Q^+ = H_Q^+ = Q \times Q' \simeq C_3 \times C_3, \quad \tilde{G}_Q = \tilde{H}_Q = Q \times Q'\langle t \rangle \simeq C_3 \times S_3,$$

where t inverts Q' . In particular, $\tilde{G}_Q \leq G_1$ and $\tilde{H}_Q \leq H_1$. Moreover, $e_Q^+ = f_Q^+ = 1, \mu_Q^+ = \sum_{\zeta \in (Q \times Q')^\vee} (\zeta \times \bar{\zeta})$ is $\Delta(\alpha_Q)$ -stable, and $\mu_Q^+(1, 1) = 9$. If $\zeta^t \neq \zeta$, then $Q' \not\leq \ker \zeta$ and $\Delta(\alpha_Q)_{\zeta \times \bar{\zeta}} = G_Q^+ \times H_Q^+$. If $\zeta^t = \zeta$, then $Q' \leq \ker \zeta$ and $\Delta(\alpha_Q)_{\zeta \times \bar{\zeta}} = \Delta(\alpha_Q)$. In this case $\zeta \times \bar{\zeta}$ extends to a character $E_Q(\zeta \times \bar{\zeta})$ containing $(Q' \times Q')\langle (t, t) \rangle$ in its kernel. It follows easily that conditions **(I1.3a)** and **(I1.3b)** hold. Moreover, $E_Q(\mu_Q^+)(t, t) = 3$, so that **(G1)** and **(G2)** also hold.

Suppose $p = 5$. Then $D \simeq 5$, $H^+ \simeq D_{10}$, and $\tilde{H} \simeq 10 : 4$. The characters $1, 9, 8_1, 8_2$ in $(G^+, e^+)^\vee$ have stabilizers $\tilde{G}, \tilde{G}, G_2, G_2$ in \tilde{G} and extend to their stabilizers. Let $E(1), E(9), E(8_1), E(8_2)$ be the extensions given in (6.3). The following is the character table of \tilde{H} :

TABLE (6.5). $\tilde{H}, p = 5$

10	2	5	5	2	2	10	2	5	5	2	2				
1A	2A	5A	B*	4B	4B	2D	2A	10A	B*	4C	4C				
1	1	1	1	:	1	1	:	1	1	1	:	1	1		
1	-1	1	1	:	i	$-i$:	1	-1	1	:	i	$-i$		
2	0	$b5$	$*$	\bullet	0	0	:	-2	0	$-b5$:	$*$	\bullet	0	0
2	0	$*$	$b5$	\bullet	0	0	:	-2	0	$*$:	$-b5$	\bullet	0	0

The characters $1_1, 1_2, 2_1, 2_2$ in $(H^+, f^+)^\vee$ have stabilizers $\tilde{H}, \tilde{H}, H_2, H_2$ in \tilde{H} and extend to their stabilizers. Let $E(1_1), E(1_2), E(2_1), E(2_2)$ be the extensions given in (6.5). The isometry $\mu^+ = (J_{\mu^+}, \epsilon_{\mu^+})$ maps

$$1 \mapsto 1_1, \quad 9 \mapsto -1_2, \quad 8_1 \mapsto -2_1, \quad 8_2 \mapsto -2_2.$$

μ^+ is clearly $\Delta(\alpha)$ -stable, and

$$E(\mu^+) = E(1) \times E(1_1) - E(9) \times E(1_2) - E(8_1) \times E(2_1) - E(8_2) \times E(2_2),$$

where the right-hand side is suitably interpreted on $\Delta(\alpha)$. Since $\Delta(\alpha_1) - (G \times H)$ and $\Delta(\alpha_3) - (G \times H)$ contain no elements centralizing a 5-element, the conditions **(I1.3a)** and **(I1.3b)** on $E(\mu^+)$ need only be checked on (g, h) in $\Delta(\alpha_2)$. This is straightforward.

The only non-trivial cyclic subgroup of D is D itself. Then

$$G_D^+ = H_D^+ = D, \quad \tilde{G}_D = \tilde{H}_D = D \times \langle t \rangle \simeq C_{10}.$$

In particular, $\tilde{G}_D \leq G_2$ and $\tilde{H}_D \leq H_2$. Moreover, $e_D^+ = f_D^+ = 1$, μ_D^+ is the identity isometry, μ_D^+ is $\Delta(\alpha_D)$ -stable, and $\mu_D^+(1, 1) = 5$. We may choose $E_Q(\mu_Q^+)$ as $\sum_{\zeta} \zeta(\zeta \times \bar{\zeta})$, where ζ runs over the characters of $D \times \langle t \rangle$ with $\langle t \rangle$ in their kernels. Then $E_D(\mu_D^+)(t, t) = 5$. The conditions **(I1)**, **(G1)**, **(G2)** now follow.

7. THE ISOTYPY CONJECTURE FOR ALTERNATING GROUPS FOR $p = 2$

Let $G = S(\Pi) = S_n$ and $G^+ = A(\Pi) = A_n$ be the symmetric group and alternating group on the set Π of size n , and let $p = 2$. The 2-blocks of \mathcal{OG} are self-dual since 2-cores are self-dual. Thus each block of \mathcal{OG}^+ is covered by a unique block of \mathcal{OG} . Suppose e is a block of \mathcal{OG} parametrized by the 2-core λ_0 . If e has defect 0, then $\text{Res}_{G^+}^G(\chi_{\lambda_0}) = \chi_{\lambda_{0,+}} + \chi_{\lambda_{0,-}}$ and e covers two blocks of \mathcal{OG}^+ of defect 0, namely the blocks e_{\pm} containing $\chi_{\lambda_{0,\pm}}$. If e has positive defect, then $(G, e)^\vee$ contains characters χ_{λ} which do not split on G^+ , and e covers a unique block e^+ of \mathcal{OG}^+ . Thus a block of \mathcal{OG} of positive defect covers a unique G -stable block of \mathcal{OG}^+ . Conversely a G -stable block of \mathcal{OG}^+ is covered by a unique block of \mathcal{OG} of positive defect.

We consider a block of \mathcal{OG}^+ with abelian defect group. If the block is not G -stable, then it has defect 0 and is e_{\pm} for some block e of \mathcal{OG} of defect 0. If the block is G -stable, then it is e^+ for a unique block e of \mathcal{OG} of positive defect. Let

w be the weight of e and let $\sum_{i \geq 1} 2^i w_i$ be the 2-adic expansion of $2w$. Then e has defect group

$$D = \prod_{i \geq 1} (T_i)^{w_i},$$

where $T_i = Z_2 \wr Z_2 \wr \dots \wr Z_2$ (i times) is a Sylow 2-subgroup of S_{2^i} , and e^+ has defect group $D^+ = D \cap G^+ < D$.

Lemma (7A). *Suppose D^+ is abelian. Then either $D^+ = 1$ and $D = T_1$, or $D^+ \simeq Z_2 \times Z_2$ and $D = T_2$.*

Proof. T_i is non-abelian for $i \geq 2$ and $[T_i, T_i]$ is non-abelian for $i \geq 3$. Thus D is T_1, T_2 , or $T_1 \times T_2$. The case $D = T_1 \times T_2$ is impossible since D^+ would have a non-abelian section isomorphic to T_2 . Thus $D = T_1$ or T_2 . \square

Let e^+ be a block of $\mathcal{O}G^+$ with abelian defect group $D^+ > 1$, let e be the unique block of $\mathcal{O}G$ covering e^+ , and let D be a defect group of e such that $D^+ = D \cap G^+$. So (7A) implies that $D = T_2$ and $D^+ \simeq Z_2 \times Z_2$. Let $H = N_G(D^+)$ and $H^+ = N_{G^+}(D^+)$. Let λ_0 be the 2-core parametrizing e . If $|\lambda_0| = 0$, then $n = 4, G = H = S_4, G^+ = H^+ = A_4$, and the isotopy conjecture holds trivially. If $|\lambda_0| = 1$, then $n = 5, G = S_5, G^+ = A_5, H = S_4, H^+ = A_4$, and the isotopy conjecture holds by [4, Appendice A1.1]. So we suppose $|\lambda_0| \geq 3$ and $n \geq 7$. Let $\Pi = \Pi_0 \cup \Pi_1$, where Π_0 and Π_1 are the fixed-point set and the support of D . Let $H_0 = S(\Pi_0)$ and $H_1 = S(\Pi_1) = S_4$. Then $H = H_0 \times H_1, H^+ = (H_0 \times H_1)^+$, and

$$\begin{aligned} C_G(D^+) &= C_H(D^+) = H_0 \times D^+, \\ C_{G^+}(D^+) &= C_{H^+}(D^+) = H_0^+ \times D^+. \end{aligned}$$

Let χ_{λ_0} be the character of H_0 parametrized by λ_0 , let 1_{H_1} be the 1-character of H_1 , and let f be the block of $\mathcal{O}H$ containing $\chi_{\lambda_0} \times 1_{H_1}$. Then f covers a unique block f^+ of $\mathcal{O}H^+$ since $\text{Res}_{H^+}^H(\chi_{\lambda_0} \times 1_{H_1})$ is irreducible, and f and f^+ are the Brauer correspondents of e and e^+ . We take as maximal e^+ -subpair $(D^+, e_{D^+}^+)$ and maximal f^+ -subpair $(D^+, f_{D^+}^+)$ the subpair $(D^+, e_{\lambda_0,+})$, where $e_{\lambda_0,+}$ is the block of $\mathcal{O}(H_0^+ \times D^+)$ containing $\chi_{\lambda_0,+} \times 1_{D^+}$.

The partitions λ with 2-core λ_0 have the following 2-quotients:

$$(2, \emptyset), \quad (1^2, \emptyset), \quad (1, 1), \quad (\emptyset, 2), \quad (\emptyset, 1^2).$$

We label the χ_λ by the 2-quotients of λ as $\chi_{(2,\emptyset)}, \chi_{(1^2,\emptyset)}, \chi_{(1,1)}, \chi_{(\emptyset,2)}, \chi_{(\emptyset,1^2)}$. In particular, $\chi_{(2,\emptyset)}^* = \chi_{(\emptyset,1^2)}, \chi_{(1^2,\emptyset)}^* = \chi_{(\emptyset,2)}, \chi_{(1,1)}^* = \chi_{(1,1)}$. So $(G^+, e^+)^{\vee}$ consists of the four characters

$$\chi_{(2,\emptyset)}^+ = \text{Res}_{G^+}^G(\chi_{(2,\emptyset)}), \quad \chi_{(1^2,\emptyset)}^+ = \text{Res}_{G^+}^G(\chi_{(1^2,\emptyset)}), \quad \chi_{(1,1),\pm},$$

where $\chi_{(1,1),\pm}$ are the constituents of $\text{Res}_{G^+}^G(\chi_{(1,1)})$.

The irreducible characters in $(H, f)^{\vee}$ are

$$\chi_{\lambda_0} \times \chi_{(4)}, \quad \chi_{\lambda_0} \times \chi_{(3,1)}, \quad \chi_{\lambda_0} \times \chi_{(2^2)}, \quad \chi_{\lambda_0} \times \chi_{(2,1^2)}, \quad \chi_{\lambda_0} \times \chi_{(1^4)},$$

where the second factors are the irreducible characters of $S(\Pi_1)$. Since the partitions $(4), (3, 1), (2^2), (2, 1^2), (1^4)$ have 2-quotients $(2, \emptyset), (1^2, \emptyset), (1, 1), (\emptyset, 2), (\emptyset, 1^2)$, we also label these characters in $(H, f)^{\vee}$ as $\psi_{(2,\emptyset)}, \psi_{(1^2,\emptyset)}, \psi_{(1,1)}, \psi_{(\emptyset,2)}, \psi_{(\emptyset,1^2)}$. In

particular, $\psi_{(2,\emptyset)}^* = \psi_{(\emptyset,1^2)}$, $\psi_{(1^2,\emptyset)}^* = \psi_{(\emptyset,2)}$, $\psi_{(1,1)}^* = \psi_{(1,1)}$. So $(H^+, f^+)^\vee$ consists of the four characters

$$\psi_{(2,\emptyset)}^+ = \text{Res}_{H^+}^H(\psi_{(2,\emptyset)}), \quad \psi_{(1^2,\emptyset)}^+ = \text{Res}_{H^+}^H(\psi_{(1^2,\emptyset)}), \quad \psi_{(1,1),\pm}$$

where $\psi_{(1,1),\pm}$ are the constituents of $\text{Res}_{H^+}^H(\psi_{(1,1)})$.

Let $\mu^+ = (J^+, \epsilon^+)$ be the isometry in $\mathcal{R}_K((G^+, e^+), (H^+, f^+))$ given by

$$\chi_{(2,\emptyset)}^+ \mapsto \psi_{(2,\emptyset)}^+, \quad \chi_{(1^2,\emptyset)}^+ \mapsto -\psi_{(1^2,\emptyset)}^+, \quad \chi_{(1,1),\pm} \mapsto -\psi_{(1,1),\pm}$$

Each non-trivial cyclic subgroup $P = \langle x \rangle$ of D^+ is $Z(D)^t$ for some t in $A(\Pi_1)$. Thus

$$G_x = H_x = S(\Pi_0) \times D^t, \quad G_x^+ = H_x^+ = (S(\Pi_0) \times D^t)^+$$

and the subpairs $(\langle x \rangle, e_x^+)$ and $(\langle x \rangle, f_x^+)$ coincide. We take $\mu_x^+ = (J_x^+, \epsilon_x^+)$ as the identity isometry. The diagram

$$(7.1) \quad \begin{array}{ccc} \text{CF}(G^+, e^+, K) & \xrightarrow{\text{R}_{\mu_{\langle 1 \rangle}^+}} & \text{CF}(H^+, f^+, K) \\ \downarrow d_{G^+}^{(x, e_x^+)} & & \downarrow d_{H^+}^{(x, f_x^+)} \\ \text{CF}_{P'}(G_P^+, e_x^+, K) & \xrightarrow{\text{R}_{\mu_x^+, p'}} & \text{CF}_{P'}(H_P^+, f_x^+, K) \end{array}$$

is then commutative. Indeed, it suffices to show that

$$\chi^+(xy) = \text{R}_{\mu_{\langle 1 \rangle}^+}(\chi^+)(xy)$$

for all χ^+ in $(G^+, e^+)^\vee$ and all y in $H_{x,2^r}^+$. Such y 's lie in $A(\Pi_0)$. Now

$$\begin{aligned} \chi_{(2,\emptyset)}^+(xy) &= \chi_{(2,\emptyset)}(xy) = \chi_{\lambda_0}(y), \\ \chi_{(1^2,\emptyset)}^+(xy) &= \chi_{(1^2,\emptyset)}(xy) = \chi_{\lambda_0}(y), \end{aligned}$$

by the Murnaghan-Nakayama formula. On the other hand,

$$\begin{aligned} \psi_{(2,\emptyset)}^+(xy) &= \psi_{(2,\emptyset)}(xy) = \chi_{\lambda_0}(y), \\ \psi_{(1^2,\emptyset)}^+(xy) &= \psi_{(1^2,\emptyset)}(xy) = -\chi_{\lambda_0}(y), \end{aligned}$$

since $\chi_{(4)}(x) = 1$ and $\chi_{(3,1)}(x) = -1$. Lastly,

$$\chi_{(1,1),\pm}(xy) = \frac{1}{2}\chi_{(1,1)}(xy) = -\chi_{\lambda_0}(y).$$

The first equality holds since xy has cycle type unequal to $\eta(\lambda)$, where λ is the partition with 2-core λ_0 and 2-quotient $(1, 1)$. The second equality holds by the Murnaghan-Nakayama formula. We may suppose with suitable labeling that

$$\begin{aligned} \psi_{(1,1),+}(xy) &= \chi_{\lambda_0,+}(y)\chi_{(2^2),+}(x) + \chi_{\lambda_0,-}(y)\chi_{(2^2),-}(x), \\ \psi_{(1,1),-}(xy) &= \chi_{\lambda_0,+}(y)\chi_{(2^2),-}(x) + \chi_{\lambda_0,-}(y)\chi_{(2^2),+}(x). \end{aligned}$$

Both equal $\chi_{\lambda_0}(y)$, since $\chi_{(2^2),\pm}(x) = 1$. Thus (7.1) is commutative. It now follows as in §5 that e^+ and f^+ are isotypic blocks.

REFERENCES

1. J. L. Alperin, M. Broué, *Local methods in block theory*, Ann. of Math. **110** (1979), 143–157. MR **80f**:20010
2. M. Broué, *Radical, hauteurs, p -sections et blocs*, Ann. of Math. **107** (1978), 89–107. MR **80a**:20014
3. M. Broué, *Les ℓ -blocs des groupes $GL(n, q)$ et $U(n, q^2)$ et leurs structures locales*, Astérisque **133–134** (1986), 159–188. MR **87e**:20021
4. M. Broué, *Isométries parfaites, types de blocs, catégories dérivées*, Astérisque **181–182** (1990), 61–91. MR **91i**:20006
5. M. Broué, L. Puig, *Characters and local structure in G -algebras*, J. Algebra **63** (1980), 306–317. MR **81j**:20021
6. O. Celik, A. Kerber, H. Pahlings, *Zur Darstellungstheorie gewisser Verallgemeinerungen der Serien von Weyl-Gruppen.*, Mitt. Math. Sem. Giessen **119** (1975), 15–90. MR **54**:7600
7. J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson, *An ATLAS of finite groups.*, Oxford University Press, Oxford, 1985. MR **88g**:20025
8. M. Enguehard, *Isométries parfaites dans les groupes symétriques*, Rapport de recherche du LMENS, DMI, Ecole Normale Supérieure **88–10** (1988), 1–19. cf. MR **91d**:20015
9. M. Enguehard, *Isométries parfaites entres blocs de groupes lineaires ou unitaires*, Math. Zeit. **204** (1991), 1–24. MR **92c**:20017
10. P. Fong, M. E. Harris, *On perfect isometries and isotopies in finite groups*, Invent. Math. **114** (1993), 139–191. MR **94i**:20012
11. G. Frobenius, *Über die Charaktere der alternirenden Gruppe*, Sitzber. Preuß. Akad. Wiss. (1901), 303–315.
12. G. James, A. Kerber, *The representation theory of the symmetric group*, Addison-Wesley, Reading, MA, 1981. MR **83k**:20003
13. L. Puig, *Local block theory in p -solvable groups*, Proc. Sympos. Pure Math., vol. 37, Amer. Math. Soc., Providence, RI, 1979, pp. 385–388. MR **82c**:20045
14. R. Rouquier, *Sur les blocs à groupe de défaut abélien dans les groupes symétriques et sporadiques*, J. Algebra **168** (1994), 648–694. MR **95h**:20015
15. J. Stembridge, *On the eigenvalues of representations of reflection groups and wreath products*, Pacific. J. of Math. **140** (1989), 353–396. MR **91a**:20022

DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, ILLINOIS 60680

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55455