

TURNPIKE PROPERTY FOR EXTREMALS
OF VARIATIONAL PROBLEMS
WITH VECTOR-VALUED FUNCTIONS

A. J. ZASLAVSKI

ABSTRACT. In this paper we study the structure of extremals $\nu: [0, T] \rightarrow R^n$ of variational problems with large enough T , fixed end points and an integrand f from a complete metric space of functions. We will establish the turnpike property for a generic integrand f . Namely, we will show that for a generic integrand f , any small $\varepsilon > 0$ and an extremal $\nu: [0, T] \rightarrow R^n$ of the variational problem with large enough T , fixed end points and the integrand f , for each $\tau \in [L_1, T - L_1]$ the set $\{\nu(t): t \in [\tau, \tau + L_2]\}$ is equal to a set $H(f)$ up to ε in the Hausdorff metric. Here $H(f) \subset R^n$ is a compact set depending only on the integrand f and $L_1 > L_2 > 0$ are constants which depend only on ε and $|\nu(0)|, |\nu(T)|$.

1. INTRODUCTION

In this paper we analyse the structure of optimal solutions of the variational problem

$$(P) \quad \int_0^T f(z(t), z'(t)) dt \rightarrow \min, \quad z(0) = x, \quad z(T) = y,$$

$z: [0, T] \rightarrow R^n$ is an absolutely continuous function

where $T > 0$, $x, y \in R^n$ and $f: R^{2n} \rightarrow R^1$ is an integrand.

An optimal solution $\nu: [0, T] \rightarrow R^n$ of the variational problem (P) always depends on the integrand f and on x, y, T . We say that the integrand f has the *turnpike property* if for large enough T the dependence on x, y, T is not essential. Namely, for any $\varepsilon > 0$ there exist constants $L_1 > L_2 > 0$ which depend only on $|x|, |y|, \varepsilon$ such that for each $\tau \in [L_1, T - L_1]$ the set

$$\{\nu(t): t \in [\tau, \tau + L_2]\}$$

is equal to a set $H(f)$ up to ε in the Hausdorff metric where $H(f) \subset R^n$ is a compact set depending only on the integrand f .

More formally we say that an integrand $f = f(x, u) \in C(R^{2n})$ has the *turnpike property* if there exists a compact set $H(f) \subset R^n$ such that for each bounded set $K \subset R^n$ and each $\varepsilon > 0$ there exist numbers $L_1 > L_2 > 0$ such that for each

Received by the editors September 29, 1995 and, in revised form, November 18, 1996.

1991 *Mathematics Subject Classification*. Primary 49J99, 58F99.

Key words and phrases. Good function, turnpike property, representation formula, minimal long-run average cost growth rate.

©1999 American Mathematical Society

$T \geq 2L_1$, each $x, y \in K$ and an optimal solution $\nu: [0, T] \rightarrow R^n$ for the variational problem (P) the relation

$$\text{dist}(H(f), \{\nu(t): t \in [\tau, \tau + L_2]\}) \leq \varepsilon$$

holds for each $\tau \in [L_1, T - L_1]$. (Here $\text{dist}(\cdot, \cdot)$ is the Hausdorff metric.)

The turnpike property is well known in mathematical economics. It was studied by many authors for optimal trajectories of a von Neumann-Gale model determined by a superlinear set-valued mapping (see Makarov and Rubinov [14] and the survey [16]) and for optimal trajectories of convex autonomous systems (see Carlson, Haurie and Leizarowitz [7, Ch. 4.6].) In the control theory the turnpike property was established by Artstein and Leizarowitz [1] for a tracking periodic problem. In all these cases we have an optimal control problem with a convex cost function and a convex set of trajectories. Asymptotic turnpike properties for optimal control problems with infinite time horizon were studied by Brock and Haurie [4], Carlson [5], Carlson, Haurie and Jabrane [6], Leizarowitz [10] and Zaslavski [19].

Our goal is to show that the turnpike property is a general phenomenon which holds for a large class of variational problems with vector-valued functions. We consider the complete metric space of integrands $\overline{\mathfrak{M}}_k$ (k is a nonnegative integer) described below and establish the existence of a set $\mathcal{F} \subset \overline{\mathfrak{M}}_k$ which is a countable intersection of open everywhere dense sets in $\overline{\mathfrak{M}}_k$ and such that each integrand $f \in \mathcal{F}$ has the turnpike property.

Moreover we show that the turnpike property holds for approximate solutions of variational problems with a generic integrand f and that the turnpike phenomenon is stable under small perturbations of a generic integrand f .

A better understanding of the general nature of the turnpike phenomenon was achieved by our recent study of discrete-time control systems [17, 18] for which we established a weak version of the turnpike property. More recently in Zaslavski [20] employing the reduction to finite rewards by Leizarowitz [11, 12] and the representation formula by Leizarowitz and Mizel [13] an analogous result was established for optimal solutions of the variational problem (P) with $x, y \in R^n$, large enough T and a generic integrand f belonging to the space of functions \mathfrak{A} described below.

In the weak version of the turnpike property established in [20] for an optimal solution of the problem (P) with $x, y \in R^n$, large enough T and a generic integrand $f \in \mathfrak{A}$ the relation

$$\text{dist}(H(f), \{\nu(t): t \in [\tau, \tau + L_2]\}) \leq \varepsilon$$

with L_2 which depends on ε and $|x|, |y|$ and a compact set $H(f) \subset R^n$ depending only on the integrand f , holds for each $\tau \in [0, T] \setminus E$ where $E \subset [0, T]$ is a measurable subset such that the Lebesgue measure of E does not exceed a constant which depends on ε and $|x|, |y|$.

The turnpike property which will be established in the present work guarantees that we may take $E = [0, L_1] \cup [T - L_1, T]$ where $L_1 > 0$ is a constant which depends on ε and $|x|, |y|$.

Denote by $|\cdot|$ the Euclidean norm in R^n and denote by \mathfrak{A} the set of continuous functions $f: R^n \times R^n \rightarrow R^1$ which satisfy the following assumptions:

- (A)(i) for each $x \in R^n$ the function $f(x, \cdot): R^n \rightarrow R^1$ is convex;
- (ii) $f(x, u) \geq \sup\{\psi(|x|), \psi(|u|)|u|\} - a$ for each $(x, u) \in R^n \times R^n$ where $a > 0$ is a constant and $\psi: [0, \infty) \rightarrow [0, \infty)$ is an increasing function such that $\psi(t) \rightarrow +\infty$ as $t \rightarrow \infty$ (here a and ψ are independent on f);

(iii) for each $M, \varepsilon > 0$ there exist $\Gamma, \delta > 0$ such that

$$|f(x_1, u_1) - f(x_2, u_2)| \leq \varepsilon \sup\{f(x_1, u_1), f(x_2, u_2)\}$$

for each $u_1, u_2, x_1, x_2 \in R^n$ which satisfy

$$|x_i| \leq M, |u_i| \geq \Gamma \ (i = 1, 2), \quad \sup\{|x_1 - x_2|, |u_1 - u_2|\} \leq \delta.$$

It is an elementary exercise to show that an integrand $f = f(x, u) \in C^1(R^{2n})$ belongs to \mathfrak{A} if f satisfies assumptions (Ai), (Aii) with a constant $a > 0$ and a function $\psi: [0, \infty) \rightarrow [0, \infty)$ and there exists an increasing function $\psi_0: [0, \infty) \rightarrow [0, \infty)$ such that for each $x, u \in R^n$

$$\sup\{|\partial f / \partial x(x, u)|, |\partial f / \partial u(x, u)|\} \leq \psi_0(|x|)(1 + \psi(|u|)|u|).$$

For the set \mathfrak{A} we consider the uniformity which is determined by the following base

$$E(N, \varepsilon, \lambda) = \{(f, g) \in \mathfrak{A} \times \mathfrak{A}: |f(x, u) - g(x, u)| \leq \varepsilon \ (u, x \in R^n, |x|, |u| \leq N), \\ (|f(x, u)| + 1)(|g(x, u)| + 1)^{-1} \in [\lambda^{-1}, \lambda] \ (x, u \in R^n, |x| \leq N)\}$$

where $N > 0, \varepsilon > 0, \lambda > 1$ (see Kelley [9]).

It was shown in Zaslavski [20] that the uniform space \mathfrak{A} is metrizable and complete. We consider functionals of the form

$$(1.1) \quad I^f(T_1, T_2, x) = \int_{T_1}^{T_2} f(x(t), x'(t)) dt$$

where $f \in \mathfrak{A}, 0 \leq T_1 \leq T_2 < +\infty$ and $x: [T_1, T_2] \rightarrow R^n$ is an absolutely continuous (a.c.) function.

For $f \in \mathfrak{A}, y, z \in R^n$ and numbers T_1, T_2 satisfying $0 \leq T_1 < T_2$ we set

$$(1.2) \quad U^f(T_1, T_2, y, z) = \inf\{I^f(T_1, T_2, x): x: [T_1, T_2] \rightarrow R^n \text{ is an a.c. function} \\ \text{satisfying } x(T_1) = y, x(T_2) = z\}.$$

It is easy to see that $-\infty < U^f(T_1, T_2, y, z) < +\infty$ for each $f \in \mathfrak{A}$, each $y, z \in R^n$ and all numbers T_1, T_2 satisfying $0 \leq T_1 < T_2$.

Let $f \in \mathfrak{A}$. For any a.c. function $x: [0, \infty) \rightarrow R^n$ we set

$$(1.3) \quad J(x) = \liminf_{T \rightarrow \infty} T^{-1} I^f(0, T, x).$$

Of special interest is the *minimal long-run average cost growth rate*

$$(1.4) \quad \mu(f) = \inf\{J(x): x: [0, \infty) \rightarrow R^n \text{ is an a.c. function}\}.$$

Clearly $-\infty < \mu(f) < +\infty$. By a simple modification of the proof of Proposition 4.4 in Leizarowitz and Mizel [13] (see [20, Theorems 8.1, 8.2]) we established the representation formula

$$(1.5) \quad U^f(0, T, x, y) = T\mu(f) + \pi^f(x) - \pi^f(y) + \theta_T^f(x, y), \\ x, y \in R^n, \quad T \in (0, \infty),$$

where $\pi^f: R^n \rightarrow R^1$ is a continuous function and $(T, x, y) \rightarrow \theta_T^f(x, y) \in R^1$ is a continuous nonnegative function defined for $T > 0, x, y \in R^n$,

$$(1.6) \quad \pi^f(x) = \inf \left\{ \liminf_{T \rightarrow +\infty} [I^f(0, T, \nu) - \mu(f)T]: \nu: [0, \infty) \rightarrow R^n \right. \\ \left. \text{is an a.c. function satisfying } \nu(0) = x \right\}, \quad x \in R^n,$$

and for every $T > 0$, every $x \in R^n$ there is $y \in R^n$ satisfying $\theta_T^f(x, y) = 0$.

Here we follow Leizarowitz [11] in defining “good functions” for the infinite horizon variational problem with the integrand f .

An a.c. function $x: [0, \infty) \rightarrow R^n$ is called an (f) -good function if the function $\Phi_x^f: T \rightarrow I^f(0, T, x) - \mu(f)T, T \in (0, \infty)$ is bounded. In [20] we showed that for each $f \in \mathfrak{A}$ and each $z \in R^n$ there exists an (f) -good function $\nu: [0, \infty) \rightarrow R^n$ satisfying $\nu(0) = z$.

Propositions 1.1 and 3.2 in Zaslavski [20] imply the following result.

Proposition 1.1. *For any a.c. function $x: [0, \infty) \rightarrow R^n$ either*

$$I^f(0, T, x) - T\mu(f) \rightarrow +\infty \text{ as } T \rightarrow \infty$$

or

$$\sup\{|I^f(0, T, x) - T\mu(f)|: T \in (0, \infty)\} < \infty.$$

Moreover any (f) -good function $x: [0, \infty) \rightarrow R^n$ is bounded.

We denote $d(x, B) = \inf\{|x - y|: y \in B\}$ for $x \in R^n, B \subset R^n$. Denote by $\text{dist}(A, B)$ the Hausdorff metric for two sets $A \subset R^n, B \subset R^n$. For every bounded a.c. function $x: [0, \infty) \rightarrow R^n$ define

$$(1.7) \quad \Omega(x) = \{y \in R^n: \text{there exists a sequence } \{t_i\}_{i=0}^\infty \subset (0, \infty) \text{ for which} \\ t_i \rightarrow \infty, x(t_i) \rightarrow y \text{ as } i \rightarrow \infty\}.$$

We say that an integrand $f \in \mathfrak{A}$ has Property B if $\Omega(\nu_2) = \Omega(\nu_1)$ for all (f) -good functions $\nu_i: [0, \infty) \rightarrow R^n, i = 1, 2$.

In Zaslavski [20, Theorem 2.1] we establish the following result which describes the limit behaviour of (f) -good functions for a generic $f \in \mathfrak{A}$.

Theorem 1.1. *There exists a set $\mathcal{F} \subset \mathfrak{A}$ which is a countable intersection of open everywhere dense subsets of \mathfrak{A} and such that each $f \in \mathcal{F}$ has Property B.*

By Proposition 1.1 for each integrand $f \in \mathfrak{A}$ which has Property B there exists a compact set $H(f) \subset R^n$ such that $\Omega(\nu) = H(f)$ for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$.

Denote by \mathfrak{M} the set of all functions $f \in C^1(R^{2n})$ satisfying the following assumptions which ensure that each solution of (P) belongs to $C^2([0, T]; R^n)$:

$$\partial f / \partial u_i \in C^1(R^{2n}) \quad \text{for } i = 1, \dots, n;$$

the matrix $(\partial^2 f / \partial u_i \partial u_j)(x, u), i, j = 1, \dots, n$, is positive definite for all $(x, u) \in R^{2n}$;

$$f(x, u) \geq \sup\{\psi(|x|), \psi(|u|)|u|\} - a \quad \text{for all } (x, u) \in R^n \times R^n;$$

there exist a number $c_0 > 1$ and monotone increasing functions $\phi_i: [0, \infty) \rightarrow [0, \infty)$, $i = 0, 1, 2$, such that

$$\phi_0(t)t^{-1} \rightarrow +\infty \text{ as } t \rightarrow +\infty, \quad f(x, u) \geq \phi_0(c_0|u|) - \phi_1(|x|), \quad x, u \in R^n;$$

$$\sup\{|\partial f/\partial x_i(x, u)|, |\partial f/\partial u_i(x, u)|\} \leq \phi_2(|x|)(1 + \phi_0(|u|)),$$

$$x, u \in R^n, \quad i = 1, \dots, n.$$

It is easy to see that $\mathfrak{M} \subset \mathfrak{A}$. We will establish the following result.

Theorem 1.2. *Assume that an integrand $f \in \mathfrak{M}$ has Property B and $\varepsilon, K > 0$. Then there exists a neighborhood \mathcal{U} of f in \mathfrak{A} and numbers $M > K$, $l_0 > l > 0$, $\delta > 0$ such that for each $g \in \mathcal{U}$, each $T \geq 2l_0$ and each a.c. function $\nu: [0, T] \rightarrow R^n$ which satisfies*

$$|\nu(0)|, |\nu(T)| \leq K, \quad I^g(0, T, \nu) \leq U^g(0, T, \nu(0), \nu(T)) + \delta$$

the relation $|\nu(t)| \leq M$ holds for all $t \in [0, T]$ and

$$(1.8) \quad \text{dist}(H(f), \{\nu(t) : t \in [\tau, \tau + l]\}) \leq \varepsilon$$

for each $\tau \in [l_0, T - l_0]$. Moreover if $d(\nu(0), H(f)) \leq \delta$, then (1.8) holds for each $\tau \in [0, T - l_0]$ and if $d(\nu(T), H(f)) \leq \delta$, then (1.8) holds for each $\tau \in [l_0, T - l]$.

Let $k \geq 1$ be an integer. Denote by \mathfrak{A}_k the set of all integrands $f \in \mathfrak{A} \cap C^k(R^{2n})$. For $p = (p_1, \dots, p_{2n}) \in \{0, \dots, k\}^{2n}$ and $f \in C^k(R^{2n})$ we set

$$|p| = \sum_{i=1}^{2n} p_i, \quad D^p f = \partial^{|p|} f / \partial y_1^{p_1} \dots \partial y_{2n}^{p_{2n}}.$$

For the set \mathfrak{A}_k we consider the uniformity which is determined by the following base.

$$E(N, \varepsilon, \lambda) = \{(f, g) \in \mathfrak{A}_k \times \mathfrak{A}_k : |D^p f(x, u) - D^p g(x, u)| \leq \varepsilon$$

$$(u, x \in R^n, |x|, |u| \leq N, p \in \{0, \dots, k\}^{2n}, |p| \leq k),$$

$$|f(x, u) - g(x, u)| \leq \varepsilon (u, x \in R^n, |x|, |u| \leq N),$$

$$(|f(x, u)| + 1)(|g(x, u)| + 1)^{-1} \in [\lambda^{-1}, \lambda](x, u \in R^n, |x| \leq N)\}$$

where $N > 0$, $\varepsilon > 0$, $\lambda > 1$ (see Kelley [9]). It is easy to verify that the uniform space \mathfrak{A}_k is metrizable and complete (see [20], Section 2).

For each integer $k \geq 1$ we define $\mathfrak{M}_k = \mathfrak{M} \cap \mathfrak{A}_k$. Set

$$\mathfrak{A}_0 = \mathfrak{A}, \quad \mathfrak{M}_0 = \mathfrak{M}.$$

Let $k \geq 0$ be an integer. Denote by $\overline{\mathfrak{M}}_k$ the closure of \mathfrak{M}_k in \mathfrak{A}_k and consider the topological subspace $\overline{\mathfrak{M}}_k \subset \mathfrak{A}_k$ with the relative topology. We will establish the following result.

Theorem 1.3. *Let $q \geq 0$ be an integer. Then there exists a set $\mathcal{F}_q \subset \overline{\mathfrak{M}}_q$ which is a countable intersection of open everywhere dense subsets of $\overline{\mathfrak{M}}_q$ and such that each $f \in \mathcal{F}_q$ has Property B and the following property:*

For each $\varepsilon, K > 0$ there exist a neighborhood \mathcal{U} of f in \mathfrak{A} and numbers $M > K$, $l_0 > l > 0$, $\delta > 0$ such that for each $g \in \mathcal{U}$, each $T \geq 2l_0$ and each a.c. function $\nu: [0, T] \rightarrow R^n$ which satisfies

$$|\nu(0)|, |\nu(t)| \leq K, \quad I^g(0, T, \nu) \leq U^g(0, T, \nu(0), \nu(T)) + \delta$$

the relation $|\nu(t)| \leq M$ holds for all $t \in [0, T]$ and

$$(1.9) \quad \text{dist}(H(f), \{\nu(t) : t \in [\tau, \tau + l]\}) \leq \varepsilon$$

for each $\tau \in [l_0, T - l_0]$. Moreover if $d(\nu(0), H(f)) \leq \delta$, then (1.9) holds for each $\tau \in [0, T - l_0]$ and if $d(\nu(T), H(f)) \leq \delta$, then (1.9) holds for each $\tau \in [l_0, T - l]$.

Theorems 1.2 and 1.3 are extensions to the class of variational problems with vector-valued functions of the main result in Zaslavski [21] established for a class of one-dimensional variational problems arising in continuum mechanics which was discussed in Leizarowitz and Mizel [13] and Coleman, Marcus and Mizel [8]. In the approach used in [21] the following property of this class of one-dimensional variational problems established in Leizarowitz and Mizel [13] played a crucial role.

Property C. In the space of integrands there exists an everywhere dense subset E such that for each $f \in E$ there exists an (f) -good periodic trajectory.

It is not clear whether Property C holds in general. In Zaslavski [20] and in the present paper we develop a more general approach based on the idea that the validity of Property B implies the weak version of the turnpike property for an integrand $f \in \mathfrak{A}$ and implies the turnpike property for an integrand $f \in \mathfrak{M}$.

2. AUXILIARY RESULTS

In [20] we established the following results.

Proposition 2.1 ([20, Proposition 3.1]). *For each $f \in \mathfrak{A}$ there exists a neighborhood \mathcal{U} of f in \mathfrak{A} and a number $M > 0$ such that for each $g \in \mathcal{U}$ and each (g) -good function $x: [0, \infty) \rightarrow R^n$*

$$\limsup_{t \rightarrow \infty} |x(t)| < M.$$

Proposition 2.2 ([20, Proposition 3.2]). *Let $f \in \mathfrak{A}$ and $M_1, M_2, c > 0$. Then there exist a neighborhood \mathcal{U} of f in \mathfrak{A} and $S > 0$ such that for each $g \in \mathcal{U}$, each $T_1 \in [0, \infty)$ and each $T_2 \in [T_1 + c, \infty)$ the following property holds.*

For each $x, y \in R^n$ satisfying $|x|, |y| \leq M_1$ and each a.c. function $\nu: [T_1, T_2] \rightarrow R^n$ satisfying $\nu(T_1) = x$, $\nu(T_2) = y$, $I^g(T_1, T_2, \nu) \leq U^g(T_1, T_2, x, y) + M_2$ the following relation holds: $|\nu(t)| \leq S$ ($t \in [T_1, T_2]$).

Proposition 2.3 ([20, Proposition 3.8]). *Let $f \in \mathfrak{A}$, $0 < c_1 < c_2 < \infty$, $D, \varepsilon > 0$. Then there is a neighborhood V of f in \mathfrak{A} such that for each $g \in V$, each $T_1, T_2 \geq 0$ satisfying $T_2 - T_1 \in [c_1, c_2]$ and each a.c. function $x: [T_1, T_2] \rightarrow R^n$ satisfying $\inf\{I^f(T_1, T_2, x), I^g(T_1, T_2, x)\} \leq D$ the relation $|I^f(T_1, T_2, x) - I^g(T_1, T_2, x)| \leq \varepsilon$ holds.*

Proposition 2.4 ([20, Proposition 3.9]). *Let $f \in \mathfrak{A}$, $0 < c_1 < c_2 < \infty$, $c_3, \varepsilon > 0$. Then there exists a neighborhood V of f in \mathfrak{A} such that for each $g \in V$, each $T_1, T_2 \geq 0$ satisfying $T_2 - T_1 \in [c_1, c_2]$ and each $z, y \in R^n$ satisfying $|y|, |z| \leq c_3$ the relation $|U^f(T_1, T_2, y, z) - U^g(T_1, T_2, y, z)| \leq \varepsilon$ holds.*

Proposition 2.5 ([20, Theorem 5.1]). *Assume that $f \in \mathfrak{A}$ and there exists a compact set $H(f) \subset R^n$ such that $\Omega(\nu) = H(f)$ for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$. Let ε be a positive number. Then there exists an integer $L \geq 1$ such that for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$*

$$\text{dist}(H(f), \{\nu(t) : t \in [T, T + L]\}) \leq \varepsilon \quad \text{for all large } T.$$

Proposition 2.6 ([20, Theorem 6.1]). *Assume that $f \in \mathfrak{A}$. Then the mapping $(T_1, T_2, x, y) \rightarrow U^f(T_1, T_2, x, y)$ is continuous for $T_1 \in [0, \infty)$, $T_2 \in (T_1, \infty)$, $x, y \in R^n$.*

Proposition 2.7 ([20, Theorem 2.3]). *Assume that $f \in \mathfrak{A}$ and there exists a compact set $H(f) \subset R^n$ such that $\Omega(\nu) = H(f)$ for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$. Let ε be a positive number. Then there exist an integer $L \geq 1$ and a neighborhood \mathcal{U} of f in \mathfrak{A} such that for each $g \in \mathcal{U}$ and each (g) -good function $\nu: [0, \infty) \rightarrow R^n$*

$$\text{dist}(H(f), \{\nu(t): t \in [T, T + L]\}) \leq \varepsilon \quad \text{for all large } T.$$

Proposition 2.8 ([20, Lemma 10.2]). *Assume that $f \in \mathfrak{A}$ and $H(f) \subset R^n$ is a compact set such that $\Omega(\nu) = H(f)$ for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$. Let $\varepsilon_0 \in (0, 1)$, $K_0, M_0 > 0$ and let l be a positive integer such that for each (f) -good function $x: [0, \infty) \rightarrow R^n$*

$$\text{dist}(H(f), \{x(t): t \in [T, T + l]\}) \leq 8^{-1}\varepsilon_0$$

for all large T (the existence of l follows from Proposition 2.5). Then there exist an integer $N \geq 10$ and a neighborhood \mathcal{U} of f in \mathfrak{A} such that for each $g \in \mathcal{U}$, each $S \in [0, \infty)$ and each a.c. function $x: [S, S + Nl] \rightarrow R^n$ satisfying

$$\begin{aligned} |x(S)|, |x(S + Nl)| &\leq K_0, \\ I^g(S, S + Nl, x) &\leq U^g(S, S + Nl, x(S), x(S + Nl)) + M_0 \end{aligned}$$

there exists an integer $i_0 \in [0, N - 8]$ such that for all $T \in [S + i_0l, S + (i_0 + 7)l]$

$$\text{dist}(H(f), \{x(t): t \in [T, T + l]\}) \leq \varepsilon_0.$$

Proposition 2.9 ([20, Lemma 10.3]). *Assume that $f \in \mathfrak{A}$ and $H(f) \subset R^n$ is a compact set such that $\Omega(\nu) = H(f)$ for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$. Let $\varepsilon > 0$. Then there exists $\delta > 0$ such that for each $x_1, x_2 \in R^n$ which satisfy $d(x_i, H(f)) \leq \delta$, $i = 1, 2$ there exists an a.c. function $\nu: [0, T] \rightarrow R^n$ for which*

$$T \geq 1, \nu(0) = x_1, \nu(T) = x_2, \quad I^f(0, T, \nu) - \pi^f(x_1) + \pi^f(x_2) - T\mu(f) \leq \varepsilon.$$

Proposition 2.10 ([20, Lemma 10.4]). *Assume that $f \in \mathfrak{A}$ and $H(f) \subset R^n$ is a compact set such that $\Omega(\nu) = H(f)$ for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$. Let $\varepsilon \in (0, 1)$ and let L be a positive integer such that for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$*

$$(2.1) \quad \text{dist}(H(f), \{\nu(t): t \in [S, S + L]\}) \leq \varepsilon$$

for all large S (the existence of L follows from Proposition 2.5).

Then there exists $\delta > 0$ such that for each $T \in [L, \infty)$ and each a.c. function $\nu: [0, T] \rightarrow R^n$ which satisfies

$$\begin{aligned} d(\nu(0), H(f)) &\leq \delta, \quad d(\nu(T), H(f)) \leq \delta, \\ I^f(0, T, \nu) - T\mu(f) - \pi^f(\nu(0)) + \pi^f(\nu(T)) &\leq \delta \end{aligned}$$

relation (2.1) holds for every $S \in [0, T - L]$.

Proposition 2.11 ([20, Lemma 9.1]). *Assume that $f \in \mathfrak{A}$. Then there exists a compact set $H^* \subset R^n$ which has the following properties:*

- there exists an (f) -good function $u: [0, \infty) \rightarrow R^n$ such that $\Omega(u) = H^*$;*
- for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$ either $\Omega(\nu) = H^*$ or $\Omega(\nu) \setminus H^* \neq \emptyset$.*

Proposition 2.12 ([20, Lemmas 9.3, 9.4]). *Let $f \in \mathfrak{A}$ and let H^* be as guaranteed in Proposition 2.11. Assume that $\phi: R^n \rightarrow [0, \infty)$ is a continuous bounded function such that $H^* = \{x \in R^n: \phi(x) = 0\}$. For $r \in (0, 1]$ we set*

$$f_r(x, u) = f(x, u) + r\phi(x), \quad x, u \in R^n.$$

Then $f_r \in \mathfrak{A}$, $r \in (0, 1]$ and for each $r \in (0, 1]$ and each (f_r) -good function $\nu: [0, \infty) \rightarrow R^n$

$$\Omega(\nu) = H^*.$$

Moreover for any neighborhood \mathcal{U} of f in \mathfrak{A} there exists $r_0 \in (0, 1)$ such that $f_r \in \mathfrak{A}$ for every $r \in (0, r_0)$.

Proposition 2.13 ([20, Proposition 3.4]). *Assume that $f \in \mathfrak{A}$, $M_1 > 0$, $0 \leq T_1 < T_2$, $x_i: [T_1, T_2] \rightarrow R^n$, $i = 1, 2, \dots$ is a sequence of a.c. functions such that $I^f(T_1, T_2, x_i) \leq M_1$, $i = 1, 2, \dots$. Then there exist a subsequence $\{x_{i_k}\}_{k=1}^\infty$ and an a.c. function $x: [T_1, T_2] \rightarrow R^n$ such that $I^f(T_1, T_2, x) \leq M_1$, $x_{i_k}(t) \rightarrow x(t)$ as $k \rightarrow \infty$ uniformly in $[T_1, T_2]$ and $x'_{i_k} \rightarrow x'$ as $k \rightarrow \infty$ weakly in $L^1(R^n; (T_1, T_2))$.*

Theorem 8.1, (8.2) and Proposition 7.3 in [20] imply the following result.

Proposition 2.14. *Let $f \in \mathfrak{A}$. Then $\pi^f(x) \rightarrow +\infty$ as $|x| \rightarrow \infty$.*

The following result was established in Aubin and Ekeland [2, Ch. 2, Sec. 3].

Proposition 2.15. *Let Ω be a closed subset of R^q . Then there exists a bounded nonnegative function $\phi \in C^\infty(R^q)$ such that $\Omega = \{x \in R^q: \phi(x) = 0\}$ and for each sequence of nonnegative integers p_1, \dots, p_q the function $\partial^{p_1}\phi/\partial x_1^{p_1}, \dots, \partial x_q^{p_q}: R^q \rightarrow R^1$ is bounded where $|p| = \sum_{i=1}^q p_i$.*

We can establish the following proposition which is a higher dimensional version of a well known result (Ball and Mizel [3], Morrey [15]).

Proposition 2.16. *Suppose that $f \in \mathfrak{M}$, $x, y \in R^n$, $T_1 \in [0, \infty)$, $T_2 > T_1$ and $w: [T_1, T_2] \rightarrow R^n$ is an a.c. function such that*

$$w(T_1) = x, \quad w(T_2) = y, \quad I^f(T_1, T_2, w) = U^f(T_1, T_2, x, y).$$

Then $w \in C^2([T_1, T_2]; R^n)$.

3. STRUCTURE OF THE PROOF OF THEOREM 1.2

Assume that $f \in \mathfrak{M}$ and $H(f) \subset R^n$ is a compact set such that $\Omega(\nu) = H(f)$ for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$.

We will describe briefly the proof of Lemma 4.4 which is established in Section 4 and which plays a crucial role in our discussion of Theorem 1.2.

For each a.c. function $u: [\tau_1, \tau_2] \rightarrow R^n$ where $\tau_1 \geq 0$, $\tau_2 > \tau_1$ and each $r_1, r_2 \in [\tau_1, \tau_2]$ satisfying $r_1 < r_2$ we set

$$\sigma(r_1, r_2, u) = I^f(r_1, r_2, u) - \pi^f(u(r_1)) + \pi^f(u(r_2)) - (r_2 - r_1)\mu(f).$$

Let $\varepsilon > 0$. To prove Lemma 4.4 we need to show that there is a number $q \geq 8$ such that for each $h_1, h_2 \in H(f)$ there exists an a.c. function $\nu: [0, q] \rightarrow R^n$ which satisfies

$$(3.1) \quad \nu(0) = h_1, \quad \nu(q) = h_2, \quad \sigma(0, q, \nu) \leq \varepsilon.$$

By Proposition 2.6 there exists a sequence of positive numbers $\{\delta_i\}_{i=0}^\infty$ such that

$$(3.2) \quad \delta_0 \in (0, 8^{-1}\varepsilon), \quad \delta_{i+1} < \delta_i, \quad i = 0, 1, \dots$$

and for each integer $i \geq 0$, each $x_1, x_2, y_1, y_2 \in H(f)$ which satisfy $|x_j - y_j| \leq \delta_i$, $j = 1, 2$ the following relations hold:

$$(3.3) \quad \begin{aligned} |U^f(0, 1, x_1, x_2) - U^f(0, 1, y_1, y_2)| &\leq 2^{-i-8}\varepsilon, \\ |\pi^f(x_j) - \pi^f(y_j)| &\leq 2^{-i-8}\varepsilon, \quad j = 1, 2. \end{aligned}$$

We will show that there exists an (f) -good function $\nu_*: [0, \infty) \rightarrow H(f)$ such that $\sigma(T_1, T_2, \nu_*) = 0$ for each $T_1 \geq 0, T_2 > T_1$. Then we will define a function $\phi: [0, \infty) \rightarrow R^1$ as

$$(3.4) \quad \phi(\tau) = I^f(0, 1, \nu_* + P_\tau) - \mu(f) - \pi^f(\nu_*(0)) + \pi^f(\nu_*(\tau)), \quad \tau \in [0, \infty),$$

where

$$(3.5) \quad P_\tau(t) = t(\nu_*(\tau) - \nu_*(1)), \quad t \in R^1, \quad \tau \in [0, \infty),$$

and verify that

$$(3.6) \quad \phi \in C^1([0, \infty); R^1), \quad \phi(1) = 0, \quad \phi(t) \geq 0, \quad t \in [0, \infty).$$

We can find a number $L \geq 10$ and a sequence of numbers $\{T_p\}_{p=1}^\infty$ such that

$$(3.7) \quad \text{dist}(H(f), \{\nu_*(t) : t \in [T, T + L]\}) \leq 4^{-1}\delta_0 \text{ for all } T \in [0, \infty),$$

$$(3.8) \quad T_p \geq 2L + 8, \quad |\nu_*(0) - \nu_*(T_p)| \leq 2^{-8}\delta_p, \quad p = 1, 2, \dots$$

Fix a positive number $\varepsilon_0 < 2^{-8}L^{-1}\varepsilon$. By (3.6) there exists a positive number Δ such that

$$(3.9) \quad \Delta < 2^{-8}, \quad |\phi'(t)| \leq 2^{-1}\varepsilon_0, \quad t \in [1 - \Delta, 1 + \Delta].$$

Choose an integer $N > 64(L + 1)\Delta^{-1}$ and set $q = \sum_{i=1}^N T_i + 8L + 8$.

Let $h_1, h_1 \in H(f)$. We will construct an a.c. function $\nu: [0, q] \rightarrow R^n$ satisfying (3.1). By the definition of L there exists numbers t_1, t_2 such that

$$(3.10) \quad t_1 \in [0, L], \quad t_2 \in [8, L + 8], \quad |h_j - \nu_*(t_j)| \leq 4^{-1}\delta_0, \quad j = 1, 2.$$

We set $\Delta_0 = (N - 1)^{-1}(8L + 8 - (t_2 - t_1))$ and verify that $\Delta_0 \in (0, \Delta)$. By using (3.8), (3.10) and the definition of $\{\delta_i\}_{i=0}^\infty$ we can construct functions $w_0: [0, T_1 - t_1] \rightarrow R^n$ and $u_0: [0, t_2] \rightarrow R^n$ such that

$$(3.11) \quad \begin{aligned} w_0(0) = h_1, \quad w_0(T_1 - t_1) = \nu_*(0), \quad \sigma(0, T_1 - t_1, w_0) &\leq 2^{-6}\varepsilon, \\ u_0(0) = \nu_*(0), \quad u_0(t_2) = h_2, \quad \sigma(0, t_2, u_0) &\leq 2^{-7}\varepsilon. \end{aligned}$$

For each integer $k \geq 1$ there exists an a.c. function $w_k: [0, \Delta_0 + T_{k+1}] \rightarrow R^n$ such that

$$\begin{aligned} w_k(t) = \nu_*(t) + P_{1-\Delta_0}(t), \quad t \in [0, 1], \quad w_k(t) = \nu_*(t - \Delta_0), \quad t \in [1, \Delta_0 + T_{k+1} - 1], \\ w_k(\Delta_0 + T_{k+1}) = \nu_*(0), \end{aligned}$$

$$I^f(\Delta_0 + T_{k+1} - 1, \Delta_0 + T_{k+1}, w_k) = U^f(0, 1, w_k(\Delta_0 + T_{k+1} - 1), w_k(\Delta_0 + T_{k+1})).$$

By using (3.4)–(3.6), (3.9), (3.8) and the definition of $\{\delta_i\}_{i=0}^\infty$ we show that

$$\sigma(0, T_{k+1} + \Delta_0, w_k) \leq 2^{-1}\varepsilon_0\Delta_0 + 2^{-k-8}\varepsilon, \quad k = 1, 2, \dots$$

We will finally define $\nu: [0, q] \rightarrow R^n$ as a concatenation of the functions $w_k, k = 0, \dots, N - 1, u_0$ and show that (3.1) holds.

The structure of the proof of Theorem 1.2. For simplicity we will only sketch the proof of the turnpike property for the integrand f and will not discuss the stability of the turnpike phenomenon under small perturbations of f . We choose a small enough number $\delta > 0$ and large enough numbers $l_0 > l > 0$ depending on ε, K .

Assume that $T \geq 2l_0$ and an a.c. function $\nu: [0, T] \rightarrow R^n$ satisfies

$$(3.12) \quad |\nu(0)|, |\nu(T)| \leq K, \quad I^f(0, T, \nu) \leq U^f(0, T, \nu(0), \nu(T)) + \delta.$$

We will show that for each $\tau \in [l_0, T - l_0]$

$$(3.13) \quad \text{dist}(H(f), \{\nu(t) : t \in [\tau, \tau + l]\}) \leq \varepsilon.$$

Assume the contrary. Then there is a number $\tau \in [l_0, T - l_0]$ for which

$$(3.14) \quad \text{dist}(H(f), \{\nu(t) : t \in [\tau, \tau + l]\}) > \varepsilon.$$

By Proposition 2.8 there are numbers $S_1, S_2 \in [0, T]$ such that

$$(3.15) \quad d(\nu(S_i), H(f)) \leq \delta, \quad i = 1, 2, \quad S_2 - \tau, \tau - S_1 \in [c_1, c_2],$$

where c_1, c_2 are some positive constants depending on ε, K .

It follows from (3.14), (3.15) and Proposition 2.10 that

$$(3.16) \quad \sigma(S_1, S_2, \nu) > \delta_0$$

where $\delta_0 > 8\delta$ is some constant depending on ε, K . By using (3.15) and Lemma 4.4 we show that there exists an a.c. function $u: [0, T] \rightarrow R^n$ such that

$$(3.17) \quad u(t) = \nu(t), \quad t \in [0, S_1] \cup [S_2, T], \quad \sigma(S_1, S_2, u) < \delta_0 - \delta.$$

It follows from (3.12), (3.16), (3.17) that

$$\delta \geq I^f(0, T, \nu) - I^f(0, T, u) = \sigma(S_1, S_2, \nu) - \sigma(S_1, S_2, u) > \delta.$$

The obtained contradiction proves that (3.13) holds for each $\tau \in [l_0, T - l_0]$.

4. PROOF OF THEOREM 1.2

Assume that $f \in \mathfrak{M}$ and $H(f) \subset R^n$ is a compact set such that $\Omega(\nu) = H(f)$ for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$.

Lemma 4.1. *Let $h \in H(f)$. Then there exists an (f) -good function $\nu: [0, \infty) \rightarrow H(f)$ such that $\nu(0) = h$ and*

$$(4.1) \quad I^f(T_1, T_2, \nu) = \mu(f)(T_2 - T_1) + \pi^f(\nu(T_1)) - \pi^f(\nu(T_2))$$

for each $T_1 \geq 0, T_2 > T_1$.

Proof. Consider any (f) -good function $w: [0, \infty) \rightarrow R^n$. Then

$$\Omega(w) = H(f).$$

By Proposition 2.1 the function w is bounded. It is easy to see that the following property holds:

(a) for each $\varepsilon > 0$ there exists $T(\varepsilon) > 0$ such that for each $T_1 \geq T(\varepsilon), T_2 > T_1$

$$I^f(T_1, T_2, w) - \mu(f)(T_2 - T_1) - \pi^f(w(T_1)) + \pi^f(w(T_2)) \leq \varepsilon.$$

There exists a sequence of numbers $\{T_p\}_{p=0}^\infty \subset [0, \infty)$ such that

$$(4.2) \quad T_{p+1} \geq T_p + 1, \quad p = 0, 1, \dots, \quad w(T_p) \rightarrow h \quad \text{as } p \rightarrow \infty.$$

For every integer $p \geq 1$ we set

$$(4.3) \quad \nu_p(t) = w(t + T_p), \quad t \in [0, \infty).$$

By Proposition 2.13, the boundness of w , (4.3) and property (a) there exists a subsequence $\{\nu_{p_j}\}_{j=1}^\infty$ and an a.c. function $\nu: [0, \infty) \rightarrow R^n$ such that for each integer $N \geq 1$

$$(4.4) \quad \begin{aligned} \nu_{p_j}(t) &\rightarrow \nu(t) \quad \text{as } j \rightarrow \infty \text{ uniformly in } [0, N], \\ I^f(0, N, \nu) &\leq \liminf_{j \rightarrow \infty} I^f(0, N, \nu_{p_j}). \end{aligned}$$

(4.2)–(4.4) imply that $\nu(0) = h$ and $\nu(t) \in H(f)$, $t \in [0, \infty)$. It follows from property (a) and (4.3), (4.4) that (4.1) holds for each $T_1 \geq 0$, $T_2 > T_1$. The lemma is proved.

By Lemma 4.1 there exists an (f) -good function $\nu_*: [0, \infty) \rightarrow H(f)$ such that

$$(4.5) \quad I^f(T_1, T_2, \nu_*) = \mu(f)(T_2 - T_1) + \pi^f(\nu_*(T_1)) - \pi^f(\nu_*(T_2))$$

for each $T_1 \geq 0$, $T_2 > T_1$.

It follows from Proposition 2.16 that

$$(4.6) \quad \nu_* \in C^2([0, \infty); R^n).$$

Lemma 4.2. *The function $\pi^f \cdot \nu_* \in C^1([0, \infty); R^1)$.*

Proof. By (4.5) for each $T \geq 0$

$$\pi^f(\nu_*(T)) = -I^f(0, T, \nu_*) + \mu(f)T + \pi^f(\nu_*(0)).$$

Together with (4.6) this implies the assertion of the lemma.

For each $\tau \in [0, \infty)$ we define

$$(4.7) \quad P_\tau(t) = t(\nu_*(\tau) - \nu_*(1)), \quad t \in R^1, \quad \psi(\tau) = I^f(0, 1, \nu_* + P_\tau).$$

Lemma 4.3. *$\psi \in C^1([0, \infty); R^1)$.*

Proof. For $\lambda, t \in [0, \infty)$ we set

$$(4.8) \quad B(\lambda, t) = (\nu_*(t) + P_\lambda(t), \nu'_*(t) + P'_\lambda(t)).$$

Let $\tau, h \in [0, \infty)$, $\tau \neq h$ and $t \in [0, 1]$. By (4.7), (4.8) there exists $\lambda_h(t) \in [\inf\{h, \tau\}, \sup\{h, \tau\}]$ such that

$$(h - \tau)^{-1}[f(B(h, t)) - f(B(\tau, t))] = \partial f / \partial x(B(\lambda_h(t), t))t\nu'_*(\lambda_h(t)) + \partial f / \partial u(B(\lambda_h(t), t))\nu'_*(\lambda_h(t)) \rightarrow \partial f / \partial x(B(\tau, t))t\nu'_*(\tau) + \partial f / \partial u(B(\tau, t))\nu'_*(\tau)$$

as $h \rightarrow \tau$ uniformly for all $t \in [0, 1]$. This implies that $\psi \in C^1([0, \infty); R^1)$. The lemma is proved.

Lemma 4.4. *Let $\varepsilon > 0$. Then there exists a number $q \geq 8$ such that for each $h_1, h_2 \in H(f)$ there exists an a.c. function $\nu: [0, q] \rightarrow R^n$ which satisfies*

$$(4.9) \quad \nu(0) = h_1, \quad \nu(q) = h_2,$$

$$(4.10) \quad I^f(0, q, \nu) \leq q\mu(f) + \pi^f(\nu(0)) - \pi^f(\nu(q)) + \varepsilon.$$

Proof. Define a function $\phi: [0, \infty) \rightarrow R^1$ as follows:

$$(4.11) \quad \phi(t) = \psi(t) - \mu(f) - \pi^f(\nu_*(0)) + \pi^f(\nu_*(t)), \quad t \in [0, \infty).$$

It follows from (4.11), (4.7), Lemmas 4.2, 4.3, (4.5) and the representation formula (see (1.5), (1.6)) that

$$(4.12) \quad \phi \in C^1([0, \infty); R^1), \quad \phi(1) = 0, \quad \phi(t) \geq 0, \quad t \in [0, \infty).$$

By Proposition 2.6 there exists a sequence of positive numbers $\{\delta_i\}_{i=0}^\infty$ such that

$$(4.13) \quad \delta_0 \in (0, 8^{-1}\varepsilon), \quad \delta_{i+1} < \delta_i, \quad i = 0, 1, \dots$$

and for each integer $i \geq 0$, each $x_1, x_2, y_1, y_2 \in H(f)$ which satisfy $|x_j - y_j| \leq \delta_i$, $j = 1, 2$ the following relations hold:

$$(4.14) \quad \begin{aligned} |U^f(0, 1, x_1, x_2) - U^f(0, 1, y_1, y_2)| &\leq 2^{-i-8}\varepsilon, \\ |\pi^f(x_j) - \pi^f(y_j)| &\leq 2^{-i-8}\varepsilon, \quad j = 1, 2. \end{aligned}$$

By the definition of ν , Proposition 2.5 there exists an integer $L \geq 10$ such that

$$(4.15) \quad \text{dist}(H(f), \{\nu_*(t) : t \in [T, T + L]\}) \leq 4^{-1}\delta_0$$

for all $T \in [0, \infty)$.

Since $\Omega(\nu_*) = H(f)$ and $\nu_*(0) \in H(f)$ there exists a sequence of numbers $\{T_p\}_{p=1}^\infty$ such that

$$(4.16) \quad T_p \geq 2L + 8, \quad |\nu_*(0) - \nu_*(T_p)| \leq 2^{-8}\delta_p, \quad p = 1, 2, \dots$$

Fix a positive number ε_0 for which

$$(4.17) \quad \varepsilon_0 < 2^{-8}L^{-1}\varepsilon.$$

It follows from (4.12) that there exists a positive number Δ such that

$$(4.18) \quad \Delta < 2^{-8}, \quad |\phi'(t)| \leq 2^{-1}\varepsilon_0, \quad t \in [1 - \Delta, 1 + \Delta].$$

Choose an integer

$$(4.19) \quad N > 64(L + 1)\Delta^{-1}$$

and set

$$(4.20) \quad q = \sum_{i=1}^N T_i + 8L + 8.$$

Let $h_1, h_1 \in H(f)$. We will construct an a.c. function $\nu: [0, q] \rightarrow R^n$ satisfying (4.9), (4.10). It follows from (4.15) which holds for each $T \in [0, \infty)$ that there exists numbers t_1, t_2 such that

$$(4.21) \quad t_1 \in [0, L], \quad t_2 \in [8, L + 8], \quad |h_j - \nu_*(t_j)| \leq 4^{-1}\delta_0, \quad j = 1, 2.$$

Set

$$(4.22) \quad \Delta_0 = (N - 1)^{-1}(8L + 8 - (t_2 - t_1)).$$

(4.22), (4.21), (4.19), (4.18) imply that

$$(4.23) \quad \Delta_0 \in (0, \Delta).$$

For each a.c. function $u: [\tau_1, \tau_2] \rightarrow R^n$ where $\tau_1 \geq 0$, $\tau_2 > \tau_1$ and each $r_1, r_2 \in [\tau_1, \tau_2]$ satisfying $r_1 < r_2$ we set

$$(4.24) \quad \sigma(r_1, r_2, u) = I^f(r_1, r_2, u) - \pi^f(u(r_1)) + \pi^f(u(r_2)) - (r_2 - r_1)\mu(f).$$

It follows from (4.16), (4.21) and Proposition 2.13 that there exists an a.c. function $w_0: [0, T_1 - t_1] \rightarrow R^n$ such that

$$(4.25) \quad \begin{aligned} w_0(0) &= h_1, \quad w_0(t) = \nu_*(t_1 + t), \quad t \in [1, T_1 - t_1 - 1], \quad w_0(T_1 - t_1) = \nu_*(0), \\ I^f(\tau, \tau + 1, w_0) &= U^f(0, 1, w_0(\tau), w_0(\tau + 1)), \quad \tau = 0, T_1 - t_1 - 1. \end{aligned}$$

By (4.24), the definition of ν_* , (4.5), (4.25), (4.21), (4.16) and the definition of $\{\delta_j\}_{j=0}^\infty$

$$\begin{aligned}
 & \sigma(0, T_1 - t_1, w_0) \\
 &= \sigma(0, 1, w_0) + \sigma(T_1 - t_1 - 1, T_1 - t_1, w_0) \\
 &= U^f(0, 1, h_1, \nu_*(t_1 + 1)) - \pi^f(h_1) + \pi^f(\nu_*(t_1 + 1)) - \mu(f) \\
 (4.26) \quad &+ U^f(0, 1, \nu_*(T_1 - 1), \nu_*(0)) - \pi^f(\nu_*(T_1 - 1)) + \pi^f(\nu_*(0)) - \mu(f) \\
 &\leq 4 \cdot 2^{-8}\varepsilon + U^f(0, 1, \nu_*(t_1), \nu_*(t_1 + 1)) - \pi^f(\nu_*(t_1)) \\
 &+ \pi^f(\nu_*(t_1 + 1)) - \mu(f) + U^f(0, 1, \nu_*(T_1 - 1), \nu_*(T_1)) \\
 &- \pi^f(\nu_*(T_1 - 1)) + \pi^f(\nu_*(T_1)) - \mu(f) \leq 2^{-6}\varepsilon.
 \end{aligned}$$

Let $k \geq 1$ be an integer. By (4.16), (4.7), (4.23), (4.18) and Proposition 2.13 there exists an a.c. function $w_k: [0, \Delta_0 + T_{k+1}] \rightarrow R^n$ such that

$$\begin{aligned}
 (4.27) \quad & w_k(t) = \nu_*(t) + P_{1-\Delta_0}(t), \quad t \in [0, 1], \quad w_k(t) = \nu_*(t - \Delta_0), \quad t \in [1, \Delta_0 + T_{k+1} - 1], \\
 & w_k(\Delta_0 + T_{k+1}) = \nu_*(0), \\
 & I^f(\Delta_0 + T_{k+1} - 1, \Delta_0 + T_{k+1}, w_k) = U^f(0, 1, w_k(\Delta_0 + T_{k+1} - 1), w_k(\Delta_0 + T_{k+1})).
 \end{aligned}$$

By (4.27), (4.7)

$$(4.28) \quad w_k(0) = \nu_*(0).$$

We will estimate $\sigma(0, T_{k+1} + \Delta_0, w_k)$. It follows from (4.27), (4.24), (4.5) that

$$(4.29) \quad \sigma(0, T_{k+1} + \Delta_0, w_k) = \sigma(0, 1, w_k) + \sigma(T_{k+1} + \Delta_0 - 1, T_{k+1} + \Delta_0, w_k).$$

(4.27), (4.24), (4.7), (4.11) imply that

$$(4.30) \quad \sigma(0, 1, w_k) = \phi(1 - \Delta_0).$$

It follows from (4.30), (4.23), (4.18), (4.12) that

$$(4.31) \quad \sigma(0, 1, w_k) \leq 2^{-1}\Delta_0\varepsilon_0.$$

By (4.27), (4.24), (4.16), the definition of ν_* , (4.5), the definition of $\{\delta_i\}_{i=0}^\infty$ (see (4.13), (4.14))

$$\begin{aligned}
 & \sigma(T_{k+1} + \Delta_0 - 1, T_{k+1} + \Delta_0, w_k) \\
 &= U^f(0, 1, \nu_*(T_{k+1} - 1), \nu_*(0)) - \pi^f(\nu_*(T_{k+1} - 1)) \\
 (4.32) \quad &+ \pi^f(\nu_*(0)) - \mu(f) \\
 &\leq U^f(0, 1, \nu_*(T_{k+1} - 1), \nu_*(T_{k+1})) - \pi^f(\nu_*(T_{k+1} - 1)) \\
 &+ \pi^f(\nu_*(T_{k+1})) - \mu(f) + 2 \cdot 2^{-k-9}\varepsilon = 2^{-k-8}\varepsilon.
 \end{aligned}$$

Combining (4.29), (4.31), (4.32) we obtain that

$$(4.33) \quad \sigma(0, T_{k+1} + \Delta_0, w_k) \leq 2^{-1}\varepsilon_0\Delta_0 + 2^{-k-8}\varepsilon.$$

By Proposition 2.13 there exists an a.c. function $u_0: [0, t_2] \rightarrow R^n$ such that

$$\begin{aligned}
 (4.34) \quad & u_0(t) = \nu_*(t), \quad t \in [0, t_2 - 1], \quad u_0(t_2) = h_2, \\
 & I^f(t_2 - 1, t_2, u_0) = U^f(0, 1, u_0(t_2 - 1), u_0(t_2)).
 \end{aligned}$$

It follows from (4.34), (4.24), (4.21), the definition of $\{\delta_i\}_{i=0}^\infty$ (see (4.13), (4.14)), the definition of ν_* , (4.5) that

$$\begin{aligned}
 \sigma(0, t_2, u_0) &= \sigma(t_2 - 1, t_2, u_0) \\
 &= U^f(0, 1, \nu_*(t_2 - 1), h_2) - \pi^f(\nu_*(t_2 - 1)) + \pi^f(h_2) - \mu(f) \\
 (4.35) \quad &\leq U^f(0, 1, \nu_*(t_2 - 1), \nu_*(t_2)) - \pi^f(\nu_*(t_2 - 1)) \\
 &\quad + \pi^f(\nu_*(t_2)) - \mu(f) + 2^{-7}\varepsilon \leq 2^{-7}\varepsilon.
 \end{aligned}$$

(4.22), (4.20) imply that

$$(4.36) \quad T_1 - t_1 + \sum_{k=1}^{N-1} (\Delta_0 + T_{k+1}) + t_2 = q.$$

By (4.36), (4.25), (4.27), (4.28), (4.34) there exists an a.c. function $\nu: [0, q] \rightarrow R^n$ such that

$$\begin{aligned}
 \nu(t) &= w_0(t), \quad t \in [0, T_1 - t_1], \quad \nu(t) = w_k \left(t - \left(\sum_{i=1}^k T_i + (k-1)\Delta_0 - t_1 \right) \right), \\
 (4.37) \quad t &\in \left[\sum_{i=1}^k T_i + (k-1)\Delta_0 - t_1, \sum_{i=1}^{k+1} T_i + k\Delta_0 - t_1 \right], \quad k = 1, \dots, N-1, \\
 \nu(t) &= u_0 \left(t - \left(\sum_{i=1}^N T_i + (N-1)\Delta_0 - t_1 \right) \right), \quad t \in \left[\sum_{i=1}^N T_i + (N-1)\Delta_0 - t_1, q \right].
 \end{aligned}$$

(4.37), (4.25), (4.36), (4.34) imply that

$$\nu(0) = h_1, \quad \nu(q) = h_2.$$

It follows from (4.24), (4.37), (4.26), (4.33), (4.35), (4.22), (4.21), (4.17) that

$$\begin{aligned}
 I^f(0, q, \nu) - \pi^f(\nu(0)) + \pi^f(\nu(q)) - q\mu(f) \\
 &= \sigma(0, T_1 - t_1, w_0) + \sum_{k=1}^{N-1} \sigma(0, T_{k+1} + \Delta_0, w_k) + \sigma(0, t_2, u_0) \\
 &\leq 2^{-6}\varepsilon + \sum_{k=1}^{N-1} (2^{-1}\varepsilon_0\Delta_0 + 2^{-k-8}\varepsilon) + 2^{-7}\varepsilon \leq 2^{-5}\varepsilon + 2^{-1}(N-1)\varepsilon_0\Delta_0 \\
 &\leq 2^{-5}\varepsilon + 2^{-1}(9L + 16)\varepsilon_0 \leq 2^{-1}\varepsilon.
 \end{aligned}$$

This completes the proof of the lemma.

Proof of Theorem 1.2. Let $\varepsilon, K > 0$. We may assume that

$$\varepsilon < 1, \quad K > \sup\{|h|: h \in H(f)\} + 4.$$

By Proposition 2.2 there exist a neighborhood \mathcal{U}_1 of f in \mathfrak{A} and a number $M > K$ such that for each $g \in \mathcal{U}_1$, each $T_1 \geq 0$, $T_2 \geq T_1 + 1$ and each a.c. function $\nu: [T_1, T_2] \rightarrow R^n$ which satisfies

$$(4.38) \quad |\nu(T_i)| \leq 2K + 4, \quad i = 1, 2, \quad I^g(T_1, T_2, \nu) \leq U^g(T_1, T_2, \nu(T_1), \nu(T_2)) + 2$$

the following relation holds:

$$(4.39) \quad |\nu(t)| \leq M \quad (t \in [T_1, T_2]).$$

By Proposition 2.6 there exists

$$(4.40) \quad \delta_1 \in (0, 8^{-1}\varepsilon)$$

such that for each $x_1, x_2, y_1, y_2 \in R^n$ which satisfy

$$(4.41) \quad |x_i|, |y_i| \leq 2M + 4 + 2 \sup\{|h|: h \in H(f)\}, \quad |x_i - y_i| \leq 4\delta_1, \quad i = 1, 2$$

the following relations hold:

$$(4.42) \quad \begin{aligned} |U^f(0, 1, x_1, x_2) - U^f(0, 1, y_1, y_2)| &\leq 2^{-8}\varepsilon, \\ |\pi^f(x_i) - \pi^f(y_i)| &\leq 2^{-8}\varepsilon, \quad i = 1, 2. \end{aligned}$$

By Proposition 2.5 there exists an integer $l \geq 1$ such that for each (f)-good function $\nu: [0, \infty) \rightarrow R^n$

$$(4.43) \quad \text{dist}(H(f), \{\nu(t): t \in [T, T + l]\}) \leq \varepsilon$$

for all large T . By Proposition 2.10 there exists

$$(4.44) \quad \delta_0 \in (0, 2^{-1}\delta_1)$$

such that for each $T \in [l, \infty)$ and each a.c. function $\nu: [0, T] \rightarrow R^n$ which satisfies

$$(4.45) \quad d(\nu(0), H(f)) \leq \delta_0, \quad d(\nu(T), H(f)) \leq \delta_0,$$

$$(4.46) \quad I^f(0, T, \nu) - T\mu(f) - \pi^f(\nu(0)) + \pi^f(\nu(T)) \leq \delta_0$$

the relation

$$(4.47) \quad \text{dist}(H(f), \{\nu(t): t \in [S, S + l]\}) \leq \varepsilon$$

holds for every $S \in [0, T - l]$. By Proposition 2.6 there exists

$$(4.48) \quad \delta \in (0, 32^{-1}\delta_0)$$

such that for each $x_1, x_2, y_1, y_2 \in R^n$ which satisfy

$$(4.49) \quad |x_i|, |y_i| \leq 2M + 4 + 2 \sup\{|h|: h \in H(f)\}, \quad |x_i - y_i| \leq 4\delta, \quad i = 1, 2,$$

the following relations hold:

$$(4.50) \quad \begin{aligned} |U^f(0, 1, x_1, x_2) - U^f(0, 1, y_1, y_2)| &\leq 2^{-8}\delta_0, \\ |\pi^f(x_i) - \pi^f(y_i)| &\leq 2^{-8}\delta_0, \quad i = 1, 2. \end{aligned}$$

By Proposition 2.5 there exists an integer $L \geq 1$ such that for each (f)-good function $\nu: [0, \infty) \rightarrow R^n$

$$(4.51) \quad \text{dist}(H(f), \{\nu(t): t \in [T, T + L]\}) \leq 8^{-1}\delta$$

for all large T .

By Proposition 2.8 there exists an integer $N \geq 10$ and a neighborhood \mathcal{U}_2 of f in \mathfrak{A} such that for each $g \in \mathcal{U}_2$, each $S \in [0, \infty)$ and each a.c. function $x: [S, S + NL] \rightarrow R^n$ which satisfies

$$(4.52) \quad \begin{aligned} |x(S)|, |x(S + NL)| &\leq 2M + 2, \\ I^g(S, S + NL, x) &\leq U^g(S, S + NL, x(S), x(S + NL)) + 4 \end{aligned}$$

there exists an integer $i_0 \in [0, N - 8]$ such that for all $T \in [S + i_0L, S + (i_0 + 7)L]$

$$(4.53) \quad \text{dist}(H(f), \{x(t): t \in [T, T + L]\}) \leq \delta.$$

By Lemma 4.4 there exists a number $q \geq 8$ such that for each $h_1, h_2 \in H(f)$ there exists an a.c. function $\nu: [0, q] \rightarrow R^n$ which satisfies

$$(4.54) \quad \nu(0) = h_1, \quad \nu(q) = h_2, \quad I^f(0, q, \nu) - q\mu(f) - \pi^f(\nu(0)) + \pi^f(\nu(q)) \leq 8^{-1}\delta.$$

By Proposition 2.3 there exists a neighborhood \mathcal{U}_3 of f in \mathfrak{A} such that for each $g \in \mathcal{U}_3$, each $T_1 \geq 0$, $T_2 \in [T_1 + 8^{-1}, T_1 + 6N(q + l + L)]$ and each a.c. function $x: [T_1, T_2] \rightarrow R^n$ which satisfies

$$(4.55) \quad \inf\{I^f(T_1, T_2, x), I^g(T_1, T_2, x)\} \\ \leq 4 + 2 \sup\{|\pi^f(h)|: h \in R^n, |h| \leq \sup\{|z|: z \in H(f)\} + 4\} \\ + 6|\mu(f)|N(q + l + L)$$

the following relation holds

$$(4.56) \quad |I^f(T_1, T_2, x) - I^g(T_1, T_2, x)| \leq 4^{-1}\delta.$$

By Proposition 2.4 there exists a neighborhood \mathcal{U}_4 of f in \mathfrak{A} such that for each $g \in \mathcal{U}_4$, each $x_1, x_2 \in R^n$ which satisfy

$$|x_1|, |x_2| \leq 2M + 4 + 2 \sup\{|z|: z \in H(f)\}$$

the following relation holds:

$$(4.57) \quad |U^f(0, 1, x_1, x_2) - U^g(0, 1, x_1, x_2)| \leq 2^{-8}\delta.$$

Set

$$(4.58) \quad l_0 = 2l + 2q + 2NL + 6,$$

$$(4.59) \quad \mathcal{U} = \bigcap_{i=1}^4 \mathcal{U}_i.$$

Assume that $g \in \mathcal{U}$, $T \geq 2l_0$ and an a.c. function $\nu: [0, T] \rightarrow R^n$ satisfies

$$(4.60) \quad |\nu(0)|, |\nu(T)| \leq K, \quad I^g(0, T, \nu) \leq U^g(0, T, \nu(0), \nu(T)) + \delta.$$

It follows from the definition of \mathcal{U}_1 (see (4.38), (4.39)) and (4.60) that

$$(4.61) \quad |\nu(t)| \leq M, \quad t \in [0, T].$$

Assume that there exist numbers $S_1, S_2 \in [0, T]$ such that

$$(4.62) \quad d(\nu(S_i), H(f)) \leq \delta, \quad i = 1, 2, \quad S_2 - S_1 \in [1 + l + q, 5N(L + l + q)].$$

We will show that for each $\tau \in [S_1, S_2 - l]$

$$(4.63) \quad \text{dist}(H(f), \{\nu(t): t \in [\tau, \tau + l]\}) \leq \varepsilon.$$

Let us assume the converse. Then there exists a number τ such that

$$(4.64) \quad \tau \in [S_1, S_2 - l], \quad \text{dist}(H(f), \{\nu(t): t \in [\tau, \tau + l]\}) > \varepsilon.$$

It follows from (4.62), (4.64), (4.48) and the definition of δ_0 (see (4.44)–(4.47)) that

$$(4.65) \quad I^f(S_1, S_2, \nu) - (S_2 - S_1)\mu(f) - \pi^f(\nu(S_1)) + \pi^f(\nu(S_2)) > \delta_0.$$

We will show that

$$(4.66) \quad I^g(S_1, S_2, \nu) - (S_2 - S_1)\mu(f) - \pi^f(\nu(S_1)) + \pi^f(\nu(S_2)) \geq 2^{-1}\delta_0.$$

Assume the contrary. Then by (4.62)

$$I^g(S_1, S_2, \nu) \leq 2 \sup\{|\pi^f(z)|: z \in R^n, d(z, H(f)) \leq 1\} + |\mu(f)|(S_2 - S_1) + 1.$$

It follows from this relation, (4.62) and the definition of \mathcal{U}_3 (see (4.55), (4.56)) that

$$|I^f(S_1, S_2, \nu) - I^g(S_1, S_2, \nu)| \leq 4^{-1}\delta.$$

Together with (4.65) this implies (4.66). The obtained contradiction proves that (4.66) holds.

By (4.62) there exist $h_1, h_2 \in H(f)$ such that

$$(4.67) \quad |\nu(S_i) - h_i| \leq \delta, \quad i = 1, 2.$$

By Lemma 4.1 there exists an (f) -good function

$$(4.68) \quad w_0: [0, \infty) \rightarrow H(f) \quad \text{such that} \quad w_0(0) = h_1,$$

$$(4.69) \quad I^f(t_1, t_2, w_0) - (t_2 - t_1)\mu(f) - \pi^f(w_0(t_1)) + \pi^f(w_0(t_2)) = 0$$

for each $t_1 \geq 0, t_2 > t_1$.

It follows from the definition of q (see (4.54), (4.62), (4.68)) that there exists an a.c. function $w_1: [0, q] \rightarrow R^n$ such that

$$(4.70) \quad \begin{aligned} w_1(0) &= w_0(S_2 - S_1 - q), \quad w_1(q) = h_2, \\ I^f(0, q, w_1) - q\mu(f) - \pi^f(w_1(0)) + \pi^f(w_1(q)) &\leq 8^{-1}\delta. \end{aligned}$$

By (4.62), Proposition 2.13, (4.68), (4.70) there exists an a.c. function $u: [0, T] \rightarrow R^n$ such that

$$(4.71) \quad \begin{aligned} u(t) &= \nu(t), \quad t \in [0, S_1] \cup [S_2, T], \\ u(t) &= w_0(t - S_1), \quad t \in [S_1 + 1, S_2 - q], \\ u(t) &= w_1(t - (S_2 - q)), \quad t \in [S_2 - q, S_2 - 1], \\ I^g(r, r + 1, u) &= U^g(0, 1, u(r), u(r + 1)), \quad r = S_1, S_2 - 1. \end{aligned}$$

For each a.c. function $y: [a, b] \rightarrow R^n$ where $a \geq 0, b > a$ and each $r_1, r_2 \in [a, b]$ satisfying $r_1 \leq r_2$ we set

$$(4.72) \quad \sigma(r_1, r_2, y) = I^g(r_1, r_2, y) - \pi^f(y(r_1)) + \pi^f(y(r_2)) - (r_2 - r_1)\mu(f).$$

It follows from (4.60), (4.71), (4.72) that

$$(4.73) \quad \begin{aligned} \delta &\geq I^g(0, T, \nu) - I^g(0, T, u) \\ &= \sigma(0, T, \nu) - \sigma(0, T, u) = \sigma(S_1, S_2, \nu) - \sigma(S_1, S_2, u). \end{aligned}$$

By (4.70), (4.68) and the definition of M (see (4.38), (4.39))

$$(4.74) \quad |w_1(t)| \leq M, \quad t \in [0, q].$$

By (4.70) there exists an a.c. function $\tilde{w}: [S_1, S_2] \rightarrow R^n$ such that

$$(4.75) \quad \begin{aligned} \tilde{w}(t) &= w_0(t - S_1), \quad t \in [S_1, S_2 - q], \\ \tilde{w}(t) &= w_1(t - (S_2 - q)), \quad t \in [S_2 - q, S_2]. \end{aligned}$$

(4.73), (4.72), (4.66), (4.71), (4.75) imply that

$$(4.76) \quad \begin{aligned} \delta &\geq 2^{-1}\delta_0 - \sigma(S_1, S_2, u) = 2^{-1}\delta_0 - \sigma(S_1, S_2, \tilde{w}) \\ &\quad + [\sigma(S_1, S_1 + 1, \tilde{w}) - \sigma(S_1, S_1 + 1, u)] \\ &\quad + [\sigma(S_2 - 1, S_2, \tilde{w}) - \sigma(S_2 - 1, S_2, u)]. \end{aligned}$$

We will estimate $\sigma(S_1, S_2, \tilde{w})$ and $\sigma(h, h + 1, \tilde{w}) - \sigma(h, h + 1, u), h = S_1, S_2 - 1$.

Let $h \in \{S_1, S_2 - 1\}$. (4.70), (4.75), (4.71), (4.74), (4.68), (4.61), (4.62), (4.67) imply that

$$\begin{aligned} |\tilde{w}(h)|, |\tilde{w}(h+1)|, |u(h)|, |u(h+1)| &\leq M + \sup\{|z|: z \in H(f)\}, \\ |\tilde{w}(h) - u(h)|, |\tilde{w}(h+1) - u(h+1)| &\leq \delta. \end{aligned}$$

It follows from these relations, (4.71), (4.72), the definition of \mathcal{U}_4 (see (4.57)) and δ (see (4.49), (4.50), (4.48)) that

$$\begin{aligned} (4.77) \quad &\sigma(h, h+1, \tilde{w}) - \sigma(h, h+1, u) \\ &\geq U^g(0, 1, \tilde{w}(h), \tilde{w}(h+1)) - \pi^f(\tilde{w}(h)) + \pi^f(\tilde{w}(h+1)) \\ &\quad - [U^g(0, 1, u(h), u(h+1)) - \pi^f(u(h)) + \pi^f(u(h+1))] \\ &\geq U^f(0, 1, \tilde{w}(h), \tilde{w}(h+1)) - \pi^f(\tilde{w}(h)) + \pi^f(\tilde{w}(h+1)) \\ &\quad - [U^f(0, 1, u(h), u(h+1)) - \pi^f(u(h)) + \pi^f(u(h+1))] - 2^{-7}\delta \\ &\geq -2^{-6}\delta_0, \quad h \in \{S_1, S_2 - 1\}. \end{aligned}$$

We will estimate $\sigma(S_1, S_2, \tilde{w})$. It follows from (4.69), (4.70), (4.75) that

$$(4.78) \quad I^f(S_1, S_2, \tilde{w}) - \pi^f(\tilde{w}(S_1)) + \pi^f(\tilde{w}(S_2)) - (S_2 - S_1)\mu(f) \leq 8^{-1}\delta.$$

By this relation, (4.62), (4.75), (4.68), (4.70) and the definition of \mathcal{U}_3 (see (4.55), (4.56))

$$|I^f(S_1, S_2, \tilde{w}) - I^g(S_1, S_2, \tilde{w})| \leq 4^{-1}\delta.$$

Together with (4.78), (4.72) this implies that

$$\sigma(S_1, S_2, \tilde{w}) \leq 3 \cdot 8^{-1}\delta.$$

It follows from this relation, (4.76), (4.77) that

$$\delta \geq 2^{-1}\delta_0 - 3 \cdot 8^{-1}\delta - 2^{-5}\delta_0.$$

This is contradictory to (4.48). The obtained contradiction proves that (4.63) holds for each $\tau \in [S_1, S_2 - l]$. Therefore we have shown that the following property holds:

Property D. For each $S_1, S_2 \in [0, T]$ which satisfy (4.62) relation (4.63) holds for each $\tau \in [S_1, S_2 - l]$.

It follows from (4.60), (4.61) and the definition of \mathcal{U}_2, N (see (4.52), (4.53)) that for each $r_0 \in [0, T - (1 + l + q + L(N + 2))]$ there exists a number r_1 such that

$$r_1 - r_0 \in [1 + l + q + 2L, 1 + l + q + L(N + 2)], \quad d(\nu(r_1), H(f)) \leq \delta.$$

This implies that there exists a finite sequence of numbers $\{S_i\}_{i=0}^Q \subset [0, T]$ such that

$$\begin{aligned} S_0 = 0, S_{i+1} - S_i &\in [1 + l + q + 2L, 1 + l + q + L(N + 2)], \quad i = 0, \dots, Q - 1, \\ T - S_Q &\leq 1 + l + q + L(N + 2), \quad d(\nu(S_i), H(f)) \leq \delta, \quad i = 1, \dots, Q. \end{aligned}$$

The assertion of the theorem follows from these relations and Property D.

5. PROOF OF THEOREM 1.3

Lemma 5.1. *Assume that an integrand $f \in \mathfrak{M}$ has Property B and $\varepsilon > 0$. Then there exists a neighborhood \mathcal{U} of f in \mathfrak{A} such that for each $g \in \mathcal{U}$ and each (g) -good function $\nu: [0, \infty) \rightarrow R^n$*

$$\text{dist}(\Omega(\nu), H(f)) \leq \varepsilon.$$

Proof. By Proposition 2.1 there exist a neighborhood \mathcal{U}_1 of f in \mathfrak{A} and a number $K > 0$ such that for each $g \in \mathcal{U}_1$ and each (g) -good function $\nu: [0, \infty) \rightarrow R^n$

$$\limsup_{t \rightarrow \infty} |\nu(t)| < K.$$

By Theorem 1.2 there exist a neighborhood \mathcal{U} of f in \mathfrak{A} which satisfies $\mathcal{U} \subset \mathcal{U}_1$ and numbers $l_0 > l > 0, \delta > 0$ such that for each $g \in \mathcal{U}$, each $T \geq 2l_0$ and each a.c. function $\nu: [0, T] \rightarrow R^n$ which satisfies

$$|\nu(0)|, |\nu(T)| \leq K, \quad I^g(0, T, \nu) \leq U^g(0, T, \nu(0), \nu(T)) + \delta$$

the relation $\text{dist}(H(f), \{\nu(t) : t \in [\tau, \tau + l]\}) \leq \varepsilon$ holds for each $\tau \in [l_0, T - l_0]$.

Assume that $g \in \mathcal{U}$ and $\nu: [0, \infty) \rightarrow R^n$ is a (g) -good function. It follows from the definition of $\mathcal{U}, \mathcal{U}_1, K$ and Proposition 1.1 that there exists a number $T_0 \geq 0$ such that

$$|\nu(t)| \leq K, \quad t \in [T_0, \infty),$$

$$I^g(t_1, t_2, \nu) \leq U^g(t_1, t_2, \nu(t_1), \nu(t_2)) + \delta \quad \text{for each } t_1 \geq T_0, t_2 > t_1.$$

It follows from these relations and the definition of $\mathcal{U}, l_0, l, \delta$ that

$$\text{dist}(H(f), \Omega(\nu)) \leq \varepsilon.$$

The lemma is proved.

Construction of the set \mathcal{F}_q . Suppose that q is a nonnegative integer. By Propositions 2.11, 2.12, 2.15 there exists a set $E_q \subset \mathfrak{M}_q$ which is an everywhere dense subset of $\overline{\mathfrak{M}}_q$ and such that each integrand $f \in E_q$ has Property B. Therefore for each $f \in E_q$ there exists a compact set $H(f) \subset R^n$ such that $\Omega(\nu) = H(f)$ for each (f) -good function $\nu: [0, \infty) \rightarrow R^n$.

By Theorem 1.2 and Lemma 5.1 for each $f \in E_q$ and each integer $p \geq 1$ there exist an open neighborhood $\mathcal{U}(f, p)$ of f in \mathfrak{A} and numbers $M(f, p) > p, l_0(f, p) > l(f, p) > 0, \delta(f, p) \in (0, p^{-1})$ such that for each $g \in \mathcal{U}(f, p)$ and each (g) -good function $\nu: [0, \infty) \rightarrow R^n$

$$\text{dist}(H(f), \Omega(\nu)) \leq 4^{-1}\delta(f, p);$$

for each $g \in \mathcal{U}(f, p)$, each $T \geq 2l_0(f, p)$ and each a.c. function $\nu: [0, T] \rightarrow R^n$ which satisfies

$$(5.1) \quad |\nu(0)|, |\nu(T)| \leq p, \quad I^g(0, T, \nu) \leq U^g(0, T, \nu(0), \nu(T)) + \delta(f, p)$$

the relation $|\nu(t)| \leq M(f, p)$ holds for all $t \in [0, T]$ and the following properties hold:

- (i) for each $\tau \in [l_0(f, p), T - l_0(f, p)]$
- (5.2) $\text{dist}(H(f), \{\nu(t) : t \in [\tau, \tau + l(f, p)]\}) \leq p^{-1};$
- (ii) if $d(\nu(0), H(f)) \leq \delta(f, p)$, then (5.2) holds for each $\tau \in [0, T - l_0(f, p)];$
- (iii) if $d(\nu(T), H(f)) \leq \delta(f, p)$, then (5.2) holds for each $\tau \in [l_0(f, p), T - l(f, p)].$

We define

$$(5.3) \quad \mathcal{F}_q = \left[\bigcap_{p=1}^{\infty} \cup \{ \mathcal{U}(f, p) : f \in E_q \} \right] \cap \overline{\mathfrak{M}}_q.$$

Clearly \mathcal{F}_q is a countable intersection of open everywhere dense subsets of $\overline{\mathfrak{M}}_q$.

Assume that $f \in \mathcal{F}_q$, $\varepsilon, K > 0$. Fix a natural number p such that

$$(5.4) \quad p > 2K + 4 + 8\varepsilon^{-1}.$$

There exists $G \in E_q$ such that

$$(5.5) \quad f \in \mathcal{U}(G, p).$$

It follows from (5.4), (5.5) and the definition of $\mathcal{U}(G, p)$, $\delta(G, p)$ that for each (f)-good function $\nu: [0, \infty) \rightarrow R^n$

$$(5.6) \quad \text{dist}(H(G), \Omega(\nu)) \leq 4^{-1}\delta(G, p) < (4p)^{-1} < 8^{-1}\varepsilon.$$

This implies that for each (f)-good function $\nu_i: [0, \infty) \rightarrow R^n$, $i = 1, 2$,

$$\text{dist}(\Omega(\nu_1), \Omega(\nu_2)) \leq \varepsilon.$$

Since ε is any positive number we conclude that f has Property B and there exists a compact set $H(f) \subset R^n$ such that $\Omega(w) = H(f)$ for each (f)-good function $w: [0, \infty) \rightarrow R^n$. It follows from (5.6) that

$$(5.7) \quad \text{dist}(H(G), H(f)) \leq 4^{-1}\delta(G, p).$$

Set

$$(5.8) \quad \begin{aligned} \mathcal{U} &= \mathcal{U}(G, p), & M &= M(G, p), \\ l_0 &= l_0(G, p), & l &= l(G, p), & \delta &= 8^{-1}\delta(G, p). \end{aligned}$$

Assume that $g \in \mathcal{U}$, $T \geq 2l_0$ and an a.c. function $\nu: [0, T] \rightarrow R^n$ satisfies

$$(5.9) \quad |\nu(0)|, |\nu(T)| \leq K, \quad I^g(0, T, \nu) \leq U^g(0, T, \nu(0), \nu(T)) + \delta.$$

It follows from (5.9), (5.8), (5.5), (5.4) and the definition of $\mathcal{U}(G, p)$, $M(G, p)$, $l_0(G, p)$, $l(G, p)$, $\delta(G, p)$ that

$$(5.10) \quad |\nu(t)| \leq M, \quad t \in [0, T],$$

and properties (i)–(iii) hold with $f = G$. Together with (5.7), (5.8), (5.4) this implies that

$$(5.11) \quad \text{dist}(H(f), \{ \nu(t) : t \in [\tau, \tau + l] \}) \leq \varepsilon$$

for each $\tau \in [l_0, T - l_0]$; if $d(\nu(0), H(f)) \leq \delta$, then (5.11) holds for each $\tau \in [0, T - l_0]$. If $d(\nu(T), H(f)) \leq \delta$, then (5.11) holds for each $\tau \in [l_0, T - l]$. This completes the proof of the theorem.

6. EXAMPLES

Fix a constant $a > 0$ and set $\psi(t) = t$, $t \in [0, \infty)$. Consider the complete metric space \mathfrak{A} of integrands $f: R^n \times R^n \rightarrow R^1$ defined in Section 1.

Example 1. Consider an integrand $f(x, u) = |x|^2 + |u|^2$, $x, u \in R^n$. It is easy to see that $f \in \mathfrak{M}_q$ for each integer $q \geq 0$ if the constant a is large enough. We can show (see [20, Section 14]) that $\Omega(\nu) = \{0\}$ for every (f)-good function $\nu: [0, \infty) \rightarrow R^n$. Therefore Theorem 1.2 holds with the integrand f .

Example 2. Fix a number $q > 0$ and consider an integrand

$$g(x, u) = q|x|^2|x - e|^2 + |u|^2, \quad x, u \in \mathbb{R}^n,$$

where $e = (1, 1, \dots, 1)$. It is easy to see that $g \in \mathfrak{M}$ if the constant a is large enough. Clearly f does not have the turnpike property.

REFERENCES

1. Z. Artstein and A. Leizarowitz, *Tracking periodic signals with overtaking criterion*, IEEE Trans. on Autom. Control AC **30** (1985), 1122–1126.
2. J. P. Aubin and I. Ekeland, *Applied nonlinear analysis*, Wiley Interscience, New York, 1984. MR **87a**:58002
3. J. M. Ball and V. J. Mizel, *One-dimensional variational problems whose minimizers do not satisfy the Euler-Lagrange equation*, Arch. Rational Mech. Anal. **90** (1985), 325–388. MR **86k**:49002
4. W. A. Brock and A. Haurie, *On existence of overtaking optimal trajectories over an infinite horizon*, Math. Oper. Res. **1** (1976), 337–346. MR **56**:14975
5. D. A. Carlson, *The existence of catching-up optimal solutions for a class of infinite horizon optimal control problems with time delay*, SIAM Journal on Control and Optimization **28** (1990), 402–422. MR **91b**:49008
6. D. A. Carlson, A. Haurie and A. Jabrane, *Existence of overtaking optimal solutions to infinite dimensional control problems on unbounded time intervals*, SIAM Journal on Control and Optimization **25** (1987), 1517–1541. MR **88j**:90063
7. D. A. Carlson, A. Haurie and A. Leizarowitz, *Infinite horizon optimal control*, Springer-Verlag, Berlin, 1991.
8. B. D. Coleman, M. Marcus and V. J. Mizel, *On the thermodynamics of periodic phases*, Arch. Rational Mech. Anal. **117** (1992), 321–347. MR **93d**:73008
9. J. L. Kelley, *General topology*, D. Van Nostrand Co., 1955. MR **51**:6681
10. A. Leizarowitz, *Existence of overtaking optimal trajectories for problems with convex integrands*, Math. Oper. Res. **10** (1985), 450–461. MR **87f**:49002
11. A. Leizarowitz, *Infinite horizon autonomous systems with unbounded cost*, Appl. Math. and Opt. **13** (1985), 19–43. MR **86g**:49002
12. A. Leizarowitz, *Optimal trajectories on infinite horizon deterministic control systems*, Appl. Math. and Opt. **19** (1989), 11–32. MR **89m**:49049
13. A. Leizarowitz and V. J. Mizel, *One dimensional infinite-horizon variational problems arising in continuum mechanics*, Arch. Rational Mech. Anal. **106** (1989), 161–194. MR **90b**:49007
14. V. L. Makarov and A. M. Rubinov, *Mathematical theory of economic dynamics and equilibria*, Nauka, Moscow, 1973; English transl., Springer-Verlag, 1977. MR **55**:11973
15. C. B. Morrey, *Multiple integrals in the calculus of variations*, Springer, Berlin Heidelberg New York, 1966. MR **34**:2380
16. A. M. Rubinov, *Economic dynamics*, in *Itogy Nauki. Sovremennye problemy mat.* **19** (1982) VINITI Moscow, 59–110; English transl., J. Soviet Math. **26** (1984), 4. MR **83k**:90034
17. A. J. Zaslavski, *Optimal programs on infinite horizon I*, SIAM Journal on Control and Optimization **33** (1995), 1643–1660. MR **96i**:49047
18. A. J. Zaslavski, *Optimal programs on infinite horizon II*, SIAM Journal on Control and Optimization **33** (1995), 1661–1686. MR **96i**:49047
19. A. J. Zaslavski, *The existence and structure of extremals for a class of second order infinite horizon variational problems*, Journal of Mathematical Analysis and Applications **194** (1995), 660–696. MR **96h**:49005
20. A. J. Zaslavski, *Dynamic properties of optimal solutions of variational problems*, Nonlinear Analysis: Theory, Methods and Applications **27** (1996), 895–931. MR **97h**:49022
21. A. J. Zaslavski, *Structure of extremals for one-dimensional variational problems arising in continuum mechanics*, Journal of Mathematical Analysis and Applications **198** (1996), 893–921. MR **97a**:49006

DEPARTMENT OF MATHEMATICS, TECHNION-ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA, 32000, ISRAEL

E-mail address: ajzasl@techunix.technion.ac.il