

ON THE TELESCOPIC HOMOTOPY THEORY OF SPACES

A. K. BOUSFIELD

ABSTRACT. In telescopic homotopy theory, a space or spectrum X is approximated by a tower of localizations $L_n^f X$, $n \geq 0$, taking account of v_n -periodic homotopy groups for progressively higher n . For each $n \geq 1$, we construct a telescopic Kuhn functor Φ_n carrying a space to a spectrum with the same v_n -periodic homotopy groups, and we construct a new functor Θ_n left adjoint to Φ_n . Using these functors, we show that the n th stable monocular homotopy category (comprising the n th fibers of stable telescopic towers) embeds as a retract of the n th unstable monocular homotopy category in two ways: one giving infinite loop spaces and the other giving “infinite L_n^f -suspension spaces.” We deduce that Ravenel’s stable telescope conjectures are equivalent to unstable telescope conjectures. In particular, we show that the failure of Ravenel’s n th stable telescope conjecture implies the existence of highly connected infinite loop spaces with trivial Johnson-Wilson $E(n)_*$ -homology but nontrivial v_n -periodic homotopy groups, showing a fundamental difference between the unstable chromatic and telescopic theories. As a stable chromatic application, we show that each spectrum is $K(n)$ -equivalent to a suspension spectrum. As an unstable chromatic application, we determine the $E(n)_*$ -localizations and $K(n)_*$ -localizations of infinite loop spaces in terms of $E(n)_*$ -localizations of spectra under suitable conditions. We also determine the $E(n)_*$ -localizations and $K(n)_*$ -localizations of arbitrary Postnikov H -spaces.

1. INTRODUCTION

With the probable failure of Ravenel’s telescope conjecture, a “telescopic” or “finite chromatic” approach to stable homotopy theory has emerged in its own right (see [Mil2], [Rav4], [MS]). In this approach, a spectrum E is approximated at a prime p by a tower $\{L_n^f E\}_{n \geq 0}$ of localizations away from the finite p -local spectra of type $n + 1$. Recall that a finite p -local spectrum W is of *type* n when $K(i)_* W = 0$ for $i < n$ and $K(n)_* W \neq 0$, where $K(n)$ is the n th Morava K -theory. By results of Mitchell, Hopkins, and Smith, such a W exists for each n and admits a self-map $\omega : \Sigma^d W \rightarrow W$ with $K(n)_* \omega$ an isomorphism and with $K(i)_* \omega = 0$ for $i \neq n$. Such an ω is called a v_n -map and leads to the v_n -periodic homotopy groups $v_n^{-1} \pi_*(E; W) = \omega^{-1}[W, E]_*$ of a spectrum E . The localization $E \rightarrow L_n^f E$ preserves the v_i -periodic homotopy groups of E for $i \leq n$ while annihilating those for $i > n$, and the fiber $M_n^f E$ of $L_n^f E \rightarrow L_{n-1}^f E$ captures the v_n -periodic homotopy groups alone. From this perspective, $\{L_n^f E\}_{n \geq 0}$ resembles a Postnikov tower. As suggested by Ravenel, we call the spectrum $M_n^f E$ *monocular*.

Received by the editors March 29, 1999.

2000 *Mathematics Subject Classification*. Primary 55P60; Secondary 55N20, 55P42, 55P65, 55U35.

Research partially supported by the National Science Foundation.

©2000 American Mathematical Society

For spaces, a similar telescopic theory is obtained using the nullification or periodization functors of [Bou7], [Bou8], or [Dro]. A pointed space X is approximated at a prime p by a tower $\{L_n^f X\}_{n \geq 0}$ of nullifications away from desuspensions of finite p -local spectra of type $n + 1$. Like its stable counterpart, the nullification $X \rightarrow L_n^f X$ preserves the v_i -periodic homotopy groups of X for $i \leq n$ while annihilating those for $i > n$, and the fiber $M_n^f X$ of $L_n^f X \rightarrow L_{n-1}^f X$ captures the v_n -periodic homotopy groups alone. We likewise call the space $M_n^f X$ *monocular*.

For $n \geq 1$, Kuhn [Kuh2] introduced a remarkable functor ϕ_n from pointed spaces to spectra with $\phi_n \Omega^\infty E \simeq L_{K(n)} E$ for a spectrum E , where $L_{K(n)}$ is the $K(n)_*$ -localization functor. This greatly extended a similar result of [Bou6] for $n = 1$. Using Kuhn's basic approach, we now construct a telescopic Kuhn functor Φ_n from pointed spaces to spectra with $\Phi_n \Omega^\infty E \simeq \widehat{M}_n^f E$ and $v_n^{-1} \pi_*(\Phi_n X; W) \cong v_n^{-1} \pi_*(X; W)$ (by Theorem 5.3) for a spectrum E and pointed space X , where \widehat{M}_n^f is a variant of the n th monocular functor M_n^f . Our key result (Theorem 5.4) provides a new functor Θ_n from spectra to pointed spaces which is homotopically left adjoint to Φ_n in the sense that

$$\text{map}_*(\Theta_n E, L_n^f X) \simeq \text{map}_*(E, \Phi_n X).$$

In particular, $\widehat{K(n)^*}(\Theta_n E) \simeq K(n)^*(E)$, so that $\Theta_n E$ behaves somewhat like an infinite desuspension of E . Moreover, we establish a natural equivalence $L_n^f \Sigma^\infty \Theta_n E \simeq M_n^f E$ showing that monocular spectra are closely related to suspension spectra. This is carried by $L_{K(n)}$ to a natural equivalence $L_{K(n)} \Sigma^\infty \Theta_n E \simeq L_{K(n)} E$ showing that every spectrum is $K(n)$ -equivalent to a suspension spectrum (see Corollary 5.8).

For $n \geq 1$, we let \mathcal{M}_n^f be the n th stable monocular homotopy category comprising all $M_n^f E$ for spectra E , and we let \mathcal{UN}_n^f be the n th unstable monocular homotopy category comprising all $(M_n^f X)\langle d_n \rangle$ for spaces X , where d_n is a certain integer increasing with n and $\langle d_n \rangle$ denotes a d_n -connected cover. In Theorem 5.14, we deduce that \mathcal{M}_n^f embeds as a retract of \mathcal{UN}_n^f in two ways given by the adjoint factorizations

$$\begin{array}{ccccc} \mathcal{M}_n^f & \xrightarrow{\Theta_n} & \mathcal{UN}_n^f & \xrightarrow{L_n^f \Sigma^\infty} & \mathcal{M}_n^f \\ \mathcal{M}_n^f & \xleftarrow{M_n^f \Phi_n} & \mathcal{UN}_n^f & \xleftarrow{\widetilde{\Omega}^\infty} & \mathcal{M}_n^f \end{array}$$

of the identity functor for \mathcal{M}_n^f , where $\widetilde{\Omega}^\infty E = (\Omega^\infty E)\langle d_n \rangle$. We note that the $\widetilde{\Omega}^\infty$ -embedding of \mathcal{M}_n^f in \mathcal{UN}_n^f consists of infinite loop spaces, while the Θ_n -embedding of \mathcal{M}_n^f in \mathcal{UN}_n^f consists of "infinite L_n^f -suspension spaces" since $\Theta_n E \simeq L_n^f \Sigma^k (\Theta_n \Sigma^{-k} E)$ for $k \geq 0$. Moreover, there is a correspondence of unstable and stable maps given by isomorphisms $[\Theta_n E, \widetilde{\Omega}^\infty G] \cong [E, G]$ for $E, G \in \mathcal{M}_n^f$.

Ravenel's original telescope conjecture involves the mapping telescopes of v_n -maps, but may be reformulated to assert that each $K(n)_*$ -equivalence of spectra is a v_n -periodic homotopy equivalence (see e.g. [Rav1], [MS], or 6.13). Since the converse is true, this would give a perfect Whitehead theorem for $K(n)_*$ -equivalences of spectra. From another standpoint, the telescope conjecture concerns the natural telescopic-to-chromatic transformation $L_n^f \rightarrow L_n$ where L_n is the $E(n)_*$ -localization functor. In fact, $L_n^f E \rightarrow L_n E$ is an equivalence for each spectrum E if and only if the m th telescope conjecture holds for each $m \leq n$. The n th telescope conjecture was proved for $n = 1$ by Mahowald [Mah] and Miller [Mil1], but has been strongly

doubted for $n \geq 2$ since 1990 when Ravenel [Rav2] sketched a refutation for $n = 2$ and $p \geq 5$. However, a completely convincing refutation has been extraordinarily hard to achieve, and Ravenel cautions us against drawing conclusions at this time.

To formulate an unstable telescope conjecture, we call a map $X \rightarrow Y$ of pointed spaces a *virtual $K(n)_*$ -equivalence* when it induces a $K(n)_*$ -equivalence $(\Omega X)\langle n+2 \rangle \rightarrow (\Omega Y)\langle n+2 \rangle$. By [Bou9], the virtual $K(n)_*$ -equivalences behave much like periodic homotopy equivalences; for instance, in a ladder of fiber sequences

$$\begin{array}{ccccc} F & \longrightarrow & X & \longrightarrow & B \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ F' & \longrightarrow & X' & \longrightarrow & B', \end{array}$$

if any two of α , β , and γ are virtual $K(n)_*$ -equivalences, then so is the third. Our unstable telescope conjecture asserts that *each virtual $K(n)_*$ -equivalence of pointed spaces is a v_n -periodic homotopy equivalence*, and we prove that this is equivalent to the stable telescope conjecture for $K(n)_*$ (see Theorem 6.5). If the conjecture fails, then we can deduce the existence of highly connected infinite loop spaces with trivial $E(n)_*$ -homology but nontrivial v_n -periodic homotopy groups (see Theorem 6.11).

Our telescopic results have important chromatic applications. They enable us to determine the $E(n)_*$ -localizations of many infinite loop spaces in terms of the $E(n)_*$ -localizations of spectra, very much as we determined the $E(1)_*$ -localizations of infinite loop spaces in [Bou4]. For a connected infinite loop space $\Omega_0^\infty E$, we show that $L_n \Omega_0^\infty E$ is given by $\Omega_0^\infty L_n E$ with low dimensional adjustments under suitable conditions (Theorem 8.1). These conditions hold, for instance, when $n \leq 2$ or when E is a BP -module spectrum. The $K(n)_*$ -localizations of spaces are very closely related to $E(n)_*$ -localizations; if Y is a connected H -space or simply connected space, then $L_{K(n)} Y \simeq (L_n Y)_p^\wedge$ (see Theorem 7.3). In particular, $L_{K(n)} \Omega_0^\infty E$ is given by $\Omega_0^\infty (L_n E)_p^\wedge$ with low dimensional adjustments under suitable conditions (see Theorem 8.2), and may differ substantially from $\Omega_0^\infty L_{K(n)} E$. We also determine $L_n Y$ and $L_{K(n)} Y$ for any Postnikov H -space Y using work of Hopkins, Ravenel, and Wilson [HRW].

Much of our effort is devoted to constructing the functors Φ_n and Θ_n using the technology of model categories. We give a much improved version of the Bousfield-Friedlander [BF] construction of new model categories from old ones with “ Q -structures,” and deduce that each nullification functor determines a proper model category for spaces (see Theorem 9.9). In particular, the telescopic functor L_n^f determines a proper model category structure for spaces, giving the n th telescopic homotopy category. We construct a chain of right Quillen functors and left Quillen equivalences from this model category to the model category of spectra, and we compose their total derived functors to define Φ_n . We then construct an adjoint chain of functors and compose their total derived functors to define Θ_n . The homotopy functors Φ_n and Θ_n can be induced by simplicial functors between the categories of spaces and spectra.

The paper is organized as follows. In Section 2 we discuss the general theory of relative homological localizations of spectra, and in Section 3 we review the telescopic homotopy theory of spectra. Although this material is more or less known, it is needed to establish notation, terminology, and background. In Section

4 we introduce the telescopic homotopy theory of spaces. In Section 5 we state our main theorems on the existence and basic properties of the functors Φ_n and Θ_n , and we derive our embedding theorem for monocular homotopy categories. In Section 6 we prove the equivalence of the stable and unstable telescope conjectures. In Section 7 we determine the $E(n)_*$ -localizations and $K(n)_*$ -localizations of Postnikov H -spaces. In Section 8 we determine the $E(n)_*$ -localizations and $K(n)_*$ -localizations of infinite loop spaces under suitable conditions. In Section 9 we show that a nullification functor, such as L_n^f , determines a proper model category structure for spaces. In Sections 10-13 we construct the functors Φ_n and Θ_n and establish their basic properties listed in Theorems 5.3 and 5.4.

We work simplicially so that a “space” is a simplicial set, and a “spectrum” is as in [BF] or [HSS, 4.2]. However, to make the presentation more accessible, we frequently work in the homotopy categories Ho_* of pointed spaces and Ho^s of spectra. We develop our results in telescopic and chromatic homotopy theory at a fixed but arbitrary prime p .

We are very grateful to the referee for critical comments and suggestions which have helped to make this work more accessible.

2. RELATIVE HOMOLOGICAL LOCALIZATIONS OF SPECTRA

To set the stage for our discussion of telescopic stable homotopy theory in Section 3, we shall establish an equivalence between two sorts of relative homological localizations of spectra. We work in the stable homotopy category of spectra and assume familiarity with

2.1. Homological localizations and acyclizations of spectra. For a spectrum E , we let $X \rightarrow L_E X$ and $C_E X \rightarrow X$ denote the E_* -localization and E_* -acyclization of a spectrum X , and we obtain a homotopy cofiber sequence $C_E X \rightarrow X \rightarrow L_E X$ (see [Bou2], [Rav1]). We let $\mathcal{L}_E \subset Ho^s$ (resp. $\mathcal{C}_E \subset Ho^s$) denote the full subcategory of E_* -local (resp. E_* -acyclic) spectra.

2.2. Ayclicity classes of spectra. The functors L_E and C_E depend only on the equivalence class $\langle E \rangle$ obtained by considering two spectra as equivalent when their homology theories have the same acyclic spectra. These *acyclicity classes* $\langle E \rangle$ are studied at length in [Bou1], [Rav1], and [HP]. They are partially ordered, where $\langle G \rangle \geq \langle E \rangle$ means that each G_* -acyclic spectrum is E_* -acyclic, and they have the usual smash and wedge operations. A class $\langle X \rangle$ may have a *complement* $\langle X \rangle^c$, which is uniquely determined by the conditions $\langle X \rangle \wedge \langle X \rangle^c = \langle 0 \rangle$ and $\langle X \rangle \vee \langle X \rangle^c = \langle S \rangle$ where S is the sphere spectrum. For a class $\langle W \rangle$ and a complemented class $\langle X \rangle$, the *difference class* $\langle W \rangle \setminus \langle X \rangle$ is defined as $\langle W \rangle \wedge \langle X \rangle^c$.

2.3. Relative homological localizations. For spectra E and G with $\langle G \rangle \geq \langle E \rangle$, there is a natural map $L_G X \rightarrow L_E X$ with homotopy fiber $C_E L_G X$ for a spectrum X . We regard $C_E L_G X$ as a *relative homological localization* of X giving its G_* -local E_* -acyclic part. When $\langle E \rangle$ is complemented, we may choose a representative $G \setminus E$ of $\langle G \rangle \setminus \langle E \rangle$ and regard $L_{G \setminus E} X$ as another sort of *relative homological localization* of X . We shall show that $C_E L_G$ and $L_{G \setminus E}$ are functorially equivalent when E is smashing.

2.4. Smashing spectra. As in [Rav1], a spectrum E is called *smashing* when the natural map $X \wedge L_E S \rightarrow L_E X$ is an equivalence for all spectra X . When E is smashing, $X \wedge C_E S \rightarrow C_E X$ is also an equivalence, and $\langle E \rangle$ is complemented with $\langle E \rangle = \langle L_E S \rangle$ and $\langle E \rangle^c = \langle C_E S \rangle$.

Theorem 2.5. *For a smashing spectrum E and spectrum G with $\langle G \rangle \geq \langle E \rangle$, the relative homological localization functors $C_E L_G$ and $L_{G \setminus E}$ are related by natural equivalences $L_{G \setminus E}(C_E L_G X) \simeq L_{G \setminus E} X$ and $C_E L_G(L_{G \setminus E} X) \simeq C_E L_G X$ for $X \in Ho^s$. Moreover, they restrict to adjoint functors $C_E L_G : Ho^s \rightleftarrows \mathcal{L}_G : L_{G \setminus E}$ which in turn restrict to adjoint equivalences $C_E L_G : \mathcal{L}_{G \setminus E} \rightleftarrows C_E \wedge \mathcal{L}_G : L_{G \setminus E}$. Furthermore, $L_{G \setminus E} X \simeq F(C_E S, L_G X) \simeq L_{S \setminus E} L_G X$ for $X \in Ho^s$.*

Cases of this result have been proved in [Bou3], [Bou8], [HPS], and [HSt]. Our proof in 2.7 will use the following categorical lemma. For a class \mathcal{W} of maps in a category \mathcal{A} , an object $J \in \mathcal{A}$ is called \mathcal{W} -local (resp. \mathcal{W} -colocal) when $f^* : \text{Hom}(B, J) \cong \text{Hom}(A, J)$ (resp. $f^* : \text{Hom}(J, A) \cong \text{Hom}(J, B)$) for each map $f : A \rightarrow B$ in \mathcal{W} . A \mathcal{W} -localization of $A \in \mathcal{A}$ consists of a map $A \rightarrow I$ in \mathcal{W} such that I is \mathcal{W} -local, and a \mathcal{W} -colocalization consists of a map $J \rightarrow A$ in \mathcal{W} such that J is \mathcal{W} -colocal.

Lemma 2.6. *If each object of \mathcal{A} has both a \mathcal{W} -localization and a \mathcal{W} -colocalization, then:*

- (i) *the \mathcal{W} -colocalization functor $C : \mathcal{A} \rightarrow \mathcal{A}$ is left adjoint to the \mathcal{W} -localization functor $L : \mathcal{A} \rightarrow \mathcal{A}$;*
- (ii) *there are natural isomorphisms $LCX \cong LX$ and $CLX \cong CX$ for $X \in \mathcal{A}$;*
- (iii) *C and L restrict to adjoint equivalences between the full subcategories of \mathcal{W} -local objects and of \mathcal{W} -colocal objects in \mathcal{A} .*

Proof. C is a left adjoint to L via the natural bijections

$$\text{Hom}(CX, Y) \cong \text{Hom}(CX, LY) \cong \text{Hom}(X, LX).$$

The lemma follows easily since C and L carry maps of \mathcal{W} to isomorphisms by adjunction arguments. □

2.7. Proof of Theorem 2.5. Let \mathcal{W} be the class of C_E -equivalences in Ho^s . Then each spectrum X has a \mathcal{W} -localization $X \rightarrow L_{S \setminus E} X$ and a \mathcal{W} -colocalization $C_E X \rightarrow X$. Thus $L_{S \setminus E} : Ho^s \rightarrow Ho^s$ is right adjoint to $C_E : Ho^s \rightarrow Ho^s$ by Lemma 2.6, and hence $L_{S \setminus E} X \simeq F(C_E S, X)$ for $X \in Ho^s$ since $C_E X \simeq X \wedge C_E S$. If $W \in Ho^s$ is $G \setminus E_*$ -acyclic, then $W \wedge (G \wedge C_E S) \simeq 0$ and thus $[W, F(C_E S, L_G X)] = 0$ for $X \in Ho^s$. Hence $F(C_E S, L_G X)$ is $G \setminus E_*$ -local and $L_{G \setminus E} X \simeq F(C_E S, L_G X) \simeq L_{S \setminus E} L_G X$. The theorem now follows easily.

Examples of smashing spectra arise from the theory of

2.8. Homotopical trivializations and cotrivialization of spectra. As in [Bou1] for $W \in Ho^s$, we say that a spectrum I is W -trivial when $[W, I]_* = 0$ and say that a spectrum J is W -cotrivial when $[J, Y]_* = 0$ for each W -trivial spectrum Y . By [Bou1, 1.7], each spectrum X has a W -trivialization $X \rightarrow L^W X$ and a W -cotrivialization $C^W X \rightarrow X$, and we obtain a homotopy (co)fiber sequence

$C^W X \rightarrow X \rightarrow L^W X$. We let $\mathcal{L}^W \subset Ho^s$ (resp. $\mathcal{C}^W \subset Ho^s$) denote the full subcategory of W -trivial (resp. W -cotrivial) spectra. By [Bou1, 1.6], \mathcal{C}^W is the smallest full subcategory of Ho^s which contains W , is closed under arbitrary wedges, and contains all three spectra in a cofiber sequence when it contains any two. We remark that homological localizations of spectra are actually examples of homotopical trivializations by [Bou2, 1.14].

The following theorem generalizes a result of [Bou2, 3.5] or [Mil2] showing that the spectrum $L^W S$ is smashing when W is a wedge of finite spectra.

Theorem 2.9. *Let W be a spectrum such that \mathcal{L}^W is closed under arbitrary wedges. Then $L^W S$ is smashing with $\langle L^W S \rangle = \langle W \rangle^c$, and the $L^W S_*$ -localization of a spectrum X is given by the W -trivialization $X \rightarrow L^W X$.*

Proof. If $E \in \mathcal{L}^W$, then $X \wedge E \in \mathcal{L}^W$ for each $X \in Ho^s$, since $\mathcal{C}^E \subset \mathcal{L}^W$ by 2.8. Using the cofiber sequence $X \wedge C^W S \rightarrow X \wedge S \rightarrow X \wedge L^W S$ with $X \wedge C^W S \in \mathcal{C}^W$ and $X \wedge L^W S \in \mathcal{L}^W$, we obtain natural equivalences $X \wedge C^W S \simeq C^W X$ and $X \wedge L^W S \simeq L^W X$ for $X \in Ho^s$. Since $W \wedge C^W S \simeq W$ and $C^W S \wedge L^W S \simeq 0$, we easily deduce $\langle W \rangle = \langle C^W S \rangle$ and $\langle W \rangle^c = \langle L^W S \rangle$ as required. \square

3. THE TELESCOPIC HOMOTOPY THEORY OF SPECTRA

We now discuss the “finite chromatic” or “telescopic” approach to stable homotopy theory in which spectra are approximated by their homotopical trivializations away from finite spectra of progressively higher types as in [Mil2], [Rav4], or [MS]. Using our knowledge of relative homological localizations, we establish an equivalence between two sorts of “monocular” stable homotopy categories. Let p be a fixed prime, and let $K(n)$ be the n th Morava K -theory spectrum at p for $n \geq 0$ where $K(0) = HQ$.

3.1. Types of spectra. The *type* of a finite p -local spectrum W is the smallest integer $n \geq 0$ such that $K(n)_* W \neq 0$, or is ∞ when there is no such n (i.e., when $W \simeq 0$). A spectrum W of type n exists for each $n \geq 0$, and any two such spectra have the same acyclicity class; moreover, this class decreases monotonely as n increases.

3.2. Telescopic towers of spectra. For $n \geq -1$, let $F(n+1)$ be a finite p -local spectrum of type $n+1$, let $\check{F}(n+1) = F(n+1) \vee \bigvee_{q \neq p} S/q$ be the wedge of $F(n+1)$

with the Moore spectra S/q for all primes $q \neq p$, and let $S(n)$ be the $\check{F}(n+1)$ -trivialization of the sphere spectrum S . We note that $S(n)$ does not depend on the choice of $F(n+1)$, and we have $S(-1) = 0$ and $S(0) = HQ$. By Theorem 2.9, $S(n)$ is smashing with $\langle S(n) \rangle = \langle \check{F}(n+1) \rangle^c$ and hence with $\langle S(m) \rangle \leq \langle S(n) \rangle$ for $m \leq n$. For a spectrum X , we write $L_n^f X = L_{S(n)} X$ and $C_n^f X = C_{S(n)} X$ and obtain the natural *telescopic tower*

$$L_0^f X \leftarrow L_1^f X \leftarrow \cdots \leftarrow L_{n-1}^f X \leftarrow L_n^f X \leftarrow \cdots$$

under X . The homotopy fiber $M_n^f X$ of $L_n^f X \rightarrow L_{n-1}^f X$ is given by $M_n^f X = C_{n-1}^f L_n^f X$ and is called the n th *monocular* part of X . It has a variant $\widehat{M}_n^f X = L_{T(n)} X$ where $T(n) = S(n) \wedge F(n)$ is the “difference spectrum” with

$$\langle T(n) \rangle = \langle S(n) \rangle \setminus \langle S(n-1) \rangle = \langle F(n) \rangle \setminus \langle F(n+1) \rangle.$$

We let $\mathcal{L}_n^f \subset Ho^s$, $\mathcal{M}_n^f \subset Ho^s$, and $\widehat{\mathcal{M}}_n^f \subset Ho^s$ respectively denote the full subcategories given by all $L_n^f X$, $M_n^f X$, and $\widehat{M}_n^f X$ for $X \in Ho^s$. By Theorem 2.5 we have

Theorem 3.3. *For $n \geq 0$, the functors M_n^f and \widehat{M}_n^f are related by natural equivalences $\widehat{M}_n^f M_n^f X \simeq \widehat{M}_n^f X$ and $M_n^f \widehat{M}_n^f X \simeq M_n^f X$ for $X \in Ho^s$. Moreover, they restrict to adjoint functors $M_n^f : Ho^s \rightleftarrows \mathcal{L}_n^f : \widehat{M}_n^f$ which in turn restrict to adjoint equivalences $M_n^f : \widehat{\mathcal{M}}_n^f \rightleftarrows \mathcal{M}_n^f : \widehat{M}_n^f$. Furthermore, $\widehat{M}_n^f X \simeq F(C_{n-1}^f S, L_n^f X) \simeq F(C_{n-1}^f S_{(p)}, L_n^f X) \simeq L_{\tilde{F}(n)} L_n^f X \simeq L_{F(n)} L_n^f X$ for $X \in Ho^s$.*

A similar result is proved in [HSt, Theorem 6.19], for the monochromatic functors M_n and \widehat{M}_n , and can also be derived from Theorem 2.5. We immediately deduce

Corollary 3.4. *For spectra $X, Y \in Ho^s$ and $n > 0$, there are isomorphisms*

$$[M_n^f X, M_n^f Y] \cong [M_n^f X, \widehat{M}_n^f Y] \cong [\widehat{M}_n^f X, \widehat{M}_n^f Y].$$

The functors L_n^f , M_n^f , and \widehat{M}_n^f are perhaps best understood in terms of

3.5. v_n -periodic homotopy groups. Let $\mathcal{F}_n \subset Ho^s$ be the full subcategory of finite p -local spectra of type $\geq n$. For a spectrum $W \in \mathcal{F}_n$ with $n > 0$, a v_n -map is a map $\omega : \Sigma^d W \rightarrow W$ with $d > 0$ such that $K(n)_* \omega$ is an isomorphism and $K(i)_* \omega = 0$ for all $i \neq n$. By the periodicity theorem of Hopkins and Smith (see [Hop], [HSm], [Rav3]), each spectrum $W \in \mathcal{F}_n$ has a v_n -map, which is unique and natural up to iteration. The v_n -periodic homotopy groups of a spectrum $X \in Ho^s$ with coefficients in a spectrum $W \in \mathcal{F}_n$ are defined by

$$v_n^{-1} \pi_*(X; W) = \omega^{-1}[W, X]_* = \operatorname{colim}_i [\Sigma^{id} W, X]_*$$

for $n > 0$ using the sequence

$$W \xleftarrow{\omega} \Sigma^d W \xleftarrow{\omega} \Sigma^{2d} W \xleftarrow{\omega} \dots$$

for a v_n -map $\omega : \Sigma^d W \rightarrow W$, and by $v_0^{-1} \pi_*(X; W) = [W, X_Q]_*$ for $n = 0$. The periodicity theorem ensures that $v_n^{-1} \pi_*(X; W)$ does not depend on the choice of ω and is functorial in X and W . In fact, $v_n^{-1} \pi_*(-; W)$ is a homology theory by

Lemma 3.6. *For a spectrum X and $W \in \mathcal{F}_n$ with $n \geq 0$, there is a natural isomorphism $v_n^{-1} \pi_*(X; W) \cong (S(n) \wedge DW)_* X$ where $DW = F(W, S_{(p)})$ is the p -local Spanier-Whitehead dual of W .*

Proof. We may assume $n > 0$. Since $D\omega : DW \rightarrow \Sigma^{-d} DW$ is a v_n -map in \mathcal{F}_n , its homotopy cofiber is in \mathcal{F}_{n+1} , and the map $D\omega \wedge C$ is nilpotent for each $C \in \mathcal{F}_{n+1}$ by the nilpotence theorem of Devinatz, Hopkins, and Smith (see [Hop], [HSm], [Rav3]). Hence, the sequence

$$DW \xrightarrow{D\omega} \Sigma^{-d} DW \xrightarrow{D\omega} \Sigma^{-2d} DW \xrightarrow{D\omega} \dots$$

has the telescope $L_n^f DW \simeq S(n) \wedge DW$ and

$$v_n^{-1} \pi_*(X; W) \cong \operatorname{colim}_i (\Sigma^{-id} DW)_* X \cong (S(n) \wedge DW)_* X.$$

□

3.7. v_n -periodic homotopy equivalences. For a finite p -local spectrum W of type n , we have $\langle S(n) \wedge DW \rangle = \langle S(n) \wedge F(n) \rangle = \langle T(n) \rangle$. Thus by Lemma 3.6, a $v_n^{-1}\pi_*(-; W)$ -equivalence of spectra is the same as a $T(n)_*$ -equivalence and does not depend on the choice of W . It will be called a $v_n^{-1}\pi_*$ -equivalence. A map which is a $v_i^{-1}\pi_*$ -equivalence for $0 \leq i \leq n$ will be called a $\bigoplus_{i=0}^n v_i^{-1}\pi_*$ -equivalence and is the same as an $S(n)_*$ -equivalence since $\langle S(n) \rangle = \langle T(0) \rangle \vee \cdots \vee \langle T(n) \rangle$.

3.8. An interpretation of the telescopic tower. For a spectrum X and $n \geq 0$, we may interpret $L_n^f X$ as the $\bigoplus_{i=0}^n v_i^{-1}\pi_*$ -localization of X , and we obtain

$$v_i^{-1}\pi_*(L_n^f X; F(i)) \cong \begin{cases} v_i^{-1}\pi_*(X; F(i)) & \text{for } i \leq n, \\ 0 & \text{for } i > n, \end{cases}$$

$$v_i^{-1}\pi_*(M_n^f X; F(i)) \cong \begin{cases} v_i^{-1}\pi_*(X; F(i)) & \text{for } i = n, \\ 0 & \text{for } i \neq n, \end{cases}$$

for $i \geq 0$ where $F(i)$ is a finite p -local spectrum of type i . Thus the telescopic tower $\{L_n^f X\}_{n \geq 0}$ resembles a Postnikov tower for X , while its monocular fibers $M_n^f X$ resemble Eilenberg-MacLane objects. We may likewise interpret $\widehat{M}_n^f X$ as the $v_n^{-1}\pi_*$ -localization of X .

4. THE TELESCOPIC HOMOTOPY THEORY OF SPACES

We now introduce a “finite chromatic” or “telescopic” approach to the homotopy theory of spaces, using the theory of homotopical nullifications (= periodizations) developed by Dror-Farjoun [Dro] and the author [Bou7], [Bou8]. For convenience, we work primarily in the homotopy category Ho_* of pointed spaces (i.e. pointed simplicial sets), and start by recalling

4.1. Nullifications and conullifications of spaces. For a space $A \in Ho_*$, we say that a space $Y \in Ho_*$ is A -null if the map $A \rightarrow *$ induces an equivalence $Y \simeq \text{map}(A, Y)$ of unpointed mapping spaces. When Y is connected, this means that the pointed mapping space $\text{map}_*(A, Y)$ is contractible. We say that a space $J \in Ho_*$ is A -conull if $\text{map}_*(J, Y) \simeq *$ for each A -null space $Y \in Ho_*$. By [Bou7] or [Dro], each space $X \in Ho_*$ has an A -nullification $X \rightarrow P_A X$ going universally to an A -null target, and has an A -conullification $\overline{P}_A X \rightarrow X$ coming universally from an A -conull source. These form a homotopy fiber sequence $\overline{P}_A X \rightarrow X \rightarrow P_A X$ by [Dro, p. 37]. We note that the terms “ A -null” and “ A -nullification” have exactly the same meanings as the earlier terms “ A -periodic” and “ A -periodization” used in [Bou7].

4.2. Nullity classes and relative nullifications. The functors P_A and \overline{P}_A depend only on the equivalence class $\langle A \rangle$ obtained by considering two spaces as equivalent when they have the same null spaces. These nullity classes $\langle A \rangle$ are studied in [Bou7] and [Dro]. They are partially ordered where $\langle A \rangle \leq \langle B \rangle$ means that each B -null space is A -null, which happens if and only if $P_B A \simeq *$. (The opposite ordering is used in [Dro].) If $\langle A \rangle \leq \langle B \rangle$, then for $X \in Ho_*$ there is a natural map $P_A X \rightarrow P_B X$ with homotopy fiber $\overline{P}_B P_A X$, and we regard $\overline{P}_B P_A X$ as a relative nullification of X , giving its A -null B -conull part.

4.3. Telescopic towers of spaces. For $n \geq 0$, we choose a suspension space $A(n+1) \in Ho_*$ such that $\Sigma^\infty A(n+1)$ is a finite p -local spectrum of type $n+1$ with the connectivity of $\tilde{H}_*(A(n+1); Z/p)$ as low as possible. For instance, we might choose $A(1) = M(Z/p, 2) = S^2 \cup_p e^3$. We let $d_n - 1$ denote the connectivity of $\tilde{H}_*(A(n+1); Z/p)$ and observe that $d_0 = 2$, $d_{n+1} \geq d_n$, and $d_n \geq n+1$ by [Bou7, 13.8]. We let

$$\check{A}(n+1) = A(n+1) \vee \bigvee_{q \neq p} M(Z/q, 2)$$

be the wedge of $A(n+1)$ with the Z/q -Moore spaces $M(Z/q, 2)$ for primes $q \neq p$. Then by [Bou7, 9.15], the nullity class $\langle \check{A}(n+1) \rangle$ does not depend on the choice of $A(n+1)$, and it decreases monotonely as n increases. For a pointed space X , we write $L_n^f X = P_{\check{A}(n+1)} X$ and $C_n^f X = \overline{P}_{\check{A}(n+1)} X$, and we obtain the natural telescopic tower

$$L_0^f X \leftarrow L_1^f X \leftarrow \dots \leftarrow L_{n-1}^f X \leftarrow L_n^f X \leftarrow \dots$$

under X . For $n \geq 1$, the homotopy fiber $M_n^f X$ of $L_n^f X \rightarrow L_{n-1}^f X$ is given by $M_n^f X = C_{n-1}^f L_n^f X$ and is called the n th *monocular* part of X . We let $\mathcal{UL}_n^f \subset Ho_*$ and $\mathcal{UM}_n^f \subset Ho_*$ respectively denote the full subcategories given by all $L_n^f X$ and $M_n^f X$ for $X \in Ho_*$. As will be explained below in 4.11, $L_n^f X$ is very closely related to the v_n -periodization $P_{v_n} X$ of [Bou7].

4.4. Simplicial functoriality. Since the functors $L_n^f : Ho_* \rightarrow Ho_*$ for $n \geq 0$ are nullifications, they are induced as in [Bou7, 2.10] by simplicial functors $L_n^f : \mathcal{SS}_* \rightarrow \mathcal{SS}_*$ which may be constructed with simplicial natural transformations giving a functorial telescopic tower $X \rightarrow \{L_n^f X\}_{n \geq 0}$ for $X \in \mathcal{SS}_*$.

The telescopic tower is perhaps best understood in terms of

4.5. v_n -periodic homotopy groups. The v_n -periodic homotopy groups of a space $X \in Ho_*$ with coefficients in a spectrum $W \in \mathcal{F}_n$ are defined for $n \geq 1$ by

$$v_n^{-1} \pi_*(X; W) = \omega^{-1}[W, X]_* = \operatorname{colim}_i [\Sigma^{id} W, X]_*$$

using the eventual desuspension to Ho_* of the sequence

$$W \xleftarrow{\omega} \Sigma^d W \xleftarrow{\omega} \Sigma^{2d} W \leftarrow \dots$$

for a v_n -map $\omega : \Sigma^d W \rightarrow W$. As in the stable case (3.5), $v_n^{-1} \pi_*(X; W)$ does not depend on the choice of ω and is functorial in X and W . A map $f : X \rightarrow Y$ in Ho_* will be called a $v_n^{-1} \pi_*$ -equivalence when $f_* : v_n^{-1} \pi_*(X; W) \cong v_n^{-1} \pi_*(Y; W)$ for some (and hence all) $W \in \mathcal{F}_n$ of type n (see [Bou7, 11.11]). It will be called a $v_0^{-1} \pi_*$ -equivalence when $f_* : Q \otimes \pi_* X \cong Q \otimes \pi_* Y$ assuming that X and Y are nilpotent.

4.6. An interpretation of the telescopic tower. For a space $X \in Ho_*$, we have

$$\pi_i L_0^f X \cong \begin{cases} \pi_i X & \text{for } i < 2, \\ \pi_i X / \tau \pi_i X & \text{for } i = 2, \\ \pi_i X \otimes Q & \text{for } i > 2 \end{cases}$$

by [Bou7, Theorem 5.2], where $\tau\pi_i X$ is the torsion subgroup of $\pi_i X$. For $n \geq 1$ and $i \geq 1$, we obtain

$$v_i^{-1}\pi_*(L_n^f X; F(i)) \cong \begin{cases} v_i^{-1}\pi_*(X; F(i)) & \text{for } i \leq n, \\ 0 & \text{for } i > n, \end{cases}$$

$$v_i^{-1}\pi_*(M_n^f X; F(i)) \cong \begin{cases} v_i^{-1}\pi_*(X; F(i)) & \text{for } i = n, \\ 0 & \text{for } i \neq n \end{cases}$$

by 4.11 below and [Bou7, 11.5] where $F(i)$ is a finite p -local spectrum of type i . Thus the telescopic tower $\{L_n^f X\}_{n \geq 0}$ resembles a Postnikov tower for X , capturing progressively higher types of periodic homotopy.

This interpretation can be strengthened for highly connected spaces. For $j \geq 0$ let $Y\langle j \rangle$ denote the j -connected cover of a space $Y \in Ho_*$, and let $Ho_*\langle j \rangle \subset Ho_*$ denote the full subcategory of j -connected spaces. By 4.11 below and Theorems 11.5, 13.1, and 13.3 of [Bou7], we have

Theorem 4.7. *For $n \geq 0$ and $X \in Ho_*\langle d_n \rangle$, the map $X \rightarrow L_n^f X$ is a $\bigoplus_{i=0}^n v_i^{-1}\pi_*$ -localization of X in $Ho_*\langle d_n \rangle$. Moreover, for $Y \in Ho_*$, there is a natural equivalence $L_n^f(Y\langle d_n \rangle) \simeq (L_n^f Y)\langle d_n \rangle$.*

Corollary 4.8. *For $n \geq 0$, a map in $Ho_*\langle d_n \rangle$ is an L_n^f -equivalence if and only if it is a $\bigoplus_{i=0}^n v_i^{-1}\pi_*$ -equivalence.*

For $n \geq 1$, we write $N_n^f X = (M_n^f X)\langle d_n \rangle$ and let $\mathcal{UN}_n^f \subset Ho_*$ denote the full subcategory given by all $N_n^f X$ for $X \in Ho_*$. We call \mathcal{UN}_n^f the n th *monocular homotopy category* of spaces.

Lemma 4.9. *For $n \geq 1$, \mathcal{UN}_n^f is contained in \mathcal{UM}_n^f .*

Proof. For $X \in Ho_*$, we have $L_{n-1}^f(N_n^f X) \simeq *$ by Corollary 4.8 and have $N_n^f X$ in \mathcal{UL}_n^f by Theorem 4.7 since $N_n^f X = (M_n^f X)\langle d_n \rangle$ with $M_n^f X \in \mathcal{UL}_n^f$. \square

Thus, \mathcal{UN}_n^f is the full subcategory of \mathcal{UM}_n^f given by its d_n -connected spaces.

Finally, we shall explain the very close relationship between $L_n^f : Ho_* \rightarrow Ho_*$ and the v_n -periodization functor $P_{v_n} : Ho_* \rightarrow Ho_*$ of [Bou7]. For a pointed space X , we let $P_{(p)}X$ denote the M -nullification of X where $M = \bigvee_{q \neq p} M(Z/q, 2)$. By [Bou7, Theorem 5.2], $P_{(p)}X$ is a partial p -localization of X with

$$\pi_i P_{(p)}X \cong \begin{cases} \pi_i X & \text{for } i < 2, \\ \pi_i X / \tau_p \pi_i X & \text{for } i = 2, \\ \pi_i X \otimes Z_{(p)} & \text{for } i > 2, \end{cases}$$

where $\tau_p \pi_i X \subset \pi_i X$ is the subgroup of torsion prime to p . Since $\check{A}(n+1) = A(n+1) \vee M$, there are natural maps $P_{(p)}P_{A(n+1)} \rightarrow L_n^f X$ and $P_{A(n+1)}P_{(p)}X \rightarrow L_n^f X$, and we have

Lemma 4.10. *For a space $X \in Ho_*$ and $n \geq 0$, there are natural equivalences $P_{(p)}P_{A(n+1)} \simeq L_n^f X$ and $P_{A(n+1)}P_{(p)}X \simeq L_n^f X$.*

Proof. Since $A(n + 1)$ is a simply connected space with p -torsion homology groups $\tilde{H}_*(A(n + 1); Z)$, the functor $P_{(p)}$ preserves the $A(n + 1)$ -null property, while $P_{A(n+1)}$ preserves the M -null property. Hence, $P_{(p)}P_{A(n+1)}X$ and $P_{A(n+1)}P_{(p)}X$ are $\dot{A}(n + 1)$ -null, and the lemma follows easily. \square

4.11. Comparison of L_n^f and P_{v_n} . For a space $X \in Ho_*$, the localizations $L_n^f X$ and $P_{v_n} X$ are closely related by natural maps $P_{A(n+1)} X \rightarrow L_n^f X$ and $P_{A(n+1)} X \rightarrow P_{v_n} X$. The first map is obtained by Lemma 4.10 and gives $\pi_i P_{A(n+1)} X \otimes Z_{(p)} \cong \pi_i L_n^f X$ for $i > 2$. The second map is obtained from the natural equivalences $P_{A(n+1)} X \simeq P_{\Sigma^j W_n} X$ and $P_{v_n} X \simeq P_{\Sigma W_n} X$ of [Bou7, 10.1 and 10.2], and gives $\pi_i P_{A(n+1)} X \cong \pi_i P_{v_n} X$ for $i > d_n$ by [Bou7, 7.2]. Consequently, $\pi_i L_n^f X \cong \pi_i P_{v_n} X \otimes Z_{(p)}$ for $i > d_n$. Our functors $L_n^f : Ho_* \rightarrow Ho_*$ are designed to be compatible with their stable counterparts (see Proposition 5.2) and to be well-behaved above the homotopy dimension d_n (see Lemma 12.5). We are generally not concerned with their action below the homotopy dimension d_n .

5. THE TELESCOPIC KUHN FUNCTOR AND ITS ADJOINT

In this section, we state our main results on the telescopic Kuhn functor $\Phi_n : Ho_* \rightarrow Ho^s$ and our new functor $\Theta_n : Ho^s \rightarrow Ho_*$, establishing strong relations between the stable and unstable telescopic homotopy theories. We begin by noting some elementary relations involving infinite loop spaces.

Proposition 5.1. *For spectra $E \in Ho^s$ and $W \in \mathcal{F}_n$ with $n \geq 1$, there is a natural isomorphism $v_n^{-1}\pi_*(\Omega^\infty E; W) \cong v_n^{-1}\pi_*(E; W)$.*

Of course, we also have $\pi_i(\Omega^\infty E) \otimes Q \cong \pi_i E \otimes Q$ for $i \geq 0$. More geometrically, the infinite loop map $\Omega^\infty E \rightarrow \Omega^\infty L_n^f E$ induces a map $\lambda : L_n^f \Omega^\infty E \rightarrow \Omega^\infty L_n^f E$ since $\Omega^\infty L_n^f E$ is in \mathcal{UL}_n^f . This provides natural comparison maps of telescopic towers $\lambda : \{L_n^f \Omega^\infty E\}_{n \geq 0} \rightarrow \{\Omega^\infty L_n^f E\}_{n \geq 0}$ and of monocular parts $\lambda : M_n^f \Omega^\infty E \rightarrow \Omega^\infty M_n^f E$.

Proposition 5.2. *For a spectrum E , there are equivalences*

$$\begin{aligned} \lambda : (L_n^f \Omega^\infty E)\langle d_n \rangle &\simeq (\Omega^\infty L_n^f E)\langle d_n \rangle && \text{for } n \geq 0, \\ \lambda : (M_n^f \Omega^\infty E)\langle d_n \rangle &\simeq (\Omega^\infty M_n^f E)\langle d_n \rangle && \text{for } n \geq 1. \end{aligned}$$

Proof. The case $n = 0$ is clear. For $n \geq 1$, these maps are equivalences since they are $\bigoplus_{i=0}^n v_i^{-1}\pi_*$ -equivalences of $\bigoplus_{i=0}^n v_i^{-1}\pi_*$ -local spaces in $Ho_*\langle d_n \rangle$ by Corollary 4.8 and Proposition 5.1. \square

In [Kuh2] for $n \geq 1$, Kuhn introduced a remarkable functor $\phi_n : Ho_* \rightarrow Ho_*^s$ with $\phi_n \Omega^\infty E \simeq L_{K(n)} E$ for $E \in Ho^s$. This greatly extended a similar result of [Bou6] for $n = 1$. We shall derive a telescopic Kuhn functor $\Phi_n : Ho_* \rightarrow Ho^s$ with $\Phi_n \Omega^\infty E \simeq \widehat{M}_n^f E$, and thus with $L_{K(n)} \Phi_n \Omega^\infty E \simeq L_{K(n)} E$.

Theorem 5.3 (Telescopic Kuhn theorem). *For $n \geq 1$, there is a functor $\Phi_n : Ho_* \rightarrow Ho^s$ such that*

- (i) $\Phi_n X$ is in $\widehat{\mathcal{M}}_n^f$ for each $X \in Ho_*$;
- (ii) for each $W \in \mathcal{F}_n$, there are isomorphisms

$$v_n^{-1}\pi_*(X; W) \cong [W, \Phi_n X]_* \cong v_n^{-1}\pi_*(\Phi_n X; W)$$

which are natural in $X \in Ho_*$;

- (iii) for $E \in Ho^s$, there is a natural equivalence $\Phi_n \Omega^\infty E \simeq \widehat{M}_n^f E$;
- (iv) for $K, X \in Ho_*$, there is a natural equivalence $\Phi_n (L_n^f X)^K \simeq (\Phi_n X)^K$;
- (v) Φ_n preserves homotopy fiber squares.

A homotopy (co)fiber square in Ho_* or Ho^s is a square represented by a homotopy (co)fiber square of pointed spaces or spectra in the usual sense [BF, A.2]. A more functorial version of 5.3(v) is provided in 5.5 below. We shall prove this theorem in 13.8, and we have the following companion theorem which is the key result of this paper.

Theorem 5.4. *For $n \geq 1$, there is a functor $\Theta_n : Ho^s \rightarrow Ho_*$ such that*

- (i) $\Theta_n E$ is in \mathcal{UN}_n^f (and hence in \mathcal{UM}_n^f) for each $E \in Ho^s$;
- (ii) for $E \in Ho^s$ and $X \in Ho_*$, there is a natural equivalence

$$\text{map}_*(\Theta_n E, L_n^f X) \simeq \text{map}_*(E, \Phi_n X);$$

- (iii) for $E \in Ho^s$, there is a natural equivalence $L_n^f \Sigma^\infty \Theta_n E \simeq M_n^f E$;
- (iv) for $E \in Ho^s$ and $K \in Ho_*$, there is a natural equivalence $\Theta_n (K \wedge E) \simeq L_n^f (K \wedge \Theta_n E)$;
- (v) Θ_n carries a homotopy cofiber square in Ho^s to an L_n^f -homotopy cofiber square in Ho_* .

An L_n^f -homotopy cofiber square of pointed spaces (in \mathcal{SS}_*) is a commutative square

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D \end{array}$$

such that the map from the homotopy pushout of $C \leftarrow A \rightarrow B$ to D is an L_n^f -equivalence; an L_n^f -homotopy cofiber square in Ho_* is a square represented by such a diagram of pointed spaces. We shall prove Theorem 5.4 in 13.9.

5.5. Simplicial functoriality. As we shall see in 12.7, the functors $\Phi_n : Ho_* \rightarrow Ho^s$ and $\Theta_n : Ho^s \rightarrow Ho_*$ for $n \geq 0$ can be induced by simplicial functors $\Phi_n : \mathcal{SS}_* \rightarrow \mathcal{S}p$ and $\Theta_n : \mathcal{S}p \rightarrow \mathcal{SS}_*$ on the categories \mathcal{SS}_* and $\mathcal{S}p$ of pointed simplicial sets and spectra (see 10.3). Moreover, the functor $\Phi_n : \mathcal{SS}_* \rightarrow \mathcal{S}p$ preserves homotopy fiber squares, while $\Theta_n : \mathcal{S}p \rightarrow \mathcal{SS}_*$ carries homotopy cofiber squares to L_n^f -homotopy cofiber squares.

We now derive some consequences of Theorems 5.3 and 5.4.

Corollary 5.6. *For $n \geq 1$, the functor $\Theta_n : Ho^s \rightarrow \mathcal{UL}_n^f$ is left adjoint to $\Phi_n : \mathcal{UL}_n^f \rightarrow Ho^s$.*

Proof. This follows since there is a natural isomorphism $[\Theta_n E, L_n^f X] \cong [E, \Phi_n X]$ for $E \in Ho^s$ and $X \in Ho_*$ by 5.4(ii). □

Corollary 5.7. *For $n \geq 1$, $G \in \mathcal{L}_n^f$, and $X \in \widehat{Ho^s}$, there is a natural isomorphism $\widetilde{G}^*(\Theta_n X) \cong (\widehat{M}_n^f G)^* X$. In particular, $\widetilde{K}(n)^*(\Theta_n X) \cong K(n)^* X$ and $\widetilde{K}(i)^*(\Theta_n X) = 0$ for $0 \leq i < n$. Likewise, $\widetilde{K}^*(\Theta_1 X; \widehat{Z}_p) \cong K^*(X; \widehat{Z}_p)$.*

Proof. There are isomorphisms

$$[\Theta_n X, \Omega^\infty G] \cong [X, \Phi_n \Omega^\infty G] \cong [X, \widehat{M}_n^f G]$$

by Corollary 5.6 and Theorem 5.3(iii). □

Using the $K(n)_*$ -localization functor $L_{K(n)}$, we obtain a closely related result:

Corollary 5.8. *There is a natural equivalence $L_{K(n)}\Sigma^\infty\Theta_n X \simeq L_{K(n)}X$ for $n \geq 1$ and $X \in Ho^s$.*

Proof. This follows by applying $L_{K(n)}$ to the equivalence $L_n^f\Sigma^\infty\Theta_n X \simeq M_n^f X$ of Theorem 5.4(iii). □

Thus each spectrum is $K(n)$ -equivalent to a suspension spectrum. The preceding results show that Θ_n behaves somewhat like an infinite desuspension, and we similarly have

Corollary 5.9. *For a space $A \in Ho_*$ with a map $\nu : \Sigma^d A \rightarrow A$ such that $\Sigma^\infty\nu$ is a v_n -map in \mathcal{F}_n , there is a canonical equivalence $\Theta_n\Sigma^\infty(\Sigma^2 A) \simeq L_n^f(\Sigma^2 A)$.*

Proof. Since $\Sigma^\infty\text{cof } \nu$ is in \mathcal{F}_{n+1} , $[\Sigma^i\text{cof } \nu, X]$ is trivial for $X \in \mathcal{UL}_n^f$ and $i \geq 1$. Hence there is an isomorphism $[\overline{A}, X] \cong v_n^{-1}\pi_0(X; \Sigma^\infty\overline{A})$ for $\overline{A} = \Sigma^2 A$, and there are natural isomorphisms

$$[L_n^f\overline{A}, X] \cong v_n^{-1}\pi_0(X; \Sigma^\infty\overline{A}) \cong [\Sigma^\infty\overline{A}, \Phi_n X] \cong [\Theta_n\Sigma^\infty\overline{A}, X]$$

for $X \in \mathcal{UL}_n^f$ by Theorem 5.3(ii) and Corollary 5.6. Consequently, $\Theta_n\Sigma^\infty\overline{A} \simeq L_n^f\overline{A}$. □

Corollary 5.10. *For $n \geq 1$, we have*

- (i) *a map $f : X \rightarrow Y$ in Ho_* is a $v_n^{-1}\pi_*$ -equivalence if and only if $\Phi_n f : \Phi_n X \simeq \Phi_n Y$;*
- (ii) *a map $g : E \rightarrow F$ in Ho^s is a $v_n^{-1}\pi_*$ -equivalence if and only if $\Theta_n g : \Theta_n E \simeq \Theta_n F$.*

Proof. Part (i) follows by 5.3(i) and (ii). For part (ii), note that $\Theta_n g : \Theta_n E \simeq \Theta_n F$ if and only if $g^* : [F, \Phi_n X] \cong [E, \Phi_n X]$ for each $X \in \mathcal{UL}_n^f$. By 5.3(i) and (iii), this is equivalent to $g^* : [F, M] \cong [E, M]$ for each $M \in \widehat{\mathcal{M}}_n^f$, and hence to $\widehat{M}_n^f g : \widehat{M}_n^f E \simeq \widehat{M}_n^f F$. □

Thus we see that $\Phi_n(Y\langle i \rangle) \simeq \Phi_n Y$ and $\Theta_n(E\langle i \rangle) \simeq \Theta_n E$ for each i . In particular, Φ_n kills Postnikov spaces and Θ_n kills Postnikov spectra. Finally, we will show that the stable monocular homotopy category \mathcal{M}_n^f is a retract of the unstable monocular homotopy category \mathcal{UN}_n^f in two different ways. For this we need

Lemma 5.11. *For $n \geq 0$ and $X \in Ho_*$, the map $X \rightarrow L_n^f X$ is an $S(n)_*$ -equivalence.*

Proof. This follows as in [Bou7, 12.1] since $L_n^f X$ is the nullification of X with respect to an $S(n)_*$ -acyclic space $\check{A}(n+1)$. □

Lemma 5.12. *For $n \geq 1$, the functor $L_n^f\Sigma^\infty : \mathcal{UN}_n^f \rightarrow \mathcal{M}_n^f$ is left adjoint to $\tilde{\Omega}^\infty : \mathcal{M}_n^f \rightarrow \mathcal{UN}_n^f$ where $\tilde{\Omega}^\infty E = (\Omega^\infty E)\langle d_n \rangle$.*

Proof. For a space $X \in \mathcal{UN}_n^f$, the spectrum $L_n^f\Sigma^\infty X$ is in \mathcal{M}_n^f because X and $\Sigma^\infty X$ are $S(n-1)_*$ -acyclic by Lemma 5.11 since $L_{n-1}^f X \simeq *$. For a spectrum $E \in \mathcal{M}_n^f$, the space $\tilde{\Omega}^\infty E$ is in \mathcal{UN}_n^f since $\tilde{\Omega}^\infty E \simeq (M_n^f\Omega^\infty E)\langle d_n \rangle$ by Proposition 5.2. The result now follows easily from the adjointness of Σ^∞ and Ω^∞ . □

Lemma 5.13. *For $n \geq 1$, the functor $\Theta_n : \mathcal{M}_n^f \rightarrow \mathcal{UN}_n^f$ is left adjoint to $M_n^f \Phi_n : \mathcal{UN}_n^f \rightarrow \mathcal{M}_n^f$.*

Proof. This follows by 5.4(i), 5.6, and 3.3. □

By 5.3(iii), 5.4(iii), 3.3, and the above lemmas, we have

Theorem 5.14. *For $n \geq 1$, the adjoint compositions*

$$\begin{array}{ccccc} \mathcal{M}_n^f & \xrightarrow{\Theta_n} & \mathcal{UN}_n^f & \xrightarrow{L_n^f \Sigma^\infty} & \mathcal{M}_n^f \\ \mathcal{M}_n^f & \xleftarrow{M_n^f \Phi_n} & \mathcal{UN}_n^f & \xleftarrow{\tilde{\Omega}^\infty} & \mathcal{M}_n^f \end{array}$$

each express \mathcal{M}_n^f as a categorical retract of \mathcal{UN}_n^f ; that is, the compositions $(L_n^f \Sigma^\infty)\Theta_n$ and $(M_n^f \Phi_n)\tilde{\Omega}^\infty$ are each equivalent to the identity functor on \mathcal{M}_n^f .

For $n \geq 1$, we let $\tilde{\Omega}^\infty \mathcal{M}_n^f \subset \mathcal{UN}_n^f$ and $\Theta_n \mathcal{M}_n^f \subset \mathcal{UN}_n^f$ denote the image subcategories of the faithful embeddings $\tilde{\Omega}^\infty : \mathcal{M}_n^f \rightarrow \mathcal{UN}_n^f$ and $\Theta_n : \mathcal{M}_n^f \rightarrow \mathcal{UN}_n^f$. Clearly, $\tilde{\Omega}^\infty \mathcal{M}_n^f$ consists of infinite loop spaces, while $\Theta_n \mathcal{M}_n^f$ consists of “infinite L_n^f -suspension spaces” since there is a natural equivalence $\Theta_n E \simeq L_n^f \Sigma^k (\Theta_n \Sigma^{-k} E)$ for $E \in Ho^s$ and $k \geq 0$ by Corollary 5.4(iv). Finally, we easily see

Proposition 5.15. *For spectra $E, G \in \mathcal{M}_n^f$ and $n \geq 1$, there is a natural isomorphism $[\Theta_n E, \tilde{\Omega}^\infty G] \cong [E, G]$.*

Thus a map from a space in $\Theta_n \mathcal{M}_n^f$ to a space in $\tilde{\Omega}^\infty \mathcal{M}_n^f$ corresponds to a stable map in \mathcal{M}_n^f .

6. THE UNSTABLE TELESCOPE CONJECTURE

In this section, we formulate the unstable telescope conjecture and prove that it is equivalent to Ravenel’s stable telescope conjecture. Our proof relies heavily on the functors Φ_n and Θ_n . We start by recalling

6.1. The stable telescope conjecture for $K(n)_*$. For $n \geq 0$, this asserts that *each $K(n)_*$ -equivalence of spectra is a $v_n^{-1} \pi_*$ -equivalence.* This is equivalent to Ravenel’s original telescope conjecture (see [Rav1], [MS], or 6.13 below). It holds trivially for $n = 0$ and holds for $n = 1$ by work of Mahowald [Mah] and Miller [Mil1], but has been strongly doubted for $n \geq 2$ since 1990 when Ravenel [Rav2] sketched a refutation for $n = 2$ and $p \geq 5$. It has the following obvious converse.

Theorem 6.2. *For $n \geq 0$, each $v_n^{-1} \pi_*$ -equivalence of spectra is a $K(n)_*$ -equivalence.*

Proof. This follows by 3.7 since $\langle T(n) \rangle \geq \langle T(n) \wedge K(n) \rangle = \langle K(n) \rangle$. □

Thus the stable telescope conjecture would provide a perfect Whitehead theorem for $K(n)_*$ -equivalences. To formulate an unstable telescope conjecture, we use the following concept from [Bou9]. For $n \geq 1$, a map $X \rightarrow Y$ in Ho_* is called a *virtual $K(n)_*$ -equivalence* if it induces a $K(n)_*$ -equivalence $(\Omega X)\langle n+2 \rangle \rightarrow (\Omega Y)\langle n+2 \rangle$. Results of [Bou9, Section 11] are combined to give

Theorem 6.3. *For $n \geq 1$ we have*

- (i) *each $K(n)_*$ -equivalence of connected H -spaces is a virtual $K(n)_*$ -equivalence;*
- (ii) *an i -connected cover $X\langle i \rangle \rightarrow X$ of a pointed space is a virtual $K(n)_*$ -equivalence for $i \geq 0$;*

- (iii) if any two of three pointed space maps $f : X \rightarrow Y$, $g : Y \rightarrow Z$ and $gf : X \rightarrow Z$ are virtual $K(n)_*$ -equivalences, then so is the third;
- (iv) for a map of homotopy fiber sequences of pointed spaces

$$\begin{array}{ccccc} F & \longrightarrow & X & \longrightarrow & B \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ F' & \longrightarrow & X' & \longrightarrow & B' \end{array}$$

- if any two of α , β , and γ are virtual $K(n)_*$ -equivalences, then so is the third;
- (v) if $f : X \rightarrow Y$ is a virtual $K(n)_*$ -equivalence of pointed spaces, then $f\langle i \rangle : X\langle i \rangle \rightarrow Y\langle i \rangle$ is a $K(n)_*$ -equivalence for $i \geq n + 2$.

Although an ordinary $K(n)_*$ -equivalence of spaces (such as the Barratt-Priddy map $K(\Sigma_\infty, 1) \rightarrow \Omega_0^\infty S$) need not be a $v_n^{-1}\pi_*$ -equivalence, we had hoped to prove

6.4. The unstable telescope conjecture for $K(n)_*$. For $n \geq 1$, this asserts that each virtual $K(n)_*$ -equivalence of pointed spaces is a $v_n^{-1}\pi_*$ -equivalence. This holds for $n = 1$ by [Bou9, 11.11], but now seems likely to fail for higher n by

Theorem 6.5. For each $n \geq 1$, the unstable telescope conjecture for $K(n)_*$ is equivalent to the stable telescope conjecture for $K(n)_*$.

This theorem will be proved in 6.10. The converse of the unstable telescope conjecture holds for $n = 1$ by [Bou9, 11.11] but fails for $n \geq 2$; for instance, $\Omega^\infty K(1)$ is $v_n^{-1}\pi_*$ -trivial but not virtually $K(n)_*$ -acyclic by Theorem 6.3(v) and [Bou10]. However, by [Bou9, 11.13], we have

Theorem 6.6. For $n \geq 1$, each $\bigoplus_{i=1}^n v_i^{-1}\pi_*$ -equivalence of spaces is a virtual $K(n)_*$ -equivalence.

In particular, the map $X \rightarrow L_n^f X$ is a virtual $K(n)_*$ -equivalence for each pointed space X by 4.6. The proof of Theorem 6.5 will be based on the following lemmas.

Lemma 6.7. For $n \geq 1$, if $h : E \rightarrow E'$ is a $K(n)_*$ -equivalence of spectra, then $\Omega^\infty M_n^f h : \Omega^\infty M_n^f E \rightarrow \Omega^\infty M_n^f E'$ is a virtual $K(n)_*$ -equivalence of spaces.

Proof. Using the natural $\bigoplus_{i=0}^n v_i^{-1}\pi_*$ -equivalence $\Sigma^\infty \Theta_n E \rightarrow M_n^f E$ of Theorem 5.4(iii), we obtain a commutative diagram

$$\begin{array}{ccc} \Omega^\infty \Sigma^\infty \Theta_n E & \longrightarrow & \Omega^\infty M_n^f E \\ \downarrow & & \downarrow \\ \Omega^\infty \Sigma^\infty \Theta_n E' & \longrightarrow & \Omega^\infty M_n^f E' \end{array}$$

whose horizontal maps are virtual $K(n)_*$ -equivalences by Theorem 6.6. The map $\Theta_n E \rightarrow \Theta_n E'$ is a $K(n)_*$ -equivalence by Corollary 5.7. Hence the left map is a $K(n)_*$ -equivalence by [Bou4, Remark after 1.2], or [Kuh1, Proposition 2.4], and is a virtual $K(n)_*$ -equivalence by Theorem 6.3(i). Therefore the right map is also a virtual $K(n)_*$ -equivalence. □

The converse of the above lemma holds by

Lemma 6.8. *For $n \geq 1$ and a map $h : E \rightarrow E'$ of spectra, if $\Omega^\infty h : \Omega^\infty E \rightarrow \Omega^\infty E'$ is a virtual $K(n)_*$ -equivalence of spaces, then $h : E \rightarrow E'$ and $\Phi_n \Omega^\infty h : \Phi_n \Omega^\infty E \rightarrow \Phi_n \Omega^\infty E'$ are $K(n)_*$ -equivalences.*

Proof. Since $(\Omega^\infty h)\langle n + 2 \rangle$ is a $K(n)_*$ -equivalence by Theorem 6.3(v), so is the map $h\langle n + 2 \rangle$ by [Bou4, Proposition 1.4]. Hence the map h is a $K(n)_*$ -equivalence since any spectrum whose homotopy groups are bounded above is $K(n)_*$ -acyclic because its rational and torsion parts are $K(n)_*$ -acyclic by [Rav1, Theorem 4.8]. Thus, since the maps $E \rightarrow \widehat{M}_n^f E$ and $E' \rightarrow \widehat{M}_n^f E'$ are also $K(n)_*$ -equivalences by Theorem 6.2, so is $\widehat{M}_n^f h$. Hence, $\Phi_n \Omega^\infty h$ is a $K(n)_*$ -equivalence by Theorem 5.3(iii). □

More generally, we have

Lemma 6.9. *For $n \geq 1$, if $g : X \rightarrow Y$ is a virtual $K(n)_*$ -equivalence of spaces, then $\Phi_n g : \Phi_n X \rightarrow \Phi_n Y$ is a $K(n)_*$ -equivalence of spectra.*

Proof. The map $L_n^f g$ is a virtual $K(n)_*$ -equivalence by Theorem 6.6, and hence $\text{map}_*(F, L_n^f g)$ is a virtual $K(n)_*$ -equivalence for each finite cell complex $F \in Ho_*$ by Theorem 6.3. Let $A \in Ho_*$ be a finite cell complex with a map $\nu : \Sigma^d A \rightarrow A$ such that $\Sigma^\infty A$ is a finite p -local spectrum of type n and $\Sigma^\infty \nu$ is a v_n -map. Then the virtual $K(n)_*$ -equivalence $\text{map}_*(\Sigma^2 A, L_n^f g)$ is an infinite loop map, where the infinite loop structure of $\text{map}_*(\Sigma^2 A, L_n^f X)$ comes from the natural equivalences

$$\text{map}_*(\Sigma^2 A, L_n^f X) \simeq \text{map}_*(\Sigma^{d+2} A, L_n^f X) \simeq \Omega^d \text{map}_*(\Sigma^2 A, L_n^f X)$$

induced by $\nu : \Sigma^d A \rightarrow A$. Hence $\text{map}_*(\Sigma^2 A, L_n^f g)$ induces a $K(n)_*$ -equivalence $\text{map}_*(\Sigma^2 A, \Phi_n g)$ by Lemma 6.8 and Theorem 5.3(iv). Using Spanier-Whitehead duality, we conclude that $\Phi_n g$ is a $K(n)_*$ -equivalence. □

6.10. Proof of Theorem 6.5. Given the unstable telescope conjecture for $K(n)_*$, let $h : E \rightarrow E'$ be a $K(n)_*$ -equivalence of spectra. Then $\Omega^\infty \widehat{M}_n^f h$ is a virtual $K(n)_*$ -equivalence by Lemma 6.7, and hence is a $v_n^{-1} \pi_*$ -equivalence. Thus $\widehat{M}_n^f h$ and h are $v_n^{-1} \pi_*$ -equivalences. Conversely, given the stable telescope conjecture for $K(n)_*$, let $f : X \rightarrow Y$ be a virtual $K(n)_*$ -equivalence of spaces. Then $\Phi_n f$ is a $K(n)_*$ -equivalence by Lemma 6.9 and hence is a $v_n^{-1} \pi_*$ -equivalence. Thus f is a $v_n^{-1} \pi_*$ -equivalence by Theorem 5.3(ii). □

More explicitly, a stable counterexample to the telescope conjecture will lead to a strong unstable counterexample.

Theorem 6.11. *Suppose that, for some $n \geq 2$, the stable telescope conjecture fails for $K(n)_*$. Then, for each $j > d_n$, there exists a j -connected infinite loop space Y such that $\widetilde{E}(n)_* Y = 0$, $v_n^{-1} \pi_* Y \neq 0$, $v_i^{-1} \pi_* Y = 0$ for each $i \neq n$, $\widetilde{S}(n)_* Y \neq 0$, and $\widetilde{S}(n-1)_* Y = 0$.*

Proof. Choose a spectrum E with $v_n^{-1} \pi_* E \neq 0$ and $K(n)_* E = 0$, and let $Y = (\Omega^\infty \widehat{M}_n^f E)\langle j \rangle$. Then clearly $v_i^{-1} \pi_* Y = 0$ for $i \neq n$ and $v_n^{-1} \pi_* Y \neq 0$. Hence, $\widetilde{S}(n-1)_* Y = 0$ by Corollary 4.8 and Lemma 5.11, and $\widetilde{S}(n)_* Y \neq 0$ by [Bou7, 13.13]. By Lemma 6.7, $\Omega^\infty \widehat{M}_n^f E$ is virtually $K(n)_*$ -acyclic, and hence $\widetilde{K}(n)_* Y = 0$

by Theorem 6.3(v) since $j > d_n \geq n + 1$. Thus $\widetilde{E(n)}_* Y = 0$ since $\langle E(n) \rangle = \langle E(n-1) \vee K(n) \rangle \leq \langle S(n-1) \vee K(n) \rangle$. □

6.12. Variants of the telescope conjecture. Since each $S(n)_*$ -equivalence of spectra is an $E(n)_*$ -equivalence for $n \geq 0$, there is a natural map $L_n^f X \rightarrow L_n X$ for $X \in Ho^s$. Moreover, since $\langle E(n) \rangle = \langle K(0) \vee \dots \vee K(n) \rangle$ and $\langle S(n) \rangle = \langle T(0) \vee \dots \vee T(n) \rangle$, the following conditions are equivalent for each $n \geq 0$:

- (i) the stable telescope conjecture for $K(m)_*$ holds for $0 \leq m \leq n$;
- (ii) each $E(n)_*$ -equivalence of spectra is an $S(n)_*$ -equivalence;
- (iii) the map $L_n^f X \rightarrow L_n X$ is an equivalence for each spectrum X .

In [Bou7, 12.3], we considered a variant of the unstable telescope conjecture asserting that, for $n \geq 0$, each $E(n)_*$ -equivalence of spaces is an $S(n)_*$ -equivalence. This holds for $n = 1$, but will fail for some $n \geq 2$ by Theorem 6.11 if the stable telescope conjecture for $K(n)_*$ fails.

We conclude by showing that the stable telescope conjecture for $K(n)_*$ is equivalent to Ravenel’s original telescope conjecture [Rav1]. For $n \geq 1$, we consider the v_n -telescope \widehat{W} given by the homotopy direct limit of

$$W \xrightarrow{\omega} \Sigma^{-d}W \xrightarrow{\omega} \Sigma^{-2d}W \longrightarrow \dots$$

for a finite p -local spectrum W of type n with a v_n -map $\omega : \Sigma^d W \rightarrow W$. Ravenel’s original telescope conjecture asserts that the map $\widehat{W} \rightarrow L_n W$ is an equivalence, and we have

Proposition 6.13. *The map $\widehat{W} \rightarrow L_n W$ is an equivalence if and only if the stable telescope conjecture holds for $K(n)_*$.*

Proof. Note that $\widehat{W} \simeq L_n^f W$ as in the proof of 3.6. First assume that the stable telescope conjecture holds for $K(n)_*$. Then the $K(n)_*$ -equivalence $W \rightarrow L_n W$ is a $v_n^{-1}\pi_*$ -equivalence, and is also an $S(n-1)_*$ -equivalence since $S(n-1)_* W = 0$ and $L_n W \simeq L_n S \wedge W$ by the smash product theorem of Hopkins and Ravenel [Rav3]. Thus $W \rightarrow L_n W$ is an $S(n)_*$ -localization and $L_n^f W \simeq L_n W$. Conversely, assume that $L_n^f W \simeq L_n W$. Then

$$\langle K(n) \rangle = \langle E(n) \wedge W \rangle = \langle L_n W \rangle = \langle L_n^f W \rangle = \langle S(n) \wedge W \rangle = \langle T(n) \rangle$$

and the stable telescope conjecture holds for $K(n)_*$. □

7. THE $E(n)_*$ -LOCALIZATIONS OF POSTNIKOV H -SPACES

We now determine the $E(n)_*$ -localizations and $K(n)_*$ -localizations of Postnikov H -spaces and of other H -spaces which are virtually $K(n)_*$ -acyclic. We hope that our results can eventually be extended from H -spaces to nilpotent spaces. They depend on work of Hopkins, Ravenel, and Wilson [HRW], and will be needed in Section 8. For a nilpotent space X , we use the p -localization $X_{(p)}$, the rationalization X_Q , and the p -completion \widehat{X}_p of [BK]. We start by recalling

7.1. Modified Postnikov sections of p -complete nilpotent spaces. Recall that a nilpotent space is p -complete if and only if its homotopy groups are Ext- p -complete in the sense of [BK]. For an Ext- p -complete nilpotent group A , there is a short exact sequence $0 \rightarrow \widehat{\tau}A \rightarrow A \rightarrow A/\widehat{\tau}A \rightarrow 0$ where $\widehat{\tau}A$ is the Ext- p -completion

of the torsion subgroup of A , and where $A/\widehat{\tau}A$ is the greatest torsion-free Ext- p -complete quotient of A . For a p -complete nilpotent space Y and $n \geq 0$, let $Y \rightarrow \widehat{P}_\tau^{n+1}Y$ denote the *modified $n + 1$ -Postnikov section* with

$$\pi_i \widehat{P}_\tau^{n+1}Y = \begin{cases} \pi_i Y & \text{if } i < n + 1, \\ \pi_{n+1} Y / \widehat{\tau} \pi_{n+1} Y & \text{if } i = n + 1, \\ 0 & \text{if } i > n + 1. \end{cases}$$

A space will be called *Postnikov* when its homotopy groups vanish above some dimension. The following key result will be proved in 7.13.

Theorem 7.2. *If X is a connected H -space which is virtually $K(n)_*$ -acyclic (e.g. Postnikov) for some $n \geq 1$, then $L_{K(n)}X \simeq \widehat{P}_\tau^{n+1}\widehat{X}_p$.*

The $K(n)_*$ -localizations of spaces are closely related to $E(n)_*$ -localizations by

Theorem 7.3. *If X is a connected H -space or a simply connected pointed space and $n \geq 1$, then $L_{K(n)}X \simeq (L_n X)_p^\wedge$ and there is a homotopy fiber square*

$$\begin{array}{ccc} L_n X & \longrightarrow & L_{K(n)} X \\ \downarrow & & \downarrow \\ X_Q & \longrightarrow & (L_{K(n)} X)_Q. \end{array}$$

This follows by [Bou10, 1.6]. For a pointed nilpotent space Y and $n \geq 0$, let $Y \rightarrow P_\tau^{n+1}Y$ denote the *modified $n + 1$ -Postnikov section* with

$$\pi_i P_\tau^{n+1}Y = \begin{cases} \pi_i Y & \text{if } i < n + 1, \\ \pi_{n+1} Y / \tau \pi_{n+1} Y & \text{if } i = n + 1, \\ 0 & \text{if } i > n + 1, \end{cases}$$

where $\tau \pi_{n+1} Y$ is the torsion subgroup of $\pi_{n+1} Y$. We can now determine the $E(n)_*$ -localization of a Postnikov H -space.

Theorem 7.4. *If X is a connected H -space which is virtually $K(n)_*$ -acyclic (e.g. Postnikov) for some $n \geq 1$, then*

$$\pi_i L_n X = \begin{cases} \pi_i X \otimes Z_{(p)} & \text{if } i < n + 1, \\ (\pi_{n+1} X / \tau \pi_{n+1} X) \otimes Z_{(p)} & \text{if } i = n + 1, \\ \pi_i X \otimes Q & \text{if } i > n + 1. \end{cases}$$

Proof. This follows since the combined square and the right square of

$$\begin{array}{ccccc} L_n X & \longrightarrow & P_\tau^{n+1} X_{(p)} & \longrightarrow & \widehat{P}_\tau^{n+1} \widehat{X}_p \\ \downarrow & & \downarrow & & \downarrow \\ X_Q & \longrightarrow & P_\tau^{n+1} X_Q & \longrightarrow & (\widehat{P}_\tau^{n+1} \widehat{X}_p)_Q \end{array}$$

are both homotopy fiber squares by 7.2, 7.3, and the arithmetic square theorem [DDK]. Hence, the left square is also a homotopy fiber square. \square

To prove Theorem 7.2, we need a series of lemmas.

Lemma 7.5. *If $X = K(A, m)$ for an abelian group A and $m \geq 1$, then $L_{K(n)}X \simeq \widehat{P}_\tau^{n+1}\widehat{X}_p$ for each $n \geq 1$.*

Proof. This follows by [Bou5, 6.3] since $K(Z/p, j)$ and $K(Z, j + 1)$ are $K(n)_*$ -nonacyclic for $1 \leq j \leq n$ and are $K(n)_*$ -acyclic for $j > n$ by [RW] and [HRW, 4.4]. □

Lemma 7.6. *For $n \geq 1$, if $X \in Ho_*$ is a virtually $K(n)_*$ -acyclic nilpotent space, then the map $X \rightarrow \widehat{P}_\tau^{n+1}\widehat{X}_p$ is a $K(n)_*$ -equivalence.*

Proof. The map $X \rightarrow P_\tau^{n+1}X$ is a $K(n)_*$ -equivalence since the spaces $X\langle n + 2 \rangle$, $K(\pi_{n+2}X, n + 2)$, and $K(\tau\pi_{n+1}X, n + 1)$ are $K(n)_*$ -acyclic by Theorem 6.3(v) and Lemma 7.5. The map $P_\tau^{n+1}X \rightarrow \widehat{P}_\tau^{n+1}\widehat{X}_p$ is a $K(n)_*$ -equivalence since it is an H/p_* -equivalence. □

Lemma 7.7. *An Ext- p -complete nilpotent group A is adjusted (i.e. $\widehat{\tau}A = A$) if and only if $\text{Hom}(A, \widehat{Z}_p) = 0$.*

Proof. The upper central series quotients of $A/\widehat{\tau}A$ are torsion-free Ext- p -complete abelian groups by [BK, p. 174] and [War, p. 6]. The lemma now follows since \widehat{Z}_p is a retract of each nontrivial torsion-free Ext- p -complete abelian group by [BK, p. 181]. □

Lemma 7.8. *For $n \geq 1$, if $f : X \rightarrow Y$ is a $K(n)_*$ -equivalence of pointed nilpotent spaces, then the homotopy fiber of $\widehat{f}_p : \widehat{X}_p \rightarrow \widehat{Y}_p$ is an $(n - 1)$ -connected p -complete space F with $\pi_n F$ adjusted.*

Proof. Since $K(Z/p, n)$ and $K(\widehat{Z}_p, n + 1)$ are $K(n)_*$ -local, by Lemma 7.5, we have $f_* : H^i(Y; Z/p) \cong H^i(X; Z/p)$ and $f_* : H^{i+1}(Y; \widehat{Z}_p) \cong H^{i+1}(X; \widehat{Z}_p)$ for $i \leq n$. Hence, $f_* : \pi_i \widehat{X}_p \rightarrow \pi_i \widehat{Y}_p$ is an isomorphism for $i < n$ and onto for $i = n$ by [Bou5, 5.2]. Thus the homotopy fiber of $\widehat{f}_p : \widehat{X}_p \rightarrow \widehat{Y}_p$ is an $(n - 1)$ -connected p -complete space F with $H^0(\widehat{Y}_p; H^n(F; \widehat{Z}_p)) = 0$ by the Serre spectral sequence. Since $H^n(F; \widehat{Z}_p) = \text{Hom}(\pi_n F, \widehat{Z}_p)$ and since $\pi_1 \widehat{X}_p$ acts nilpotently on $\pi_n F$ by [BK, pp. 60, 61], we conclude that $H^n(F; \widehat{Z}_p) = 0$. Thus the Ext- p -complete nilpotent group $\pi_n F$ is adjusted by Lemma 7.7, and the result follows. □

For a nilpotent space $X \in Ho_*$ and $j \geq 1$, let $X\langle j \rangle_\tau$ denote the homotopy fiber of the map $X \rightarrow P_\tau^j X$. Generalizing a result of Hopkins, Ravenel, and Wilson [HRW], we obtain

Lemma 7.9. *If X is a connected Postnikov H -space, then the map $X\langle j \rangle_\tau \rightarrow X$ is a $K(n)_*$ -monomorphism for $j, n \geq 1$.*

Proof. Since X is a connected H -space, it is a retract of $\Omega\Sigma X$, and thus is a retract of ΩY where Y is a sufficiently large Postnikov section of ΣX . Hence it suffices by naturality to show that the map $(\Omega Y)\langle j \rangle_\tau \rightarrow \Omega Y$ is a $K(n)_*$ -monomorphism for each simply connected Postnikov space Y and $j, n \geq 1$. We can assume that Y has trivial 1-skeleton and view Y as the homotopy direct limit of Postnikov sections of its finite subcomplexes. Thus by a limit argument, we can assume that Y is a simply connected Postnikov space with finitely generated homotopy groups, and by [HRW, 2.3], we can also assume that $\pi_2 Y$ is torsion. After replacing Y by the homotopy fiber of $Y \rightarrow Y[1/p]$, we obtain the desired result from Theorem 2.1 of [HRW]. □

For a p -complete nilpotent space $X \in Ho_*$ and $j \geq 1$, let $X\langle \widehat{j} \rangle_\tau$ denote the homotopy fiber of the map $X \rightarrow \widehat{P}_\tau^j X$.

Lemma 7.10. *If X is a p -complete connected Postnikov H -space, then $X\langle j \rangle_\tau \rightarrow X$ is a $K(n)_*$ -monomorphism for $j, n \geq 1$.*

Proof. This follows from Lemma 7.9 since $X\langle j \rangle_\tau$ is the p -completion of $X\langle j \rangle_\tau$. \square

Lemma 7.11. *If $u : A \rightarrow B$ is a monomorphism of p -torsion abelian groups, then $K(u, j) : K(A, j) \rightarrow K(B, j)$ is a $K(n)_*$ -monomorphism for $j, n \geq 1$.*

Proof. The result holds for an inclusion $Z/p^s \rightarrow Z/p^\infty$ with $1 \leq s < \infty$ by [RW, 13.1] and [HRW, 4.4], and hence holds for a finite direct sum of such inclusions by the Kunnetth theorem. When B is finite, we may choose such finite direct sums $\alpha : A \rightarrow I$ and $\beta : B \rightarrow J$, and then use the divisibility of I and J to construct a splittable monomorphism $\bar{u} : I \rightarrow J$ with $\bar{u}\alpha = \beta u$. Thus the result holds when B is finite, and holds in general by a direct limit argument. \square

Lemma 7.12. *For $n \geq 1$, if X is a p -complete connected H -space with $X \simeq \widehat{P}_\tau^{n+1}X$, then X is $K(n)_*$ -local.*

Proof. The $K(n)_*$ -localization $X \rightarrow L_{K(n)}X$ is a virtual $K(n)_*$ -equivalence by Theorem 6.3(i), and thus $L_{K(n)}X \rightarrow \widehat{P}_\tau^{n+1}L_{K(n)}X$ is a $K(n)_*$ -equivalence by Lemma 7.6. Hence $L_{K(n)}X$ is a retract of $\widehat{P}_\tau^{n+1}L_{K(n)}X$ and $L_{K(n)}X \simeq \widehat{P}_\tau^{n+1}L_{K(n)}X$. By Lemma 7.8, the homotopy fiber sequence $F \rightarrow X \rightarrow L_{K(n)}X$ has $\widehat{P}_\tau^n F \simeq *$ and is therefore covered by a homotopy fiber sequence $F \rightarrow X\langle n \rangle_\tau \rightarrow (L_{K(n)}X)\langle n \rangle_\tau$ which is of the form $K(A', n)_p^\wedge \rightarrow K(A, n)_p^\wedge \rightarrow K(A'', n)_p^\wedge$ for a short exact sequence $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ of p -torsion abelian groups obtained as follows. For any p -complete nilpotent space Y with $\pi_1 Y$ adjusted, there is a natural equivalence $Y \simeq (t_p Y)_p^\wedge$ by [Bou7, 14.1], where $t_p Y$ is the “ p -torsion part” of Y given by the homotopy fiber of the rationalization map $Y \rightarrow Y_Q$. Moreover, if $\pi_* Y$ is trivial except for an adjusted group $\pi_n Y$ and a torsion-free group $\pi_{n+1} Y$, then $t_p Y \simeq K(\pi_n t_p Y, n)$ where $\pi_n t_p Y$ is a p -torsion group. Thus the desired homotopy fiber sequence $K(A', n) \rightarrow K(A, n) \rightarrow K(A'', n)$ may be obtained by applying t_p to $F \rightarrow X\langle n \rangle_\tau \rightarrow (L_{K(n)}X)\langle n \rangle_\tau$. Since $X \rightarrow L_{K(n)}X$ is a $K(n)_*$ -equivalence, $K(A, n) \rightarrow K(A'', n)$ is a $K(n)_*$ -monomorphism by Lemma 7.10, and $K(A', n) \rightarrow K(A, n)$ is a $K(n)_*$ -monomorphism by Lemma 7.11. Thus $K(A', n)$ is $K(n)_*$ -acyclic and $A' = 0$ by Lemma 7.5. Hence, $F \simeq K(A', n)_p^\wedge \simeq *$ and $X \simeq L_{K(n)}X$. \square

7.13. Proof of Theorem 7.2. By Lemmas 7.6 and 7.12, the map $X \rightarrow \widehat{P}_\tau^{n+1}\widehat{X}_p$ is a $K(n)_*$ -equivalence to a $K(n)_*$ -local space. \square

8. $E(n)_*$ -LOCALIZATIONS OF INFINITE LOOP SPACES

For a spectrum X , we study the $E(n)_*$ -localization $L_n \Omega_0^\infty X$ of the connected infinite loop space $\Omega_0^\infty X = (\Omega^\infty X)(0)$. There is a natural comparison map $L_n \Omega_0^\infty X \rightarrow \Omega_0^\infty L_n X$ since the space $\Omega_0^\infty L_n X$ is $E(n)_*$ -local. In [Bou4], we showed that $L_1 \Omega_0^\infty X$ is given by $\Omega_0^\infty L_1 X$ with low dimensional modifications. We now obtain a similar result for $L_2 \Omega_0^\infty X$, and more generally for $L_n \Omega_0^\infty X$ when $L_{n-1}^f X \simeq L_{n-1} X$. The latter condition would be automatic if the telescope conjecture were valid, and holds when X is a BP -module spectrum or any other BP -cotrivial spectrum (see 6.12 and 8.9).

Theorem 8.1. For $n \geq 1$ and a spectrum X with $L_{n-1}^f X \simeq L_{n-1} X$, the natural map

$$L_n \Omega_0^\infty X \rightarrow \Omega_0^\infty L_n X \times_{P_\tau^{n+1} \Omega_0^\infty L_n X} P_\tau^{n+1} \Omega_0^\infty X_{(p)}$$

is a π_i -isomorphism for $i \neq n + 1$ and is a π_{n+1} -monomorphism. The condition $L_{n-1}^f X \simeq L_{n-1} X$ holds for all spectra when $n \leq 2$, and the natural map is always an equivalence when $n = 1$.

This will be proved in 8.6.

Corollary 8.2. For $n \geq 1$ and a spectrum X with $L_{n-1}^f X \simeq L_{n-1} X$, the natural map

$$L_{K(n)} \Omega_0^\infty X \rightarrow (\Omega_0^\infty L_n X)_p^\wedge \times_{\widehat{P}_\tau^{n+1}(\Omega_0^\infty L_n X)_p^\wedge} \widehat{P}_\tau^{n+1}(\Omega_0^\infty X)_p^\wedge$$

is a π_i -isomorphism for $i \neq n + 1, n + 2$ and is a π_{n+2} -monomorphism. The natural map is always an equivalence when $n = 1$.

This follows from Theorems 8.1 and 7.3. To prove Theorem 8.1, we need three lemmas.

Lemma 8.3. For $n \geq 1$, if $h : X \rightarrow Y$ is a $K(n)_*$ -equivalence and an L_{n-1}^f -equivalence of spectra, then $\Omega^\infty h : \Omega^\infty X \rightarrow \Omega^\infty Y$ is a virtual $K(n)_*$ -equivalence of spaces.

Proof. In the map of fiber sequences

$$\begin{array}{ccccc} \Omega^\infty M_n^f X & \longrightarrow & \Omega^\infty L_n^f X & \longrightarrow & \Omega^\infty L_{n-1}^f X \\ \downarrow & & \downarrow & & \downarrow \simeq \\ \Omega^\infty M_n^f Y & \longrightarrow & \Omega^\infty L_n^f Y & \longrightarrow & \Omega^\infty L_{n-1}^f Y \end{array}$$

the vertical arrows are virtual $K(n)_*$ -equivalences by Lemma 6.7 and Theorem 6.3(iv). Since the maps $\Omega^\infty X \rightarrow \Omega^\infty L_n^f X$ and $\Omega^\infty Y \rightarrow \Omega^\infty L_n^f Y$ are also virtual $K(n)_*$ -equivalences by Theorem 6.6, so is the map $\Omega^\infty X \rightarrow \Omega^\infty Y$. \square

Lemma 8.4. For $n \geq 1$, if X is a spectrum with $L_{n-1}^f X \simeq L_{n-1} X$, then the map $L_n \Omega_0^\infty X \rightarrow \Omega_0^\infty L_n X$ is a π_i -isomorphism for $i > n + 1$ and a π_{n+1} -monomorphism.

Proof. Since $S(n - 1)$ and $E(n)$ are smashing with $\langle E(n - 1) \rangle = \langle S(n - 1) \wedge E(n) \rangle$, we have $L_{n-1} X \simeq L_{n-1}^f L_n X$ and hence $L_{n-1}^f X \simeq L_{n-1}^f L_n X$. Thus Lemma 8.3 shows that the map of spectra $X \rightarrow L_n X$ induces a virtual $K(n)_*$ -equivalence of spaces $\Omega_0^\infty X \rightarrow \Omega_0^\infty L_n X$. Since the $E(n)_*$ -localization $\Omega_0^\infty X \rightarrow L_n \Omega_0^\infty X$ is also a virtual $K(n)_*$ -equivalence by Theorem 6.3(i), so is the map $L_n \Omega_0^\infty X \rightarrow \Omega_0^\infty L_n X$. Thus its homotopy fiber F is virtually $K(n)_*$ -acyclic. Since F is an $E(n)_*$ -local, p -torsion, infinite loop space by [Bou4], we have $F \simeq P_\tau^{n+1} F$ by Theorem 7.4, and $\pi_i F = 0$ for $i \geq n + 1$. \square

Lemma 8.5. For $n \geq 1$, each $E(n)_*$ -equivalence of p -local connected H -spaces is a P_τ^{n+1} -equivalence.

Proof. Let $\mathcal{C} \subset Ho_*$ be the full subcategory of all p -local connected H -spaces Y with $P_\tau^{n+1} Y \simeq Y$. Since each $Y \in \mathcal{C}$ is $E(n)_*$ -local by Theorem 7.4, an $E(n)_*$ -equivalence $f : A \rightarrow B$ in Ho_* induces a bijection $f^* : [B, Y] \cong [A, Y]$, and thus induces a bijection $(P_\tau^{n+1} f)^* : [P_\tau^{n+1} B, Y] \cong [P_\tau^{n+1} A, Y]$ for all $Y \in \mathcal{C}$. When

A and B are p -local connected H -spaces, so are $P_\tau^{n+1}A$ and $P_\tau^{n+1}B$, and thus $P_\tau^{n+1}f : P_\tau^{n+1}A \rightarrow P_\tau^{n+1}B$ is an equivalence in \mathcal{C} by the Yoneda lemma. \square

8.6. Proof of Theorem 8.1. The theorem follows for $n = 1$ by [Bou4]. Since $L_1^f X \simeq L_1 X$ holds for all X (see 6.12), we may assume $n \geq 2$ with $L_{n-1}^f X \simeq L_{n-1} X$. Thus the map $L_n \Omega_0^\infty X \rightarrow \Omega_0^\infty L_n X$ is a π_i -isomorphism for $i > n + 1$ and a π_{n+1} -monomorphism by Lemma 8.4, and the map $\Omega_0^\infty X_{(p)} \rightarrow L_n \Omega_0^\infty X$ is a P_τ^{n+1} -equivalence by Lemma 8.5. The theorem now follows easily. \square

8.7. A counterexample. Unfortunately, when $n = 2$, the natural map of Theorem 8.1 may actually fail to be a π_3 -isomorphism. To see this, note that for $1 \leq j \leq \infty$, the map $\Sigma^\infty M(Z/p^j, 3)\langle 3 \rangle \rightarrow \Sigma^\infty M(Z/p^j, 3)$ is carried to an equivalence by $\Omega_0^\infty L_2$, $P_\tau^3 \Omega_0^\infty L_2$, and $P_\tau^3 \Omega_0^\infty$, but not by $L_2 \Omega_0^\infty$, because we have

Lemma 8.8. *For $1 \leq j \leq \infty$, the map $QM(Z/p^j, 3)\langle 3 \rangle \rightarrow QM(Z/p^j, 3)$ of spaces is not a $K(2)_*$ -monomorphism.*

Proof. In the principal fiber sequence

$$QM(Z_{p^\infty}, 2) \longrightarrow K(Z_{p^\infty}, 2) \longrightarrow QM(Z_{p^\infty}, 3)\langle 3 \rangle,$$

$K(2)_*QM(Z_{p^\infty}, 2) \cong K(2)_*QS^3$ is an exterior algebra on odd dimensional generators by Kashiwabara [Kas]. Thus since $K(2)_*K(Z_{p^\infty}, 2)$ is concentrated in even dimensions, a bar spectral sequence argument shows that the map $K(Z_{p^\infty}, 2) \rightarrow QM(Z_{p^\infty}, 3)\langle 3 \rangle$ is a $K(2)_*$ -monomorphism. Since the map $K(Z_{p^j}, 2) \rightarrow K(Z_{p^\infty}, 2)$ is also a $K(2)_*$ -monomorphism by Lemma 7.11, we see that the map $K(Z/p^j, 2) \rightarrow QM(Z/p^j, 3)\langle 3 \rangle$ is a nontrivial $K(2)_*$ -monomorphism for $1 \leq j \leq \infty$. Hence, the map $QM(Z/p^j, 3)\langle 3 \rangle \rightarrow QM(Z/p^j, 3)$ cannot be a $K(2)_*$ -monomorphism. \square

Finally, we show that our results on $L_n \Omega^\infty X$ and $L_{K(n)} \Omega^\infty X$ apply whenever X is a BP -cotrivial spectrum in the sense of 2.8. We note that the BP -cotrivial spectra include the BP -module spectra and the BP -nilpotent spectra in the sense of [Bou2, 3.7].

Lemma 8.9. *If X is a BP -cotrivial spectrum and $n \geq 0$, then $L_n^f X \simeq L_n X$.*

Proof. Let \mathcal{C} be the class of all spectra X with $L_n^f X \simeq L_n X$. Since $L_n^f X \simeq X \wedge L_n^f S$, $L_n X \simeq X \wedge L_n S$, and $BP \wedge L_n^f S \simeq BP \wedge L_n S$ by [Rav4, 2.7], we have (i) \mathcal{C} contains BP ; (ii) \mathcal{C} is closed under arbitrary wedges; and (iii) \mathcal{C} contains all three spectra in a cofiber sequence when it contains any two. Hence, \mathcal{C} contains the BP -cotrivial spectra by 2.8. \square

9. MODEL CATEGORIES AND NULLIFICATIONS

To prepare for the proofs of Theorems 5.3 and 5.4, we now give a much improved version of the Bousfield-Friedlander [BF] construction of new model categories from old ones with “ Q -structures,” and then show that each nullification functor determines a proper model category structure on spaces. By a model category, we mean a closed model category in Quillen’s original sense [Qui]. We refer the reader to [DS], [GJ], [Hir], and [Hov] for good recent or forthcoming treatments of model categories, and we note that our results are fully compatible with the slightly refined definitions of a “model category” used in [Hir], [Hov], and elsewhere. We shall need

9.1. Proper model categories. A model category \mathcal{C} is called:

- (i) *right proper* if every pullback of a weak equivalence along a fibration is a weak equivalence;
- (ii) *left proper* if every pushout of a weak equivalence along a cofibration is a weak equivalence;
- (iii) *proper* if it is both right proper and left proper.

As explained in [BF] or [Hir], an elementary theory of homotopy pullbacks (resp. pushouts) is available in a right (resp. left) proper model category. All model categories used in this paper will turn out to be proper.

9.2. Q -structures on proper model categories. Let \mathcal{C} be a proper model category with a functor $Q : \mathcal{C} \rightarrow \mathcal{C}$ and natural transformation $\alpha : 1 \rightarrow Q$ satisfying the following axioms:

- (A1) if $f : X \rightarrow Y$ is a weak equivalence in \mathcal{C} , then so is $Qf : QX \rightarrow QY$;
- (A2) for each $X \in \mathcal{C}$, the maps $\alpha, Q\alpha : QX \rightarrow QQX$ are weak equivalences;
- (A3) for a pullback square

$$\begin{array}{ccc} V & \xrightarrow{k} & X \\ \downarrow g & & \downarrow f \\ W & \xrightarrow{h} & Y \end{array}$$

in \mathcal{C} , if f is a fibration of fibrant objects such that $\alpha : X \rightarrow QX, \alpha : Y \rightarrow QY$, and $Qh : QW \rightarrow QY$ are weak equivalences, then $Qk : QV \rightarrow QX$ is a weak equivalence.

A map $f : X \rightarrow Y$ in \mathcal{C} will be called a *Q -equivalence* if $Qf : QX \rightarrow QY$ is a weak equivalence, a *Q -cofibration* if f is a cofibration, and a *Q -fibration* if f has the right lifting property for Q -trivial cofibrations (the Q -cofibrations which are Q -equivalences). We let \mathcal{C}^Q denote the category \mathcal{C} equipped within its Q -equivalences, Q -cofibrations, and Q -fibrations. The following theorem is a much improved version of Theorem A.7 from [BF]. Our present axiom (A3) replaces the hard-to-verify axiom (A.6), which involved general Q -fibrations and included a dual pushout condition.

Theorem 9.3. *If \mathcal{C} is a proper model category with a functor $Q : \mathcal{C} \rightarrow \mathcal{C}$ and transformation $\alpha : 1 \rightarrow Q$ satisfying (A1), (A2), and (A3), then \mathcal{C}^Q is also a proper model category. Moreover, a map $f : X \rightarrow Y$ in \mathcal{C} is a Q -fibration if and only if f is a fibration and*

$$\begin{array}{ccc} X & \xrightarrow{\alpha} & QX \\ \downarrow f & & \downarrow Qf \\ Y & \xrightarrow{\alpha} & QY \end{array}$$

is a homotopy fiber square in \mathcal{C} .

Proof. The proof proceeds as in [BF, Appendix A] with a new construction of factorizations into Q -trivial cofibrations and Q -fibrations, and with a new verification

of properness. To factor a map $f : X \rightarrow Y$, take the commutative diagram

$$\begin{CD} X @>\alpha>> QX @>u>> \widehat{QX} \\ @VfVV @VQfVV @V\widehat{Qf}VV \\ Y @>\alpha>> QY @>v>> \widehat{QY} \end{CD}$$

where u and v are trivial cofibrations and $\widehat{Qf} : \widehat{QX} \rightarrow \widehat{QY}$ is a fibration of fibrant objects. Then \widehat{Qf} is a Q -fibration by [BF, (A.8)], and \widehat{Qf} pulls back along the Q -equivalence $Y \rightarrow \widehat{QY}$ to give a Q -fibration $g : E \rightarrow Y$ such that the map $E \rightarrow \widehat{QX}$ is a Q -equivalence by (A3). Hence the map $X \rightarrow E$ is a Q -equivalence and has a factorization into a Q -trivial cofibration $j : X \rightarrow D$ and a trivial fibration $k : D \rightarrow E$. Now $f = (kj)j$ is the desired factorization of f . The resulting model category \mathcal{C}^Q is left proper since the pushout of a Q -equivalence along a cofibration may be obtained by successively pushing out a Q -trivial cofibration and a weak equivalence. It is right proper by (A3) and the following lemma. \square

Lemma 9.4. *A model category \mathcal{D} is right proper if every pullback of a weak equivalence along a fibration of fibrant objects in \mathcal{D} is a weak equivalence.*

Proof. Let

$$\begin{CD} V @>k>> X \\ @VgVV @VfVV \\ W @>h>> Y \end{CD}$$

be a pullback of a weak equivalence h along a fibration f in \mathcal{D} . Construct a fibration $\widehat{f} : \widehat{X} \rightarrow \widehat{Y}$ of fibrant objects \widehat{X} and \widehat{Y} with trivial cofibrations $u : X \rightarrow \widehat{X}$ and $v : Y \rightarrow \widehat{Y}$ such that $vf = \widehat{f}u$. Let $f' : X' \rightarrow Y$ and $g' : V' \rightarrow W$ be the induced fibrations of \widehat{f} . Since the maps $X' \rightarrow \widehat{X}$ and $V' \rightarrow \widehat{X}$ are weak equivalences by hypothesis, so are the maps $V' \rightarrow X'$ and $X \rightarrow X'$. By Ken Brown's lemma (see [DS, 9.9]), the functor $h^* : \mathcal{D} \downarrow Y \rightarrow \mathcal{D} \downarrow W$ preserves weak equivalences of fibrant objects. Thus h^* carries the weak equivalence $f \rightarrow f'$ to a weak equivalence $g \rightarrow g'$. Hence the maps $V \rightarrow V'$ and $V \rightarrow X$ are weak equivalences. \square

Remark 9.5. Theorem 9.3 remains valid when “proper” is replaced by “right proper.” However, the proof then requires another new argument to verify Lemma A.8(iii) of [BF]. This can be given using a lemma of Dan Kan proved in [Hir, 11.1.16].

9.6. Simplicial model categories. By a *simplicial model category*, we mean a closed simplicial model category in Quillen's original sense [Qui, II §2], but without any finiteness conditions on simplicial sets in the axiom SM0. In particular, a simplicial model category \mathcal{C} is enriched over the category \mathcal{SS} of simplicial sets and is both tensored and cotensored (see e.g. [Bor]). For $X, Y \in \mathcal{C}$ and $K \in \mathcal{SS}$, we write $\text{map}(X, Y) \in \mathcal{SS}$, $X \otimes K \in \mathcal{C}$, and $X^K = \text{map}(K, X) \in \mathcal{C}$ for the mapping space, tensor, and cotensor. Theorem 9.3 now extends to

Theorem 9.7. *Under the hypotheses of Theorem 9.3, if \mathcal{C} is a proper simplicial model category, then so is \mathcal{C}^Q .*

Proof. It suffices to verify Quillen’s axiom SM7(b) for \mathcal{C}^Q . The only nontrivial step is to show that if $A \rightarrow B$ is a Q -trivial cofibration, then so is

$$F_n : B \otimes \partial\Delta^n \coprod_{A \otimes \partial\Delta^n} A \otimes \Delta^n \longrightarrow B \otimes \Delta^n$$

for $n \geq 0$. Let $\Lambda_0^n \subset \Delta^n$ and $\partial\Delta^n \subset \Delta^n$ be the 0-horn and boundary of the standard n -simplex. In the pushout square

$$\begin{array}{ccc} B \otimes \partial\Delta^{n-1} \coprod_{A \otimes \partial\Delta^{n-1}} A \otimes \Delta^{n-1} & \longrightarrow & B \otimes \Lambda_0^n \coprod_{A \otimes \Lambda_0^n} A \otimes \Delta^n \\ \downarrow F_{n-1} & & \downarrow G_n \\ B \otimes \Delta^{n-1} & \longrightarrow & B \otimes \partial\Delta^n \coprod_{A \otimes \partial\Delta^n} A \otimes \Delta^n \end{array}$$

we can assume inductively that F_{n-1} is a Q -trivial cofibration and deduce that G_n is also a Q -trivial cofibration. Since \mathcal{C} is a simplicial model category and $\Lambda_0^n \rightarrow \Delta^n$ is a trivial cofibration, we see that $F_n G_n$ is a trivial cofibration. Likewise, F_n is a cofibration. Hence F_n is a Q -trivial cofibration. \square

9.8. The nullification functors. We now work in the proper simplicial model category $\mathcal{S}\mathcal{S}_*$ of pointed spaces (i.e. pointed simplicial sets). For each pointed space W , we consider the W -nullification (= W -periodization) functor $P_W : \mathcal{S}\mathcal{S}_* \rightarrow \mathcal{S}\mathcal{S}_*$ and transformation $\alpha : 1 \rightarrow P_W$ of [Bou7, 2,10] or [Dro], which induce the W -nullification functor $P_W : Ho_* \rightarrow Ho_*$ and transformation $\alpha : 1 \rightarrow P_W$ of 4.1. Using the terminology of 9.2, we have

Theorem 9.9. *If W is a pointed space, then $\mathcal{S}\mathcal{S}_*^{P_W}$ is a proper simplicial model category. Moreover, a map $f : X \rightarrow Y$ in $\mathcal{S}\mathcal{S}_*$ is a P_W -fibration if and only if f is a fibration and*

$$\begin{array}{ccc} X & \xrightarrow{\alpha} & P_W X \\ \downarrow f & & \downarrow P_W f \\ Y & \xrightarrow{\alpha} & P_W Y \end{array}$$

is a homotopy fiber square in $\mathcal{S}\mathcal{S}_$.*

Proof. It suffices by Theorem 9.7 to verify our three axioms for $P_W : \mathcal{S}\mathcal{S}_* \rightarrow \mathcal{S}\mathcal{S}_*$. Axioms (A1) and (A2) follow since P_W induces an idempotent functor on Ho_* . For (A3), we must show that if

$$\begin{array}{ccc} A & \xrightarrow{k} & C \\ \downarrow g & & \downarrow f \\ B & \xrightarrow{h} & D \end{array}$$

is a pullback square in $\mathcal{S}\mathcal{S}_*$ where f is a fibration of W -null fibrant spaces and where h is a P_W -equivalence, then k is a P_W -equivalence. When C and D are connected, this follows by [Bou7, 4.8(i)] since the homotopy fiber of h is P_W -acyclic by [Bou7, 4.3]. The result now follows in general since the functor $P_W : \mathcal{S}\mathcal{S}_* \rightarrow \mathcal{S}\mathcal{S}_*$ comes from a functor $P_W : \mathcal{S}\mathcal{S} \rightarrow \mathcal{S}\mathcal{S}$ which preserves disjoint unions when W is connected, and which takes contractible values when W is not connected. \square

9.10. The homotopy category \mathcal{P}_W . For a pointed space W , we let $\mathcal{P}_W \subset Ho_*$ be the full subcategory of W -null spaces (see 4.1). This is the homotopy category of fibrant cofibrant objects in $\mathcal{S}\mathcal{S}_*^{P_W}$, and is thus canonically equivalent to the Quillen homotopy category $Ho(\mathcal{S}\mathcal{S}_*^{P_W})$ obtained by inverting the P_W -equivalences in $\mathcal{S}\mathcal{S}_*$.

Remark 9.11. The W -nullification functors P_W are special cases of the f -localization functors L_f constructed for arbitrary maps f of pointed spaces in [Bou7], [Bou9], [Dro], and [Hir]. In general, there is a left proper simplicial model category structure on $\mathcal{S}\mathcal{S}_*$ whose weak equivalences are the L_f -equivalences and whose cofibrations are the usual injections (see [Bou9, 4.6] or [Hir]). However, this model category will not be right proper unless L_f is equivalent to a nullification, as we now explain. By [Bou9, Theorem 4.4], for an arbitrary map f of pointed spaces, there exists a pointed space $A(f)$ such that $P_{A(f)}$ and L_f have the same acyclic spaces. Moreover, each $P_{A(f)}$ -equivalence of pointed spaces is an L_f -equivalence. Now suppose that the model category structure for L_f on $\mathcal{S}\mathcal{S}_*$ is right proper. Then for a pointed connected space X , the homotopy fiber of $X \rightarrow L_f X$ is L_f -acyclic, since a path fibration over $L_f X$ is an L_f -fibration by [BF, A.3(iii)]. Hence, this homotopy fiber is also $P_{A(f)}$ -acyclic, and $X \rightarrow L_f X$ is a $P_{A(f)}$ -equivalence by [Bou7, Corollary 4.8(i)]. Thus each L_f -equivalence of pointed connected spaces is a $P_{A(f)}$ -equivalence, and this generalizes to nonconnected spaces as in the proof of Theorem 9.9. Hence L_f is equivalent to $P_{A(f)}$.

10. THE FUNCTORS Φ_A AND Θ_A

To prepare for the proofs of Theorems 5.3 and 5.4, we now introduce a Kuhn functor $\Phi_A : \mathcal{P}_W \rightarrow Ho^s$ and its left adjoint $\Theta_A : Ho^s \rightarrow \mathcal{P}_W$ for a “ P_W -cospectrum” A , where W is a fixed pointed space. These functors will be constructed more generally in Section 11. To establish notation and terminology, we first discuss

10.1. Pointed simplicial Quillen functors. A pointed simplicial model category \mathcal{C} is automatically enriched, tensored, and cotensored over $\mathcal{S}\mathcal{S}_*$. For $X, Y \in \mathcal{C}$ and $K \in \mathcal{S}\mathcal{S}_*$, we let $\text{map}_*(X, Y) \in \mathcal{S}\mathcal{S}_*$, $X \wedge K \in \mathcal{C}$, and $X^K = \text{map}_*(K, X) \in \mathcal{C}$ denote the pointed mapping space, tensor, and cotensor in \mathcal{C} . The Quillen homotopy category $Ho\mathcal{C}$ is consequently enriched, tensored, and cotensored over $Ho(\mathcal{S}\mathcal{S}_*) = Ho_*$. For pointed simplicial model categories \mathcal{C} and \mathcal{D} , a pair of functors $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ are called *pointed simplicial left and right Quillen functors* if F is pointed simplicially left adjoint to G and the following equivalent conditions are satisfied:

- (i) F preserves cofibrations and G preserves fibrations;
- (ii) F preserves cofibrations and trivial cofibrations;
- (iii) G preserves fibrations and trivial fibrations.

Thus, in the language of enriched category theory (see e.g. [Bor, p. 340]), F and G are $\mathcal{S}\mathcal{S}_*$ -functors such that F is left $\mathcal{S}\mathcal{S}_*$ -adjoint to G and they satisfy conditions (i)–(iii). By adjunction arguments, there are natural isomorphisms $F(X \wedge K) \cong F(X) \wedge K$ and $G(Y^K) \cong (GY)^K$ for $X \in \mathcal{C}$, $Y \in \mathcal{D}$, and $K \in \mathcal{S}\mathcal{S}_*$. Moreover, F (resp. G) preserves weak equivalences of cofibrant (resp. fibrant) objects by Ken Brown’s lemma [DS, 9.9]. Thus, F preserves homotopy cofiber squares of cofibrant objects, while G preserves homotopy fiber squares of fibrant objects.

By [Qui, I.4], F has a *total left derived functor* $\mathbf{L}F : Ho\mathcal{C} \rightarrow Ho\mathcal{D}$, and G has a *total right derived functor* $\mathbf{R}G : Ho\mathcal{D} \rightarrow Ho\mathcal{C}$, where $\mathbf{L}F$ is left adjoint to

RG. Moreover, there are natural equivalences $\mathbf{LF}(X \wedge K) \simeq \mathbf{LF}(X) \wedge K$ and $\mathbf{RG}(Y^K) \simeq (\mathbf{RG}Y)^K$ for $X \in Ho\mathcal{C}$, $Y \in Ho\mathcal{D}$, and $K \in Ho_*$. The functor \mathbf{LF} preserves homotopy cofiber squares, while \mathbf{RG} preserves homotopy fiber square.

The pointed simplicial left and right Quillen functors $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ are called *Quillen equivalences* when, for each cofibrant $X \in \mathcal{C}$ and fibrant $Y \in \mathcal{D}$, a map $FX \rightarrow Y$ is a weak equivalence if and only if its adjoint $X \rightarrow GY$ is a weak equivalence. This implies that the functors $\mathbf{LF} : Ho\mathcal{C} \rightleftarrows Ho\mathcal{D} : \mathbf{RG}$ are adjoint equivalences. To construct our Quillen functors, we use

10.2. P_W -cospectra. A *cospectrum* A consists of spaces $A^n \in \mathcal{SS}_*$ for $n \geq 0$ and structure maps $\sigma : A^{n+1} \wedge S^1 \rightarrow A^n$ where $S^1 = \Delta^1/\partial\Delta^1$. We call A a *P_W -cospectrum* if $\sigma : A^{n+1} \wedge S^1 \rightarrow A^n$ is a P_W -equivalence for $n \geq 0$. This implies that A^0 is an “infinite suspension space” in P_W -homotopy theory. We remark that a P_W -equivalence $J \wedge S^d \rightarrow J$, for a space J and $d > 0$, can be used to construct a P_W -cospectrum with successive spaces $J, J \wedge S^{d-1}, \dots, J \wedge S^1, J, \dots$ where $S^d = S^1 \wedge \dots \wedge S^1$.

10.3. Spectra. A *spectrum* E consists of spaces $E_n \in \mathcal{SS}_*$ for $n \geq 0$ with structure maps $\sigma : S^1 \wedge E_n \rightarrow E_{n+1}$. It has homotopy groups $\pi_*E = \text{colim}_n \pi_{*+n}E_n$. By [BF] or [HSS], there is a proper pointed simplicial model category \mathcal{Sp} of spectra, where a map $f : E \rightarrow F$ is a *weak equivalence* when $f_* : \pi_*E \cong \pi_*F$, a *cofibration* when f induces cofibrations $E_0 \rightarrow F_0$ and $E_{n+1} \coprod_{S^1 \wedge E_n} S^1 \wedge F_n \rightarrow F_{n+1}$ for $n \geq 0$, and a *fibration* when f induces fibrations $E_n \rightarrow F_n$ and homotopy fiber squares

$$\begin{array}{ccc} E_n & \longrightarrow & (QE)_n \\ \downarrow & & \downarrow \\ F_n & \longrightarrow & (QF)_n \end{array}$$

in \mathcal{SS}_* for $n \geq 0$ using an omegafication functor $Q : \mathcal{Sp} \rightarrow \mathcal{Sp}$. A map $f : E \rightarrow F$ is a trivial fibration in \mathcal{Sp} if and only if it induces trivial fibrations $E_n \rightarrow F_n$ in \mathcal{SS}_* for $n \geq 0$. We may identify $Ho(\mathcal{Sp})$ with the stable homotopy category Ho^s .

10.4. The functor $\phi_A : \mathcal{SS}_*^{P_W} \rightarrow \mathcal{Sp}$. For a P_W -cospectrum A , there is a pointed simplicial functor $\phi_A : \mathcal{SS}_*^{P_W} \rightarrow \mathcal{Sp}$ where $\phi_A(X)$ is the spectrum with $\phi_A(X)_n = X^{A^n}$ for $n \geq 0$ and with the induced structure maps. The functor ϕ_A has a left adjoint pointed simplicial functor $\theta_A : \mathcal{Sp} \rightarrow \mathcal{SS}_*^{P_W}$ carrying a spectrum E to the pointed space $\theta_A(E)$, which may be constructed as the quotient of $\bigvee_{n \geq 0} (A^n \wedge E_n)$

by the relations given by

$$A^{n+1} \wedge E_{n+1} \xleftarrow{1 \wedge \sigma} A^{n+1} \wedge S^1 \wedge E_n \xrightarrow{\sigma \wedge 1} A^n \wedge E_n$$

for $n \geq 0$. To show that θ_A and ϕ_A are left and right Quillen functors, we need

Lemma 10.5. *Let $T : \mathcal{C} \rightarrow \mathcal{D}$ be a functor of model categories. If \mathcal{C} is right proper and T preserves pullbacks, trivial fibrations, and fibrations of fibrant objects, then T preserves arbitrary fibrations.*

Proof. For a fibration $f : X \rightarrow Y$ in \mathcal{C} , construct a square

$$\begin{array}{ccc} X & \xrightarrow{u} & \widehat{X} \\ \downarrow f & & \downarrow \widehat{f} \\ Y & \xrightarrow{v} & \widehat{Y} \end{array}$$

where u and v are trivial cofibrations and \widehat{f} is a fibration of fibrant objects. Let $f' : X' \rightarrow Y$ be the pullback of \widehat{f} along v , and note that the maps $X \rightarrow X' \rightarrow \widehat{X}$ are weak equivalences since \mathcal{C} is right proper. Factor $X \rightarrow X'$ as the composite of a trivial cofibration $i : X \rightarrow X''$ and a trivial fibration $j : X'' \rightarrow X'$. Then the fibration $f : X \rightarrow Y$ is a retract of $jj' : X'' \rightarrow Y$ by the lifting axiom. Since T preserves the fibrations j and f' , it also preserves f . \square

Lemma 10.6. *For a P_W -cospectrum A , the functors $\theta_A : Sp \rightarrow \mathcal{SS}_*^{P_W}$ and $\phi_A : \mathcal{SS}_*^{P_W} \rightarrow Sp$ are pointed simplicial left and right Quillen functors.*

Proof. It suffices by Lemma 10.5 to show that ϕ_A preserves trivial fibrations and fibrations of fibrant objects. If $f : X \rightarrow Y$ is a trivial fibration in $\mathcal{SS}_*^{P_W}$ (and hence in \mathcal{SS}_*), then $\phi_A(f) : \phi_A(X) \rightarrow \phi_A(Y)$ is a levelwise trivial fibration of spectra, and hence is a trivial fibration in Sp . If $f : X \rightarrow Y$ is a fibration of fibrant objects in $\mathcal{SS}_*^{P_W}$, then $\phi_A(f) : \phi_A(X) \rightarrow \phi_A(Y)$ is a levelwise fibration of omega spectra, and hence is a fibration in Sp . \square

We let $\Theta_A : Ho^s \rightarrow \mathcal{P}_W$ and $\Phi_A : \mathcal{P}_W \rightarrow Ho^s$ denote the total left and right derived functors $\mathbf{L}\theta_A : Ho(Sp) \rightarrow Ho(\mathcal{SS}_*^{P_W})$ and $\mathbf{R}\phi_A : Ho(\mathcal{SS}_*^{P_W}) \rightarrow Ho(Sp)$. Combining Lemma 10.6 with 10.1, we have

Theorem 10.7. *For a P_W -cospectrum A , the functor $\Theta_A : Ho^s \rightarrow \mathcal{P}_W$ is left adjoint to $\Phi_A : \mathcal{P}_W \rightarrow Ho^s$ with $\text{map}_*(\Theta_A E, X) \simeq \text{map}_*(E, \Phi_A X)$ for $E \in Ho^s$ and $X \in \mathcal{P}_W$. Moreover, there are natural equivalences $\Theta_A(K \wedge E) \simeq P_W(K \wedge \Theta_A E)$ and $\Phi_A(X^K) \simeq (\Phi_A X)^K$ for $K \in Ho_*$, $E \in Ho^s$, and $X \in \mathcal{P}_W$. Furthermore, the functor $\Phi_A : \mathcal{P}_W \rightarrow Ho^s$ preserves homotopy fiber squares, while $\Theta_A : Ho^s \rightarrow \mathcal{P}_W$ carries homotopy cofiber squares to P_W -homotopy cofiber squares.*

10.8. The simplicial functoriality of Θ_A and Φ_A . The total derived functors $\Theta_A = \mathbf{L}\theta_A : Ho(Sp) \rightarrow Ho(\mathcal{SS}_*^{P_W})$ and $\Phi_A = \mathbf{R}\phi_A : Ho(\mathcal{SS}_*^{P_W}) \rightarrow Ho(Sp)$ are induced by simplicial functors $\Theta_A = \theta_A I : Sp \rightarrow \mathcal{SS}_*^{P_W}$ and $\Phi_A = \phi_A J : \mathcal{SS}_*^{P_W} \rightarrow Sp$, which preserve weak equivalences, where $I : Sp \rightarrow Sp$ is a simplicial cofibrant replacement functor giving a cofibrant spectrum $I(E)$ with a natural trivial fibration $I(E) \rightarrow E$ for $E \in Sp$, and where $J : \mathcal{SS}_*^{P_W} \rightarrow \mathcal{SS}_*^{P_W}$ is a simplicial fibrant replacement functor giving a fibrant (i.e. W -null Kan) complex $J(X) \in \mathcal{SS}_*^{P_W}$ with a natural trivial cofibration (i.e. monic P_W -equivalence) $X \rightarrow J(X)$ for $X \in \mathcal{SS}_*^{P_W}$. The functor $\Theta_A : Sp \rightarrow \mathcal{SS}_*^{P_W}$ preserves homotopy cofiber squares, while $\Phi_A : \mathcal{SS}_*^{P_W} \rightarrow Sp$ preserves homotopy fiber squares.

11. THE GENERAL FUNCTORS Φ_A AND Θ_A

Extending the preceding work, we now introduce a general Kuhn functor $\Phi_A : \mathcal{P}_W \rightarrow Ho^s$ and its left adjoint $\Theta_A : Ho^s \rightarrow \mathcal{P}_W$ for a “spectral P_W -cospectrum” A , where W is a fixed pointed space. We first discuss

11.1. Spectral P_W -cospectra. A *spectral P_W -cospectrum* A consists of P_W -cospectra A_n for $n \geq 0$ with structure maps $\sigma : S^1 \wedge A_n \rightarrow A_{n+1}$ which are levelwise cofibrations of cospectra (in the sense that $\sigma : S^1 \wedge A_n^k \rightarrow A_{n+1}^k$ is a cofibration in \mathcal{SS}_* for $k, n \geq 0$).

11.2. The functor $\bar{\phi}_A : \mathcal{SS}_*^{Pw} \rightarrow \mathcal{CspSp}$. A *cospectrum of spectra* E consists of spectra E^n for $n \geq 0$ with structure maps $\sigma : E^{n+1} \wedge S^1 \rightarrow E^n$. There is a proper pointed simplicial model category \mathcal{CspSp} of cospectra of spectra, where a map $f : E \rightarrow F$ in \mathcal{CspSp} is a *weak equivalence* (resp. *cofibration*) when f induces weak equivalences (resp. cofibrations) $E^n \rightarrow F^n$ in \mathcal{Sp} for $n \geq 0$, and a *fibration* (resp. *trivial fibration*) when f induces fibrations (resp. trivial fibrations) $E^0 \rightarrow F^0$ and

$$E^{n+1} \rightarrow F^{n+1} \times_{(F^n)^{S^1}} (E^n)^{S^1}$$

in \mathcal{Sp} for $n \geq 0$. For a spectral P_W -cospectrum A , there is a pointed simplicial functor $\bar{\phi}_A : \mathcal{SS}_*^{Pw} \rightarrow \mathcal{CspSp}$ where $\bar{\phi}_A(X)$ is the cospectrum of spectra with $\bar{\phi}_A(X)^n = \phi_{A_n}(X)$ for $n \geq 0$ and with the induced structure maps. Thus $\bar{\phi}_A(X)^n_m = \text{map}_*(A_n^m, X)$ for $m, n \geq 0$, and $\bar{\phi}_A$ has a pointed simplicial left adjoint $\bar{\theta}_A : \mathcal{CspSp} \rightarrow \mathcal{SS}_*^{Pw}$ carrying a cospectrum of spectra E to the pointed space $\bar{\theta}_A(E)$, which may be constructed as the quotient of $\bigvee_{m,n \geq 0} (A_n^m \wedge E_m^n)$ by the

relations given by

$$\begin{aligned} A_n^{m+1} \wedge E_{m+1}^n &\xleftarrow{1 \wedge \sigma} A_n^{m+1} \wedge S^1 \wedge E_m^n \xrightarrow{\sigma \wedge 1} A_n^m \wedge E_m^n \\ A_{n+1}^m \wedge E_m^{n+1} &\xleftarrow{\tau(1 \wedge \sigma)} E_m^{n+1} \wedge S^1 \wedge A_n^m \xrightarrow{\tau(\sigma \wedge 1)} A_n^m \wedge E_m^n \end{aligned}$$

for $m, n \geq 0$ where τ is the twisting map.

Lemma 11.3. *For a spectral P_W -cospectrum A , the functors $\bar{\theta}_A : \mathcal{CspSp} \rightarrow \mathcal{SS}_*^{Pw}$ and $\bar{\phi}_A : \mathcal{SS}_*^{Pw} \rightarrow \mathcal{CspSp}$ are pointed simplicial left and right Quillen functors.*

Proof. By Lemma 10.5, it suffices to show that $\bar{\phi}_A(f)$ is a trivial fibration (resp. fibration) in \mathcal{CspSp} whenever $f : X \rightarrow Y$ is a trivial fibration (resp. fibration of fibrant objects) in \mathcal{SS}_*^{Pw} . This follows since the induced maps $\phi_{A_0}(X) \rightarrow \phi_{A_0}(Y)$ and

$$\phi_{A_{n+1}}(X) \rightarrow \phi_{A_{n+1}}(Y) \times_{\phi_{A_n}(Y)^{S^1}} \phi_{A_n}(X)^{S^1}$$

are levelwise trivial fibrations of spectra (resp. levelwise fibrations of omega spectra), and thus are trivial fibrations (resp. fibrations) in \mathcal{CspSp} . \square

11.4. The functors $\beta : \mathcal{CspSp} \rightarrow \mathcal{TowSp}$. A *tower of spectra* $E = \{E^n\}_{n \geq 0}$ consists of spectra E^n with maps $E^{n+1} \rightarrow E^n$ for $n \geq 0$. There is a proper pointed simplicial model category \mathcal{TowSp} of towers of spectra, where a map $f : E \rightarrow F$ in \mathcal{TowSp} is a *weak equivalence* (resp. *cofibration*) when f induces weak equivalences (resp. cofibrations) $E^n \rightarrow F^n$ in \mathcal{Sp} for $n \geq 0$, and a *fibration* (resp. *trivial fibration*) when f induces fibrations (resp. trivial fibrations) $E^0 \rightarrow F^0$ and $E^{n+1} \rightarrow F^{n+1} \times_{F^n} E^n$ in \mathcal{Sp} for $n \geq 0$. There is a pointed simplicial functor $\beta : \mathcal{CspSp} \rightarrow \mathcal{TowSp}$ where $\beta(X)$ is the tower of spectra with $\beta(X)^n = X^n \wedge S^n$ for $n \geq 0$. The functor β has a pointed simplicial right adjoint $\gamma : \mathcal{TowSp} \rightarrow \mathcal{CspSp}$ where $\gamma(Y)$ is the cospectrum of spectra with $\gamma(Y)^n = (Y^n)^{S^n}$ for $n \geq 0$. It is now straightforward to prove the following.

Lemma 11.5. *The functors $\beta : \mathcal{C}sp\mathcal{S}p \rightarrow \mathcal{T}ow\mathcal{S}p$ and $\gamma : \mathcal{T}ow\mathcal{S}p \rightarrow \mathcal{C}sp\mathcal{S}p$ are pointed simplicial left and right Quillen equivalences.*

Lemma 11.6. *The constant tower functor $c^* : \mathcal{S}p \rightarrow \mathcal{T}ow\mathcal{S}p$ and the inverse limit functor $\lim : \mathcal{T}ow\mathcal{S}p \rightarrow \mathcal{S}p$ are pointed simplicial left and right Quillen functors.*

Finally, for a spectral P_W -cospectrum A , we let $\Theta_A : Ho^s \rightarrow \mathcal{P}_W$ and $\Phi_A : \mathcal{P}_W \rightarrow Ho^s_*$ denote the compositions of total derived functors

$$\begin{array}{ccccccc} Ho(\mathcal{S}p) & \xrightarrow{\mathbf{L}c^*} & Ho(\mathcal{T}ow\mathcal{S}p) & \xrightarrow{\mathbf{R}\gamma} & Ho(\mathcal{C}sp\mathcal{S}p) & \xrightarrow{\mathbf{L}\bar{\theta}_A} & Ho(\mathcal{S}\mathcal{S}^{P_W}_*) \\ Ho(\mathcal{S}p) & \xleftarrow{\mathbf{R}\lim} & Ho(\mathcal{T}ow\mathcal{S}p) & \xleftarrow{\mathbf{L}\beta} & Ho(\mathcal{C}sp\mathcal{S}p) & \xleftarrow{\mathbf{R}\bar{\phi}_A} & Ho(\mathcal{S}\mathcal{S}^{P_W}_*) \end{array}$$

Noting that $\mathbf{R}\gamma$ and $\mathbf{L}\beta$ are adjoint equivalences, we obtain

Theorem 11.7. *Theorem 10.7 remains valid when “a P_W -cospectrum A ” is replaced by “a spectral P_W -cospectrum A .”*

11.8. The simplicial functoriality of Φ_A and Θ_A . In each of the model categories $\mathcal{S}\mathcal{S}^{P_W}_*$, $\mathcal{S}p$, $\mathcal{C}sp\mathcal{S}p$, and $\mathcal{T}ow\mathcal{S}p$, it is straightforward to construct a simplicial (co)fibrant replacement functor. Thus since $\Theta_A : Ho(\mathcal{S}p) \rightarrow Ho(\mathcal{S}\mathcal{S}^{P_W}_*)$ and $\Phi_A : Ho(\mathcal{S}\mathcal{S}^{P_W}_*) \rightarrow Ho(\mathcal{S}p)$ are composites of total derived functors of Quillen functors linking these categories, they are induced by simplicial functors $\Theta_A : \mathcal{S}p \rightarrow \mathcal{S}\mathcal{S}_*$ and $\Phi_A : \mathcal{S}\mathcal{S}^{P_W}_* \rightarrow \mathcal{S}p$ which preserve weak equivalences, as in 10.8. Moreover, the functor $\Theta_A : \mathcal{S}p \rightarrow \mathcal{S}\mathcal{S}^{P_W}_*$ preserves homotopy cofiber squares, while $\Phi_A : \mathcal{S}\mathcal{S}^{P_W}_* \rightarrow \mathcal{S}p$ preserves homotopy fiber squares.

Finally, we establish a key property of Φ_A .

Theorem 11.9. *For a spectral P_W -cospectrum A and a $\Sigma^\infty W$ -trivial spectrum $E \in Ho^s$, there is a natural equivalence $\Phi_A(\Omega^\infty E) \simeq F(A^0_*, E)$ in Ho^s where A^0_* is the edge-spectrum of A .*

This will be proved in 11.11 using

11.10. Bispectra. A bispectrum X consists of spaces $X_{m,n} \in \mathcal{S}\mathcal{S}_*$ for $m, n \geq 0$ together with structure maps $\sigma_1 : S^1 \wedge X_{m,n} \rightarrow X_{m+1,n}$ and $\sigma_2 : S^1 \wedge X_{m,n} \rightarrow X_{m,n+1}$ such that the compositions

$$\sigma_1(1 \wedge \sigma_2), \sigma_2(1 \wedge \sigma_1)(\tau \wedge 1) : S^1 \wedge S^1 \wedge X_{m,n} \rightarrow X_{m+1,n+1}$$

are equal for $m, n \geq 0$ where τ is the twisting map. A bispectrum X is called *fibrant* when the spaces $X_{m,n}$ are fibrant and the structure maps $X_{m,n} \rightarrow (X_{m+1,n})^{S^1}$ and $X_{m,n} \rightarrow (X_{m,n+1})^{S^1}$ are weak equivalences for $m, n \geq 0$. If X is a fibrant bispectrum, then the edge spectra $X_{0,*}$ and $X_{*,0}$ are related by a chain of natural weak equivalences

$$\begin{array}{ccccccc} X_{0,*} & \longrightarrow & \text{Sing}|X_{0,*}| & \longleftarrow & \text{Sing Tel}|X_{0,*}| & \longrightarrow & \text{Sing Tel}^{(2)}|X_{*,*}| \\ & & \longleftarrow & \text{Sing Tel}|X_{*,0}| & \longrightarrow & \text{Sing}|X_{*,0}| & \longleftarrow & X_{*,0} \end{array}$$

in $\mathcal{S}p$, where Sing is the singular functor, $|-|$ is the geometric realization functor, Tel is the telescope functor of Adams [Ada, p. 171], and $\text{Tel}^{(2)}$ is the double telescope functor of Adams [Ada, pp. 173-176]. More generally, if X is a cospectrum of fibrant bispectra, then the edge objects $X_{0,*} \in \mathcal{C}sp\mathcal{S}p$ and $X_{*,0} \in \mathcal{C}sp\mathcal{S}p$ are related by a chain of natural weak equivalences in $\mathcal{C}sp\mathcal{S}p$ as displayed above.

11.11. Proof of Theorem 11.9. Let $\phi(A, E)$ be the cospectrum of bispectra with $\phi(A, E)_{m,n}^q = \text{map}_*(A_q^m, E_n)$ for $m, n, q \geq 0$. We may assume that the spectrum $E \in \mathcal{S}p$ is fibrant and conclude that $\phi(A, E)$ is a cospectrum of fibrant bispectra. Thus by 11.10 there is a natural equivalence $\phi(A_*^*, E_0) \simeq \phi(A_*^0, E_*)$ in $Ho(\mathcal{C}sp\mathcal{S}p)$ which may be written as $\bar{\phi}_A(\Omega^\infty E) \simeq \phi(A_*^0, E_*)$ since $E_0 = \Omega^\infty E$. This is carried to a natural equivalence $\Phi_A(\Omega^\infty E) \simeq F(A_*^0, E)$ by the functors

$$Ho(\mathcal{C}sp\mathcal{S}p) \xrightarrow{L\beta} Ho(\mathcal{T}ow\mathcal{S}p) \xrightarrow{\mathbf{R}\lim} Ho(\mathcal{S}p).$$

12. THE FUNCTORS Φ_n AND Θ_n

In final preparation for the proofs of Theorems 5.3 and 5.4, we now construct the required functors $\Phi_n : Ho_* \rightarrow Ho^s$ and $\Theta_n : Ho^s \rightarrow Ho_*$, and establish some of their formal properties. Throughout this section, we let $n \geq 1$ be a fixed integer, and we use the nullification functor $L_n^f = P_{\tilde{A}(n+1)}^f$. A spectral L_n^f -cospectrum B will be called *admissible* if the edge spectrum B_*^0 is equivalent to $C_{n-1}^f S_{(p)}$ in Ho^s and if the spectra $\Sigma^\infty B_i^j$ are in \mathcal{F}_n for $i, j \geq 0$.

Theorem 12.1. *For $n \geq 1$, there exists an admissible spectral L_n^f -cospectrum B .*

Proof. As in 3.2, let $F(n)$ be a finite p -local spectrum of type n . The $F(n)$ -trivialization of the p -local sphere spectrum $S_{(p)} \rightarrow L_{n-1}^f S_{(p)}$ is given by the mapping telescope of a sequence $S_{(p)} = Y_0 \rightarrow \dots \rightarrow Y_k \rightarrow Y_{k+1} \rightarrow \dots$ where the homotopy cofiber of $Y_k \rightarrow Y_{k+1}$ is a finite wedge of integral suspensions of $F(n)$. Thus, since $C_{n-1}^f S_{(p)}$ is the homotopy fiber of $S_{(p)} \rightarrow L_{n-1}^f S_{(p)}$, it is given by the mapping telescope of a sequence $X_0 \xrightarrow{u_0} X_1 \rightarrow \dots \rightarrow X_i \xrightarrow{u_i} X_{i+1} \rightarrow \dots$ of spectra in \mathcal{F}_n . Using the periodicity theorem of Hopkins and Smith (see 3.5), we may choose v_n -maps $\omega_i : X_i \wedge S^{d_i} \rightarrow X_i$ in Ho^s with $d_i | d_{i+1}$ and with $\omega_{i+1} u_i = u_i (\omega_i)^{q_i}$ for $i \geq 0$ and $q_i = d_{i+1}/d_i$. We may then desuspend these spectra and maps to give: spaces $\tilde{X}_i \in Ho_*$ for $i \geq 0$ with $\Sigma^\infty \tilde{X}_i = S^{r_i} \wedge X_i$ where $r_{i+1} \geq r_i \geq 0$; maps $\tilde{\omega}_i : \tilde{X}_i \wedge S^{d_i} \rightarrow \tilde{X}_i$ in Ho_* for $i \geq 0$ which are equivalent to double suspensions with $\Sigma^\infty \tilde{\omega}_i = S^{r_i} \wedge \omega_i$; and maps $\tilde{u}_i : S^{r_{i+1}-r_i} \wedge \tilde{X}_i \rightarrow \tilde{X}_{i+1}$ in Ho_* for $i \geq 0$ with $\Sigma^\infty \tilde{u}_i = S^{r_{i+1}} \wedge u_i$ and with

$$\begin{array}{ccc} S^{r_{i+1}-r_i} \wedge \tilde{X}_i \wedge S^{d_{i+1}} & \xrightarrow{\tilde{u}_i \wedge S^{d_{i+1}}} & \tilde{X}_{i+1} \wedge S^{d_{i+1}} \\ \downarrow S^{r_{i+1}-r_i} \wedge (\tilde{\omega}_i)^{q_i} & & \downarrow \tilde{\omega}_{i+1} \\ S^{r_{i+1}-r_i} \wedge \tilde{X}_i & \xrightarrow{\tilde{u}_i} & \tilde{X}_{i+1} \end{array}$$

commuting in Ho_* . The maps $\tilde{\omega}_i : \tilde{X}_i \wedge S^{d_i} \rightarrow \tilde{X}_i$ are L_n^f -equivalences by [Bou7, 11.12–11.14]. We next rigidify this homotopy system by choosing a representative sequence of fibrant spaces \tilde{X}_i and cofibrations $\tilde{u}_i : S^{r_{i+1}-r_i} \wedge \tilde{X}_i \rightarrow \tilde{X}_{i+1}$ in $\mathcal{S}\mathcal{S}_*$, and then choosing a representative sequence of maps $\tilde{\omega}_{i+1} : \tilde{X}_{i+1} \wedge S^{d_{i+1}} \rightarrow \tilde{X}_{i+1}$ in $\mathcal{S}\mathcal{S}_*$ making the above-displayed square commute for each i . For $i \geq 0$ and $m \geq d_i$ with $d_i | m$, we let $B(\tilde{X}_i, \tilde{\omega}_i, m)$ be the obvious L_n^f -cospectrum with $B(\tilde{X}_i, \tilde{\omega}_i, m)^k = \tilde{X}_i \wedge S^r$ for $k \equiv -r \pmod m$ and $0 \leq r < m$. We inductively construct L_n^f -cospectra $B(\tilde{X}_i)$ for $i \geq 0$, together with maps $B(\tilde{X}_i, \tilde{\omega}_i, d_i) \rightarrow B(\tilde{X}_i)$ and $S^{r_{i+1}-r_i} \wedge$

$B(\tilde{X}_i) \rightarrow B(\tilde{X}_{i+1})$, by taking the pushouts

$$\begin{array}{ccc} S^{r_{i+1}-r_i} \wedge B(\tilde{X}_i, \tilde{\omega}_i, d_{i+1}) & \longrightarrow & B(\tilde{X}_{i+1}, \tilde{\omega}_{i+1}, d_{i+1}) \\ \downarrow & & \downarrow \\ S^{r_{i+1}-r_i} \wedge B(\tilde{X}_i) & \longrightarrow & B(\tilde{X}_{i+1}). \end{array}$$

We note that $B(\tilde{X}_i)^0 = \tilde{X}_i$ for each $i \geq 0$. Finally, we let B be the associated spectral L_n^f -cospectrum with $B_{r_i+j}^* = S^j \wedge B(\tilde{X}_i)$ for $0 \leq j < r_{i+1} - r_i$. \square

12.2. The functors Φ_n and Θ_n . Holding $n \geq 1$ fixed, we choose an admissible spectral L_n^f -cospectrum B , together with an equivalence $B_*^0 \simeq C_{n-1}^f S_{(p)}$ and obtain the adjoint functors $\Theta_B : Ho^s \rightarrow \mathcal{UL}_n^f$ and $\Phi_B : \mathcal{UL}_n^f \rightarrow Ho^s$. Using the reflection $L_n^f : Ho_* \rightarrow \mathcal{UL}_n^f \subset Ho_*$, we extend these functors to give $\Theta_n = \Theta_B : Ho^s \rightarrow Ho_*$ and $\Phi_n = \Phi_B L_n^f : Ho_* \rightarrow Ho^s$. We now have

Theorem 12.3. *For $n \geq 1$, the functors $\Phi_n : Ho_* \rightarrow Ho^s$ and $\Theta_n : Ho^s \rightarrow Ho_*$ satisfy the conditions of Theorem 5.3(iv), (v) and Theorem 5.4(ii), (iv), (v).*

Proof. Condition 5.3(v) follows from Lemma 12.6 below, while the other conditions follow from Theorem 11.7. \square

Condition 5.3(v) requires additional work to show that $\Phi_n : Ho_* \rightarrow Ho^s$ preserves homotopy fiber squares in Ho_* , not merely \mathcal{UL}_n^f .

Lemma 12.4. *If $f : X \rightarrow Y$ is a map in \mathcal{UL}_n^f with $f_* : \pi_i X \cong \pi_i Y$ for sufficiently large i , then $\Phi_B f : \Phi_B X \simeq \Phi_B Y$.*

Proof. It suffices to show that $\text{map}_*(B_m^k, f)$ is an equivalence for $k, m \geq 0$. This follows since $\text{map}_*(B_m^{k+j} \wedge S^j, f)$ is an equivalence for sufficiently large j and since there is an L_n^f -equivalence $B_m^{k+j} \wedge S^j \rightarrow B_m^k$ for all j .

Lemma 12.5. *For a homotopy fiber sequence $F \rightarrow X \rightarrow Y$ of pointed spaces and $n \geq 0$, the map $L_n^f F \rightarrow \text{fib}(L_n^f X \rightarrow L_n^f Y)$ has homotopy fiber with vanishing homotopy groups in dimensions $\geq d_n$.*

Proof. Since $L_n^f X \simeq P_{(p)} P_{A(n+1)} X$ by Lemma 4.10, this follows from [Bou7, 8.1]. \square

Lemma 12.6. *The functor $\Phi_n : Ho_* \rightarrow Ho^s$ preserves homotopy fiber squares.*

Proof. For a homotopy pullback $X \times_Y Z$, the map $L_n^f(X \times_Y Z) \rightarrow L_n^f X \times_{L_n^f Y} L_n^f Z$ is a π_i -equivalence for $i > d_n + 1$ by Lemma 12.5, and is consequently a Φ_B -equivalence by Lemma 12.4. The result now follows since Φ_B preserves homotopy fiber squares in \mathcal{UL}_n^f by Theorem 11.7. \square

12.7. The simplicial functoriality of Φ_n and Θ_n . The functors $\Phi_n : Ho_* \rightarrow Ho^s$ and $\Theta_n : Ho^s \rightarrow Ho_*$ for $n \geq 1$ are induced by simplicial functors

$$\begin{aligned} \Phi_n &= \Phi_B L_n^f : \mathcal{SS}_* \longrightarrow Sp \\ \Theta_n &= L_n^f \Theta_B : Sp \longrightarrow \mathcal{SS}_* \end{aligned}$$

which are constructed using the simplicial functors Φ_B , Θ_B , and L_n^f of 11.8 and 4.4. By 11.8 and the proof of Lemma 12.6, $\Phi_n : \mathcal{SS}_* \rightarrow Sp$ preserves homotopy fiber squares, while $\Theta_n : Sp \rightarrow \mathcal{SS}_*$ carries homotopy cofiber squares to L_n^f -homotopy cofiber squares.

13. PROOFS OF THEOREMS 5.3 AND 5.4

We now proceed to prove Theorems 5.3 and 5.4 by establishing the remaining properties of Φ_n and Θ_n not covered by Theorem 12.3. As above, we hold $n \geq 1$ fixed and let B be an admissible spectral L_n^f -cospectrum with a given equivalence $B_*^0 \simeq C_{n-1}^f S_{(p)}$ in Ho^s .

Theorem 13.1. *For $n \geq 1$ and $E \in Ho^s$, there is a natural equivalence $\Phi_n \Omega^\infty E \simeq \widehat{M}_n^f E$ in Ho^s .*

Proof. Since the map $L_n^f \Omega^\infty E \rightarrow \Omega^\infty L_n^f E$ is a π_i -equivalence for $i > d_n$ by Proposition 5.2, there are natural equivalences

$$\begin{aligned} \Phi_n \Omega^\infty E &\simeq \Phi_B L_n^f \Omega^\infty E \simeq \Phi_B \Omega^\infty L_n^f E \\ &\simeq F(B_*^0, L_n^f E) \simeq F(C_{n-1}^f S_{(p)}, L_n^f E) \simeq \widehat{M}_n^f E \end{aligned}$$

by Lemma 12.4, Theorem 11.9, and Theorem 3.3. □

Theorem 13.2. *For $n \geq 1$, let A be an L_n^f -cospectrum with $\Sigma^\infty A^i \in \mathcal{F}_n$ for $i \geq 0$. Then for each space $X \in \mathcal{UL}_n^f$, the spectrum $\Phi_A X$ belongs to $\mathcal{M}_n^f \cap \widehat{\mathcal{M}}_n^f$.*

Proof. $\Phi_A X$ belongs to \mathcal{L}_n^f since it is an Ω -spectrum whose spaces X^{A^i} belong to \mathcal{UL}_n^f for $i \geq 0$. Moreover,

$$v_k^{-1} \pi_*(\Phi_A X; F(k)) \cong v_k^{-1} \pi_*(X^{A^0}; F(k)) \cong v_k^{-1} \pi_*(X; A^0 \wedge F(k)) = 0$$

for $1 \leq k < n$ since $A^0 \wedge F(k) \in \mathcal{F}_n$, and $\pi_* \Phi_A X$ is p -torsion since $[\Sigma^j A^i, \Sigma^j A^i]$ is a finite p -torsion group for $i \geq 0$ and $j \geq 2$. Hence, $L_{n-1}^f \Phi_A X = 0$ by 3.8, and $\Phi_A X$ belongs to \mathcal{M}_n^f . For $1 \leq k < n$ choose a map $\omega_k : \Sigma^d W_k \rightarrow W_k$ of pointed spaces such that $\omega_k \wedge A^0$ is trivial and $\Sigma^\infty \omega_k$ is a v_k -map of finite p -local spectra of type k , and also let $\omega_0 : \Sigma^d W_0 \rightarrow W_0$ denote a map $S_{(p)}^m \rightarrow S_{(p)}^m$ of degree p^j for some $m, j > 0$ such that $\omega_0 \wedge A^0$ is trivial. Let $\omega_k^\# : (\Phi_A X)^{W_k} \rightarrow (\Phi_A X)^{\Sigma^d W_k}$ be the induced map in \mathcal{M}_n^f for $0 \leq k < n$. Then $\widehat{M}_n^f \omega_k^\#$ is trivial since $\widehat{M}_n^f \omega_k^\# = \Phi_n \Omega^\infty \omega_k^\#$ by Theorem 13.1 where $\Omega^\infty \omega_k^\#$ is trivial. Thus $\omega_k^\#$ is trivial by Theorem 3.3, and $\Phi_A X$ is $F(n)_*$ -local by [Bou8, 6.6]. Hence $\Phi_A X$ belongs to $\widehat{\mathcal{M}}_n^f$ since $\widehat{M}_n^f = L_{F(n)} L_n^f$ by Theorem 3.3. □

Theorem 13.3. *For $n \geq 1$ and $X \in Ho_*$, the spectrum $\Phi_n X$ belongs to $\widehat{\mathcal{M}}_n^f$.*

Proof. This follows since $\Phi_n X$ is a homotopy inverse limit of spectra $S^m \wedge \Phi_{B_*^m} L_n^f X$ which are in $\widehat{\mathcal{M}}_n^f$ by Theorem 13.2. □

Theorem 13.4. *For $n \geq 1$ and $W \in \mathcal{F}_n$, there are isomorphisms*

$$v_n^{-1} \pi_*(X; W) \cong [W, \Phi_n X]_* \cong v_n^{-1} \pi_*(\Phi_n X; W)$$

which are natural in $X \in Ho_*$.

Proof. By desuspending a v_n -map for W , we may obtain an L_n^f -cospectrum A such that $\Sigma^\infty A^i \in \mathcal{F}_n$ for $i \geq 0$, $\Sigma^\infty A^0 \simeq \Sigma^j W$ for some j , and $\pi_* \Phi_A X \cong v_n^{-1} \pi_*(X; A^0)$ naturally for $X \in \mathcal{UL}_n^f$. Then for $X \in Ho_*$ there are natural equivalences

$$\Phi_A L_n^f X \simeq \widehat{M}_n^f \Phi_A L_n^f X \simeq \Phi_n \Omega^\infty \Phi_A L_n^f X \simeq \Phi_n (L_n^f X)^{A^0} \simeq (\Phi_n X)^{A^0}$$

by Theorems 13.1, 13.2, and 12.3. Hence for $X \in Ho_*$ there are natural isomorphisms $\pi_* \Phi_A L_n^f X \cong \pi_*(\Phi_n X)^{A^0}$ giving isomorphisms

$$v_n^{-1} \pi_*(L_n^f X; A^0) \cong [A^0, \Phi_n X]_*,$$

and the result follows by 4.5 and 4.6. \square

Corollary 13.5. *For $n \geq 1$, if $f : X \rightarrow Y$ is a $v_n^{-1} \pi_*$ -equivalence in Ho_* , then $\Phi_n f : \Phi_n X \simeq \Phi_n Y$.*

Proof. Since $\Phi_n f$ is a $v_n^{-1} \pi_*$ -equivalence in $\widehat{\mathcal{M}}_n^f$ by 13.3 and 13.4, it is an equivalence by 3.8. \square

Theorem 13.6. *For $n \geq 1$ and $E \in Ho^s$, $\Theta_n E$ belongs to \mathcal{UN}_n^f .*

Proof. The maps $L_{n-1}^f \Theta_n E \rightarrow *$ and $(\Theta_n E)\langle d_n \rangle \rightarrow \Theta_n E$ are Φ_n -equivalences by Corollary 13.5, and thus are $[\Theta_n E, -]$ -equivalences since $\Theta_n : Ho^s \rightarrow \mathcal{UL}_n^f$ is left adjoint to $\Phi_n : \mathcal{UL}_n^f \rightarrow Ho^s$ by Theorem 12.3. Hence $\Theta_n E$ belongs to \mathcal{UM}_n^f since the localization map $\Theta_n E \rightarrow L_{n-1}^f \Theta_n E$ is trivial, and $\Theta_n E$ belongs to \mathcal{UN}_n^f since its identity map factors through $(\Theta_n E)\langle d_n \rangle$. \square

Theorem 13.7. *For $n \geq 1$ and $E \in Ho^s$, there is a natural equivalence $L_n^f \Sigma^\infty \Theta_n E \simeq M_n^f E$.*

Proof. Since the functors

$$Ho^s \xrightarrow{\Theta_n} \mathcal{UL}_n^f \xrightarrow{L_n^f \Sigma^\infty} \mathcal{L}_n^f$$

are left adjoint to

$$Ho^s \xleftarrow{\Phi_n} \mathcal{UL}_n^f \xleftarrow{\Omega^\infty} \mathcal{L}_n^f,$$

and since the functors $M_n^f : Ho^s \rightleftarrows \mathcal{L}_n^f : \widehat{M}_n^f$ are also adjoint, the natural equivalence $\Phi_n \Omega^\infty E \simeq \widehat{M}_n^f E$ of Theorem 13.1 implies a natural equivalence $L_n^f \Sigma^\infty \Theta_n E \simeq M_n^f E$. \square

13.8. Proof of Theorem 5.3. Parts (i), (ii), and (iii) follow by Theorems 13.3, 13.4, and 13.1, while parts (iv) and (v) follow by Theorem 12.3. \square

13.9. Proof of Theorem 5.4. Parts (i) and (iii) follow by Theorems 13.6 and 13.7, while parts (ii), (iv), and (v) follow by Theorem 12.3. \square

REFERENCES

- [Ada] J. F. Adams, *Stable Homotopy and Generalized Homology*, University of Chicago Press, 1974. MR **53**:6534
- [Bor] F. Borceux, *Handbook of Categorical Algebra 2, Categories and Structures*, Encyclopedia of Mathematics and its Applications, vol. 51, Cambridge University Press, 1994. MR **96g**:18001b
- [Bou1] A.K. Bousfield, *Boolean algebra of spectra*, Comment. Math. Helv. **54** (1979), 368-377; Correction, Comment Math. Helv. **58** (1983), 599-600. MR **81a**:55015; MR **85h**:55013
- [Bou2] ———, *The localization of spectra with respect to homology*, Topology **18** (1979), 257-281. MR **80m**:55006
- [Bou3] ———, *Cohomological localizations of spaces and spectra*, preprint, 1979.
- [Bou4] ———, *K-localizations and K-equivalences of infinite loop spaces*, Proc. London Math. Soc. **44** (1982), 291-311. MR **83g**:55008
- [Bou5] ———, *On homology equivalences and homological localizations of spaces*, Amer. J. Math. **104** (1982), 1025-1042. MR **84g**:55014

- [Bou6] ———, *Uniqueness of infinite deloopings for K -theoretic spaces*, Pacific J. Math. **129** (1987), 1-31. MR **89g**:55017
- [Bou7] ———, *Localization and periodicity in unstable homotopy theory*, J. Amer. Math. Soc. **7** (1994), 831-873. MR **95c**:55010
- [Bou8] ———, *Unstable localization and periodicity*, Algebraic Topology: New Trends in Localization and Periodicity, Progress in Mathematics, vol. 136, Birkhauser-Verlag, 1996, pp. 33-50. MR **98c**:55014
- [Bou9] ———, *Homotopical localizations of spaces*, Amer. J. Math. **119** (1997), 1321-1354. MR **98m**:55009
- [Bou10] ———, *On $K(n)$ -equivalences of spaces*, Homotopy invariant algebraic structures: a conference in honor of J. Michael Boardman, Contemp. Math., vol. 239. CMP 2000:03
- [BF] A.K. Bousfield and E.M. Friedlander, *Homotopy theory of Γ -spaces, spectra, and bisimplicial sets*, Lecture Notes in Math., vol. 658, Springer-Verlag, 1978, pp. 80-130. MR **80e**:55021
- [BK] A.K. Bousfield and D.M. Kan, *Homotopy Limits, Completions and Localizations*, Lecture Notes in Math., vol. 304, Springer-Verlag, 1972. MR **51**:1825
- [Dro] E. Dror Farjoun, *Cellular spaces, null spaces and homotopy localizations*, Lecture Notes in Math., vol. 1622, Springer-Verlag, 1996. MR **98f**:55010
- [DDK] E. Dror Farjoun, W.G. Dwyer, and D.M. Kan, *An arithmetic square for virtually nilpotent spaces*, Illinois J. Math. **21** (1977), 242-254. MR **55**:11246
- [DS] W.G. Dwyer and J. Spalinski, *Homotopy theories and model categories*, Handbook of Algebraic Topology, North-Holland, 1995, pp. 73-126. MR **96h**:55014
- [GJ] P.G. Goerss and J.F. Jardine, *Simplicial Homotopy Theory*, Progress in Mathematics, vol. 174, Birkhauser-Verlag, 1999. CMP 2000:02
- [Hir] P.S. Hirschhorn, *Localization in Model Categories*, in preparation.
- [Hop] M.J. Hopkins, *Global methods in homotopy theory*, Homotopy Theory – Proceedings of the Durham Symposium 1985, London Math. Soc. Lecture Note Series, vol. 117, Cambridge Univ. Press, 1987, pp. 73-96. MR **89g**:55022
- [HRW] M.J. Hopkins, D.C. Ravenel, and W.S. Wilson, *Morava Hopf algebras and spaces $K(n)$ equivalent to finite Postnikov systems*, Stable and Unstable Homotopy, Fields Institute Communications, vol. 19, 1998, pp. 137-163. MR **99e**:55005
- [HSm] M.J. Hopkins and J.H. Smith, *Nilpotence and stable homotopy theory II*, Ann. Math. **148** (1998), 1-49. MR **99h**:55009
- [Hov] M. Hovey, *Model Categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, 1998. MR **99h**:55031
- [HP] M. Hovey and J.H. Palmieri, *The structure of the Bousfield lattice*, Homotopy invariant algebraic structures: a conference in honor of J. Michael Boardman, Contemp. Math., vol. 239. CMP 2000:03
- [HPS] M. Hovey, J.H. Palmieri, and N.P. Strickland, *Axiomatic Stable Homotopy Theory*, Mem. Amer. Math. Soc. (1997). MR **98a**:55017
- [HSS] M. Hovey, B. Shipley, and J.H. Smith, *Symmetric spectra*, J. Amer. Math. Soc. **13** (2000), 149-208. CMP 2000:02
- [HSt] M. Hovey and N.P. Strickland, *Morava K -theories and localization*, Mem. Amer. Math. Soc. **139** (1999). MR **99b**:55017
- [Kas] T. Kashiwabara, *On Brown-Peterson cohomology of QX* (to appear).
- [Kuh1] N.J. Kuhn, *Suspension spectra and homology equivalences*, Trans. Amer. Math. Soc. **283** (1984), 303-313. MR **85g**:55014
- [Kuh2] ———, *Morava K -theories and infinite loop spaces*, Lecture Notes in Math., vol. 1370, Springer-Verlag, 1989, pp. 243-257. MR **90d**:55014
- [Mah] M.E. Mahowald, *The image of J in the EHP sequences*, Ann. Math. **116** (1982), 65-112. MR **86d**:55018
- [MS] M. Mahowald and H. Sadofsky, *v_n -telescopes and the Adams spectral sequence*, Duke Math. J. **78** (1995), 101-129. MR **96h**:55006
- [Mil1] H.R. Miller, *On relations between Adams spectral sequences with an application to stable homotopy theory*, J. Pure Appl. Algebra **20** (1981), 287-312. MR **82f**:55029
- [Mil2] ———, *Finite localizations*, Bol. Soc. Mat. Mexicana (Homenaje a Jose Adem) **37** (1992), 383-390. MR **96h**:55009

- [Qui] D.G. Quillen, *Homotopical Algebra*, Lecture Notes in Math., vol. 43, Springer-Verlag, 1967. MR **36**:6480
- [Rav1] D.C. Ravenel, *Localization with respect to certain periodic homology theories*, Amer. J. Math. **106** (1984), 351-414. MR **85k**:55009
- [Rav2] ———, *Progress report on the telescope conjecture*, London Math. Soc. Lecture Note Series 176, Adams Memorial Symposium on Algebraic Topology, Cambridge Univ. Press, 1992, pp. 1-21. MR **94h**:55023
- [Rav3] ———, *Nilpotence and Periodicity in Stable Homotopy Theory*, Ann. of Math. Studies 128, Princeton Univ. Press, 1992. MR **94b**:55015
- [Rav4] ———, *Life after the telescope conjecture*, Algebraic K-Theory and Algebraic Topology (P.G. Goerss and J.F. Jardine, eds.), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci. 407, Kluwer, 1993, pp. 205-222. MR **96i**:55016
- [RW] D.C. Ravenel and W.S. Wilson, *The Morava K-theories of Eilenberg-Mac Lane spaces and the Conner-Floyd conjecture*, Amer. J. Math. **102** (1980), 691-748. MR **81i**:55005
- [War] R.B. Warfield, Jr., *Nilpotent Groups*, Lecture Notes in Math., vol. 513, Springer-Verlag, 1976. MR **53**:13413

DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE (M/C 249), UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, ILLINOIS 60607

E-mail address: bous@uic.edu