

POLYNOMIALS NONNEGATIVE ON A GRID AND DISCRETE OPTIMIZATION

JEAN B. LASSERRE

ABSTRACT. We characterize the real-valued polynomials on \mathbb{R}^n that are nonnegative (not necessarily strictly positive) on a grid \mathbb{K} of points of \mathbb{R}^n , in terms of a weighted sum of squares whose degree is bounded and known in advance. We also show that the minimization of an arbitrary polynomial on \mathbb{K} (a discrete optimization problem) reduces to a convex continuous optimization problem of fixed size. The case of concave polynomials is also investigated. The proof is based on a recent result of Curto and Fialkow on the \mathbb{K} -moment problem.

1. INTRODUCTION

This paper is concerned with the characterization of real-valued polynomials on \mathbb{R}^n that are nonnegative (and not necessarily strictly positive) on a pre-defined grid of points of \mathbb{R}^n . That is, we consider the polynomials $p(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $p(x) \geq 0$ on \mathbb{K} , where \mathbb{K} is the subset of \mathbb{R}^n defined by

$$(1.1) \quad \mathbb{K} := \{x \in \mathbb{R}^n \mid g_k(x) = 0, k = 1, \dots, n\},$$

where the polynomials $g_k(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ are given by

$$x \mapsto g_k(x) := \prod_{i=1}^{2r_k} (x_k - a_{ki}), \quad k = 1, \dots, n.$$

\mathbb{K} defines a grid in \mathbb{R}^n with $s := \prod_{k=1}^n 2r_k$ points.

We obtain a result in the spirit of the “linear” representation of polynomials positive on a compact semi-algebraic set, obtained by Putinar [10], Jacobi and Prestel [4, 5] in a general framework. Namely, we show that every polynomial $p(x)$ of degree $2r_0$ or $2r_0 - 1$, nonnegative on \mathbb{K} , can be written as a sum of squares of polynomials weighted by the polynomials $g_k(x)$ defining the set \mathbb{K} , and whose degree is bounded by $r + v$ with $r := \sum_{k=1}^n (2r_k - 1)$ and $v := \max\{r_0 - r, \max_{k=1}^n r_k\}$, independently of the grid points. The important thing is that in this case, the degree of the polynomials in that representation is *bounded* and *known* in advance.

To prove this result, we use a detour and first consider the associated discrete optimization problem

$$\mathbb{P} \rightarrow p^* := \min_{x \in \mathbb{K}} p(x),$$

Received by the editors November 11, 2000.

2000 *Mathematics Subject Classification.* Primary 13P99, 30C10; Secondary 90C10, 90C22.

Key words and phrases. Algebraic geometry; positive polynomials; \mathbb{K} -moment problem; semi-definite programming.

which is known to be NP-hard in general. However, we show that \mathbb{P} is equivalent to a continuous convex optimization problem whose size depends on the number of points, but not on the points in the grid. The idea is to define a sequence of refined so-called convex *semidefinite (SDP) relaxations* $\{\mathbb{Q}_i\}$ of \mathbb{P} . For the general optimization problem where one minimizes a polynomial $p(x)$ on a compact semi-algebraic set \mathbb{K} , we have shown in Lasserre [7] that the SDP relaxations \mathbb{Q}_i “converge”, that is, $\inf \mathbb{Q}_i \uparrow p^*$ as $i \rightarrow \infty$, provided the set \mathbb{K} has some property that permits a “linear” representation of polynomials positive on \mathbb{K} (a particular case of the more general Schmüdgen’s representation [11]), as proved in Putinar [10], Jacobi and Prestel [4, 5]. This approach is also valid for 0-1 optimization problems (see Lasserre [8]). Other sequences of SDP relaxations have been provided for 0-1 optimization problems, notably the *lift and project* procedure of Lovász and Schrijver [9] (see also extensions by Kojima and Tunçel [6]) that eventually yields the convex hull of the feasible set of \mathbb{P} whenever a weak separation oracle is available for the homogeneous cone associated with the constraint set. The approach developed in Lasserre [7, 8]) is different and directly relates results of algebraic geometry with optimization. In particular, the relaxations $\{\mathbb{Q}_i\}$ and their dual $\{\mathbb{Q}_i^*\}$ perfectly match the duality between the \mathbb{K} -moment problem and the theory of polynomials, positive on a compact semi-algebraic set \mathbb{K} .

However, in the present context, we may even further extend the properties of the above relaxations. We use a recent result of Curto and Fialkow [2] on the \mathbb{K} -moment problem, to show that the optimal value p^* is obtained at most at the relaxation \mathbb{Q}_{r+v} . Next, by using a standard “strong duality” result in convex optimization, we show that the dual optimization problem \mathbb{Q}_{r+v}^* is solvable and yields the coefficients of the polynomials in the representation of $p(x) - p^*$ as a sum of squares, weighted by the polynomials $g_k(x)$ defining \mathbb{K} . In a sense, the order of the relaxation \mathbb{Q}_i for which the optimum value p^* is obtained, measures the “hardness” of problem \mathbb{P} , as it expresses the “effort” (in degree) needed to represent $p(x) - p^*$ as a weighted sum of squares. In the present case, this “effort” is finite and bounded by $r + v$. It also follows that any discrete polynomial optimization problem on \mathbb{K} , with additional arbitrary polynomial constraints $h_j(x) \geq 0$, also reduces to a convex continuous optimization problem of fixed size. Finally, we also obtain a representation of concave polynomials.

The paper is organized as follows. We first introduce the notation as well as some definitions. In Section 3, we present the results on the discrete optimization problem \mathbb{P} . In Section 4, we then present the result on the representation of $p(x) - p^*$ as a weighted sum of squares.

2. NOTATION AND DEFINITIONS

We adopt the following notation. Given any two real-valued symmetric matrices A, B let $\langle A, B \rangle$ denote the usual scalar product $\text{trace}(AB)$ and let $A \succeq B$ (resp. $A \succ B$) stand for $A - B$ positive semidefinite (resp. $A - B$ positive definite). Let

$$(2.1) \quad 1, x_1, x_2, \dots, x_n, x_1^2, x_1x_2, \dots, x_1x_n, x_2^2, x_2x_3, \dots, x_n^2, \dots, x_1^r, \dots, x_n^r$$

be a basis for the space \mathcal{A}_r of real-valued polynomials of degree at most r , and let $s(r)$ be its dimension. Therefore, an r -degree polynomial $p(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ is written

$$p(x) = \sum_{\alpha} p_{\alpha} x^{\alpha}, \quad x \in \mathbb{R}^n,$$

where

$$x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \dots x_n^{\alpha_n}, \quad \text{with } \sum_{i=1}^n \alpha_i = k,$$

is a monomial of degree k with coefficient p_α . Denote by $p = \{p_\alpha\} \in \mathbb{R}^{s(r)}$ the coefficients of the polynomial $p(x)$ in the basis (2.1). Hence, the respective vectors of coefficients of the polynomials $g_k(x)$, $k = 1, \dots, n$, in (1.1), are denoted $\{(g_k)_\alpha\} = g_k \in \mathbb{R}^{s(2r_k)}$, $k = 1, \dots, n$.

We next define the important notions of moment matrix and localizing matrix already introduced in Curto and Fialkow [2], Berg [1].

2.1. Moment matrix. Given an $s(2r)$ -sequence $(1, y_1, \dots)$, let $M_r(y)$ be the **moment** matrix of dimension $s(r)$ (denoted $M(r)$ in Curto and Fialkow [2]), with rows and columns labelled by (2.1). For instance, for illustration purposes, and for clarity of exposition, consider the 2-dimensional case. The moment matrix $M_r(y)$ is the block matrix $\{M_{i,j}(y)\}_{0 \leq i,j \leq r}$ defined by

$$(2.2) \quad M_{i,j}(y) = \begin{bmatrix} y_{i+j,0} & y_{i+j-1,1} & \dots & y_{i,j} \\ y_{i+j-1,1} & y_{i+j-2,2} & \dots & y_{i-1,j+1} \\ \dots & \dots & \dots & \dots \\ y_{j,i} & y_{i+j-1,1} & \dots & y_{0,i+j} \end{bmatrix}.$$

To fix ideas, with $n = 2$ and $r = 2$, one obtains

$$M_2(y) = \begin{bmatrix} 1 & | & y_{10} & y_{01} & | & y_{20} & y_{11} & y_{0,2} \\ \hline y_{10} & | & y_{20} & y_{11} & | & y_{30} & y_{21} & y_{12} \\ y_{01} & | & y_{11} & y_{02} & | & y_{21} & y_{12} & y_{03} \\ \hline y_{20} & | & y_{30} & y_{21} & | & y_{40} & y_{31} & y_{22} \\ y_{11} & | & y_{21} & y_{12} & | & y_{31} & y_{22} & y_{13} \\ y_{02} & | & y_{12} & y_{03} & | & y_{22} & y_{13} & y_{04} \end{bmatrix}.$$

Another more intuitive way of constructing $M_r(y)$ is as follows. If $M_r(y)(1, i) = y_\alpha$ and $M_r(y)(j, 1) = y_\beta$, then

$$(2.3) \quad M_r(y)(i, j) = y_{\alpha+\beta}, \quad \text{with } \alpha + \beta = (\alpha_1 + \beta_1, \dots, \alpha_n + \beta_n).$$

$M_r(y)$ defines a bilinear form $\langle \cdot, \cdot \rangle_y$ on \mathcal{A}_r by

$$\langle q(x), v(x) \rangle_y := \langle q, M_r(y)v \rangle, \quad q(x), v(x) \in \mathcal{A}_r,$$

and if y is a sequence of moments of some measure μ_y , then

$$(2.4) \quad \langle q, M_r(y)q \rangle = \int q(x)^2 \mu_y(dx) \geq 0,$$

so that $M_r(y) \succeq 0$.

2.2. Localizing matrix. If the entry (i, j) of the matrix $M_r(y)$ is y_β , let $\beta(i, j)$ denote the subscript β of y_β . Next, given a polynomial $\theta(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ with coefficient vector θ , we define the matrix $M_r(\theta y)$ by

$$(2.5) \quad M_r(\theta y)(i, j) = \sum_{\alpha} \theta_\alpha y_{\{\beta(i,j)+\alpha\}}.$$

For instance, with

$$M_1(y) = \begin{bmatrix} 1 & y_{10} & y_{01} \\ y_{10} & y_{20} & y_{11} \\ y_{01} & y_{11} & y_{02} \end{bmatrix} \quad \text{and } x \mapsto \theta(x) = a - x_1^2 - x_2^2,$$

we obtain

$$M_1(\theta y) = \begin{bmatrix} a - y_{20} - y_{02}, & ay_{10} - y_{30} - y_{12}, & ay_{01} - y_{21} - y_{03} \\ ay_{10} - y_{30} - y_{12}, & ay_{20} - y_{40} - y_{22}, & ay_{11} - y_{31} - y_{13} \\ ay_{01} - y_{21} - y_{03}, & ay_{11} - y_{31} - y_{13}, & ay_{02} - y_{22} - y_{04} \end{bmatrix}.$$

In a manner similar to what we have in (2.4), if y is a sequence of moments of some measure μ_y , then

$$(2.6) \quad \langle q, M_r(\theta y)q \rangle = \int \theta(x)q(x)^2 \mu_y(dx),$$

for every polynomial $q(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ with coefficient vector $q \in \mathbb{R}^{s(r)}$. Therefore, $M_r(\theta y) \succeq 0$ whenever μ_y has its support contained in the set $\{\theta(x) \geq 0\}$. In Curto and Fialkow [2], $M_r(\theta y)$ is called a *localizing* matrix (denoted by $M_\theta(r + v)$ if $\deg \theta = 2v$ or $2v - 1$).

The \mathbb{K} -moment problem identifies those sequences y that are moment-sequences of a measure with support contained in the semi-algebraic set \mathbb{K} . In duality with the theory of moments is the theory of representation of positive polynomials, which dates back to Hilbert's 17th problem. This fact will be reflected in the semidefinite relaxations proposed later. For details and recent results, the interested reader is referred to Curto and Fialkow [2], Jacobi [3], Jacobi and Prestel [4, 5], Simon [12], Schmüdgen [11] and the many references therein.

3. THE ASSOCIATED DISCRETE OPTIMIZATION PROBLEM \mathbb{P}

Consider the discrete optimization problem \mathbb{P}

$$(3.1) \quad \mathbb{P} \rightarrow p^* := \min_{x \in \mathbb{R}^n} \{p(x) \mid g_k(x) = 0, k = 1, \dots, n\},$$

where the polynomials $g_k(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ are defined by

$$(3.2) \quad g_k(x) := \prod_{i=1}^{2r_k} (x - a_{ki}), \quad k = 1, \dots, n,$$

with $r_k \in \mathbb{N}$, $k = 1, \dots, n$, and the $\{a_{ki}\}$, $i = 1, \dots, r_k$, are given real numbers such that for every $k = 1, \dots, n$, $a_{ki} \neq a_{kj}$ whenever $i \neq j$. For homogeneity in notation, we have chosen to assume that all the polynomials $g_k(x)$ have an even degree. The results presented below are also valid when the polynomials have arbitrary degree.

Let

$$(3.3) \quad \mathbb{K} := \{x \in \mathbb{R}^n \mid g_k(x) = 0, k = 1, \dots, n\}$$

be the feasible set associated with \mathbb{P} .

As we minimize $p(x)$ on \mathbb{K} , we could assume that the degree $2r_0$ (or $2r_0 - 1$) of $p(x)$ is not larger than $r := \sum_{k=1}^n (2r_k - 1)$, since otherwise, using the equations $g_k(x) = 0$, we may replace $p(x)$ with another polynomial $\tilde{p}(x)$ of degree not larger than r , and identical to $p(x)$ on \mathbb{K} . However, as for the representation of $p(x)$ in §4, we will consider an arbitrary r_0 , we do not make this assumption.

When needed below, for $i \geq \max_k r_k$, the vectors $g_k \in \mathbb{R}^{s(2r_k)}$ are extended to vectors of $\mathbb{R}^{s(2i)}$ by completing with zeros. As we minimize $p(x)$ we may and will assume that its constant term is zero, that is, $p(0) = 0$.

3.1. SDP relaxations of \mathbb{P} . For $i \geq \max_{k \in \{0, n\}} r_k$, consider the following family $\{\mathbb{Q}_i\}$ of convex positive semidefinite (psd) programs (or semidefinite programming (SDP) relaxations of \mathbb{P})

$$(3.4) \quad \mathbb{Q}_i \begin{cases} \min_y \sum_{\alpha} p_{\alpha} y_{\alpha} \\ M_i(y) \succeq 0, \\ M_{i-r_k}(g_k y) = 0, \quad k = 1, \dots, n, \end{cases}$$

with respective dual problems

$$(3.5) \quad \mathbb{Q}_i^* \begin{cases} \max_{X \succeq 0, Z_k} -X(1, 1) - \sum_{k=1}^n g_k(0) Z_k(1, 1) \\ \langle X, B_{\alpha} \rangle + \sum_{k=1}^n \langle Z_k, C_{\alpha}^k \rangle = p_{\alpha}, \quad \forall \alpha \neq 0 \end{cases}$$

where X, Z_k are real-valued symmetric matrices, the “dual variables” associated with the constraints $M_i(y) \succeq 0$ and $M_{i-r_k}(g_k y) \succeq 0$ respectively, and where we have written

$$M_i(y) = \sum_{\alpha} B_{\alpha} y_{\alpha}; \quad M_{i-r_k}(g_k y) = \sum_{\alpha} C_{\alpha}^k y_{\alpha}, \quad k = 1, \dots, n,$$

for appropriate real-valued symmetric matrices $B_{\alpha}, C_{\alpha}^k, k = 1, \dots, n$.

In the standard terminology, the constraint $M_i(y) \succeq 0$ is called a “linear matrix inequality” (LMI) and \mathbb{Q}_i and its dual \mathbb{Q}_i^* are so-called positive semidefinite (psd) programs, the semidefinite (SDP) relaxations of \mathbb{P} . Both are convex optimization problems that can be solved efficiently via *interior points methods* and nowadays, several software packages (like e.g. the LMI toolbox of MATLAB) are available. The reader interested in more details on *semidefinite programming* is referred to Vandenberghe and Boyd [14] and the many references therein.

Note that the localizing matrices $M_{i-r_k}(g_k y)$ are easily obtained from the data $\{g_k\}$ of the problem \mathbb{P} by (2.5).

Interpretation of \mathbb{Q}_i . The linear matrix inequalities (LMI) constraints of \mathbb{Q}_i state (only) necessary conditions for y to be the vector of moments up to order $2i$, of some probability measure μ_y with support contained in \mathbb{K} . This clearly implies that $\inf \mathbb{Q}_i \leq p^*$, for all i , since the vector of moments of the Dirac measure at a feasible point of \mathbb{P} is feasible for \mathbb{Q}_i .

Interpretation of \mathbb{Q}_i^* . Let $X \succeq 0$ and $\{Z_k\}, k = 1, \dots, n$, be a feasible solution of \mathbb{Q}_i^* with value ρ . From the spectral decomposition of the symmetric matrices X and Z_k , write

$$X = \sum_j u_j u_j' \quad \text{and} \quad Z_k = \sum_j v_{kj} v_{kj}' - \sum_l w_{kl} w_{kl}', \quad k = 1, \dots, n,$$

where the vectors $\{u_j\}$ correspond to the positive eigenvalues of X , and the vectors $\{v_{kj}\}$ (resp. $\{w_{kl}\}$) correspond to the positive (resp. negative) eigenvalues of Z_k . Consider the polynomials $\{u_j(x)\}$ and $\{v_{kj}(x), w_{kl}(x)\}$ with respective coefficient vectors $\{u_j\}$ and $\{v_{kj}, w_{kl}\}$ in the basis (2.1).

Let $x \in \mathbb{R}^n$ be fixed, arbitrary. Then, from the feasibility of (X, Z_k) in \mathbb{Q}_i^* ,

$$(3.6) \quad \langle X, B_\alpha x^\alpha \rangle + \sum_{k=1}^n \langle Z_k, C_\alpha^k x^\alpha \rangle = p_\alpha x^\alpha, \quad \forall \alpha \neq 0,$$

and

$$(3.7) \quad X(1, 1) + \sum_{k=1}^n g_k(0)Z_k(1, 1) = -\rho.$$

Using the notation

$$y^x = (x_1, \dots, x_n, x_1^2, x_1x_2, \dots, x_1^i, \dots, x_n^i),$$

and summing up over all α , yields

$$(3.8) \quad \langle X, M_i(y^x) \rangle + \sum_{k=1}^n \langle Z_k, M_{i-r_k}(g_k y^x) \rangle = p(x) - \rho,$$

or, equivalently,

$$(3.9) \quad \sum_j \langle u_j, M_i(y^x)u_j \rangle + \sum_{k=1}^n \left[\sum_j \langle v_{kj}, M_{i-r_k}(g_k y^x)v_{kj} \rangle - \sum_l \langle w_{kl}, M_{i-v_k}(g_k y^x)w_{kl} \rangle \right] = p(x) - \rho.$$

Therefore, from the definition of the matrices $M_i(y), M_{i-r_k}(g_k y)$ (with $y = y^x$) and using (2.4)-(2.6),

$$(3.10) \quad p(x) - \rho = \sum_j u_j(x)^2 + \sum_{k=1}^n g_k(x) \left[\sum_j v_{kj}(x)^2 - \sum_l w_{kl}(x)^2 \right].$$

As $\rho \leq p^*$, $p(x) - \rho$ is nonnegative on \mathbb{K} (strictly positive if $\rho < p^*$), and one recognizes in (3.10) a “linear” representation into a weighted sum of squares of the polynomial $p(x) - p^*$, nonnegative on \mathbb{K} , as in the theory of representation of polynomials, strictly positive on a compact semi-algebraic set \mathbb{K} (see e.g., Schmüdgen [11], Putinar [10], Jacobi [3], Jacobi and Prestel [4, 5]). Indeed, when the set \mathbb{K} has a certain property, the “linear” (in the sense that no product term like $g_k(x)g_l(x)$ is needed) representation (3.10) holds. This property states that there is some polynomial $U(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $U(x)$ has the representation (3.10) and $\{U(x) \geq 0\}$ is compact (see Putinar [10], Jacobi and Prestel [4, 5]).

It will be shown below that indeed, the polynomial $p(x) - p^*$, which is only nonnegative (and not strictly positive) on \mathbb{K} , has the representation (3.10).

Hence, both psd programs \mathbb{Q}_i and \mathbb{Q}_i^* perfectly match the duality between the \mathbb{K} -moment problem and the theory of polynomials that are positive on \mathbb{K} .

3.2. Simplified SDP relaxations. The SDP relaxation \mathbb{Q}_i has a much simpler form that can be derived as follows. For convenience, write

$$(3.11) \quad g_k(x) := x_k^{2r_k} + \sum_{j=0}^{2r_k-1} \gamma_{kj} x_k^j, \quad k = 1, \dots, n.$$

Observe that in view of the construction of the localizing matrices in (2.5), and the form of the polynomials $g_k(x), k = 1, \dots, n$, the constraints $M_{i-r_k}(g_k y) = 0$

generate linear relationships between “moments” that can be translated into the moment matrix $M_i(y)$ as follows.

Given a monomial x^α , use (3.11) to replace $x_k^{2r_k}$ by

$$\sum_{j=0}^{2r_k-1} \gamma_{kj} x_k^j, \quad k = 1, \dots, n,$$

every time $\alpha_k \geq 2r_k$, until x^α is expressed as a linear combination $\sum_l h_l x^{\beta_l}$ of monomials x^{β_l} with $(\beta_l)_k < 2r_k$ for all $k = 1, \dots, n$ and all l , that is,

$$x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \dots x_n^{\alpha_n} = \sum_l h_l x^{\beta_l}.$$

Now, if $M_i(y)(s, t) = y_\alpha (= y_{\alpha_1, \alpha_2, \dots, \alpha_n})$ is the entry (s, t) of the matrix $M_i(y)$, then formally replace y_α in $M_i(y)(s, t)$ by $\sum_l h_l y_{\beta_l}$. Hence, the LMI constraints $M_i(y) \succeq 0$ and $M_{i-r_k}(g_k y) = 0$, $k = 1, \dots, n$, reduce to the single LMI constraint $\widehat{M}_i(y) \succeq 0$ obtained from $M_i(y) \succeq 0$ by making the above substitution in all the entries of $M_i(y)$. Observe that no matter how large i is, $\widehat{M}_i(y)$ contains only “moments” y_β with $\beta_k < 2r_k$ for all $k = 1, \dots, n$.

Therefore, $r := \sum_{k=1}^n (2r_k - 1)$ is the maximum degree of distinct monomials in the sense that whenever $i > r$ a polynomial of degree i can be expressed as a linear combination of monomials of degree less than r , after the simplification induced by the constraints $g_k(x) = 0$, $k = 1, \dots, n$.

Thus, with $s := \prod_{k=1}^n 2r_k$, there are no more than $s - 1$ variables y_β in all relaxations \mathbb{Q}_i (that is, s is the number of monomials in (2.1) of degree less than r), and the relaxation \mathbb{Q}_i has the simplified form

$$(3.12) \quad \begin{cases} \min_y \sum_\alpha \tilde{p}_\alpha y_\alpha \\ \widehat{M}_i(y) \succeq 0, \end{cases}$$

where we only have variables y_α with $\alpha_k < 2r_k$ for all $k = 1, \dots, n$, and $\{\tilde{p}_\alpha\}$ is the vector of coefficients of a polynomial $\tilde{p}(x)$ identical to $p(x)$ on \mathbb{K} .

For instance, in \mathbb{R}^2 ($n = 2$), let $g_k(x) = x_k^2 - x_k$ ($r_k = 1$) so that the set \mathbb{K} defines the 4 grid points $(0, 0), (0, 1), (1, 0), (1, 1)$ in \mathbb{R}^2 . Then,

$$(3.13) \quad M_2(y) = \begin{bmatrix} 1 & | & y_{10} & y_{01} & | & y_{20} & y_{11} & y_{02} \\ \hline y_{10} & | & y_{10} & y_{11} & | & y_{30} & y_{21} & y_{12} \\ y_{01} & | & y_{11} & y_{02} & | & y_{21} & y_{12} & y_{03} \\ \hline y_{20} & | & y_{30} & y_{21} & | & y_{40} & y_{31} & y_{22} \\ y_{11} & | & y_{21} & y_{12} & | & y_{3,1} & y_{22} & y_{13} \\ y_{02} & | & y_{12} & y_{03} & | & y_{22} & y_{13} & y_{04} \end{bmatrix}$$

is replaced with

$$(3.14) \quad \widehat{M}_2(y) = \left[\begin{array}{c|cc|cc|c} 1 & & & & & & & \\ \hline & y_{10} & y_{01} & & y_{10} & y_{11} & y_{01} & \\ \hline y_{10} & & & & & & & \\ y_{01} & & & & & & & \\ \hline & & & & & & & \\ y_{10} & & & & & & & \\ y_{11} & & & & & & & \\ y_{01} & & & & & & & \end{array} \right]$$

and only the variables y_{10}, y_{01}, y_{11} appear in all the relaxations \mathbb{Q}_i .

3.3. Main result. Before stating the main result of this section, begin with the following crucial result:

Proposition 3.1. *Let $r := \sum_{k=1}^n (2r_k - 1)$ and $s := \prod_{k=1}^n 2r_k$.*

- (a) *All the simplified relaxations \mathbb{Q}_i involve at most $s - 1$ variables y_α .*
- (b) *For all the relaxations \mathbb{Q}_i with $i > r$, one has*

$$(3.15) \quad \text{rank } M_i(y) = \text{rank } M_r(y).$$

Proof. (a) is just a consequence of the comment preceding Proposition 3.1.

To get (b) observe that with $i > r$, one may write

$$M_i(y) = \left[\begin{array}{c|c} M_r(y) & B(y) \\ \hline B'(y) & C(y) \end{array} \right]$$

for appropriate matrices B, C , and we next prove that each column of $\begin{bmatrix} B(y) \\ C(y) \end{bmatrix}$ is a linear combination of some columns of $\begin{bmatrix} M_r(y) \\ B'(y) \end{bmatrix}$.

Indeed, remember from (2.3) how an element $M_i(y)(k, p)$ can be obtained. Let $y_\delta = M_i(y)(k, 1)$ and $y_\alpha = M_i(y)(1, p)$. Then

$$(3.16) \quad M_i(y)(k, p) = y_\eta \quad \text{with } \eta_i = \delta_i + \alpha_i, \quad i = 1, \dots, n,$$

that is, $M_i(y)(k, p)$ is the “moment” which corresponds to the monomial $x^\delta \times x^\alpha$ (denoted $M_i(k, p) \leftrightarrow x^\delta \times x^\alpha$). Now, consider a column $\begin{bmatrix} B(y)(\cdot, m) \\ C(y)(\cdot, m) \end{bmatrix}$ of $\begin{bmatrix} B \\ C \end{bmatrix}$, that is, some column $M_i(y)(\cdot, p)$ of $M_i(y)$, with first element $y_\alpha = B(y)(1, m) = M_i(y)(1, p)$. Therefore, the element $\begin{bmatrix} B(y) \\ C(y) \end{bmatrix}(k, m)$ (or, equivalently, the element $M_i(y)(k, p)$) is the variable y_η in (3.16). Note that α corresponds to the monomial $x^{\alpha_1 \dots \alpha_n}$ in the basis (2.1), of degree larger than r . We have seen that in view of the constraint $M_{i-r_k}(g_k y) = 0$, this element satisfies

$$y_\alpha = \sum_j \gamma_j y_{\beta^{(j)}},$$

that is, y_α is a linear combination of variables $y_{\beta^{(j)}}$ with

$$(3.17) \quad \beta^{(j)} = \beta_1^{(j)} \dots \beta_n^{(j)} \quad \text{and } \beta_k^{(j)} < 2r_k, \quad \forall j = 1, \dots, n.$$

Now, as $y_{\beta^{(j)}}$ corresponds to a monomial in the basis (2.1) of degree less than r , to each $y_{\beta^{(j)}}$ corresponds a column $M_r(y)(\cdot, p_j)$ (more precisely, $M_r(y)(1, p_j) \leftrightarrow$

$y_{\beta^{(j)}}$) and thus, $B(y)(1, m)$ is a linear combination of the elements $M_i(y)(1, p_j)$ with coefficients γ_j as in (3.17). Next, by construction, the element $M_i(y)(k, p)$ corresponds to the monomial $x^\delta \times x^\alpha$, and we have the correspondences

$$\begin{aligned} y_\eta = M_i(y)(k, p) \leftrightarrow x^\delta \times x^\alpha &= x^\delta \times \sum_j \gamma_j x^{\beta^{(j)}} \\ &= \sum_j \gamma_j (x^\delta \times x^{\beta^{(j)}}) \leftrightarrow \sum_j \gamma_j \begin{bmatrix} M_r(y) \\ B' \end{bmatrix} (k, p_j), \end{aligned}$$

and thus,

$$M_i(y)(k, p) = \sum_j \gamma_j \begin{bmatrix} M_r(y) \\ B'(y) \end{bmatrix} (k, p_j),$$

which states that the column $\begin{bmatrix} B(y) \\ C(y) \end{bmatrix} (., m)$ is a linear combination of the columns $\begin{bmatrix} M_r(y) \\ B'(y) \end{bmatrix} (., p_j)$. From, this it immediately follows that $\text{rank } M_i(y) = \text{rank } M_r(y)$ whenever $i > r$. □

We now are in position to state the following result.

Theorem 3.2. *Let \mathbb{P} be as in (3.1) with \mathbb{K} as in (3.3) and let $r := \sum_{k=1}^n (2r_k - 1)$, $v := \max\{r_0 - r, \max_{k=1, \dots, n} r_k\}$. Then, whenever $i \geq r + v$,*

$$(3.18) \quad p^* = \min \mathbb{Q}_i,$$

and every optimal solution y^* of \mathbb{Q}_i is the vector of moments of a probability measure μ_{y^*} supported on $s = \text{rank } M_r(y)$ optimal solutions of \mathbb{P} .

Proof. First let $i := r + v$. Let y be any feasible solution of \mathbb{Q}_i . From Proposition 3.1, it follows that $\text{rank } M_{r+1}(y) = \text{rank } M_r(y)$. Therefore, from a deep result of Curto and Fialkow, namely [2, Th. 1.1, p. 4], $M_r(y)$ has positive flat extensions $M_{r+j}(y)$ for all $j = 1, 2, \dots$, that is, $M_{r+j}(y) \succeq 0$ and $\text{rank } M_{r+j}(y) = \text{rank } M_r(y)$ for all $j = 1, 2, \dots$. This implies that y is the vector of moments of some rank $M_r(y)$ -atomic probability measure μ_y . Moreover, from $M_{i-r_k}(g_k y) = 0$ (equivalently $M_{i-r_k}(g_k y) \succeq 0$ and $M_{i-r_k}(-g_k y) \succeq 0$), it also follows that μ_y has its support contained in $\bigcap_{k=1}^n \{g_k(x) = 0\}$, that is, the set \mathbb{K} . In the terminology of Curto and Fialkow [2], $M_{r+v-v_k}(g_k y)$ is well-defined relative to the unique flat extension $M_{r+v}(y)$ of $M_r(y)$. Theorem 1.1 in Curto and Fialkow [2, p. 4] is stated for the complex \mathbb{K} -moment problem, but is valid for the \mathbb{K} -moment problem in several real or complex variables (see [2, p. 2]).

But then, for every admissible solution y of \mathbb{Q}_i , we have

$$\begin{aligned} \sum_\alpha p_\alpha y_\alpha &= \int_{\mathbb{K}} p(x) \mu_y(dx) \\ &\geq p^* \mu_y(\mathbb{K}) = p^* \end{aligned}$$

so that $p^* \geq \inf \mathbb{Q}_i$, which combined with $\inf \mathbb{Q}_i \leq p^*$ for all i , yields $\inf \mathbb{Q}_i = p^*$ when $i = r + v$. For $i > r + v$, it suffices to notice that $p^* \geq \inf \mathbb{Q}_{i+1} \geq \inf \mathbb{Q}_i$ for all i , to get $\inf \mathbb{Q}_i = p^*$ whenever $i > r + v$. Moreover, to a global minimizer $x^* \in \mathbb{K}$ of \mathbb{P} , corresponds the admissible solution

$$y^* = (x_1^*, \dots, x_n^*, \dots, (x_1^*)^{2i}, \dots, (x_n^*)^{2i})$$

of \mathbb{Q}_i , with value p^* , which implies $\inf \mathbb{Q}_i = \min \mathbb{Q}_i = p^*$ whenever $i > r + v$. \square

In Theorem 3.2, we have $i \geq r + v$, and v depends on r_0 . Indeed, to be consistent, \mathbb{Q}_i must be such that $2i \geq 2r_0$ to contain all the moments up to $2r_0$. However, we can make the following remark.

Remark 3.3. We have seen that \mathbb{Q}_i has the simplified equivalent form (3.12) with the single LMI constraint $\widehat{M}_i(y) \succeq 0$ after the simplification induced by the constraints $g_k(x) = 0$. But, since $M_r(y) \succeq 0$ implies that $M_{r+j}(y) \succeq 0$ and $\text{rank } M_{r+j}(y) = \text{rank } M_r(y)$ for all $j = 1, \dots$, it follows that $\widehat{M}_r(y) \succeq 0$ also implies $\widehat{M}_{r+j}(y) \succeq 0$ and $\text{rank } \widehat{M}_{r+j}(y) = \text{rank } \widehat{M}_r(y)$ for all $j = 1, \dots$. Therefore, p^* is obtained by solving the SDP relaxation

$$(3.19) \quad \min_y \left\{ \sum_{\alpha} \tilde{p}_{\alpha} y_{\alpha} \mid \widehat{M}_r(y) \succeq 0 \right\},$$

in which we only have variables y_{α} such that $\alpha_k < 2r_k$ for all $k = 1, \dots, n$.

Therefore, no matter its degree, the global minimization of a polynomial $p(x)$ on \mathbb{K} reduces to a **convex continuous optimization problem of fixed dimension**, namely the convex psd program (3.19). This is because even if $p(x)$ has degree larger than r , the $\{\tilde{p}_{\beta}\}$ are the coefficients of a polynomial $\tilde{p}(x)$ of degree less than r , identical to $p(x)$ on \mathbb{K} . However, this simplification is formally obtained at the relaxation \mathbb{Q}_{r+v} that we will need for the representation of $p(x)$ in §4.

Despite its theoretical interest, Theorem 3.2 is of little value for solving discrete optimization problems for the dimension of the psd program \mathbb{Q}_{r+v} (or, equivalently, (3.19)) is exponential in the problem size.

Fortunately, in many cases, the optimal value p^* is obtained at relaxations \mathbb{Q}_i with $i \ll r + v$. For instance, this will be the case whenever $p(x) - p^*$ has the representation (3.10) with polynomials $\{u_j(x)\}$ of maximum degree $a_0 < r + v$, and polynomials $\{v_{kj}(x), w_{kl}(x)\}$ of maximum degree $a_k < r + v - r_k$, $k = 1, \dots, n$. Indeed, let $\{u_j, v_{kj}, w_{kl}\}$ be their vectors of coefficients in $\mathbb{R}^{s(i)}, \mathbb{R}^{s(i-r_k)}$ respectively, with $i := \max[a_0, \max_k(r_k + a_k)]$. Then, the real-valued symmetric matrices $X \in \mathbb{R}^{s(a_0) \times s(a_0)}$ and $Z_k \in \mathbb{R}^{s(a_k) \times s(a_k)}$, $k = 1, \dots, n$, defined by

$$X := \sum_j u_j u'_j; \quad Z_k := \sum_j v_{kj} v'_{kj} - \sum_l w_{kl} w'_{kl},$$

form an admissible solution of \mathbb{Q}_i^* with value p^* (just redo backward the derivations (3.6)-(3.9)). Since from weak duality we must have $\sup \mathbb{Q}_i^* \leq \inf \mathbb{Q}_i \leq p^*$, it follows that $\inf \mathbb{Q}_i = p^*$, and thus, $\min \mathbb{Q}_i = p^*$ (since the vector y^* of moments of the Dirac measure δ_{x^*} at a global minimizer x^* of \mathbb{P} , is obviously an admissible solution of \mathbb{Q}_i with value p^*).

For instance, consider the so-called MAX-CUT discrete optimization problem

$$\min_{x \in \{0,1\}^n} x' M x,$$

where $M \in \mathbb{R}^{n \times n}$ is a real-valued symmetric matrix with a null diagonal, which is known to be NP-hard. We have solved a sample of MAX-CUT problems in \mathbb{R}^{10} with the the nondiagonal entries of M randomly generated, uniformly between 0 and 1. In all cases, the SDP relaxation \mathbb{Q}_2 provided the optimal value p^* , with no need to solve \mathbb{Q}_{11} , as predicted by Theorem 3.2 (since in this case, with $r_k = 1$, $r = \sum_k (2r_k - 1) = 10$). The computational savings are significant since the simplified

form $\widehat{\mathbb{Q}}_2$ of \mathbb{Q}_2 is a psd program with 385 variables y_α and a single LMI constraint of dimension 56×56 , whereas \mathbb{Q}_{11} would involve $2^1 - 1 = 1023$ variables and a matrix $\widehat{M}_{10}(y)$ of size 1024×1024 !

Remark 3.4. If $p(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ is concave, then its minimum p^* on the grid \mathbb{K} is attained at some “corner” of \mathbb{K} . Therefore, only the corner points are useful and one may replace the initial grid \mathbb{K} , by the new grid \mathbb{K}' defined by the quadratic equality constraints

$$g'_k(x) := (x_k - a_{k1}) \times (x_k - a_{k2r_k}) = 0, \quad k = 1, \dots, n.$$

The resulting grid \mathbb{K}' has only 2^n points and the optimal value p^* is obtained at most at the \mathbb{Q}_{n+v} relaxation (since $r = n$).

3.4. Constrained discrete optimization. Consider now the problem \mathbb{P} with the additional constraints $h_j(x) \geq 0, j = 1, \dots, m$, for some real-valued polynomials $h_j(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ of degree $2v_j$ or $2v_j - 1$ (depending on the parity), that we may assume to be not larger than r . The new set \mathbb{K} is now $\mathbb{K}' := \mathbb{K} \cap [\bigcap_{j=1}^m \{h_j(x) \geq 0\}]$ and the new associated relaxations \mathbb{Q}_i now include the additional constraints $M_{i-v_j}(h_j y) \succeq 0, j = 1, \dots, m$. Theorem 3.2 also holds, that is, the (new) optimal value p^* is also obtained at the relaxation \mathbb{Q}_{r+v} with now $v := \max\{r_0 - r, \max_j v_j, \max_k r_k\}$. Indeed, as in the proof of Theorem 3.2, we may invoke Theorem 1.1 of Curto and Fialkow [2] to show that $M_r(y)$ admits flat positive extensions $M_{r+l}(y), l = 1, \dots$, and as $M_{r+v-v_j}(h_j y)$ are all well defined, relative to the unique flat extension $M_{r+v}(y)$, every admissible solution y of \mathbb{Q}_{r+v} is the the vector of moments of some probability measure μ_y supported on \mathbb{K}' , and the result follows with same arguments.

4. REPRESENTATION OF POLYNOMIALS NONNEGATIVE ON \mathbb{K}

We now investigate the representation of polynomials that are nonnegative on \mathbb{K} . We will see that the previous results obtained for the discrete optimization problem \mathbb{P} will help us to get the desired representation of $p(x) - p^*$, and thus, the representation of every polynomial $p(x)$, nonnegative on \mathbb{K} . We then present results for concave polynomials.

4.1. Representation of polynomials, nonnegative on \mathbb{K} . For a polynomial $g(x)$, nonnegative on \mathbb{K} , let $\tilde{g}(x) := g(x) - g(0)$ and let $\tilde{p}^* := \min_{x \in \mathbb{K}} \tilde{g}(x)$ and $0 \leq p^* := \min_{x \in \mathbb{K}} g(x)$. Then, $g(x) = \tilde{g}(x) - \tilde{p}^* + p^*$. Therefore, if one obtains a representation of $\tilde{g}(x) - \tilde{p}^*$ as a weighted sum of squares, one obtains the same representation for the polynomial $g(x)$, positive on \mathbb{K} , by adding the nonnegative constant term p^* .

Therefore, with no loss of generality, we may and will assume that the constant term of $p(x)$ vanishes, that is, $p(0) = 0$. Consider the polynomial $p(x) - p^*$, with $p^* := \min_{x \in \mathbb{K}} p(x)$, which is nonnegative on \mathbb{K} and let $2r_0$ (resp. $2r_0 - 1$) be its degree if even (resp. odd). Before proceeding further, we need the following remark.

Remark 4.1. We have seen that with

$$v := \max\{r_0 - r, \max_{k=1, \dots, n} r_k\},$$

we have $\min \mathbb{Q}_{r+v} = p^*$ and \mathbb{Q}_{r+v} is equivalent to solving the psd program

$$\min_y \left\{ \sum_{\alpha} \tilde{p}_{\alpha} y_{\alpha} \mid \widehat{M}_r(y) \succeq 0 \right\},$$

where $\widehat{M}_r(y)$ is the “simplified” form of $M_r(y)$ after the simplifications in \mathbb{Q}_{r+v} induced by the constraints $g_k(x) = 0, k = 1, \dots, n$ (cf. (3.19) in Remark 3.3), so that we only have variables y_{α} with $\alpha_k < 2r_k$ for all k .

In fact, $\widehat{M}_r(y)$ can be further simplified and reduced in size. Indeed, all the columns of $\widehat{M}_r(y)$ that correspond to monomials x^{α} in the basis (2.1), and simplified as a linear combination of monomials x^{β} with $\beta_k < 2r_k$ for all k , can be deleted (and the corresponding rows as well). Indeed, by rearranging rows and columns, write

$$\widehat{M}_r(y) = \left[\begin{array}{c|c} H_r(y) & B(y) \\ \hline B'(y) & C(y) \end{array} \right],$$

where the elements of $H_r(y)(1, \cdot)$ correspond to the monomials x^{β} in the basis (2.1) with $\beta_k < 2r_k$ for all k . Thus, $\begin{bmatrix} B(y) \\ C(y) \end{bmatrix}$ is the submatrix of $\widehat{M}_r(y)$ whose columns are precisely the columns of $\widehat{M}_r(y)$ that are linear combinations of the columns of $\begin{bmatrix} H_r(y) \\ B'(y) \end{bmatrix}$. Therefore, we can write $B(y) = H_r(y)W$ and $C(y) = B'(y)W$ for some matrix W . Next, let y be such that $H_r(y) \succeq 0$. From $C(y) = W'H_r(y)W \succeq 0$, it follows that $\widehat{M}_r(y) \succeq 0$ and, in addition, $\text{rank } \widehat{M}_r(y) = \text{rank } H_r(y)$. Therefore, y is admissible for \mathbb{Q}_{r+v} with same value, so that solving \mathbb{Q}_{r+v} is equivalent to solving the new psd program of reduced size

$$(4.1) \quad \widehat{\mathbb{Q}}_r(y) \left\{ \begin{array}{l} \min \sum_{\alpha} \tilde{p}_{\alpha} y_{\alpha} \\ H_r(y) \succeq 0. \end{array} \right.$$

With $s := \prod_{k=1}^n 2r_k$, $H_r(y)$ is a matrix of size $s \times s$, (s is precisely the number of monomials x^{β} with $\beta_k < 2r_k$ for all $k = 1, \dots, n$).

To illustrate the above Remark 4.1, consider the matrix $\widehat{M}_2(y)$ in (3.14) (for the case $n = 2$ and $g_k(x) = x_k^2 - x_k$), which can be written

$$\widehat{M}_2(y) = \left[\begin{array}{cccc|cc} 1 & y_{10} & y_{01} & y_{11} & y_{10} & y_{01} \\ y_{10} & y_{10} & y_{11} & y_{11} & y_{10} & y_{11} \\ y_{01} & y_{11} & y_{01} & y_{11} & y_{11} & y_{01} \\ y_{11} & y_{11} & y_{11} & y_{11} & y_{11} & y_{11} \\ - & - & - & - & - & - \\ y_{10} & y_{10} & y_{11} & y_{11} & y_{10} & y_{11} \\ y_{01} & y_{11} & y_{01} & y_{11} & y_{11} & y_{01} \end{array} \right],$$

with the last two columns being identical to the second and third columns respectively. Therefore, in this case, solving \mathbb{Q}_{2+v} reduces to solving the psd program

$$\widehat{\mathbb{Q}}_2 \left\{ \begin{array}{l} \min_y \sum_{\alpha} \tilde{p}_{\alpha} y_{\alpha} \\ H_2(y) = \begin{bmatrix} 1 & y_{10} & y_{01} & y_{11} \\ y_{10} & y_{10} & y_{11} & y_{11} \\ y_{01} & y_{11} & y_{01} & y_{11} \\ y_{11} & y_{11} & y_{11} & y_{11} \end{bmatrix} \succeq 0. \end{array} \right.$$

We need the simplified psd program $\widehat{\mathbb{Q}}_r$ to prove the absence of a duality gap between $\mathbb{Q}_{r+v}(y)$ and its dual \mathbb{Q}_{r+v}^* , which in turn, will yield the desired result on the representation of polynomials $p(x)$, nonnegative on \mathbb{K} .

Theorem 4.2. *Let $p(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ be an arbitrary polynomial of degree $2r_0$ or $2r_0 - 1$, with $p(0) = 0$. Let \mathbb{P} be as in (3.1) with \mathbb{K} as in (3.3) and let $r := \sum_{k=1}^n (2r_k - 1)$ and $s := \prod_{k=1}^n 2r_k$. Then with $v := \max\{r_0 - r, \max_{k=1}^n r_k\}$.*

(i) *There is no duality gap between the psd program \mathbb{Q}_{r+v} and its dual \mathbb{Q}_{r+v}^* . Moreover, \mathbb{Q}_{r+v}^* is solvable, that is,*

$$(4.2) \quad \max \mathbb{Q}_{r+v}^* = \min \mathbb{Q}_{r+v} = p^*.$$

(ii) *$p(x) - p^*$ has the following representation:*

$$(4.3) \quad p(x) - p^* = \sum_{j=1}^{m_0} u_j(x)^2 + \sum_{k=1}^n g_k(x) \left[\sum_{j=1}^{m_k} v_{kj}(x)^2 - \sum_{l=1}^{n_k} w_{kl}(x)^2 \right],$$

for some $m_0 (\leq s(r + v))$ polynomials $\{u_j(x)\}$ of degree at most $r + v$ and some polynomials $\{v_{kj}(x), w_{kl}(x)\}$ of degree at most $r + v - r_k$, with $m_k + n_k \leq s(r + v - r_k)$, $k = 1, \dots, n$.

Proof. From Remark 4.1, we know that $\min \widehat{\mathbb{Q}}_r = p^*$ and the dimension of the matrix $H_r(y)$ is $s \times s$ with $s := \prod_{k=1}^n 2r_k$. Next, to each of the s (grid) points of \mathbb{K} , labelled $x(k)$, $k = 1, \dots, s$, corresponds an admissible solution $y(k)$ of \mathbb{Q}_{r+v} with

$$(4.4) \quad y(k) = (x(k)_1, \dots, x(k)_n, \dots, x(k)_1^{2(r+v)}, \dots, x(k)_n^{2(r+v)}),$$

and with

$$M_{r+v}(y(k)) \succeq 0, \quad k = 1, \dots, s.$$

Let $\tilde{y}(k)$ be the vector obtained from $y(k)$ by keeping the only variables $y(k)_\alpha$ corresponding to the monomials x^α with $\alpha_k < 2r_k$ for all k , then the vector $(1, \tilde{y}(k))$ satisfies $H_r(\tilde{y}(k)) \succeq 0$ and $\tilde{y}(k)$ is admissible for $\widehat{\mathbb{Q}}_r(y)$ in (4.1).

Moreover, the s points $(1, \tilde{y}(k)) \in \mathbb{R}^s$ are linearly independent (see a proof in the Appendix). Consider a convex combination $z \in \mathbb{R}^s$ defined by

$$z := \sum_{k=1}^s z_k (1, \tilde{y}(k)), \quad \text{with } z_k > 0 \text{ and } \sum_{k=1}^s z_k = 1.$$

From the linearity of $H_r(z)$ it follows that

$$H_r(z) = \sum_{k=1}^s z_k H_r(\tilde{y}(k)) \succeq 0.$$

Moreover, from the definition of the moment matrix $M_{r+v}(y)$, all the matrices $M_{r+v}(y(k))$ are the rank-one matrices

$$M_{r+v}(y(k)) = \begin{bmatrix} 1 \\ y(k) \end{bmatrix} \times [1, y(k)], \quad k = 1, \dots, s.$$

From this and the definition of the simplified moment matrix $H_r(y)$, it also follows that all the matrices $H_r(\tilde{y}(k))$ are the rank-one matrices

$$H_r(\tilde{y}(k)) = \begin{bmatrix} 1 \\ \tilde{y}(k) \end{bmatrix} \times [1, \tilde{y}(k)], \quad k = 1, \dots, s.$$

As the points $(1, \tilde{y}(k))$ are linearly independent in \mathbb{R}^s , it follows that $H_r(z) \succ 0$, that is, $H_r(z)$ is positive definite, and thus, z is a strictly admissible solution of $\widehat{\mathbb{Q}}_r$. In other words, Slater’s interior point condition holds for the convex psd program $\widehat{\mathbb{Q}}_r$. As $\inf \widehat{\mathbb{Q}}_r = p^* > -\infty$, by a standard (strong duality) result in convex optimization, we have that $\widehat{\mathbb{Q}}_r^*$ is solvable and there is no duality gap between $\widehat{\mathbb{Q}}_r$ and $\widehat{\mathbb{Q}}_r^*$ (see e.g. Sturm [13, Th. 2.24]). That is,

$$\max \widehat{\mathbb{Q}}_r^* = \min \widehat{\mathbb{Q}}_r = p^*.$$

If we now remember how $\widehat{\mathbb{Q}}_r$ was obtained from \mathbb{Q}_{r+v} , it also follows that \mathbb{Q}_{r+v}^* is also solvable and there is no duality gap between \mathbb{Q}_{r+v} and \mathbb{Q}_{r+v}^* , which proves the first assertion of Theorem 4.2.

Next, as \mathbb{Q}_{r+v}^* is solvable with optimal value p^* , let $X^* \succeq 0$ and $Z_k^*, k = 1, \dots, n$, be an optimal solution of \mathbb{Q}_{r+v}^* . From the spectral decomposition of the symmetric matrices X^* and Z_k^* , write

$$X^* = \sum_{j=1}^{m_0} u_j u_j' \quad \text{and} \quad Z_k^* = \sum_{j=1}^{m_k} v_{kj} v_{kj}' - \sum_{l=1}^{n_k} w_{kl} w_{kl}', \quad k = 1, \dots, n,$$

where the vectors $\{u_j\}$ correspond to the m_0 positive eigenvalues of X^* , and the vectors $\{v_{kj}\}$ (resp. $\{w_{kl}\}$) correspond to the m_k positive (resp. the n_k negative) eigenvalues of Z_k^* . Consider the polynomials $\{u_j(x)\}$ and $\{v_{kj}(x), w_{kl}(x)\}$ with respective coefficient vectors $\{u_j\}$ and $\{v_{kj}, w_{kl}\}$, in the basis (2.1).

Fix $x \in \mathbb{R}^n$, arbitrary. From the feasibility of (X^*, Z_k^*) in \mathbb{Q}_{r+v}^* ,

$$\langle X^*, B_\alpha x^\alpha \rangle + \sum_{k=1}^n \langle Z_k^*, C_\alpha^k x^\alpha \rangle = p_\alpha x^\alpha, \quad \forall \alpha \neq 0,$$

and from $\max \mathbb{Q}_{r+v}^* = p^*$

$$X^*(1, 1) + \sum_{k=1}^n g_k(0) Z_k^*(1, 1) = -p^*.$$

Using the notation $y^x = (x_1, \dots, x_n, x_1^2, x_1 x_2, \dots, x_1^{2(r+v)}, \dots, x_n^{2(r+v)})$, and summing up over all α , yields

$$\langle X^*, M_{r+v}(y^x) \rangle + \sum_{k=1}^n \langle Z_k^*, M_{r+v-r_k}(g_k y^x) \rangle = p(x) - p^*,$$

or, equivalently,

$$p(x) - p^* = \sum_{j=1}^{m_0} \langle u_j, M_{r+v}(y^x) u_j \rangle + \sum_{k=1}^n \left[\sum_{j=1}^{m_k} \langle v_{kj}, M_{r+v-r_k}(g_k y^x) v_{kj} \rangle - \sum_{l=1}^{n_k} \langle w_{kl}, M_{r+v-r_k}(g_k y^x) w_{kl} \rangle \right].$$

Therefore, using (2.4)-(2.6) (with y^x),

$$p(x) - p^* = \sum_{j=1}^{m_0} u_j(x)^2 + \sum_{k=1}^n g_k(x) \left[\sum_{j=1}^{m_k} v_{kj}(x)^2 - \sum_{l=1}^{n_k} w_{kl}(x)^2 \right]$$

which is (4.3). □

4.2. Concave polynomials. As a consequence of Theorem 4.2, we obtain a representation of concave polynomials. Remember that the global minimum of a concave polynomial on a grid \mathbb{K} is attained at some “corner” point of the grid \mathbb{K} (see Remark 3.4). Therefore, let $p(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ be a concave polynomial of degree $2r_0$ or $2r_0 - 1$, with $p^* := \min_{x \in \mathbb{K}} p(x)$. Then,

(4.5)

$$p(x) - p^* = \sum_{j=1}^{m_0} u_j(x)^2 + \sum_{k=1}^n (x_k - a_{k1})(a_{k2r_k} - x_k) \left[\sum_{j=1}^{m_k} v_{kj}(x)^2 - \sum_{l=1}^{n_k} w_{kl}(x)^2 \right]$$

for some polynomials $\{u_j(x)\}$ of degree at most $n + v$, and some polynomials $\{v_{kj}(x)\}$ and $\{w_{kl}(x)\}$ of degree at most $n + v - 1$.

This is because, we may replace the initial grid \mathbb{K} with the coarser grid \mathbb{K}' defined by the 2^n corner points, so that we have $r = n$ and $v = \max[1, r_0 - n]$. Moreover, from the concavity of $p(x)$,

$$p^* = \min_{x \in \mathbb{K}'} p(x) = \min\{p(x) \mid a_{k1} \leq x_k \leq a_{k2r_k}, \quad k = 1, \dots, n\},$$

that is, p^* is also the global minimum of $p(x)$ on the box $S := \prod_{k=1}^n [a_{k1}, a_{k2r_k}]$ (equivalently, the convex hull of \mathbb{K}). But, observe that (4.5) is also a “linear” representation with respect to the constraints $(x - a_{k1})(a_{k2r_k} - x) \geq 0, k = 1, \dots, n$, that are necessary conditions for $x \in S$. Therefore, we have

$$(4.6) \quad \begin{aligned} p(x) - \min_{x \in S} p(x) &= \sum_{j=1}^{m_0} u_j(x)^2 \\ &+ \sum_{k=1}^n (x_k - a_{k1})(a_{k2r_k} - x_k) \left[\sum_{j=1}^{m_k} v_{kj}(x)^2 - \sum_{l=1}^{n_k} w_{kl}(x)^2 \right]. \end{aligned}$$

Remark 4.3. The reader will easily convince himself that Theorem 3.2 of this paper is also valid if instead of a grid, \mathbb{K} is now the more general variety $\bigcap_{k=1}^n \{g_k(x) = 0\}$, with polynomials $g_k(x)$ of the form

$$x \mapsto g_k(x) := x_k^{r_k} + h_k(x), \quad k = 1, \dots, n,$$

and where $h_k(x)$ is a polynomial of degree not larger than $r_k - 1$.

4.3. Concave polynomials on a simplex. We end up this section with the characterization of concave polynomials on a simplex. Let $\Delta \subset \mathbb{R}^n$ be the canonical simplex

$$(4.7) \quad \Delta := \{x \in \mathbb{R}_+^n \mid \sum_{i=1}^n x_i = 1\}.$$

Let $p(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ be a concave polynomial of degree $2r_0$ or $2r_0 - 1$. Its minimum on Δ is attained at some vertex of Δ . Therefore,

$$(4.8) \quad \begin{aligned} \mathbb{P} \rightarrow p^* = \min_{x \in \Delta} p(x) &= \min_{x \in \Delta \cap \{0,1\}^n} p(x) \\ &= \min_{x \in \{0,1\}^n} \{p(x) \mid \theta(x) := 1 - \sum_{i=1}^n x_i = 0\}. \end{aligned}$$

The constraints $x \in \{0, 1\}^n$ are modelled with the polynomial constraints

$$g_k(x) := x_k - x_k^2 = 0, \quad k = 1, 2, \dots, n.$$

As we already did before, we also assume with no loss of generality that $p(0) = 0$. In view of Section 3.4 and Remark 3.3, (and as $r = n$, $v := \max[1, r_0 - n]$) the optimal value p^* is obtained at the \mathbb{Q}_{n+v} SDP relaxation

$$\mathbb{Q}_{n+v} \left\{ \begin{array}{l} \min \sum_{\alpha} p_{\alpha} y_{\alpha} \\ M_{n+v}(y) \succeq 0, \\ M_{n+v-1}(g_k y) = 0, \quad k = 1, \dots, n. \\ M_{n+v-1}(\theta y) = 0, \end{array} \right.$$

The constraints $M_{n+v-1}(g_k y) = 0$ and $M_{n+v-1}(\theta y) = 0$ generate linear relationships between the variables y_{α} and in view of those constraints, there are only $n - 1$ independent variables that may be chosen to be

$$y_{10\dots 0}, y_{010\dots 0}, \dots, y_{0\dots 010},$$

corresponding to the monomials x_1, \dots, x_{n-1} . Indeed, for instance, the constraint $M_{n+v-1}(\theta y)(1, 1) = 0$ states that

$$y_{10\dots 0} + y_{010\dots 0} + \dots + y_{0\dots 01} = 1,$$

which yields $n - 1$ independent variables among the variables y_{α} corresponding to first-order moments, that is, with $|\alpha| = 1$ (where $|\alpha| = \sum_i \alpha_i$). The constraint $M_{n+v-1}(\theta y)(1, 2) = 0$ generates

$$y_{10\dots 0} - y_{10\dots 0} - y_{110\dots 0} - y_{101\dots 0} - \dots - y_{10\dots 01} = 0,$$

which (as $y_{\alpha} \geq 0$) yields $y_{\alpha} = 0$ for all variables y_{α} corresponding to monomials $x_1 x_i, i = 2, \dots, n$, etc. so that finally, all the variables y_{α} with $|\alpha| = 2$ vanish. From the other constraints induced by $M_{n+v-1}(\theta y) = 0$, it follows that all the variables y_{α} with $|\alpha| > 1$, vanish. This corresponds to the fact that, on $\Delta \cap \{0, 1\}^n$, all the monomials x^{α} with $\alpha_i < 2$ and $|\alpha| > 1$ vanish. Thus, after these substitutions, \mathbb{Q}_{n+v} is the SDP relaxation,

$$\mathbb{Q}_{n+v} \left\{ \begin{array}{l} \min \sum_{|\alpha| \leq 1} \tilde{p}_{\alpha} y_{\alpha} \\ \left[\begin{array}{ccccc} 1 & y_{10\dots 0} & y_{010\dots 0} & \dots & y_{0\dots 01} \\ y_{10\dots 0} & y_{10\dots 0} & 0 & \dots & 0 \\ y_{01\dots 0} & 0 & y_{010\dots 0} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ y_{0\dots 01} & 0 & \dots & 0 & y_{0\dots 01} \end{array} \right] \begin{array}{l} | \\ | \\ | \\ | \\ | \end{array} \begin{array}{l} 0 \\ 0 \\ 0 \\ \dots \\ 0 \end{array} \succeq 0 \end{array} \right.$$

with

$$(4.9) \quad y_{10\dots 0} + y_{010\dots 0} + \dots + y_{0\dots 01} = 1.$$

Therefore, using (4.9) to remove one variable, say $y_{0\dots 01}$, the SDP \mathbb{Q}_{n+v} is strictly equivalent to the simplified SDP relaxation

$$\widehat{\mathbb{Q}}_{n+v} \left\{ \begin{array}{l} \tilde{p}_{0\dots 01} + \min \sum_{|\alpha| \leq 1} (\tilde{p}_{\alpha} - \tilde{p}_{0\dots 01}) y_{\alpha} \\ H(y) := \left[\begin{array}{ccccc} 1 & y_{10\dots 0} & y_{010\dots 0} & \dots & y_{0\dots 010} \\ y_{10\dots 0} & y_{10\dots 0} & 0 & \dots & 0 \\ y_{01\dots 0} & 0 & y_{010\dots 0} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ y_{0\dots 01} & 0 & \dots & 0 & y_{0\dots 010} \end{array} \right] \succeq 0 \end{array} \right.$$

with $\{y_\alpha\} \in \mathbb{R}^{n-1}$. Next, let $\{x(k)\}$, $k = 1, \dots, n$, be the n vertices of Δ , that is, $x(k) = e_k \in \mathbb{R}^n$ with $e_{ki} = \delta_{ki}$. To each $x(k) \in \mathbb{R}^n$ corresponds the vector $(1, y(k)) \in \mathbb{R}^n$ with $y(k)_j = \delta_{kj}$, $j = 1, \dots, n - 1$, for all $k = 1, \dots, n - 1$, and $y(n) = 0$. Therefore, the n -vectors $(1, y(k))$ are linearly independent, and, in addition,

$$H(y(k)) = \begin{bmatrix} 1 \\ y(k) \end{bmatrix} \times [1, y(k)] \succeq 0, \quad k = 1, \dots, n,$$

so that every $y(k)$ is admissible for $\widehat{\mathbb{Q}}_{n+v}$. Moreover, let $z := \sum_{k=1}^n z_k(1, y(k))$ with $z_k > 0$ and $\sum_{k=1}^n z_k = 1$. From the linear independence of the vectors $(1, y(k))$, it follows that $H(z) \succ 0$, and thus, z is a strictly admissible solution of $\widehat{\mathbb{Q}}_{n+v}$, that is, Slater’s interior point condition holds for \mathbb{Q}_{n+v} . As $\min \widehat{\mathbb{Q}}_{n+v} = \min \mathbb{Q}_{n+v} = p^*$, from a standard result in convex optimization, it follows that there is no duality gap between $\widehat{\mathbb{Q}}_{n+v}$ and its dual $\widehat{\mathbb{Q}}_{n+v}^*$, and $\sup \widehat{\mathbb{Q}}_{n+v} = \max \widehat{\mathbb{Q}}_{n+v}^* = p^*$. If we remember how $\widehat{\mathbb{Q}}_{n+v}$ was obtained from \mathbb{Q}_{n+v} , this in turn implies that $\max \mathbb{Q}_{n+v}^* = p^*$.

Now, from the solvability of the dual \mathbb{Q}_{n+v}^* with $\max \mathbb{Q}_{n+v}^* = p^*$, and proceeding as in the proof of Theorem 4.2, we obtain that

$$(4.10) \quad p(x) - p^* = \sum_{j=1}^{m_0} q_j(x)^2 + \sum_{k=1}^n (x_k - x_k^2) \left[\sum_j v_{kj}(x)^2 - \sum_l w_{kl}(x)^2 \right] + \left(1 - \sum_{k=1}^n x_k \right) \left[\sum_j y_{kj}(x)^2 - \sum_l z_{kl}(x)^2 \right]$$

for some polynomials $\{q_j(x)\}$ of degree at most $n + v$, and some polynomials $\{v_{kj}(x)\}$, $\{w_{kl}(x)\}$ and $\{y_{kj}(x), z_{kl}(x)\}$ of degree at most $n + v - 1$.

Hence, every concave polynomial $p(x)$, nonnegative on Δ , has the representation (4.10) (write it $p(x) - p^* + a$ for some scalar $a > 0$).

5. CONCLUSION

We have provided a representation of polynomials that are nonnegative on a grid \mathbb{K} of points of \mathbb{R}^n . This representation is a sum of squares “linearly” weighted by the polynomials defining the grid \mathbb{K} as in Putinar’s representation. However, the degree of the polynomials in this representation is bounded and known in advance and depends on the size of the grid, not on the points of the grid. A related result is that every discrete optimization problem (on a grid) is also equivalent to a continuous convex optimization problem whose size depends on the size of the grid but not on the points of the grid.

6. APPENDIX

Let $\{y(k)\}$, $k = 1, \dots, s$, be the vectors in (4.4), that is, $y(k)$ is the vector $\{x(k)^\beta\}$ of all monomials x^β with $\beta_k < 2r_k$, evaluated at the point $x(k)$ of the grid \mathbb{K} . We show by induction on the dimension of the space \mathbb{R}^n , that the s points $\{(1, y(k))\}$ in \mathbb{R}^s are linearly independent.

Denote by \mathbb{K}_n the grid in \mathbb{R}^n and let $S_n \in \mathbb{R}^{s_n \times s_n}$ be the matrix formed by the s_n points $\{(1, y(k))\}$ in \mathbb{R}^{s_n} where $s_n = \prod_{k=1}^n (2r_k)$, that is,

$$S_n = \begin{bmatrix} 1 & 1 & \dots & 1 \\ a_{11} & a_{11} & \dots & a_{1r_1} \\ a_{21} & a_{21} & \dots & a_{2r_2} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nr_n} \\ a_{11}^2 & a_{11}^2 & \dots & a_{1r_1}^2 \\ a_{11}a_{21} & a_{11}a_{21} & \dots & a_{1r_1}a_{2r_2} \\ \dots & \dots & \dots & \dots \end{bmatrix}.$$

Assume that the property is true (that is, S_n is nonsingular) whenever the dimension is $1, 2, \dots, n$. Consider a grid \mathbb{K}_{n+1} in \mathbb{R}^{n+1} defined by $g_k(x) = 0$ for all $k = 1, \dots, n + 1$, and where

$$g_k(x) := \prod_{j=1}^{2r_k} (x - a_{kj}), \quad k = 1, 2, \dots, n + 1.$$

With $s_{n+1} := \prod_{k=1}^{n+1} 2r_k$, we also define the s_{n+1} points $\{(1, \tilde{y}(k))\}$ in $\mathbb{R}^{s_{n+1}}$ and the associated matrix S_{n+1} .

From its definition, and for an easy use of the induction argument, we can rearrange S_{n+1} to rewrite it as

$$S_{n+1} = \left[\begin{array}{c|c|c|c} S_n & S_n & \dots & S_n \\ \hline - & - & - & - \\ \hline C_1 & C_2 & \dots & C_{2r_{n+1}} \end{array} \right],$$

with

$$C_1 = \begin{bmatrix} a_{n+11} & a_{n+11} & \dots & a_{n+11} \\ a_{11}a_{n+11} & a_{11}a_{n+11} & \dots & a_{1r_1}a_{n+1,1} \\ a_{11}^2a_{n+11} & \dots & \dots & a_{1r_1}^2a_{n+11} \\ \dots & \dots & \dots & \dots \end{bmatrix}$$

that is, the first column $C_1(:, 1)$ of C_1 contains all the monomials x^β with $1 < \beta_{n+1} < 2r_{n+1}$ evaluated at the point $\tilde{y}(k) = (a_{11}, a_{21}, \dots, a_{n1}, a_{n+11})$ of the grid \mathbb{K}_{n+1} . Similarly, the second column $C_1(:, 2)$ of C_1 contains all the monomials x^β with $1 < \beta_{n+1} < 2r_{n+1}$ evaluated at the point $(a_{11}, a_{21}, \dots, a_{n2}, a_{n+11})$, etc.

C_i is a verbatim copy of C_1 with a_{n+11} replaced with a_{n+1i} , for all $i = 1, \dots, 2r_{n+1}$. Next, by the induction hypothesis, S_n is full rank, and thus the only way for S_{n+1} to be singular, is that at least one column of $\begin{bmatrix} S_n \\ C_i \end{bmatrix}$ matches exactly one

column of $\begin{bmatrix} S_n \\ C_j \end{bmatrix}$ for some pair (i, j) . But this is not possible. Indeed, assume that $C_i(:, m) = C_j(:, p)$ for some pair (m, p) . Note that because of S_n , we must have $m = p$. An element $C_i(k, p)$ is of the form $x_1^{\beta_1} \dots x_n^{\beta_n} a_{n+1i}^{\beta_{n+1}}$, whereas the element $C_j(k, p)$ is $x_1^{\beta_1} \dots x_n^{\beta_n} a_{n+1j}^{\beta_{n+1}}$. Therefore, it suffices to consider a row k corresponding to a monomial x^β in the basis (2.1), with $\beta_i = 0$ for all $i = 1, \dots, n$, to yield the contradiction $a_{n+1i}^{\beta_{n+1}} = a_{n+1j}^{\beta_{n+1}}$ since the a_{n+1k} 's are distinct.

It remains to prove that the induction hypothesis is true for $n = 1$. With $n = 1$ and a grid \mathbb{K}_1 of $2r_1$ points, the matrix S_1 reads

$$\begin{bmatrix} 1 & 1 & \dots & 1 \\ x_1 & x_2 & \dots & x_{2r_1} \\ x_1^2 & x_2^2 & \dots & x_{2r_1}^2 \\ \dots & \dots & \dots & \dots \\ x_1^{2r_1-1} & x_2^{2r_1-1} & \dots & x_{2r_1}^{2r_1-1} \end{bmatrix}.$$

If S_1 is singular, then there is a linear combination $\{\gamma_k\}$ of the rows such that

$$\sum_{k=0}^{2r_1-1} \gamma_k x^k = 0, \quad \forall x \in \mathbb{K}_1,$$

that is, the univariate polynomial $x \mapsto h(x) := \sum_{k=0}^{2r_1-1} \gamma_k x^k$ has $2r_1$ distinct real zeros, which is impossible.

REFERENCES

- [1] C. BERG, *The multidimensional moment problem and semi-groups*, in: *Moment in Mathematics* (1987), pp. 110–124, ed. H.J. Landau, Amer. Math. Soc. MR **89k**:44013
- [2] R.E. CURTO AND L. A. FIALKOW, *The truncated complex K -moment problem*, *Trans. Amer. Math. Soc.* **352** (2000), pp. 2825–2855. MR **2000j**:47027
- [3] T. JACOBI, *A representation theorem for certain partially ordered commutative rings*, *Mathematische Zeitschrift*, **237** (2001), pp. 259–273. CMP 2001:14
- [4] T. JACOBI AND A. PRESTEL, *On special representations of strictly positive polynomials*, Technical report, Konstanz University, Konstanz, Germany, January 2000.
- [5] T. JACOBI AND A. PRESTEL, *Distinguished representations of strictly positive polynomials*, *J. Reine Angew. Math.* **532** (2001), 223–235. CMP 2001:09
- [6] M. KOJIMA AND L. TUNÇEL, *Cones of matrices and successive convex relaxations of non convex sets*, *SIAM J. Optim.* **10** (2000), 750–778. MR **2001i**:90062
- [7] J.B. LASSERRE, *Global optimization with polynomials and the problem of moments*, *SIAM J. Optim.* **11** (2001), 796–817.
- [8] J.B. LASSERRE, *Optimality conditions and LMI relaxations for 0-1 programs*, LAAS-CNRS Technical report #00099, submitted.
- [9] L. LOVÁSZ AND A. SCHRIJVER, *Cones of matrices and set-functions and 0-1 optimization*, *SIAM J. Optim.* **1** (1991), 166–190. MR **92c**:52016
- [10] M. PUTINAR, *Positive polynomials on compact semi-algebraic sets*, *Ind. Univ. Math. J.* **42** (1993), pp. 969–984. MR **95h**:47014
- [11] K. SCHMÜDGEN, *The K -moment problem for compact semi-algebraic sets*, *Math. Ann.* **289** (1991), pp. 203–206. MR **92b**:44011
- [12] B. SIMON, *The classical moment problem as a self-adjoint finite difference operator*, *Adv. Math.* **137** (1998), 82–203. MR **2001e**:47020
- [13] J. F. STURM, *Theory and algorithms of semidefinite programming*, in *High Performance Optimization methods*, H. Frenk, K. Roos and S. Zhang eds., Kluwer Academic Publishers, Dordrecht, 2000, pp. 3–20. MR **2001e**:90001
- [14] L. VANDENBERGHE AND S. BOYD, *Semidefinite programming*, *SIAM Review* **38** (1996), pp. 49–95. MR **96m**:90005

LAAS-CNRS, 7 AVENUE DU COLONEL ROCHE, 31077 TOULOUSE CÉDEX, FRANCE
E-mail address: lasserre@laas.fr