

THE SUPER ORDER DUAL OF AN ORDERED VECTOR SPACE AND THE RIESZ–KANTOROVICH FORMULA

CHARALAMBOS D. ALIPRANTIS AND RABEE TOURKY

ABSTRACT. A classical theorem of F. Riesz and L. V. Kantorovich asserts that if L is a vector lattice and f and g are order bounded linear functionals on L , then their supremum (least upper bound) $f \vee g$ exists in L^\sim and for each $x \in L_+$ it satisfies the so-called *Riesz–Kantorovich formula*:

$$[f \vee g](x) = \sup\{f(y) + g(z) : y, z \in L_+ \text{ and } y + z = x\}.$$

Related to the Riesz–Kantorovich formula is the following long-standing problem: *If the supremum of two order bounded linear functionals f and g on an ordered vector space exists, does it then satisfy the Riesz–Kantorovich formula?*

In this paper, we introduce an extension of the order dual of an ordered vector space and provide some answers to this long-standing problem. The ideas regarding the Riesz–Kantorovich formula owe their origins to the study of the fundamental theorems of welfare economics and the existence of competitive equilibrium. The techniques introduced here show that the existence of decentralizing prices for efficient allocations is closely related to the above-mentioned problem and to the properties of the Riesz–Kantorovich formula.

1. INTRODUCTION

Consider two ordered (real) vector spaces L and M , and let $T: L \rightarrow M$ be a linear operator. The operator T is said to be *positive*, in symbols $T \geq 0$, if $Tx \geq 0$ holds for each $x \geq 0$. Also, T is a *regular operator* if T can be written as a difference of two positive operators. Similarly, T is called *order bounded* if it carries order bounded subsets of L to order bounded subsets of M . If $\mathcal{L}^\sim(L, M)$ and $\mathcal{L}^r(L, M)$ denote the vector spaces of all order bounded and regular operators from L to M , respectively, then we have $\mathcal{L}^r(L, M) \subseteq \mathcal{L}^\sim(L, M)$. Moreover, $\mathcal{L}^r(L, M)$ and $\mathcal{L}^\sim(L, M)$ are both ordered vector spaces when equipped with the ordering \geq defined by $S \geq T$ whenever $S - T \geq 0$. (The inequality $S \geq T$ is also expressed by saying that S *dominates* T .) The ordered vector space $L^\sim = \mathcal{L}^\sim(L, \mathbb{R})$ is called the *order dual* of L . The vector space of all *regular* linear functionals on L is denoted $L^r = \mathcal{L}^r(L, \mathbb{R})$.

Received by the editors April 20, 2000 and, in revised form, August 16, 2001.

2000 *Mathematics Subject Classification*. Primary 46A40, 46E99, 47B60; Secondary 91B50.

Key words and phrases. Ordered vector space, super order dual, Riesz–Kantorovich formula, decentralizing prices.

The research of C. D. Aliprantis is supported by NSF Grant EIA-007506, and the research of R. Tourky is funded by Australian Research Council Grant A00103450.

As mentioned in the abstract, this work is associated with the following famous theorem due to F. Riesz [28] and L. V. Kantorovich [20]; for a proof see [11, Theorem 1.13, p. 12]. (Recall that the *Riesz Decomposition Property* asserts that: If $0 \leq y \leq x_1 + x_2$ and $x_1, x_2 \in L_+$, then there exist $y_1, y_2 \in L_+$ satisfying $y = y_1 + y_2$, $0 \leq y_1 \leq x_1$, and $0 \leq y_2 \leq x_2$.)

Theorem 1.1 (Riesz–Kantorovich). *If L is an ordered vector space with the Riesz Decomposition Property and M is a Dedekind complete vector lattice, then the ordered vector space $\mathcal{L}^\sim(L, M)$ is a Dedekind complete vector lattice. Moreover, if $S, T \in \mathcal{L}^\sim(L, M)$, then for each $x \in L_+$ we have*

$$\begin{aligned} [S \vee T](x) &= \sup\{Sy + T(x - y) : 0 \leq y \leq x\}, \\ [S \wedge T](x) &= \inf\{Sy + T(x - y) : 0 \leq y \leq x\}. \end{aligned}$$

The above formulas describing the lattice operations in $\mathcal{L}^\sim(L, M)$ are known as the *Riesz–Kantorovich formulas*. Related to these formulas is the following long-standing unsolved question.

- *If L and M are vector lattices with M not Dedekind complete and the supremum (least upper bound) $S \vee T$ of two operators $S, T \in \mathcal{L}^\sim(L, M)$ exists in $\mathcal{L}^\sim(L, M)$, does it then satisfy the Riesz–Kantorovich formula?*

This question and some related ones have been studied extensively in the past by many investigators; see for instance [1, 2, 3, 4, 5, 6, 30]. Although some progress has been made and some partial answers have been given over the last fifty years, a satisfactory solution to this problem has not yet been obtained.

This paper provides some answers to this question when L is an ordered topological vector space and $M = \mathbb{R}$. Notably, many of the ideas presented here regarding the Riesz–Kantorovich formulas owe their origins to recent advances in the theory of general economic equilibrium. We shall illustrate briefly below the connection between economic theory and the Riesz–Kantorovich formula.¹

For the rest of the paper, L will denote an ordered vector space with a generating positive cone L_+ , i.e., $L = L_+ - L_+$. Let f and g be a pair of order bounded linear functionals on L (i.e., $f, g \in L^\sim$) for which the supremum $f \vee g$ exists in L^\sim . We shall say that $f \vee g$ *satisfies the Riesz–Kantorovich formula* if for any $\omega \in L_+$ we have

$$[f \vee g](\omega) = \sup_{0 \leq x \leq \omega} [f(x) + g(\omega - x)].$$

In this case, the long-standing open problem regarding the Riesz–Kantorovich formula can be restated as follows:

- *Are there an ordered vector space L and two order bounded linear functionals f and g for which the supremum $f \vee g$ exists in L^\sim and fails to satisfy the Riesz–Kantorovich formula?*

To answer this question, we introduce a new collection of real-valued functions defined on the positive cone L_+ that includes all order bounded linear functionals on L . This collection (which we shall denote \mathcal{H}_s^\sim and call the *super order dual* of L) consists of all order upper bounded, positively homogeneous, and super-additive functions from L_+ to \mathbb{R} . The super order dual \mathcal{H}_s^\sim inherits the partial ordering

¹L. V. Kantorovich and T. C. Koopmans shared the 1975 Nobel Prize in Economics “for their contributions to the theory of optimum allocation of resources.”

of \mathbb{R}^{L_+} . That is, if $\psi, \phi \in \mathcal{H}_s^\sim$, then ψ dominates ϕ if $\psi(x) \geq \phi(x)$ holds for all $x \in L_+$. This partial ordering has the following basic property.

Proposition 1. *The collection of all positive functions in \mathcal{H}_s^\sim is a lattice, and the supremum of every pair of positive functions satisfies the Riesz–Kantorovich formula, i.e., for $0 \leq \psi, \phi \in \mathcal{H}_s^\sim$ and each $\omega \in L_+$ we have*

$$[\psi \vee \phi](\omega) = \sup_{0 \leq x \leq \omega} [\psi(x) + \phi(\omega - x)].$$

If we also assume that L is equipped with a Hausdorff locally convex topology τ for which the order intervals are τ -bounded and the positive cone L_+ is closed, then we can distinguish one more collection of functions from L_+ to \mathbb{R} . This is the collection \mathcal{H}'_s of all upper τ -semicontinuous, positively homogeneous, and super-additive functions from L_+ to \mathbb{R} . We call this collection the *super topological dual* of L . Clearly, $L' \subseteq \mathcal{H}'_s$, where $L' = (L, \tau)'$ is the topological dual of (L, τ) . Also, $\mathcal{H}'_s \subseteq \mathcal{H}_s^\sim$. We shall establish the following result on the embedding of L' into \mathcal{H}'_s .

Proposition 2. *The canonical embedding of L' into \mathcal{H}'_s preserves finite supremum, i.e., if f and g are continuous linear functionals on L and the supremum $f \vee g$ exists in L' , then*

$$(f \vee g)|_{L_+} = f|_{L_+} \vee g|_{L_+},$$

where the supremum on the right is taken in \mathcal{H}'_s .

Recently, in the study of market equilibrium there has been some attention on weakening the pervasive requirement that commodity spaces be vector lattices.² Indeed, as shown in [14], one can drop all vector lattice theoretic considerations if linear prices are replaced by the possibly non-linear Riesz–Kantorovich functionals. For any m -tuple of functions $f = (f_1, \dots, f_m)$ in \mathcal{H}_s^\sim we define its *Riesz–Kantorovich functional* $\mathcal{R}_f: L_+ \rightarrow \mathbb{R}$ via the formula

$$\mathcal{R}_f(\omega) = \sup \left\{ f_1(x_1) + \dots + f_m(x_m) : x_i \in L_+ \text{ for each } i \text{ and } \sum_{i=1}^m x_i \leq \omega \right\}.$$

By using the Riesz–Kantorovich functionals as the generalized measures of economic value (i.e., as prices) one can extend the fundamental results on the Arrow–Debreu–McKenzie model of general competitive equilibrium beyond vector lattices and to general ordered vector spaces.³ Notice that the Riesz–Kantorovich functional is always an element of \mathcal{H}_s^\sim ; in fact, it is the supremum of the set $\{0, f_1, \dots, f_m\}$ in \mathcal{H}_s^\sim . This paper also establishes the following result, which connects Proposition 1 on the lattice structure of \mathcal{H}_s^\sim and the embedding result for \mathcal{H}'_s in Proposition 2.

² Infinite dimensional equilibrium analysis in economics is typically done in a vector lattice setting. This is in sharp contrast to the coordinate-free finite dimensional analysis of G. Debreu [17]. In fact, all the important results in the literature that establish the fundamental theorems of welfare economics or the existence of market equilibrium utilize in a non-trivial way the lattice structure of the commodity space and its dual. For example, Mas-Colell in [24] uses the local solidness and the Riesz Decomposition Property of the Riesz commodity space; for more on these techniques see [8, 9].

³ In [14] it is also proposed to interpret the Riesz–Kantorovich functional as the revenue function from a discriminatory price auction, which is a generalization of the U.S. Treasury Bill Auction. See also [13].

Proposition 3. *Let $f = (f_1, f_2)$ be a pair of positive continuous linear functionals on L . If the supremum $f_1 \vee f_2$ exists in L' , then it satisfies the Riesz–Kantorovich formula if and only if the Riesz–Kantorovich functional \mathcal{R}_f is upper τ -semicontinuous.*

In particular, if the order intervals of L are weakly compact and either

- a. L_+ has an interior point, or
- b. L is an ordered Fréchet space,

then $f_1 \vee f_2$ must satisfy the Riesz–Kantorovich formula. Moreover, if order intervals of L are not weakly compact, then $f_1 \vee f_2$ must satisfy the Riesz–Kantorovich formula in the algebraic dual $(L')^*$ of L' .

As we shall see, continuity conditions for the Riesz–Kantorovich functionals offer a rich insight into the structure of partially ordered vector spaces that are not vector lattices. For example, it turns out that the order dual of L coincides with the regular dual of L (as in the case of vector lattices) if and only if every Riesz–Kantorovich functional is upper semicontinuous. Furthermore, if the Riesz–Kantorovich functionals are upper semicontinuous, then the order intervals in L are $\sigma((L')^*, L')$ -dense in the order intervals of $(L')^*$; this extends an important property of locally convex-solid vector lattices.

It has become apparent that there is a remarkable relationship between the fundamental theorems of welfare economics and the properties of the Riesz–Kantorovich functional. These fundamental theorems state that an allocation of resources is efficient if and only if the allocation can be decentralized by a price system, i.e., by a continuous linear functional. The welfare theorems are the basic results that provide conceptual foundations for our understanding of efficiency and decentralization of resources. Moreover, all applied economic policy analysis relies implicitly on the foundations provided by these theorems. For details about the welfare theorems, see [9].

This paper also establishes the following result, which has many consequences in economic theory.

Proposition 4. *If L is an ordered Fréchet space with weakly compact intervals, then for any m -tuple $f = (f_1, \dots, f_m)$ of linear functionals in L^\sim and any $\omega \in L_+$ we have*

$$(\star) \quad \inf_{g \geq f_1, \dots, g \geq f_m} g(\omega) = \mathcal{R}_f(\omega).$$

Surprisingly enough, it can be shown (see [12]) that the so-called third fundamental theorem of welfare economics is basically equivalent to the statement that for any m -tuple of linear functionals $f = (f_1, \dots, f_m)$ in L^\sim and any $\omega \in L_+$ a solution to the minimization problem in (\star) exists, i.e., there exists some $g^* \in L^\sim$ satisfying $g^* \geq f_i$ for each i and $g^*(\omega) = \mathcal{R}_f(\omega)$.

The material in this paper is organized as follows. The mathematical preliminaries are included in Section 2. Section 3 introduces the super order dual of an ordered vector space, which is an extension of the order dual. The lattice properties of the super order dual are discussed in Section 4. Our main embedding result and its consequences can be found in Sections 5, 6, and 7. Examples and final remarks are in the concluding section of this paper.

2. MATHEMATICAL PRELIMINARIES

For notation and terminology not explained below regarding ordered vector spaces we refer the reader to the monographs [19, 23, 25, 31]. In this work the letter L will denote exclusively an *ordered vector space*. That is, L is a real vector space equipped with an order relation \geq such that $x \geq y$ implies $x + z \geq y + z$ for all $z \in L$ and $\alpha x \geq \alpha y$ for each $\alpha \geq 0$. The convex set $L_+ = \{x \in L: x \geq 0\}$ is called the *positive cone* of L and its members are referred to as *positive vectors*.⁴ The positive cone L_+ satisfies the following properties:

1. $L_+ + L_+ \subseteq L_+$,
2. $\alpha L_+ \subseteq L_+$ for each $\alpha \geq 0$, and
3. $L_+ \cap (-L_+) = \{0\}$.

Any subset C of a vector space X that satisfies the above properties (1), (2), and (3) is called a *convex cone*. Every convex cone C induces a natural order \geq on X by letting $x \geq y$ whenever $x - y \in C$. This order makes X an ordered vector space satisfying $X_+ = C$. In other words, an ordered vector space is completely characterized by its positive cone. Any subset C of a vector space that satisfies the above properties (1) and (2) is called a *wedge*.

The cone L_+ is said to be *generating* if for each $x \in L$ there exist $y, z \in L_+$ such that $x = y - z$. Equivalently, L_+ is generating if every vector of L is dominated by some positive vector. In this work we shall assume that:

- *The cone L_+ is always generating.*

A (nonempty) subset A of L is *bounded from above* (resp. *from below*) if there exists some $x \in L$ satisfying $a \leq x$ (resp. $x \leq a$) for all $a \in A$; the vector x is called an *upper bound* (resp. *lower bound*) of A . A set is *order bounded* if it is bounded from above and below. Sets of the form $[x, y] = \{z \in L: x \leq z \leq y\}$ are called *order intervals* or simply *intervals*. Clearly, a subset of L is order bounded if and only if it is included in an interval.

A subset A of L has a *least upper bound* (or a *supremum*), denoted $\sup A$, if $\sup A$ is an upper bound of A and whenever x is an upper bound of A , then $\sup A \leq x$. The *greatest lower bound* (or *infimum*) is defined analogously. Following the standard lattice terminology, the supremum and infimum of a set $\{x, y\}$ will be denoted by $x \vee y$ and $x \wedge y$, respectively. That is,

$$x \vee y = \sup\{x, y\} \quad \text{and} \quad x \wedge y = \inf\{x, y\}.$$

For an m -tuple x_1, x_2, \dots, x_m of vectors we shall write $\bigvee_{i=1}^m x_i$ for the supremum of the set $\{x_1, x_2, \dots, x_m\}$ and $[\bigvee_{i=1}^m x_i]^+$ for the supremum of the set $\{0, x_1, \dots, x_m\}$. That is,

$$\bigvee_{i=1}^m x_i = \sup\{x_1, x_2, \dots, x_m\} \quad \text{and} \quad \left[\bigvee_{i=1}^m x_i\right]^+ = \sup\{0, x_1, \dots, x_m\}.$$

An ordered vector space L is called a *Riesz space* (or a *vector lattice*) if $x \vee y$ and $x \wedge y$ exist in L for all $x, y \in L$. For extensive and detailed treatments of Riesz spaces see [10, 11] and [23].

The symbol L^* denotes the *algebraic dual* of L , i.e., the real vector space of all linear functionals on L . A linear functional $f \in L^*$ is said to be *positive* if $f(x) \geq 0$ holds for all $x \in L_+$. The collection of all positive linear functionals on L is denoted

⁴ The inequality $x \geq y$ is also expressed by saying that x *dominates* y .

by L_+^* and is known as the *cone of positive linear functionals*. Since the cone L_+ is generating, it follows that L^* is automatically an ordered vector space by letting $f \geq g$ whenever $f - g \in L_+^*$ or $f(x) \geq g(x)$ for each $x \in L_+$.

A linear functional f is said to be *order bounded* if it carries order bounded sets of L to bounded subsets of \mathbb{R} . The collection L^\sim of all order bounded linear functionals on L is a vector subspace of L^* and is called the *order dual* of L . Clearly, every positive linear functional is order bounded, and so every *regular linear functional* (i.e., every linear functional that can be written as a difference of two positive linear functionals) is likewise order bounded. If (as usual) L^r denotes the vector space of all regular linear functionals, then we have the following vector subspace inclusions:

$$L^r \subseteq L^\sim \subseteq L^* .$$

It should be noticed that if L^\sim is a Riesz space, then $L^r = L^\sim$.

We close the section with a useful result from the work of T. Andô [15].

Lemma 2.1. *Let (L, τ) be a complete metrizable ordered topological vector space such that L_+ is generating and τ -closed. If $\{U_n : n \in \mathbb{N}\}$ is a τ -neighborhood base of zero consisting of τ -closed circled sets such that $U_{n+1} + U_{n+1} \subseteq U_n$ holds for each n , then the countable collection $\{V_n\}$, where*

$$V_n = U_n \cap L_+ - U_n \cap L_+ ,$$

is also a τ -neighborhood base of zero.

Proof. Let τ_1 be the linear topology of L with basis $\{V_n\}$. By [29, Lemma 2, p. 221], (L, τ_1) is a complete metrizable topological vector space. Moreover, the identity operator $I: (L, \tau_1) \rightarrow (L, \tau)$ is continuous. But then, by the Open Mapping Theorem (see, for instance [18, Theorem 8, p. 41]), I is a homeomorphism. This implies $\tau_1 = \tau$. \square

3. THE SUPER ORDER DUAL OF AN ORDERED VECTOR SPACE

In this section, we shall introduce a collection of real-valued functions defined on the positive cone L_+ of an ordered vector space L that provides an extension of the topological and algebraic duals of L . We shall refer to this collection of real-valued functions as the *super algebraic dual* of L . Before we introduce this dual, let us isolate some properties of functions which will play an important role in our study.

Definition 3.1. A function $f: L_+ \rightarrow \mathbb{R}$ is said to be:

1. **Positive**, if $f(x) \geq 0$ holds for all $x \in L_+$.
2. **Super-additive**, if $f(x) + f(y) \leq f(x + y)$ holds for all $x, y \in L_+$.
3. **Positively homogeneous**, if $f(\alpha x) = \alpha f(x)$ holds for every $\alpha \geq 0$ and all $x \in L_+$.
4. **Monotone**, if $0 \leq x \leq y$ implies $f(x) \leq f(y)$.
5. **Upper order bounded**, if it carries order intervals of the form $[0, x]$ to bounded from above subsets in \mathbb{R} .

If $f \in L^*$, then $f|_{L_+}$ (the restriction of f to L_+) is additive and positively homogeneous. In particular, $f|_{L_+}$ is super-additive and positively homogeneous. On the other hand, if $f: L_+ \rightarrow \mathbb{R}$ is additive and positively homogeneous, then the formula

$$\hat{f}(x) = f(x_1) - f(x_2) ,$$

where $x = x_1 - x_2$ and $x_1, x_2 \in L_+$, defines a linear functional on L (i.e., $\hat{f} \in L^*$) that extends f (uniquely) to all of L .⁵ Therefore, the members of L^* can be identified with the additive positively homogeneous real functions on L_+ .⁶ This observation leads us to the definition of the super algebraic dual of L , which will play a central role in this paper.

Definition 3.2. The collection of all super-additive and positively homogeneous real functions defined on L_+ is called the **super algebraic dual** of L and is denoted \mathcal{H}_s^* . That is,

$$\mathcal{H}_s^* = \{f \in \mathbb{R}^{L_+} : f \text{ is super-additive and positively homogeneous}\}.$$

We also isolate and name some important subsets of the super algebraic dual. First, we define the *super dual cone* \mathcal{H}_s^+ of L , which is the collection of all functions in \mathcal{H}_s^* that are positive. That is,

$$\mathcal{H}_s^+ = \{f \in \mathcal{H}_s^* : f(x) \geq 0 \text{ for all } x \in L_+\}.$$

Second, the *super order dual* \mathcal{H}_s^\sim of L is the collection of all upper order bounded functions in \mathcal{H}_s^* . That is,

$$\mathcal{H}_s^\sim = \{f \in \mathcal{H}_s^* : f([0, x]) \text{ is bounded from above in } \mathbb{R} \text{ for all } x \in L_+\}.$$

The above collections of functions will be referred to collectively as the *super duals* of L . A few basic properties of functions in the super order dual are included in the next result.

Lemma 3.3. *Every function in \mathcal{H}_s^* is concave and vanishes at zero.*

Corollary 3.4. *If $f \in \mathcal{H}_s^*$, then for each $x \in L_+$ the **upper contour set***

$$U_f^x = \{y \in X : f(y) \geq f(x)\}$$

of f at x is convex. Moreover, if $f \in \mathcal{H}_s^+$, then for each $x \in L_+$ the upper contour set U_f^x is convex and comprehensive, i.e., $U_f^x + L_+ \subseteq U_f^x$.

The functions in \mathcal{H}_s^+ are monotone.

Lemma 3.5. *Every function in \mathcal{H}_s^+ is monotone and upper order bounded.*

Proof. Let $f \in \mathcal{H}_s^+$, and assume that $x, y \in L_+$ satisfy $x \geq y$. Since $x - y \geq 0$, the super-additivity and positivity of f imply

$$f(x) = f(x - y + y) \geq f(x - y) + f(y) \geq f(y).$$

Thus, f is monotone and $f([0, x]) \subseteq [0, f(x)]$. □

Lemma 3.6. *The super duals \mathcal{H}_s^+ , \mathcal{H}_s^\sim , and \mathcal{H}_s^* are all wedges in \mathbb{R}^{L_+} , and they are closed under finite pointwise infima. Moreover, \mathcal{H}_s^+ is a convex cone in \mathbb{R}^{L_+} .*

Proof. We show that \mathcal{H}_s^* is a wedge and that it is closed under finite pointwise infima. The other cases are trivial. So, let $f, g \in \mathcal{H}_s^*$. Clearly, $f + g$ is positively homogeneous. Now if $x, y \in L_+$, then we have

$$\begin{aligned} (f + g)(x + y) &= f(x + y) + g(x + y) \geq [f(x) + f(y)] + [g(x) + g(y)] \\ &= [f(x) + g(x)] + [f(y) + g(y)] = (f + g)(x) + (f + g)(y). \end{aligned}$$

⁵ When L is Archimedean, L. V. Kantorovich has shown that a function $f: L_+ \rightarrow \mathbb{R}_+$ extends to a positive linear functional on L if and only if f is additive; see [11, Theorem 1.7, p. 7].

⁶ Notice here that this is possible since the cone of L_+ is generating. If L_+ is not generating, then our analysis must be confined to the vector subspace generated by L_+ , i.e., to $L_+ - L_+$.

Therefore, $f + g$ is super-additive, i.e., $f + g \in \mathcal{H}_s^*$. Thus, \mathcal{H}_s^* is a wedge in \mathbb{R}^{L_+} .

To see that \mathcal{H}_s^* is closed under finite pointwise infima, let $f, g \in \mathcal{H}_s^*$ again, and define $h: L_+ \rightarrow \mathbb{R}$ by $h(x) = \min\{f(x), g(x)\}$. Clearly, h is positively homogeneous. If $x \in L_+$, then

$$h(x) + h(y) \leq f(x) + f(y) \leq f(x + y) \quad \text{and} \quad h(x) + h(y) \leq g(x + y).$$

Therefore, $h(x) + h(y) \leq \min\{f(x + y), g(x + y)\} = h(x + y)$. This shows that the pointwise infimum h of f and g belongs to \mathcal{H}_s^* . \square

Since the super duals of L are subsets of \mathbb{R}^{L_+} , they inherit the natural pointwise order of \mathbb{R}^{L_+} . That is, for $f, g: L_+ \rightarrow \mathbb{R}$ we write $f \geq g$ (and say that f dominates g) whenever $f(x) \geq g(x)$ holds for each $x \in L_+$. We have the following characterization of domination amongst positively homogeneous functions.

Lemma 3.7. *Assume that $f, g: L_+ \rightarrow \mathbb{R}_+$ are positively homogeneous functions. If $f(x_0) > 0$ holds for some $x_0 \in L_+$, then the following statements are equivalent.*

1. *The function g dominates f , i.e., $g(y) \geq f(y)$ holds for each $y \in L_+$.*
2. *For each $y \in L_+$ satisfying $f(y) = f(x_0)$ we have $g(y) \geq f(x_0)$, i.e.,*

$$\{y \in L_+ : f(y) = f(x_0)\} \subseteq \{y \in L_+ : g(y) \geq f(x_0)\}.$$

Proof. (1) \implies (2) If $g \geq f$ and a vector $y \in L_+$ satisfies $f(y) = f(x_0)$, then $g(y) \geq f(y) = f(x_0)$.

(2) \implies (1) Let $y \in L_+$. If $f(y) = 0$, then $g(y) \geq f(y)$ is trivially true. Therefore, we can assume that $f(y) > 0$. In this case, $f(\frac{f(x_0)}{f(y)}y) = f(x_0)$, and so our hypothesis implies $\frac{f(x_0)}{f(y)}g(y) = g(\frac{f(x_0)}{f(y)}y) \geq f(x_0)$ or $g(y) \geq f(y)$. So, $g(y) \geq f(y)$ holds for each $y \in L_+$, i.e., g dominates f . \square

4. LATTICE PROPERTIES OF THE SUPER DUALS

We start by introducing some notation. If f_1, \dots, f_m are m functions in \mathcal{H}_s^\sim , then we shall denote the supremum in \mathcal{H}_s^\sim (if it exists) of the set $\{0, f_1, \dots, f_m\}$ by $[\bigvee_{i=1}^m f_i]^+$, i.e., we shall write

$$\left[\bigvee_{i=1}^m f_i\right]^+ = \sup\{0, f_1, \dots, f_m\}.$$

In particular, if $f \in \mathcal{H}_s^\sim$, then we have $f^+ = f \vee 0 = \sup\{0, f\}$.

We are now ready to introduce the notion of a Riesz–Kantorovich functional.

Definition 4.1. For each $i = 1, 2, \dots, m$ let $f_i: L_+ \rightarrow (-\infty, \infty]$ be an extended real-valued function. Then the **Riesz–Kantorovich functional** of the m -tuple $f = (f_1, f_2, \dots, f_m)$ is the extended real-valued function $\mathcal{R}_f: L_+ \rightarrow (-\infty, \infty]$ defined by

$$\mathcal{R}_f(x) = \sup\left\{\sum_{i=1}^m f_i(y_i) : y_i \in [0, x] \text{ for each } i \text{ and } \sum_{i=1}^m y_i \leq x\right\}.$$

It should be immediate that if each f_i belongs to the super order dual \mathcal{H}_s^\sim , then the Riesz–Kantorovich functional \mathcal{R}_f is a positive real-valued function. In addition, it should be noticed that if one of the functions f_i is monotone, then

$$\mathcal{R}_f(x) = \sup\left\{\sum_{i=1}^m f_i(y_i) : y_i \in [0, x] \text{ for each } i \text{ and } \sum_{i=1}^m y_i = x\right\}.$$

In particular, if $f \in \mathcal{H}_s^+$, then Lemma 3.5 implies that

$$\mathcal{R}_f(x) = \sup_{0 \leq y \leq x} f(y) = f(x)$$

for each $x \in L_+$. That is, $\mathcal{R}_f = f$ for each $f \in \mathcal{H}_s^+$.

The Riesz–Kantorovich formula of an m -tuple (f_1, \dots, f_m) of functions in \mathcal{H}_s^\sim is an element of \mathcal{H}_s^+ . In fact, it is the supremum in \mathcal{H}_s^\sim of the set $\{0, f_1, \dots, f_m\}$.

Lemma 4.2. *The Riesz–Kantorovich functional of an m -tuple $f = (f_1, f_2, \dots, f_m)$ of functions in \mathcal{H}_s^\sim is the supremum of the set $\{0, f_1, \dots, f_m\}$ in \mathcal{H}_s^\sim . That is, the least upper bound $[\bigvee_{i=1}^n f_i]^+$ of the set $\{0, f_1, \dots, f_m\}$ exists in \mathcal{H}_s^\sim and for each $x \in L_+$ we have*

$$\left[\bigvee_{i=1}^n f_i\right]^+(x) = \mathcal{R}_f(x) = \sup \left\{ \sum_{i=1}^m f_i(y_i) : y_i \in [0, x] \text{ for each } i \text{ and } \sum_{i=1}^m y_i \leq x \right\}.$$

Proof. Let $f = (f_1, f_2, \dots, f_m)$ be an m -tuple of functions in \mathcal{H}_s^\sim . We shall establish first that the Riesz–Kantorovich functional \mathcal{R}_f is positive, super-additive, and positively homogeneous. Start by observing that since each f_i is order upper bounded and satisfies $f_i(0) = 0$, the Riesz–Kantorovich functional \mathcal{R}_f takes values in \mathbb{R}_+ .

The positive homogeneity of \mathcal{R}_f is a consequence of the fact that each f_i is positively homogeneous and that for each $\alpha > 0$ and $x \in L_+$, a vector (y_1, \dots, y_m) in L_+^m satisfies $\sum_{i=1}^m y_i \leq x$ if and only if $\sum_{i=1}^m \alpha y_i \leq \alpha x$.

For the super-additivity of \mathcal{R}_f , fix $x, y \in L_+$, and let $(x_1, \dots, x_m) \in L_+^m$ satisfy $\sum_{i=1}^m x_i \leq x$ and $(y_1, \dots, y_m) \in L_+^m$ satisfy $\sum_{i=1}^m y_i \leq y$. Then, we have

$$(x_1 + y_1, \dots, x_m + y_m) \in L_+^m \quad \text{and} \quad \sum_{i=1}^m (x_i + y_i) \leq x + y.$$

Now the super-additivity of each f_i yields

$$\begin{aligned} & [f_1(x_1) + \dots + f_m(x_m)] + [f_1(y_1) + \dots + f_m(y_m)] \\ &= [f_1(x_1) + f_1(y_1)] + \dots + [f_m(x_m) + f_m(y_m)] \\ &\leq f_1(x_1 + y_1) + \dots + f_m(x_m + y_m) \\ &\leq \mathcal{R}_f(x + y). \end{aligned}$$

This easily implies $\mathcal{R}_f(x) + \mathcal{R}_f(y) \leq \mathcal{R}_f(x + y)$. Therefore, \mathcal{R}_f is super-additive, and thus $\mathcal{R}_f \in \mathcal{H}_s^+$.

Next, it should be obvious that $\mathcal{R}_f(x) \geq f_i(x)$ holds for all $x \in L_+$ and each i . That is, \mathcal{R}_f is an upper bound in \mathcal{H}_s^\sim for the set $\{0, f_1, \dots, f_m\}$. To see that \mathcal{R}_f is the least upper bound of the set $\{0, f_1, \dots, f_m\}$ in \mathcal{H}_s^\sim , assume that some $g \in \mathcal{H}_s^\sim$ satisfies $g \geq 0$ and $g \geq f_i$ for $i = 1, \dots, m$. If $x \in L_+$ and $(y_1, \dots, y_m) \in L_+^m$ satisfies $\sum_{i=1}^m y_i \leq x$, then, using the super-additivity and monotonicity of g (see Lemma 3.5), we obtain

$$\sum_{i=1}^m f_i(y_i) \leq \sum_{i=1}^m g(y_i) \leq g\left(\sum_{i=1}^m y_i\right) \leq g(x).$$

This clearly implies $\mathcal{R}_f(x) \leq g(x)$. Consequently, $\mathcal{R}_f(x) \leq g(x)$ for each $x \in L_+$, and thus $\mathcal{R}_f = \sup\{0, f_1, \dots, f_m\}$ in \mathcal{H}_s^\sim . \square

Remarkably, the supremum of any m -tuple of order bounded linear functionals in \mathcal{H}_s^\sim exists. This is a special case of the following result.

Theorem 4.3. *If f_1, \dots, f_m are functions in \mathcal{H}_s^\sim and at least one of them either is positive or belongs to L^\sim , then the supremum $\bigvee_{i=1}^m f_i$ exists in \mathcal{H}_s^\sim and is given by the Riesz–Kantorovich formula. That is, for each $x \in L_+$ we have*

$$(\star) \quad \left[\bigvee_{i=1}^m f_i \right](x) = \sup \left\{ \sum_{i=1}^m f_i(y_i) : y_i \in [0, x] \text{ for each } i \text{ and } \sum_{i=1}^m y_i = x \right\}.$$

Proof. By Lemma 4.2, we know that $\sup\{0, f_1, \dots, f_m\} = \mathcal{R}_f$ in \mathcal{H}_s^\sim . Now assume that f_j is positive for some j . Then, $\sup\{0, f_1, \dots, f_m\} = \sup\{f_1, \dots, f_m\}$ is clearly true, and therefore

$$\left[\bigvee_{i=1}^m f_i \right](x) = \sup \left\{ \sum_{i=1}^m f_i(y_i) : y_i \in [0, x] \text{ for each } i \text{ and } \sum_{i=1}^m y_i \leq x \right\}.$$

Now the validity of (\star) easily follows from the monotonicity of f_j .

Next, assume that $f_i \in L^\sim$ holds for some i ; we can suppose that $f_1 \in L^\sim$. Then, each $f_i - f_1$ belongs to \mathcal{H}_s^\sim . Let $g = (f_2 - f_1, f_3 - f_1, \dots, f_m - f_1)$ and consider the Riesz–Kantorovich functional \mathcal{R}_g . From Lemma 4.2, it follows that

$$\mathcal{R}_g = \left[\bigvee_{i=2}^m (f_i - f_1) \right]^+.$$

Since \mathcal{H}_s^\sim is a wedge, we have $\mathcal{R}_g + f_1 \in \mathcal{H}_s^\sim$. Clearly, $\mathcal{R}_g + f_1 \geq f_1$. Now let $x \in L_+$, and note that for each $2 \leq j \leq m$ we have

$$\begin{aligned} \mathcal{R}_g(x) + f_1(x) &= \sup \left\{ \sum_{i=2}^m (f_i - f_1)(y_i) : y \in [0, x]^{m-1} \text{ and } \sum_{i=2}^m y_i \leq x \right\} + f_1(x) \\ &\geq (f_j - f_1)(x) + f_1(x) = f_j(x). \end{aligned}$$

Therefore, $\mathcal{R}_g + f_1$ is an upper bound of the set $\{f_1, \dots, f_m\}$ in \mathcal{H}_s^\sim .

Now we shall prove that $\mathcal{R}_g + f_1$ is the least upper bound of the set $\{f_1, \dots, f_m\}$ in \mathcal{H}_s^\sim . To this end, suppose that some $h \in \mathcal{H}_s^\sim$ dominates each f_i . Once again, $h - f_1 \in \mathcal{H}_s^\sim$. Furthermore, $h - f_1 \geq 0$ and $h - f_1$ dominates $f_i - f_1$ for $2 \leq i \leq m$. Therefore, by Lemma 4.2, $\mathcal{R}_g \leq h - f_1$, and so $\mathcal{R}_g + f_1 \leq h$. This shows that $\mathcal{R}_g + f_1$ is the least upper bound in \mathcal{H}_s^\sim of the set $\{f_1, \dots, f_m\}$.

Finally, it easily follows from the additivity of f_1 that for each $x \in L_+$ we have

$$\left[\bigvee_{i=1}^m f_i \right](x) = (\mathcal{R}_g + f_1)(x) = \left\{ \sum_{i=1}^m f_i(y_i) : y_i \in [0, x] \text{ for each } i \text{ and } \sum_{i=1}^m y_i = x \right\}.$$

This completes the proof. \square

Corollary 4.4. *The set \mathcal{H}_s^+ equipped with the pointwise ordering is a lattice. Moreover, for any pair $f_1, f_2 \in \mathcal{H}_s^+$ and any $x \in L_+$ we have*

$$\begin{aligned} [f_1 \vee f_2](x) &= \sup\{f_1(y_1) + f_2(y_2) : y_1, y_2 \in L_+ \text{ and } y_1 + y_2 = x\}, \\ [f_1 \wedge f_2](x) &= \min\{f_1(x), f_2(x)\}. \end{aligned}$$

It should be noticed that the finite suprema in \mathcal{H}_s^+ need not coincide with the pointwise suprema in \mathbb{R}^{L+} . For instance, if we let $L = \mathbb{R}^2$ and $L_+ = \mathbb{R}_+^2$, and consider the “linear” functions $f_1, f_2 \in \mathcal{H}_s^+$ defined by

$$f_1(x, y) = x \quad \text{and} \quad f_2(x, y) = y,$$

then $\mathcal{R}_{(f_1, f_2)}(x, y) = x + y$, while the pointwise supremum of f_1 and f_2 in \mathbb{R}^{L+} satisfies

$$\max\{f_1(x, y), f_2(x, y)\} = \max\{x, y\},$$

which is different than $\mathcal{R}_{(f_1, f_2)}$. This phenomenon is analogous to the idea of a lattice-subspace; see [26, 27]. Moreover, notice that Theorem 4.3 *does not* tell us that the vector space $\mathcal{H}_s^+ - \mathcal{H}_s^+$ is a vector lattice under the partial ordering induced by \mathcal{H}_s^+ . In fact, $(\mathcal{H}_s^+ - \mathcal{H}_s^+) \cap \mathbb{R}^{L+} \neq \mathcal{H}_s^+$, and the order induced by the cone \mathcal{H}_s^+ is not the pointwise ordering of \mathbb{R}^{L+} .

5. EMBEDDING DUALS IN THE SUPER ORDER DUAL

Having established that the super dual cone is a lattice, we are now interested in the circumstances under which the canonical embedding of the duals in the super dual preserves suprema. To this end, assume from now to the end of the paper that L is also an *ordered topological vector space*. That is, L is also equipped with a Hausdorff locally convex topology τ for which L_+ is τ -closed and every order interval is τ -bounded.

As usual, L' denotes the topological dual of (L, τ) and L'_+ denotes the collection of all positive τ -continuous linear functionals on L , i.e.,

$$L'_+ = \{f \in L' : f(x) \geq 0 \text{ holds for each } x \in L_+\}.$$

Since we assume that the cone L_+ is generating, it follows that L'_+ is a $\sigma(L', L)$ -closed cone of L' . Moreover, since the order intervals of L are τ -bounded, it should be clear that $L' \subseteq L^\sim$.

We now isolate and name the collection of all upper τ -semicontinuous functions⁷ of \mathcal{H}_s^* .

Definition 5.1. The **super topological dual** \mathcal{H}'_s of L is the collection of all super-additive, positively homogeneous, and upper τ -semicontinuous functions from L_+ to \mathbb{R} . That is,

$$\mathcal{H}'_s = \{f \in \mathcal{H}_s^* : f \text{ is upper } \tau\text{-semicontinuous}\}.$$

The subset of all positive functions in \mathcal{H}'_s will be denoted by $(\mathcal{H}'_s)_+$.

The super topological dual \mathcal{H}'_s is closed under finite pointwise infima. Indeed, since for any family of functions $\{f_i\}_{i \in I}$, where $f_i : X \rightarrow [-\infty, \infty]$, and any $c \in \mathbb{R}$ we have

$$\{x \in X : \inf_{i \in I} f_i(x) \geq c\} = \bigcap_{i \in I} \{x \in X : f_i(x) \geq c\},$$

⁷ Recall that a function $f : X \rightarrow \mathbb{R}$ on a topological space is said to be *upper semicontinuous* if for each $c \in \mathbb{R}$ the set $\{x \in L_+ : f(x) \geq c\}$ is closed. It is well known that a function $f : X \rightarrow \mathbb{R}$ is upper τ -semicontinuous if and only if $x_\alpha \rightarrow x$ in X implies $\limsup_\alpha f(x_\alpha) \leq f(x)$; see [7, Lemma 2.39, p. 43].

it follows that the pointwise infimum of a family of upper τ -semicontinuous functions is an upper τ -semicontinuous function—which is also super-additive and positively homogeneous if each f_i is likewise so.

The next result characterizes the functions in $(\mathcal{H}'_s)_+$ that are upper τ -semicontinuous.

Lemma 5.2. *A function $f \in \mathcal{H}_s^+$ is upper τ -semicontinuous, i.e., $f \in (\mathcal{H}'_s)_+$, if and only if for each $x \in L_+$ we have*

$$f(x) = \inf\{h(x) : h \in L' \text{ and } h \geq f\}.$$

Proof. Fix a function $f \in \mathcal{H}_s^+$. If for each $x \in L_+$ we have

$$f(x) = \inf\{h(x) : h \in L' \text{ and } h \geq f\},$$

then f is clearly upper τ -semicontinuous; see the discussion following Definition 5.1.

For the converse, suppose that f is upper τ -semicontinuous. Let $x_0 \in L_+$ and fix $\epsilon > 0$. We must establish that there exists some $h \in L'$ satisfying $h \geq f$ and $f(x_0) \leq h(x_0) < f(x_0) + \epsilon$.

To this end, assume first that $f(x_0) > 0$. Put $\alpha = \frac{f(x_0)}{f(x_0) + \epsilon}$ and notice that $0 < \alpha < 1$. By the positive homogeneity of f we have $f(\alpha x_0) = \alpha f(x_0) < f(x_0)$. That is, $\alpha x_0 \notin U_f^{x_0}$, where $U_f^{x_0}$ is the upper contour set of f at x_0 , i.e., $U_f^{x_0} = \{x \in L_+ : f(x) \geq f(x_0)\}$. Since $U_f^{x_0}$ is τ -closed and convex, there exist (by the Separation Theorem) a τ -continuous linear functional g and a real number η satisfying $g(\alpha x_0) < \eta$ and $g(x) > \eta$ for each $x \in U_f^{x_0}$. The comprehensiveness of the set $U_f^{x_0}$ (see Corollary 3.7) guarantees that g is also a positive linear functional. This implies that $\eta > g(\alpha x_0) \geq 0$. Letting $h = \frac{f(x_0)}{\eta}g \in L'$, we note that $h(y) = \frac{f(x_0)}{\eta}g(y) > f(x_0)$ for each $y \in U_f^{x_0}$. By Lemma 3.7 we get $h \geq f$. Finally, notice that $h(\alpha x_0) = \frac{f(x_0)}{\eta}g(\alpha x_0) < f(x_0)$ implies $h(x_0) < \frac{f(x_0)}{\alpha} = f(x_0) + \epsilon$. This completes the proof when $f(x_0) > 0$.

We now consider the case $f(x_0) = 0$. If $f = 0$, then f is linear and the conclusion holds trivially. Thus, it remains to consider the case where $f(y) > 0$ for some $y \in L_+$. For each $0 < \alpha < 1$ let $z_\alpha = \alpha x_0 + (1 - \alpha)y$. Since f is super-additive and positively homogeneous, it must be the case that $f(z_\alpha) > 0$ for each $0 < \alpha < 1$. From $\lim_{\alpha \uparrow 1} z_\alpha = x_0$ and the upper τ -semicontinuity of f , it follows that $0 \leq \limsup_{\alpha \uparrow 1} f(z_\alpha) \leq f(x_0) = 0$. This implies $\lim_{\alpha \uparrow 1} f(z_\alpha) = 0$. In particular, there exists some $\frac{1}{2} \leq \alpha_0 < 1$ such that $0 < f(z_\alpha) < \frac{\epsilon}{2}$ for all $\alpha_0 \leq \alpha < 1$. According to the preceding case, for each $\alpha_0 \leq \alpha < 1$ there exists some $h_\alpha \in L'$ satisfying $h_\alpha \geq f \geq 0$ and $0 < f(z_\alpha) \leq h_\alpha(z_\alpha) < \frac{\epsilon}{2}$. Now observe that for each $\alpha_0 \leq \alpha < 1$ we have

$$\frac{1}{2}h_\alpha(x_0) \leq \alpha h_\alpha(x_0) \leq \alpha h_\alpha(x_0) + (1 - \alpha)h_\alpha(y) = h_\alpha(z_\alpha) < \frac{\epsilon}{2},$$

and so $f(x_0) \leq h_\alpha(x_0) < \epsilon$ holds for all $\alpha_0 \leq \alpha < 1$, and the proof is finished. \square

The next result asserts that $\mathcal{H}'_s \subseteq \mathcal{H}_s^\sim$. That is, every super-additive, positively homogeneous, and upper τ -semicontinuous function from L_+ to \mathbb{R} is upper order bounded.⁸

⁸Note that since for each function $f \in \mathcal{H}_s^*$ the sets $\{x \in L_+ : f(x) \geq c\}$ are all convex, it follows that if f is upper τ -semicontinuous, then f is also upper semicontinuous for every locally convex topology on L that is consistent with the dual pair $\langle L, L' \rangle$.

Lemma 5.3. *Every function in \mathcal{H}'_s carries τ -bounded subsets of L to bounded from above subsets in \mathbb{R} . In particular, we have $\mathcal{H}'_s \subseteq \mathcal{H}^\sim_s$.*

Proof. Suppose by way of contradiction that some $f \in \mathcal{H}'_s$ carries a τ -bounded subset B of L to an unbounded from above subset of \mathbb{R} . This means that there exists a sequence $\{x_n\} \subseteq B$ such that $f(x_n) > n$ holds for every n . Since B is τ -bounded, we have $\frac{1}{n}x_n \xrightarrow{\tau} 0$. But then, the upper τ -semicontinuity and positive homogeneity of f imply

$$0 = f(0) \geq \limsup_{n \rightarrow \infty} f\left(\frac{1}{n}x_n\right) = \limsup_{n \rightarrow \infty} \frac{1}{n}f(x_n) \geq 1,$$

which is impossible. □

There is a natural mapping $J: L^* \rightarrow \mathbb{R}^{L_+}$ defined by

$$J(f)(x) = (f|_{L_+})(x) = f(x)$$

for all $x \in L_+$. That is, $J(f) = f|_{L_+}$, the restriction of f to L_+ . It is easy to see that this is a linear mapping that preserves the order, i.e., $f \leq g$ in L^* implies $J(f) \leq J(g)$. In other words, we have the following.

Lemma 5.4. *The mapping $J: L^* \rightarrow \mathbb{R}^{L_+}$ is an injective (i.e., one-to-one) positive operator. Moreover, J maps L^* into \mathcal{H}^*_s , L^*_+ into $\mathcal{H}^+_{s,+}$, and L' and L^\sim into \mathcal{H}^\sim_s .*

It is easy to see that $(\mathcal{H}'_s)_+ \subseteq \mathcal{H}^+_{s,+}$ and that the operator $J: L^* \rightarrow \mathbb{R}^{L_+}$ carries L' into \mathcal{H}'_s . We are now in a position to show that the embedding $J: L' \rightarrow \mathcal{H}'_s$ also preserves finite suprema. Notice that we have yet to establish sufficient conditions for the suprema in \mathcal{H}'_s to coincide with suprema in \mathcal{H}^\sim_s , i.e., the suprema in \mathcal{H}'_s need not satisfy the Riesz–Kantorovich formula.

Theorem 5.5 (The Embedding Theorem). *If the supremum $\bigvee_{i=1}^m f_i$ of the continuous linear functionals f_1, f_2, \dots, f_m on L exists in L' , then $\bigvee_{i=1}^m J(f_i)$ exists in \mathcal{H}'_s and*

$$J\left(\bigvee_{i=1}^m f_i\right) = \bigvee_{i=1}^m J(f_i).$$

That is, the embedding $J: L' \rightarrow \mathcal{H}'_s$ preserves finite suprema.

Proof. Let $f_1, \dots, f_m \in L'$ and assume that $g = \bigvee_{i=1}^m f_i$ exists in L' . Identifying each $f \in L'$ with its image $J(f) = f|_{L_+}$ in \mathcal{H}'_s , what needs to be shown is that if $h \in \mathcal{H}'_s$ satisfies $h \geq f_i$ for each $i = 1, \dots, m$, then $h \geq g$. So, assume that some function $h \in \mathcal{H}'_s$ satisfies $h \geq f_i$ for each i .

Replacing each f_i with $f_i - f_1$, h with $h - f_1 \in (\mathcal{H}'_s)_+$, and g with $g - f_1 = \bigvee_{i=1}^m (f_i - f_1) \in L_+$, we can assume without loss of generality that $f_1 = 0$. Clearly, $g \geq 0$ and $h \geq 0$.

Fix a vector $x \in L_+$. From Lemma 5.2, for each $\epsilon > 0$ there exists some $\phi_\epsilon \in L'$ satisfying $\phi_\epsilon \geq h$ and $\phi_\epsilon(x) < h(x) + \epsilon$. Clearly, $\phi_\epsilon \geq f_i$ for each i . Since g is the supremum of the set $\{f_1, \dots, f_m\}$ in L' , we infer that $\phi_\epsilon \geq g$. Therefore, $g(x) \leq \phi_\epsilon(x) < h(x) + \epsilon$. This implies $g(x) < h(x) + \epsilon$ for every $\epsilon > 0$. Consequently, $g(x) \leq h(x)$ for each $x \in L_+$, as desired. □

6. THE EMBEDDING AND THE RIESZ–KANTOROVICH FORMULA

In this section, we turn our attention to the *algebraic dual* $(L')^*$ of L' . Recall that every vector $x \in L$ can be considered as a vector in $(L')^*$ by letting $x(h) = h(x)$ for each $h \in L'$. Also, if $x, y \in (L')^*$, then $x \geq y$ means that $x(h) \geq y(h)$ for all $h \in L'_+$. Thus, for each $x \in L_+$, besides the order interval $[0, x]$ in L , there is another “order interval” associated with x in $(L')^*$. This is the order interval

$$\llbracket 0, x \rrbracket = \{h \in (L')^*: 0 \leq h \leq x\}.$$

Clearly $[0, x] \subseteq \llbracket 0, x \rrbracket$ holds for each $x \in L_+$.⁹

Now, if $f = (f_1, f_2, \dots, f_m)$ is an m -tuple of τ -continuous linear functionals on L , i.e., $f_i \in L'$ for each i , then we can define another *generalized Riesz–Kantorovich function* $\mathcal{R}_f^*: L_+ \rightarrow [0, \infty]$ via the formula

$$\mathcal{R}_f^*(x) = \sup \left\{ \sum_{i=1}^m h_i(f_i) : h_i \in \llbracket 0, x \rrbracket \text{ for each } i \text{ and } \sum_{i=1}^m h_i \leq x \right\}.$$

Clearly, \mathcal{R}_f^* is positive, super-additive, and positively homogeneous. Therefore, if \mathcal{R}_f^* is also real valued, then $\mathcal{R}_f^* \in \mathcal{H}_s^+$.

Lemma 6.1. *If $f = (f_1, f_2, \dots, f_m)$ is an m -tuple of τ -continuous linear functionals on L , then for each $x \in L_+$ we have $\mathcal{R}_f(x) \leq \mathcal{R}_f^*(x)$. Moreover, if some $g \in L'_+$ satisfies $g \geq f_i$ for each i , then*

$$\mathcal{R}_f(x) \leq \mathcal{R}_f^*(x) \leq g(x)$$

for every $x \in L_+$ (and hence, in this case, $\mathcal{R}_f^* \in \mathcal{H}_s^+$).

Proof. Fix $x \in L_+$ and let $g \in L'_+$ satisfy $g \geq f_i$ for each i . The inequality $\mathcal{R}_f(x) \leq \mathcal{R}_f^*(x)$ follows immediately from the inclusion $[0, x] \subseteq \llbracket 0, x \rrbracket$. Now assume that the vector (h_1, \dots, h_m) satisfies $h_i \in \llbracket 0, x \rrbracket$ for each i and $\sum_{i=1}^m h_i \leq x$. Since g dominates each f_i , we get

$$\sum_{i=1}^m h_i(f_i) \leq \sum_{i=1}^m h_i(g) = \left[\sum_{i=1}^m h_i \right](g) \leq x(g) = g(x) < \infty.$$

This implies that $\mathcal{R}_f^*(x) \leq g(x) < \infty$, and completes the proof. \square

The next result informs us when the finite supremum of an m -tuple of continuous linear functionals satisfies the Riesz–Kantorovich formula.

Theorem 6.2. *Assume that the supremum $g = [\bigvee_{i=1}^m f_i]^+$ of an m -tuple of τ -continuous linear functionals $f = (f_1, f_2, \dots, f_m)$ exists in L' . Then we have:*

1. $[\bigvee_{i=1}^m f_i]^+ = \mathcal{R}_f$ if and only if $\mathcal{R}_f \in \mathcal{H}'_s$.
2. $[\bigvee_{i=1}^m f_i]^+ = \mathcal{R}_f^*$ if and only if $\mathcal{R}_f^* \in \mathcal{H}'_s$.

Proof. (1) If $g(x) = \mathcal{R}_f(x)$ holds for all $x \in L_+$, then \mathcal{R}_f is τ -continuous on L_+ , and hence \mathcal{R}_f is also upper τ -semicontinuous. For the converse, assume that \mathcal{R}_f is upper τ -semicontinuous, that is, $\mathcal{R}_f \in \mathcal{H}'_s$. From the Embedding Theorem 5.5, we know that g is the supremum of the set $\{0, f_1, f_2, \dots, f_m\}$ in \mathcal{H}'_s . Also, from Lemma 4.2, we know that \mathcal{R}_f is the supremum of the set $\{0, f_1, f_2, \dots, f_m\}$ in \mathcal{H}'_s .

⁹ Note that the collection $(L')^*_+ = \{f \in (L')^*: f \geq 0\}$ of all positive linear functionals on L' is a wedge of $(L')^*$, which is not necessarily a cone. However, when the cone L'_+ is generating, $(L')^*_+$ is a cone in $(L')^*$.

Since $\mathcal{H}'_s \subseteq \mathcal{H}_s^\sim$ and \mathcal{R}_f is upper τ -semicontinuous, it follows that \mathcal{R}_f is also the supremum of the set $\{0, f_1, f_2, \dots, f_m\}$ in \mathcal{H}'_s . This implies $g = \mathcal{R}_f$.

(2) Assume first that \mathcal{R}_f^* is upper τ -semicontinuous. As in the proof of part (1) of Lemma 4.2, we see that $\mathcal{R}_f^* \in \mathcal{H}_s^+$. Moreover, our assumption implies that $\mathcal{R}_f^* \in (\mathcal{H}'_s)_+$. By the Embedding Theorem 5.5, it must be the case that $g \leq \mathcal{R}_f^*$. That is, $g(x) \leq \mathcal{R}_f^*(x)$ for all $x \in L_+$. Now a glance at Lemma 6.1 tells us that $g(x) = \mathcal{R}_f^*(x)$ for every $x \in L_+$. If, on the other hand, $g(x) = \mathcal{R}_f^*(x)$ for every $x \in L_+$, then clearly \mathcal{R}_f^* is τ -continuous—and hence it is also upper τ -semicontinuous. \square

Surprisingly, when \mathcal{R}_f is upper semicontinuous it coincides with \mathcal{R}_f^* .

Lemma 6.3. *If $f = (f_1, f_2, \dots, f_m)$ is an m -tuple of τ -continuous linear functionals on L and \mathcal{R}_f is upper τ -semicontinuous, then $\mathcal{R}_f = \mathcal{R}_f^*$.*

Proof. Take an arbitrary $x \in L_+$. We claim that $\mathcal{R}_f^*(x) = \mathcal{R}_f(x)$. To see this, suppose by way of contradiction that $\mathcal{R}_f^*(x) > \mathcal{R}_f(x)$. By Lemma 5.2, there exists some $g \in L'_+$ satisfying $g \geq \mathcal{R}_f$ and $\mathcal{R}_f^*(x) > g(x)$. However, since $g \geq f_i$ holds for each i , it follows from Lemma 6.1 that $\mathcal{R}_f^*(x) \leq g(x)$, which is a contradiction. Therefore, $\mathcal{R}_f^*(x) = \mathcal{R}_f(x)$. \square

Recall that if L is a locally convex-solid Riesz space, then the order intervals in L are $\sigma(L'', L')$ -dense in the order intervals of L'' (see [11, Lemma 11.12, p. 169]). An analogous property holds whenever the Riesz–Kantorovich functionals of functions in L' are upper τ -semicontinuous.

Theorem 6.4. *If \mathcal{R}_f is upper τ -semicontinuous for every $f \in L'$, then each order interval $[0, x]$ is $\sigma((L')^*, L')$ -dense in $\llbracket 0, x \rrbracket$.*

Proof. Fix $x_0 \in L_+$ and suppose by way of contradiction that there exists some $z_0 \in \llbracket 0, x_0 \rrbracket$ satisfying $z_0 \notin \overline{\llbracket 0, x_0 \rrbracket}$, where the bar denotes $\sigma((L')^*, L')$ -closure. Since $\sigma((L')^*, L')$ is locally convex, it follows from the Separation Theorem that there exist a linear functional $f \in L'$ and a constant c such that $f(z_0) > c$ and $f(x) < c$ for all $x \in \llbracket 0, x_0 \rrbracket$. Therefore, $\mathcal{R}_f(x_0) \leq c$. On the other hand, by Lemma 6.3 we have $\mathcal{R}_f(x_0) = \mathcal{R}_f^*(x_0) \geq f(z_0) > c > 0$, a contradiction. This contradiction establishes that $\llbracket 0, x \rrbracket$ is $\sigma((L')^*, L')$ -dense in $\llbracket 0, x \rrbracket$. \square

We close this section by showing that the continuity of the Riesz–Kantorovich functionals at zero is equivalent to the identity $L^r = L^\sim$.

Theorem 6.5. *Every order bounded linear functional is regular (i.e., $L^r = L^\sim$) if and only if for every $f \in (L^\sim)^m$ its Riesz–Kantorovich functional \mathcal{R}_f is continuous at zero for some Hausdorff locally convex topology on L .*

Proof. Assume first that $L^r = L^\sim$. From $L' \subseteq L^\sim$, we know that $\sigma(L, L^\sim)$ is a Hausdorff locally convex topology on L . Suppose by way of contradiction that for some m -tuple $f = (f_1, f_2, \dots, f_m)$ of linear functionals in L^\sim the Riesz–Kantorovich functional \mathcal{R}_f is not $\sigma(L, L^\sim)$ -continuous at zero. This means that there exist a net $\{x_\alpha\} \subseteq L_+$ and some $c > 0$ such that $x_\alpha \xrightarrow{\sigma(L, L^\sim)} 0$ while $\mathcal{R}_f(x_\alpha) \geq c > 0$ holds for every α . Since $L^r = L^\sim$, there exists some $g \in L_+^\sim \subseteq L^\sim$ that dominates f_1, f_2, \dots, f_m , and therefore $\mathcal{R}_f(x) \leq g(x)$ for every $x \in L_+$. This implies that $g(x_\alpha) \geq c > 0$ holds for each α . This contradicts the fact that $g \in L^\sim$ and $x_\alpha \xrightarrow{\sigma(L, L^\sim)} 0$.

For the converse, suppose that for every $f \in (L^\sim)^m$ the Riesz–Kantorovich functional \mathcal{R}_f is continuous at zero for some Hausdorff locally convex topology τ_f on L . Since $L^r \subseteq L^\sim$ is always true, it suffices to show that for any $f \in L^\sim$ there exists a positive linear functional g such that $g \geq f$. To this end, fix $f \in L^\sim$. If $f \leq 0$, then we are done. So, we can assume that there exists some $x \in L_+$ such that $f(x) > 0$. Clearly, $\mathcal{R}_f(x) > 0$. Since \mathcal{R}_f is τ_f -continuous at zero, it follows that zero does not belong to the τ_f -closure of $U_{\mathcal{R}_f}^x = \{y \in L_+ : \mathcal{R}_f(y) \geq \mathcal{R}_f(x)\}$. By the Separation Theorem, there exist a non-zero τ_f -continuous linear functional h and a real number $c > 0$ such that $z \in U_{\mathcal{R}_f}^x$ implies $h(z) \geq c$. Since $U_{\mathcal{R}_f}^x$ is comprehensive, it must be the case that $h \geq 0$; and so $h \in L^\sim$. From $\frac{\mathcal{R}_f(x)}{c}h(y) \geq \mathcal{R}_f(x)$ for each $y \in U_{\mathcal{R}_f}^x$ and Lemma 3.7, it follows that the positive linear functional $\frac{\mathcal{R}_f(x)}{c}h$ dominates \mathcal{R}_f , and thus it also dominates f . \square

7. CONTINUITY OF THE RIESZ–KANTOROVICH FUNCTIONALS

Recall again that L is assumed to be equipped with a Hausdorff locally convex topology τ for which L_+ is τ -closed and every order interval is τ -bounded. In this section we shall also assume either that

- a. L_+ has a τ -interior point, or else that
- b. L is a complete metrizable topological vector space.

We have highlighted in the previous section the importance of the upper semi-continuity of the Riesz–Kantorovich functional. In this section, we shall investigate the continuity properties of functions in \mathcal{H}_s^+ and state some basic consequences of the embedding theorem.

We start with a generalization of a well known result from the theory of Banach lattices.

Lemma 7.1. *Every τ -convergent sequence in L has a relatively uniformly convergent subsequence. That is, if $x_n \xrightarrow{\tau} x$, then there exist a subsequence $\{u_n\}$ of $\{x_n\}$ and some $u \in L_+$ satisfying*

$$-\frac{1}{n}u \leq u_n - x \leq \frac{1}{n}u$$

for each n . In particular, every τ -convergent sequence of L has an order bounded subsequence.

Proof. Assume $x_n \xrightarrow{\tau} 0$ in L . Suppose first that e is an interior point of L_+ . In this case the conclusion follows from the fact that for each n the interval $[-\frac{1}{n}e, \frac{1}{n}e]$ is a neighborhood of zero.

Next, suppose that L is a Fréchet space. Choose a countable neighborhood base $\{U_n\}$ of zero consisting of τ -closed circled sets satisfying $U_{n+1} + U_{n+1} \subseteq U_n$ for each n . Let

$$V_n = U_n \cap L_+ - U_n \cap L_+.$$

By Lemma 2.1, the sets $\{V_n\}$ form a τ -neighborhood base at zero. So, there exists a subsequence $\{u_n\}$ of $\{x_n\}$ such that $u_n \in \frac{1}{n}V_n$ (or $nu_n \in V_n$) holds for each n . For each n pick $y_n, z_n \in U_n \cap L_+$ such that $nu_n = y_n - z_n$.

Now let $w_n = \sum_{i=1}^n y_i \in L_+$ and $v_n = \sum_{i=1}^n z_i \in L_+$. Clearly,

$$w_{n+p} - w_n \in U_{n+1} + U_{n+2} + \cdots + U_{n+p} \subseteq U_n$$

for each n . This implies that $\{w_n\}$ is a τ -Cauchy sequence and hence $\{w_n\}$ converges to some w . Since L_+ is closed, $w \in L_+$. Letting $p \rightarrow \infty$ in $w_{n+p} - w_n \in L_+$, we see that $w - w_n \in L_+$, and so $w \geq w_n \geq y_n$ for each n . Similarly, there exists some $v \in L_+$ such that $v_n \xrightarrow{\tau} v$ and $z_n \leq v$ for each n . If we let $u = w + v \in L_+$, then we have

$$-u \leq -v \leq -z_n \leq y_n - z_n = nu_n \leq y_n \leq w \leq u,$$

and so $\frac{1}{n}u \leq u_n \leq \frac{1}{n}u$ holds for each n , as desired. □

Our next result deals with the continuity at zero of the functions in \mathcal{H}_s^+ . It is well known that for a Fréchet lattice (i.e., a locally convex-solid Riesz space with a metrizable and complete topology) the identity $L' = L^r = L^\sim$ holds true; see [7, Theorem 8.11, p. 306].¹⁰ The next result informs us that we can do without the lattice property.

Theorem 7.2. *Each function in \mathcal{H}_s^+ is continuous at zero. In particular, we have $L' = L^r = L^\sim$.*

Proof. Fix a function $f \in \mathcal{H}_s^+$, and assume by way of contradiction that f is not τ -continuous at zero. This means that there exist a sequence $\{x_n\}$ of L_+ and some $\epsilon > 0$ satisfying $x_n \xrightarrow{\tau} 0$ and $f(x_n) \geq \epsilon > 0$ for each n . By Lemma 7.1, we can suppose (by passing to a subsequence if necessary) that $0 \leq x_n \leq \frac{1}{n}u$ holds for each n and some $u \in L_+$. But then, the monotonicity and homogeneity of f imply $\frac{1}{n}f(u) = f(\frac{1}{n}u) \geq f(x_n) \geq \epsilon > 0$ for every n , which is impossible. This contradiction establishes that f is continuous at zero.

The preceding conclusion implies that the Riesz–Kantorovich functional of any m -tuple of functionals in L^\sim is continuous at zero. So, by Theorem 6.5, we have $L^r = L^\sim$. Moreover, the above conclusion also shows that $L^r \subseteq L' \subseteq L^\sim$. Therefore, $L' = L^r = L^\sim$ holds true, as desired. □

Corollary 7.3. *The dual cone L'_+ generates the vector space L' , and the order intervals of $(L')^*$ are all $\sigma((L')^*, L')$ -compact.*

Proof. Theorem 7.2 implies $L^r = L'$. So, the cone L'_+ is generating in L' . Therefore, the vector space $(L')^*$ equipped with the topology $\sigma((L')^*, L')$ is an ordered topological vector space such that its order intervals are $\sigma((L')^*, L')$ -closed and $\sigma((L')^*, L')$ -bounded—and hence they are also $\sigma((L')^*, L')$ -compact. □

Sufficient conditions for the upper semicontinuity of the Riesz–Kantorovich functionals are presented in the next result.

Lemma 7.4. *For an m -tuple $f = (f_1, f_2, \dots, f_m)$ of order bounded linear functionals on L we have the following.*

1. *The generalized Riesz–Kantorovich functional \mathcal{R}_f^* is upper τ -semicontinuous.*
2. *If the order intervals in L are weakly compact, then:*
 - (a) $\mathcal{R}_f = \mathcal{R}_f^*$,
 - (b) *the Riesz–Kantorovich functional \mathcal{R}_f is upper τ -semicontinuous, and*
 - (c) $[0, x] = \llbracket 0, x \rrbracket$ *holds for each $x \in L_+$.*

¹⁰ It is also true that a given Riesz space admits at most one complete metrizable locally solid topology; see [7, Corollary 8.10, p. 306]. In particular, a Riesz space admits at most one (up to an equivalence) lattice norm under which it is a Banach lattice.

Proof. Notice first that $L^\sim = L'$. Therefore each f_i is τ -continuous.

(1) Take a sequence $\{x_n\} \subseteq L_+$ satisfying $x_n \xrightarrow{\tau} x$ for some $x \in L_+$, and suppose that $\mathcal{R}_f^*(x_n) \geq c$ for all n . We need to show that $\mathcal{R}_f^*(x) \geq c$.

From Lemma 7.1, we can assume that $\{x_n\}$ is order bounded. That is, we can assume that there exists some $e \in L_+$ such that $\{x_n\} \subseteq [0, e] \subseteq \llbracket 0, e \rrbracket$. Fix $\epsilon > 0$ and for each n pick some m -tuple $(z_1^n, \dots, z_m^n) \in \llbracket 0, e \rrbracket^m$ satisfying $\sum_{i=1}^m z_i^n \leq x_n$ and

$$(*) \quad \sum_{i=1}^m f_i(z_i^n) > \mathcal{R}_f^*(x_n) - \epsilon \geq c - \epsilon.$$

Now, according to Corollary 7.3 the order interval $\llbracket 0, e \rrbracket$ is $\sigma((L')^*, L')$ -compact. So, there exists a subnet $\{(z_1^{n_\alpha}, \dots, z_m^{n_\alpha})\}$ of the sequence $\{(z_1^n, \dots, z_m^n)\}$ such that $z_i^{n_\alpha} \xrightarrow{\sigma((L')^*, L')} z_i$ holds in $(L')^*$ for each i . Clearly, $z_i \in \llbracket 0, e \rrbracket$ for each i , and $\sum_{i=1}^m z_i \leq x$.

Now using $(*)$ and the $\sigma((L')^*, L')$ -continuity of each f_i , we see that

$$c - \epsilon \leq \limsup_{\alpha} \left[\sum_{i=1}^m f_i(z_i^{n_\alpha}) \right] \leq \sum_{i=1}^m \limsup_{\alpha} f_i(z_i^{n_\alpha}) = \sum_{i=1}^m f_i(z_i) \leq \mathcal{R}_f^*(x).$$

Since $\epsilon > 0$ is arbitrary, we get $\mathcal{R}_f^*(x) \geq c$, as desired. This completes the proof.

(2) In this case, we can modify the preceding proof to show that \mathcal{R}_f is also upper τ -continuous. The rest follows from Lemma 6.3 and Theorem 6.4. \square

The next result is a simple consequence of Lemma 7.4 and Theorem 6.2.

Theorem 7.5. *If the supremum $g = [\bigvee_{i=1}^m f_i]^+$ of an m -tuple of order bounded linear functionals $f = (f_1, f_2, \dots, f_m)$ exists in L^\sim , then for each $x \in L_+$ we have*

$$\left[\bigvee_{i=1}^m f_i \right]^+(x) = \sup \left\{ \sum_{i=1}^m f_i(y_i) : y_i \in \llbracket 0, x \rrbracket \text{ for each } i \text{ and } \sum_{i=1}^m y_i \leq x \right\}.$$

If, in addition, the order intervals in L are weakly compact, then

$$\left[\bigvee_{i=1}^m f_i \right]^+(x) = \sup \left\{ \sum_{i=1}^m f_i(y_i) : y_i \in [0, x] \text{ for each } i \text{ and } \sum_{i=1}^m y_i \leq x \right\}.$$

for each $x \in L_+$.

We close this section with a result that has many implications in economic theory.

Theorem 7.6. *If $f = (f_1, f_2, \dots, f_m)$ is an m -tuple of order bounded linear functionals, then for each $x \in L_+$ we have*

$$\inf \{g(x) : g \in L^\sim \text{ and } g \geq f_i \text{ for each } 1 \leq i \leq m\} = \mathcal{R}_f^*(x).$$

If, in addition, the order intervals in L are weakly compact, then

$$\inf \{g(x) : g \in L^\sim \text{ and } g \geq f_i \text{ for each } 1 \leq i \leq m\} = \mathcal{R}_f(x).$$

Proof. By Lemma 7.4 we know that \mathcal{R}_f^* is upper τ -semicontinuous. Therefore, by Lemmas 5.2 and 7.2 we have

$$\mathcal{R}_f^*(x) = \inf \{g(x) : g \in L^\sim \text{ and } g \geq \mathcal{R}_f^*\},$$

for each $x \in L_+$. Furthermore, Lemma 6.1 guarantees that some $g \in L^\sim$ satisfies $g \geq \mathcal{R}_f^*$ if and only if $g \geq f_i$ for each i . This concludes the first part of the theorem.

Now if the order intervals in L are weakly compact, then by Lemma 7.4(2) we have $\mathcal{R}_f = \mathcal{R}_f^*$, and the proof is finished. \square

8. EXAMPLES

Recall that the *half-ray* generated by a vector x is the set $\mathcal{L}_x = \{\alpha x : \alpha \geq 0\}$. A vector $x > 0$ is called *extremal* for the cone L_+ if $[0, x] = \{\alpha x : 0 \leq \alpha \leq 1\}$, i.e., whenever the order interval $[0, x]$ coincides with the line segment joining 0 and x . A half-ray generated by an extremal vector is called an *extremal ray* of the cone L_+ . Clearly, if x and y are extremal vectors, then either x and y are linearly independent, or else $\mathcal{L}_x = \mathcal{L}_y$.

Lemma 8.1. *In an ordered vector space any three extremal vectors that generate three distinct extremal rays are linearly independent.*

Proof. Let x_1, x_2 , and x_3 be three extremal vectors in the cone of an ordered vector space that generate three distinct extremal rays. Assume by way of contradiction that the vectors x_1, x_2, x_3 are linearly dependent. Relabeling, we can suppose that there exist scalars λ and μ such that $x_1 = \lambda x_2 + \mu x_3$.

If $\lambda = 0$, then $x_1 = \mu x_3$ and so $\mu \neq 0$. This implies $\mu > 0$ and $\mathcal{L}_{x_1} = \mathcal{L}_{x_3}$, a contradiction. Hence, $\lambda \neq 0$ and $\mu \neq 0$. Obviously, the case $\lambda < 0$ and $\mu < 0$ cannot happen. If $\lambda > 0$ and $\mu > 0$, then from $0 < \lambda x_2 \leq x_1$, it follows that $\lambda x_2 = \alpha x_1$ for some $\alpha > 0$. This implies $\mathcal{L}_{x_2} = \mathcal{L}_{x_1}$, which is a contradiction. Finally, we consider the case $\lambda > 0$ and $\mu < 0$. In this case, we have $x_1 < \lambda x_2$, and hence for some $\beta > 0$ we must have $x_1 = \beta \lambda x_2$. This implies $\mathcal{L}_{x_1} = \mathcal{L}_{x_2}$, which is also a contradiction. Hence, the vectors x_1, x_2, x_3 must be linearly independent. \square

The conclusion of the preceding result is not true for four or more extremal vectors. That is, we have the following.

- *If $\mathcal{L}_{x_1}, \mathcal{L}_{x_2}, \dots, \mathcal{L}_{x_k}$ are distinct extremal rays of a closed cone in an ordered vector space L with $\dim L \geq k \geq 4$, then the vectors x_1, x_2, \dots, x_k need not be linearly independent.*

For an example, consider the closed (convex) cone L_+ in \mathbb{R}^4 generated by the four vectors

$$x_1 = (1, 0, 0, 0), \quad x_2 = (0, 1, 0, 0), \quad x_3 = (0, 0, 1, 0), \quad x_4 = (1, -1, 1, 0).$$

That is,

$$\begin{aligned} L_+ &= \left\{ \sum_{i=1}^4 \lambda_i x_i : \lambda_i \geq 0 \text{ for each } i \right\} \\ &= \{(\lambda_1 + \lambda_4, \lambda_2 - \lambda_4, \lambda_3 + \lambda_4, 0) : \lambda_i \geq 0 \text{ for each } i\}. \end{aligned}$$

It is a routine matter to verify that L_+ is indeed a cone and that the four half-rays $\mathcal{L}_{x_1}, \mathcal{L}_{x_2}, \mathcal{L}_{x_3}$, and \mathcal{L}_{x_4} are the only distinct extremal rays of the cone L_+ . However, notice that we have $(1, -1, 1, 0) = (1, 0, 0, 0) - (0, 1, 0, 0) + (0, 0, 1, 0)$.

Definition 8.2. A **base** for the cone L_+ is a non-empty convex subset of $L_+ \setminus \{0\}$ such that for every $x > 0$ there exist a scalar $\lambda > 0$ and some $b \in B$, both uniquely determined, such that $x = \lambda b$.

Given a base B for L_+ , it is easy to see that a vector $x \in B$ is an extreme point of B if and only if x an extremal vector of the cone L_+ . Regarding the existence

of extreme rays, we have the following result of V. Klee [21, 22]. (For a proof see [19, Theorem 3.12.8, p. 144 and Corollary 3.12.9, p. 145].)

Lemma 8.3 (Klee). *If L is finite dimensional and L_+ is closed, then L_+ has a compact base. In particular, L_+ coincides with the closed convex hull of its extremal vectors.*

Two vectors $x, y \in L$ are said to be *non-comparable* if $x \not\leq y$ and $y \not\leq x$. The next theorem describes an interesting lattice property of non-comparable vectors in \mathbb{R}^3 .

Theorem 8.4. *If L is three dimensional and L_+ is a closed generating cone, then there are two non-comparable points $x, y \in L$ for which $x \vee y$ exists in L if and only if L_+ has at most four extremal rays.*

Proof. It is well known that in a Hausdorff finite dimensional ordered topological vector space with a closed generating cone the order intervals are compact—for a proof see, for instance, [16, §5.4]. This implies that in our case we have $L' = L^\sim$.

It is also well known that the positive cone L_+ has at most four extremal rays if and only if L'_+ also has at most four extremal rays, and that $L_+ = L''_+$. Therefore, in order to prove the theorem we need only to show that there are two order non-comparable functionals $f, g \in L'$ for which $f \vee g$ exists in L' if and only if L_+ has at most four extremal rays.

To this end, fix a compact base \mathcal{B} for L_+ and let \mathcal{E} be the set of all extreme points of \mathcal{B} . Therefore, aside from scalar multiples, the extremal rays of L_+ can be identified with the vectors of \mathcal{E} . Since L is three dimensional and L_+ is generating, it follows from Lemma 8.3 that $|\mathcal{E}| \geq 3$, where as usual $|S|$ denotes the cardinality of the set S .

Now assume that there is a functional $f \notin -L'_+$ and $f \notin L'_+$ for which f^+ exists. By Theorem 7.5 we know that f^+ satisfies the Riesz–Kantorovich formula. That is, for each $x \in L_+$ we have

$$f^+(x) = \sup \{f(y) : 0 \leq y \leq x\}.$$

We want to show that $|\mathcal{E}| \leq 4$. Let

$$\mathcal{E}_1 = \{x \in \mathcal{E} : f(x) \geq 0\} \quad \text{and} \quad \mathcal{E}_2 = \{x \in \mathcal{E} : f(x) < 0\}.$$

Since f^+ satisfies the Riesz–Kantorovich formula, it must be the case that $f^+(x) = f(x)$ for all $x \in \mathcal{E}_1$ and $f^+(x) = 0$ for all $x \in \mathcal{E}_2$. Now if $|\mathcal{E}_1| \geq 3$, then $f^+ = f$, since by Lemma 8.1 the set \mathcal{E}_1 spans L . This implies $f \geq 0$ or $f \in L'_+$, which is a contradiction. Therefore, $|\mathcal{E}_1| \leq 2$. Similarly, if $|\mathcal{E}_2| \geq 3$, then $f^+ = 0$ and $f \leq 0$ or $f \in -L'_+$, which is also a contradiction. Consequently, $|\mathcal{E}_2| \leq 2$ and hence $|\mathcal{E}| \leq 4$. We can also assume that \mathcal{E} is a subset of a plane. Indeed, if we choose some interior vector $p \in L'_+$, then we have $p \cdot x > 0$ for each $x > 0$, and so if for each $x \in \mathcal{E}$ we choose some $\lambda_x > 0$ such that $\lambda_x p \cdot x = 1$, then the convex hull of the set $\mathcal{E}' = \{\lambda_x x : x \in \mathcal{E}\}$ is a planar base of L_+ ; in fact the convex hull of \mathcal{E}' is either a triangle or a quadrilateral.

We shall now prove the converse. Thus, we assume that $|\mathcal{E}| \leq 4$. If $|\mathcal{E}| = 3$, then L is a vector lattice and the conclusion is trivially true. So, we can assume that $|\mathcal{E}| = 4$. That is, let $\mathcal{E} = \{x_1, x_2, x_3, x_4\}$. Therefore,

$$L_+ = \left\{ \sum_{i=1}^4 \lambda_i x_i : \lambda_i \geq 0 \text{ for each } i = 1, 2, 3, 4 \right\}.$$

By relabeling the x_i appropriately, it is not difficult to see that there exist two linear functionals g and h such that:

$$g(x_1) = g(x_2) = 0, \quad g(x_3) > 0, \quad g(x_4) > 0$$

and

$$h(x_3) = h(x_4) = 0, \quad h(x_1) < 0, \quad h(x_2) < 0.$$

Clearly, $g > 0$ and $h < 0$. Put $f = g + h$ and notice that $f \not\leq 0$ and $f \not\geq 0$. Moreover, an easy argument shows that

$$\begin{aligned} \mathcal{R}_f(x_1) &= g(x_1) = 0, & \mathcal{R}_f(x_2) &= g(x_2) = 0, \\ \mathcal{R}_f(x_3) &= f(x_3) = g(x_3) > 0, & \mathcal{R}_f(x_4) &= f(x_4) = g(x_4) > 0. \end{aligned}$$

So, if $z = \sum_{i=1}^4 \lambda_i x_i \in L_+$, then $g(z) = \sum_{i=1}^4 \lambda_i g(x_i) \geq \sum_{i=1}^4 \lambda_i f(x_i) = f(z)$, and thus $g \geq f$. But then, the super-additivity and homogeneity of \mathcal{R}_f for each $z = \sum_{i=1}^4 \lambda_i x_i \in L_+$ yields

$$g(z) \geq \mathcal{R}_f(z) \geq \sum_{i=1}^4 \lambda_i \mathcal{R}_f(x_i) = \sum_{i=1}^4 \lambda_i g(x_i) = g(z).$$

Consequently, $g(z) = \mathcal{R}_f(z)$ for every $z \in L_+$, and this shows that $g = f^+$. □

The next example illustrates Theorem 8.4.

Example 8.5. Consider the following four positive vectors in $L = \mathbb{R}^3$:

$$x_1 = (0, 1, 0), \quad x_2 = (1, 1, 0), \quad x_3 = (0, 1, 1), \quad x_4 = (1, 1, 1).$$

Let L_+ be the (convex) cone with vertex 0 generated by these points. That is,

$$L_+ = \left\{ \sum_{i=1}^4 \lambda_i x_i : \lambda_i \geq 0 \text{ for each } i = 1, 2, 3, 4 \right\}.$$

This cone (which is a closed subcone of the standard cone $L_+ = \mathbb{R}_+^3$) is not a lattice cone, but it is generating.

Now consider the vector $f = (2, -1, 0)$ as a linear functional on \mathbb{R}^3 . From

$$f \cdot x_1 = -1 < 0 \quad \text{and} \quad f \cdot x_2 = 1 > 0,$$

it follows that f and 0 are non-comparable linear functionals. Next, we claim that the linear functional $g = (1, 0, 0)$ is the least upper bound of f and 0, i.e., $g = f^+$ holds in $L' = \mathbb{R}^3$ with respect to the ordering induced by the dual cone of L'_+ .

To see this, fix an arbitrary point

$$z = \sum_{i=1}^4 \lambda_i x_i = (\lambda_2 + \lambda_4, \lambda_1 + \lambda_2 + \lambda_3 + \lambda_4, \lambda_3 + \lambda_4) \in L_+.$$

From $g - f = (-1, 1, 0)$, we see that

$$(g - f) \cdot z = (-1, 1, 0) \cdot (\lambda_2 + \lambda_4, \lambda_1 + \lambda_2 + \lambda_3 + \lambda_4, \lambda_3 + \lambda_4) = \lambda_1 + \lambda_3 \geq 0.$$

This shows that $g \geq f$ holds in L' . Moreover, $g \geq 0$ should be obvious.

To see that g is the least upper bound of f and 0, notice that $[0, x_i] = \{\alpha x_i : 0 \leq \alpha \leq 1\}$ holds for $i = 1, 2, 3, 4$. This implies

$$\begin{aligned} \mathcal{R}_f(x_1) &= g \cdot x_1 = 0, & \mathcal{R}_f(x_2) &= g \cdot x_2 = 1, \\ \mathcal{R}_f(x_3) &= g \cdot x_3 = 0, & \mathcal{R}_f(x_4) &= g \cdot x_4 = 1. \end{aligned}$$

So, if $z = \sum_{i=1}^4 \lambda_i x_i \in L_+$, then $g \geq f$, and the super-additivity and homogeneity of \mathcal{R}_f imply

$$g \cdot z \geq \mathcal{R}_f(z) \geq \sum_{i=1}^4 \lambda_i \mathcal{R}_f(x_i) = \sum_{i=1}^4 \lambda_i g \cdot x_i = g \cdot z.$$

Consequently, $g \cdot z = \mathcal{R}_f(z)$ for every $z \in L_+$, and so $g = f^+$ holds in L' .

There is another (geometrical) way of proving that $g = f^+$. Notice that the dual cone L'_+ of L_+ is the cone in \mathbb{R}^3 having the four extremal rays generated by the vectors

$$(1, 0, 0), (-1, 1, 0), (0, 0, 1), (0, 1, -1).$$

It is not difficult to see that $[(2, -1, 0) + L'_+] \cap L'_+ = (1, 0, 0) + L'_+$. This implies $g = f^+$. \square

Theorem 8.4 is not true for spaces of dimension greater than three.

Example 8.6. Let L_+ be an “ice-cream” cone in \mathbb{R}^3 . For instance, let

$$L_+ = \left\{ (x, y, z) \in \mathbb{R}^3 : z \geq \sqrt{x^2 + y^2} \right\}.$$

We consider \mathbb{R}^4 partially ordered by the cone $L_+ \times \mathbb{R}_+$. Notice that for any extremal ray \mathcal{L} of L_+ , the set $\mathcal{L} \times \{0\}$ is also an extremal ray of $L_+ \times \mathbb{R}_+$. This implies that the cone $L_+ \times \mathbb{R}_+$ has infinitely many extremal rays.

Now let h be a non-zero L_+ -negative functional on \mathbb{R}^3 and put $f = (h, 1)$. It is easy to see that $f \not\leq 0$ and $f \not\geq 0$. However, notice that $f^+ = (0, 1)$. \square

REFERENCES

1. Y. A. Abramovich, Injective envelopes of normed lattices, *Dokl. Akad. Nauk USSR* **197** (1971), 743–745. MR **44**:7257
2. Y. A. Abramovich, When each continuous operator is regular, *Functional Analysis, Optimization, and Mathematical Economics*, A memorial volume for L. V. Kantorovich, Oxford Univ. Press 1990, pp. 133–140. MR **91m**:47051
3. Y. A. Abramovich and V. A. Geiler, On a question of Fremlin concerning order bounded and regular operators, *Colloq. Math.* **46** (1982), 15–17. MR **84b**:46008
4. Y. A. Abramovich and A. W. Wickstead, Regular operators from and into a small Riesz space, *Indag. Math. N.S.* **2** (1991), 257–274. MR **93i**:47052
5. Y. A. Abramovich and A. W. Wickstead, The regularity of order bounded operators into $C(K)$, II, *Quart. J. Math. Oxford Ser. (2)* **44** (1993), 257–270. MR **94h**:47067
6. Y. A. Abramovich and A. W. Wickstead, When each continuous operator is regular, II, *Indag. Math.* **8** (1997), 281–294. MR **97c**:47046
7. C. D. Aliprantis and K. C. Border, *Infinite Dimensional Analysis: A Hitchhikers Guide*, 2nd Edition, Springer-Verlag, New York and Berlin, 1999. MR **2000k**:16001
8. C. D. Aliprantis, D. J. Brown, and O. Burkinshaw, Edgeworth equilibria, *Econometrica* **55** (1987), 1109–1137. MR **89c**:90025
9. C. D. Aliprantis, D. J. Brown, and O. Burkinshaw, *Existence and Optimality of Competitive Equilibria*, Springer-Verlag, Heidelberg and New York, 1990. MR **92i**:90023
10. C. D. Aliprantis and O. Burkinshaw, *Locally Solid Riesz Spaces*, Academic Press, New York and London, 1978. MR **58**:12271
11. C. D. Aliprantis and O. Burkinshaw, *Positive Operators*, Academic Press, New York and London, 1985. MR **87h**:47086
12. C. D. Aliprantis, M. Florenzano, and R. Tourky, Economic analysis in ordered vector spaces, in preparation.
13. C. D. Aliprantis, R. Tourky, and N. C. Yannellis, The Riesz–Kantorovich formula and general equilibrium theory, *J. Math. Econom.* **34** (2000), 55–76. MR **2001i**:91088

14. C. D. Aliprantis, R. Tourky, and N. C. Yannelis, A theory of value with non-linear prices: equilibrium analysis beyond vector lattices, *J. Econ. Theory* **100** (2001), 22–72. CMP 2002:01
15. T. Andô, On fundamental properties of a Banach space with a cone, *Pacific J. Math.* **12** (1962), 1163–1169. MR **27**:568
16. G. Debreu, *Theory of Value: An Axiomatic Analysis of Economic Equilibrium*, Cowles Foundation Monograph, Yale University Press, New Haven, 1959. MR **22**:1447
17. G. Debreu, New concepts and techniques for equilibrium analysis, *Int. Econ. Rev.* **3** (1962), 257–273.
18. T. Husain, *The Open Mapping and Closed Graph Theorems in Topological Vector Spaces*, Oxford Mathematical Monographs, Oxford University Press, London, 1965. MR **31**:2589
19. G. J. O. Jameson, *Ordered Linear Spaces*, Springer-Verlag Lecture Notes in Mathematics, **141**, Heidelberg and New York, 1970. MR **55**:10996
20. L. V. Kantorovich, Concerning the general theory of operations in partially ordered spaces, *DAN SSSR* **1** (1936), 271–274 (in Russian).
21. V. L. Klee, Jr., Extremal structure of convex sets, *Arch. Math.* **8** (1957), 234–240. MR **19**:1065a
22. V. L. Klee, Jr., Extremal structure of convex sets, II, *Math. Z.* **69** (1958), 90–104. MR **19**:1065b
23. W. A. J. Luxemburg and A. C. Zaanen, *Riesz Spaces I*, North-Holland, Amsterdam, 1971. MR **58**:23483
24. A. Mas-Colell, The price equilibrium existence problem in topological vector lattices, *Econometrica* **54** (1986), 1039–1053. MR **87m**:90083
25. A. L. Peressini, *Ordered Topological Vector Spaces*, Harper and Row, New York and London, 1967. MR **37**:3315
26. I. A. Polyakis, Lattice-subspaces of $C[0, 1]$ and positive bases, *J. Math. Anal. Appl.* **184** (1994), 1–18. MR **95g**:46040
27. I. A. Polyakis, Finite-dimensional lattice-subspaces of $C(\Omega)$ and curves of \mathbb{R}^n , *Trans. Amer. Math. Soc.* **348** (1996), 2793–2810. MR **96k**:46031
28. F. Riesz, Sur quelques notions fondamentales dans la theorie générale des opérations linéaires, *Ann. of Math.* **41** (1940), 174–206. MR **1**:147d
29. H. H. Schaefer, *Topological Vector Spaces*, Springer-Verlag, New York, Heidelberg, Berlin 1970. MR **33**:1689 (first ed.)
30. A. C. M. van Rooij, On the space of all regular operators between two Riesz spaces, *Indag. Math.* **47** (1985), 95–98. MR **86k**:46011
31. B. Z. Vulikh, *Introduction to the Theory of Partially Ordered Spaces*, Wolters-Noordhoff, Groningen, 1967. MR **37**:121

DEPARTMENT OF ECONOMICS AND DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA 47907–1310

E-mail address: `aliprantis@mgmt.purdue.edu`

DEPARTMENT OF ECONOMICS, UNIVERSITY OF MELBOURNE, PARKVILLE, VICTORIA 3052, AUSTRALIA

E-mail address: `rtourky@unimelb.edu.au`