

## QUADRATIC ITERATIONS TO $\pi$ ASSOCIATED WITH ELLIPTIC FUNCTIONS TO THE CUBIC AND SEPTIC BASE

HENG HUAT CHAN, KOK SENG CHUA, AND PATRICK SOLÉ

ABSTRACT. In this paper, properties of the functions  $A_d(q)$ ,  $B_d(q)$  and  $C_d(q)$  are derived. Specializing at  $d = 1$  and  $2$ , we construct two new quadratic iterations to  $\pi$ . These are analogues of previous iterations discovered by the Borweins (1987), J. M. Borwein and F. G. Garvan (1997), and H. H. Chan (2002). Two new transformations of the hypergeometric series  ${}_2F_1(1/3, 1/6; 1; z)$  are also derived.

### 1. INTRODUCTION

The famous Jacobi identity states that if

$$(1.1) \quad \vartheta_2(q) = \sum_{n=-\infty}^{\infty} q^{(n+1/2)^2},$$

$$(1.2) \quad \vartheta_3(q) = \sum_{n=-\infty}^{\infty} q^{n^2},$$

and

$$(1.3) \quad \vartheta_4(q) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n^2},$$

then [1, p. 40, Entry 25 (vii)]

$$(1.4) \quad \vartheta_3^4(q) = \vartheta_4^4(q) + \vartheta_2^4(q).$$

Around 1991, J. M. Borwein and P. B. Borwein discovered the following cubic analogue of (1.4) [6]:

$$(1.5) \quad a^3(q) = b^3(q) + c^3(q),$$

---

Received by the editors January 15, 2002 and, in revised form, August 21, 2002.

2000 *Mathematics Subject Classification*. Primary 11Y60, 33C05, 33E05, 11F03.

The first author was funded by National University of Singapore Academic Research Fund, Project Number R14000027112.

where

$$a(q) = \sum_{m,n=-\infty}^{\infty} q^{m^2+mn+n^2},$$

$$b(q) = \sum_{m,n=-\infty}^{\infty} \omega^{n-m} q^{n^2+mn+m^2},$$

and

$$c(q) = \sum_{m,n=-\infty}^{\infty} q^{(n+1/3)^2+(n+1/3)(m+1/3)+(m+1/3)^2},$$

$\omega$  being a primitive cube root of unity. The Borwein functions  $a(q)$ ,  $b(q)$  and  $c(q)$  and the Jacobi theta functions  $\vartheta_1(q)$ ,  $\vartheta_2(q)$  and  $\vartheta_3(q)$  share many similar features. For example, if we set

$$(1.6) \quad \alpha(q) = \frac{\vartheta_2^4(q)}{\vartheta_3^4(q)}$$

and

$$(1.7) \quad x_3(q) = \frac{c^3(q)}{a^3(q)},$$

then [1, p. 125]

$$(1.8) \quad \alpha(q^2) = \left( \frac{1 - \sqrt{1 - \alpha(q)}}{1 + \sqrt{1 - \alpha(q)}} \right)^2$$

and [2, (3.2)]

$$(1.9) \quad x_3(q^3) = \left( \frac{1 - \sqrt[3]{1 - x_3(q)}}{1 + 2\sqrt[3]{1 - x_3(q)}} \right)^3.$$

The duplication formula (1.8) for  $\alpha(q)$  plays an important role in proving the following iteration:

**Iteration 1.1.** Let  $t_0 = \frac{1}{2}$ ,  $s_0 = \frac{1}{\sqrt{2}}$ ,

$$s_{n+1} = \frac{1 - \sqrt{1 - s_n^2}}{1 + \sqrt{1 - s_n^2}}, \quad \text{and} \quad t_{n+1} = (1 + s_{n+1})^2 t_n - 2^{n+1} s_{n+1}.$$

Then  $t_n^{-1}$  converges quadratically to  $\pi$ .

In a similar way, the triplication formula (1.9) leads to

**Iteration 1.2.** Let  $t_0 = \frac{1}{3}$ ,  $s_0 = \frac{\sqrt{3}-1}{2}$ ,

$$s_n = \frac{1 - (1 - s_{n-1}^3)^{1/3}}{1 + 2(1 - s_{n-1}^2)^{1/3}}, \quad \text{and} \quad t_n = (1 + 2s_n)^2 t_{n-1} - 3^{n-1} ((1 + 2s_n)^2 - 1).$$

Then  $t_n^{-1}$  converges cubically to  $\pi$ .

For an excellent account and proofs of Iterations 1.1 and 1.2, see the paper by J. M. Borwein and F. G. Garvan [8].

Recently, motivated by the approach given in [8], H. H. Chan [10] devised a new method for deriving iterations to  $\pi$ . He succeeded in proving the following respective analogues of Iterations 1.1 and 1.2:

**Iteration 1.3.** Let  $k_0 = 0$  and  $s_0 = \frac{1}{\sqrt{2}}$ . Set

$$s_n = \frac{1 - \sqrt{1 - s_{n-1}^2}}{1 + \sqrt{1 - s_{n-1}^2}} \quad \text{and} \quad k_n = (1 + s_n)^2 k_{n-1} + 2^n (1 - s_n) s_n.$$

Then  $k_n^{-1}$  tends to  $\pi$  quadratically.

**Iteration 1.4.** Let  $k_0 = 0$  and  $s_0 = \frac{1}{\sqrt[3]{2}}$ . Set

$$s_n = \frac{1 - \sqrt[3]{1 - s_{n-1}^3}}{1 + 2\sqrt[3]{1 - s_{n-1}^3}} \quad \text{and} \quad k_n = (1 + 2s_n)^2 k_{n-1} + 8 \cdot 3^{n-2} \sqrt{3} s_n \frac{1 - s_n^3}{1 + 2s_n}.$$

Then  $k_n^{-1}$  tends to  $\pi$  cubically.

At the end of [10], Chan compared his method with that of Borwein and Garvan. He remarked that Borwein and Garvan’s method appeared to work only when  $N = p$ , since their function  $\epsilon_{N,p}$  [8, (3.13)] simplifies only in this case. In other words, for a function of “base  $p$ ” (or a function associated with the modular group  $\Gamma(p)$ ), the only iteration they obtained was a  $p$ -th order iteration to  $\pi$ . Chan’s method, on the other hand, yields (at least theoretically) an  $N$ -th order iteration to  $\pi$  for suitably chosen functions of “base  $p$ ”. Unfortunately, except for those when  $N = p$ , the iterations obtained using Chan’s method for  $N \neq p$  are not as simple as those of Iterations 1.1-1.4. An example of a quadratic iteration to the septic base was recently given by Song Heng Chan [13]. It therefore appears that we have exhausted all possible simple iterations analogous to Iterations 1.1-1.4 using the functions motivated by Ramanujan’s theories of elliptic functions to alternative bases [2], as well as the functions obtained from [8].

In this article, we introduce three new functions:

$$(1.10) \quad A_d(q) = \sum_{m,n \in \mathbb{Z}} q^{2(m^2+mn+dn^2)},$$

$$(1.11) \quad B_d(q) = \sum_{m,n \in \mathbb{Z}} (-1)^{m-n} q^{m^2+mn+dn^2},$$

and

$$(1.12) \quad C_d(q) = \sum_{m,n \in \mathbb{Z}} q^{2((m+1/2)^2+(m+1/2)n+dn^2)}.$$

Note that when  $d = 1$ , we are working with the function  $a(q^2)$ , which is associated with the cubic base. Surprisingly, in this case, we are able to obtain an analogue of Iterations 1.1-1.4, namely,

**Iteration 1.5.** Let  $k_0 = 0$  and  $y_0 = 8/9$ . Set

$$y_n = 2 \frac{(6y_{n-1}^2 - 5y_{n-1} + \sqrt{y_{n-1}(4 - y_{n-1})})}{9y_{n-1}^2 - 6y_{n-1} + 1}$$

and

$$k_n = 2^n \sqrt{3} \frac{y_{n-1}(1 - y_{n-1})}{\sqrt{4 - 3y_{n-1}}} + (4 - 3y_{n-1})k_{n-1}.$$

Then  $k_n^{-1}$  tends to  $\pi$  quadratically.

Note that this is a quadratic iteration in the cubic base. For  $d = 2$ , we have the following quadratic iteration associated with the septic base:

**Iteration 1.6.** Let  $y_0 = \frac{4}{5}, k_0 = 0$ . Let

$$y_n = \frac{2y_{n-1}^2 - y_{n-1} + \sqrt{4y_{n-1} - 3y_{n-1}^2}}{1 + y_{n-1}^2}.$$

Set

$$k_n = \frac{2^n y_{n-1}(1 - y_{n-1})}{\sqrt{7} (2 - y_{n-1})} \sqrt{(y_{n-1} + 1)(4 - 3y_{n-1})(y_{n-1}^2 - 3y_{n-1} + 4)} + (2 - y_{n-1})^2 k_{n-1}.$$

Then  $k_n^{-1}$  tends to  $\pi$  quadratically.

In Section 2, we quote several identities satisfied by  $A_d(q), B_d(q)$  and  $C_d(q)$ , and we prove that

$$(1.13) \quad A_d^2(q) = B_d^2(q) + C_d^2(q).$$

The special case of (1.13) when  $d = 2$  is proved in [11] using the existence of a unique 7-modular lattice of dimension 4 over  $\mathbb{Z}$ . Note that (1.13) is an analogue of (1.4) and (1.5).

In Section 3, we set  $d = 2$ . Motivated by (1.6) and (1.7), we define

$$(1.14) \quad Y(q) := B_2(q)/A_2(q).$$

We then establish a duplication formula satisfied by  $Y(q)$ , which is given by

$$(1.15) \quad Y(q^2) = \frac{2Y^2(q) - Y(q) + \sqrt{4Y(q) - 3Y^2(q)}}{1 + Y^2(q)}.$$

We end the section by deriving Iteration 1.6.

In Section 4, we set  $d = 1$  and establish the duplication formula

$$(1.16) \quad \mathcal{Y}(q^2) = 2 \left( \frac{6\mathcal{Y}^2(q) - 5\mathcal{Y}(q) + \sqrt{\mathcal{Y}(q)(4 - 3\mathcal{Y}(q))}}{9\mathcal{Y}^2(q) - 6\mathcal{Y}(q) + 1} \right),$$

where

$$(1.17) \quad \mathcal{Y}(q) := \frac{B_1^2(q)}{A_1^2(q)}.$$

We then prove several identities which eventually lead to Iteration 1.5. We also establish two transformation formulas for the hypergeometric series  ${}_2F_1(1/3, 1/6; 1; z)$ . In Section 5, we give another proof of the transformation formulas established in Section 4, using one of C. W. Borchardt’s mean iterations studied by the Borweins and Garvan [5], [7].

2. IDENTITIES SATISFIED BY  $A_d, B_d$  AND  $C_d$

Let  $\vartheta_i(q), i = 2, 3, 4$ , be defined as in (1.1)-(1.3). We have

**Lemma 2.1.**

$$(2.1) \quad A_d(q) = \sum_{m,n=-\infty}^{\infty} q^{2(m^2+mn+dn^2)} = \vartheta_3(q^2)\vartheta_3(q^{8d-2}) + \vartheta_2(q^2)\vartheta_2(q^{8d-2}),$$

$$(2.2) \quad B_d(q) = \sum_{m,n=-\infty}^{\infty} (-1)^{m-n} q^{m^2+mn+dn^2} = \vartheta_4(q)\vartheta_4(q^{4d-1}),$$

and

$$(2.3) \quad C_d(q) = \sum_{m,n=-\infty}^{\infty} q^{2((m+1/2)^2+(m+1/2)n+dn^2)} \\ = \vartheta_2(q^2)\vartheta_3(q^{8d-2}) + \vartheta_3(q^2)\vartheta_2(q^{8d-2}).$$

*Proof.* The identity (2.1) can be found, for example, in [12, p. 1738]. For the proofs of the subsequent identities, we will need the following simple identity, namely, for any odd integer  $n$ ,

$$(2.4) \quad \sum_{m=-\infty}^{\infty} (-1)^m q^{(m+\frac{n}{2})^2} = 0.$$

This identity follows immediately from [15, p. 464].

Next, note that

$$\begin{aligned} B_d(q) &= \sum_{m,n=-\infty}^{\infty} (-1)^{m-n} q^{m^2+mn+dn^2} \\ &= \sum_{m,n=-\infty}^{\infty} (-1)^{m+n} q^{(m+\frac{n}{2})^2+\frac{D}{4}n^2}, \quad \text{with } D = 4d - 1, \\ &= \sum_{n=-\infty}^{\infty} (-1)^n q^{Dn^2/4} \sum_{m=-\infty}^{\infty} (-1)^m q^{(m+\frac{n}{2})^2} \\ &= \sum_{\substack{n=-\infty \\ n \text{ even}}}^{\infty} (-1)^n q^{Dn^2/4} \sum_{m=-\infty}^{\infty} (-1)^m q^{(m+\frac{n}{2})^2} \\ &\quad + \sum_{\substack{n=-\infty \\ n \text{ odd}}}^{\infty} (-1)^n q^{Dn^2/4} \sum_{m=-\infty}^{\infty} (-1)^m q^{(m+\frac{n}{2})^2} \\ &= \sum_{\substack{n=-\infty \\ n \text{ even}}}^{\infty} (-1)^n q^{Dn^2/4} \sum_{m=-\infty}^{\infty} (-1)^m q^{(m+\frac{n}{2})^2}, \quad \text{by (2.4),} \\ &= \vartheta_4(q^D)\vartheta_4(q). \end{aligned}$$

Finally, rewrite  $C_d(q)$  as

$$\begin{aligned} C_d(q) &= \sum_{m,n=-\infty}^{\infty} q^{2((m+\frac{n}{2})^2+\frac{D}{4}n^2)} \\ &= \sum_{\substack{m,n=-\infty \\ n \text{ even}}}^{\infty} q^{2((m+\frac{n}{2})^2+\frac{D}{4}n^2)} + \sum_{\substack{m,n=-\infty \\ n \text{ odd}}}^{\infty} q^{2((m+\frac{n}{2})^2+\frac{D}{4}n^2)} \\ &= \vartheta_2(q^2)\vartheta_3(q^{8d-2}) + \vartheta_3(q^2)\vartheta_2(q^{8d-2}). \end{aligned}$$

□

Our next task is to establish relations between  $A_d(q)$ ,  $B_d(q)$ , and  $C_d(q)$ . We have

**Lemma 2.2.**

- (i)  $A_d(q^2) = B_d(q) + C_d(q^2)$ .
- (ii)  $A_d^2(q) = B_d^2(q) + C_d^2(q)$ .

*Proof.* From (2.1) and (2.3), we find, after some simplification, that

$$\begin{aligned} A_d(q^2) - C_d(q^2) &= (\vartheta_3(q^4) - \vartheta_2(q^4))(\vartheta_3(q^{16d-4}) - \vartheta_2(q^{16d-4})) \\ &= \vartheta_4(q)\vartheta_4(q^{4d-1}) = B_d(q). \end{aligned}$$

This completes the proof of (i).

Next, by (2.1) and (2.3), we find that

$$(2.5) \quad A_d^2(q) - C_d^2(q) = (\vartheta_3^2(q^{8d-2}) - \vartheta_2^2(q^{8d-2}))(\vartheta_3^2(q^2) - \vartheta_2^2(q^2)).$$

On the other hand,

$$(2.6) \quad \begin{aligned} \vartheta_3^2(q^4) - \vartheta_2^2(q^4) &= (\vartheta_3(q^4) - \vartheta_2(q^4))(\vartheta_3(q^4) + \vartheta_2(q^4)) \\ &= \vartheta_3(q)\vartheta_4(q) = \vartheta_4^2(q^2), \end{aligned}$$

where the last equality follows from a well-known identity (see, for example, [1, p. 40, Entry 25(iii)]). Substituting (2.6) into (2.5) completes the proof of (ii). □

**Lemma 2.3.**

$$(2.7) \quad A_d(q) + B_d(q^2) = 2 \sum_{m,n=-\infty}^{\infty} q^{2(dm^2+2mn+4n^2)}.$$

*Proof.* We have

$$\begin{aligned} A_d(q) + B_d(q^2) &= \sum_{m,n=-\infty}^{\infty} (1 + (-1)^{m-n})q^{2(m^2+mn+dn^2)} \\ &= 2 \left( \left\{ \sum_{\substack{m,n=-\infty \\ m,n \text{ both even}}}^{\infty} + \sum_{\substack{m,n=-\infty \\ m,n \text{ both odd}}}^{\infty} \right\} q^{2(m^2+mn+dn^2)} \right) \\ &= 2 \left( \left\{ \sum_{n=-\infty}^{\infty} \sum_{\substack{m=-\infty \\ m \text{ even}}}^{\infty} + \sum_{\substack{m,n=-\infty \\ m,n \text{ both odd}}}^{\infty} - \sum_{\substack{n=-\infty \\ n \text{ odd}}}^{\infty} \sum_{\substack{m=-\infty \\ m \text{ even}}}^{\infty} \right\} q^{2(m^2+mn+dn^2)} \right). \end{aligned}$$

From (2.4), we deduce that

$$\begin{aligned} & \left\{ \sum_{\substack{m,n=-\infty \\ m,n \text{ both odd}}}^{\infty} - \sum_{\substack{n=-\infty \\ n \text{ odd}}}^{\infty} \sum_{\substack{m=-\infty \\ m \text{ even}}}^{\infty} \right\} q^{2(m^2+mn+dn^2)} \\ &= - \sum_{\substack{n=-\infty \\ n \text{ odd}}}^{\infty} \sum_{m=-\infty}^{\infty} (-1)^m q^{2(m^2+mn+dn^2)} \\ &= - \sum_{\substack{n=-\infty \\ n \text{ odd}}}^{\infty} \sum_{m=-\infty}^{\infty} (-1)^m q^{2(m+\frac{n}{2})^2+\frac{d}{4}n^2} = 0. \end{aligned}$$

□

### 3. THE CASE $d = 2$ AND ITERATION 1.6

We now specialize our results in the previous section to the case  $d = 2$ . Set  $A(q) = A_2(q), B(q) = B_2(q)$  and  $C(q) = C_2(q)$ . We have

**Corollary 3.1.**

- (i)  $A^2(q) = B^2(q) + C^2(q)$ .
- (ii)  $A(q) = 2A(q^2) - B(q^2)$ .
- (iii)  $B(q) = A(q^2) - C(q^2)$ .

Now, let  $Y(q)$  be given as in (1.14). From Corollary 3.1(ii), we find that

$$(3.1) \quad \frac{A(q)}{A(q^2)} = 2 - Y(q^2).$$

From Corollary 3.1(iii), we deduce that

$$\begin{aligned} Y(q) &= \frac{A(q^2)}{A(q)} - \frac{C(q^2)}{A(q)} \\ &= \frac{A(q^2)}{A(q)} - \frac{\sqrt{A^2(q^2) - B^2(q^2)}}{A(q)} \quad \text{by Corollary 3.1(i)} \\ &= \frac{A(q^2)}{A(q)} \left( 1 - \sqrt{1 - Y^2(q^2)} \right) \\ &= \frac{1}{2 - Y(q^2)} \left( 1 - \sqrt{1 - Y^2(q^2)} \right), \quad \text{by (3.1)}. \end{aligned}$$

Solving for  $Y(q^2)$ , we obtain the duplication formula (1.15).

We are now ready to establish Iteration 1.6. It is well known that  $A(q)$  satisfies the relation [14, p. 205, (5)]

$$\sqrt{n}A(e^{-\pi\sqrt{n}/\tau}) = A(e^{-\pi/\sqrt{\tau n}}).$$

Following the method illustrated in [10], define

$$\mathcal{K} := \frac{1}{\pi A^2(e^{-\pi\sqrt{n}/\tau})} - 2\sqrt{\frac{n}{7}} \frac{\tilde{A}(e^{-\pi\sqrt{n}/\tau})}{A^3(e^{-\pi\sqrt{n}/\tau})},$$

where

$$\tilde{f}(q) := q \frac{df(q)}{dq}.$$

As in [10], we find that

$$(3.2) \quad r\mathcal{K}\left(\frac{1}{r}\right) + \mathcal{K}(r) = 0$$

and, for any positive integer  $N$ ,

$$(3.3) \quad \mathcal{K}(N^2r) = m_N^2(q) \left( \frac{2}{A^2(e^{-\pi\sqrt{r/7}})} \sqrt{\frac{r}{7}} \frac{\widetilde{m}_N}{m_N}(e^{-\pi\sqrt{r/7}}) + \mathcal{K}(r) \right),$$

where

$$m_N(q) = \frac{A(q)}{A(q^N)}.$$

Now set  $N = 2$ . In order to derive Iteration 1.6, we need to convert the derivative  $\widetilde{m}_2$  with respect to  $q$  to a derivative with respect to  $Y_2 := Y_2(q) = Y(q^2)$ . As such, we set

$$F(q) := q \frac{dY_2}{dq},$$

and we find from (3.3) and (3.1) that

$$(3.4) \quad \mathcal{K}(2^2r) = -\frac{2}{A^2(e^{-\pi\sqrt{r/7}})} \sqrt{\frac{r}{7}} (2 - Y_2(e^{-\pi\sqrt{r/7}})) F(e^{-\pi\sqrt{r/7}}) + (2 - Y_2(e^{-\pi\sqrt{r/7}}))^2 \mathcal{K}(r).$$

It remains now to evaluate  $F(q)$ , which we will complete in a few steps. First, we observe that

$$(3.5) \quad 2\eta^3(\tau)\eta^3(7\tau) = B^2(q^2)(A(q^2) - B(q^2)),$$

where

$$\eta(\tau) = q^{1/12} \prod_{k=1}^{\infty} (1 - q^{2k}), \quad q = e^{\pi i\tau}.$$

Identity (3.5) is obtained from the two representations of the theta series associated with the lattice  $A_6^2$  (see [11, Section 5.1]). Dividing by  $A^3(q^2)$ , we find that

$$Y_2^2(1 - Y_2) = 2 \frac{\eta^3(\tau)\eta^3(7\tau)}{A^3(q^2)} = 2 \frac{A^3(q)}{A^3(q^2)} \frac{\eta^3(\tau)\eta^3(7\tau)}{A^3(q)}.$$

By (3.1) and the identity (see, for example, [12, Lemma 2.2])

$$\frac{A^3(q)}{\eta^3(\tau)\eta^3(7\tau)} = 49 \frac{\eta^4(7\tau)}{\eta^4(\tau)} + 13 + \frac{\eta^4(\tau)}{\eta^4(7\tau)},$$

we find that

$$(3.6) \quad Y_2^2(1 - Y_2) = 2(2 - Y_2)^3 \frac{\chi}{7 - \chi},$$

where

$$(3.7) \quad \chi := \chi(q) = x(1 - x),$$

with

$$\frac{1}{x} = \frac{1}{x(q)} = 1 + \frac{1}{7} \frac{\eta^4(\tau)}{\eta^4(7\tau)}.$$

Solving for  $\chi$  yields

$$(3.8) \quad \chi = 7 \frac{Y_2^2(Y_2 - 1)}{(3Y_2 - 4)(Y_2^2 - 3Y_2 + 4)}.$$

Differentiating (3.8) with respect to  $q$ , we find that

$$(3.9) \quad q \frac{d\chi}{dq} = -14 \frac{Y_2(5Y_2 - 4)(Y_2 - 2)^2}{(3Y_2 - 4)^2(Y_2^2 - 3Y_2 + 4)^2} q \frac{dY_2}{dq}.$$

On the other hand, from (3.7), we find that

$$(3.10) \quad q \frac{d\chi}{dq} = (1 - 2x)q \frac{dx}{dq} = 2(1 - 2x)A^2(q)\chi,$$

using the identity [12, (2.23)]

$$q \frac{dx}{dq} = 2A^2(q)x(1 - x).$$

Since

$$x = \frac{1}{2} \left( 1 - \sqrt{1 - 4\chi} \right),$$

we deduce from (3.10) that

$$(3.11) \quad q \frac{d\chi}{dq} = 2A^2(q)\chi\sqrt{1 - 4\chi}.$$

Substituting (3.11) into (3.9) and simplifying using (3.8), we conclude that

$$(3.12) \quad F(q) = q \frac{dY_2}{dq} = A^2(q) \frac{Y_2(1 - Y_2)}{(Y_2 - 2)^2} \sqrt{(Y_2 + 1)(4 - 3Y_2)(Y_2^2 - 3Y_2 + 4)}.$$

Now, substituting (3.12) into (3.4), and setting  $r = 1$ ,  $k_n = \mathcal{K}(4^n)$  and  $y_n = Y_2(e^{-2^{n-1}\pi/\sqrt{7}})$ , we deduce that

$$k_n = \frac{2^n}{\sqrt{7}} \frac{y_{n-1}(1 - y_{n-1})}{2 - y_{n-1}} \sqrt{(y_{n-1} + 1)(4 - 3y_{n-1})(y_{n-1}^2 - 3y_{n-1} + 4)} + (2 - y_{n-1})^2 k_{n-1}.$$

To complete our proof, it remains to compute  $k_0$  and  $y_0$ .

By (3.2),  $k_0 = \mathcal{K}(1) = 0$ . To compute  $y_0$ , note that  $x(e^{-\pi/\sqrt{7}}) = 1/2$  [12, proof of (1.9)]. Hence  $\chi(e^{-\pi/\sqrt{7}}) = 1/4$ , and, solving the equation (3.6), we deduce that  $y_0 = 4/5$ .

We end this section by giving another proof of the key identity (3.5). The second proof connects our work with Ramanujan’s modular equations of degree 7.

First, we observe that (3.5) is equivalent to

$$(3.13) \quad 4\eta^3(\tau)\eta^3(7\tau) = B(q)(B(q) - A(q))^2, \quad q = e^{\pi i\tau}.$$

This follows from the identity

$$2B^2(q^2)(A(q^2) - B(q^2)) = B(q)(A(q) - B(q))^2,$$

which is a consequence of the duplication formula (1.15).

By using the identities [1, p. 122, Entry 10(i), (vii), p. 123, Entry 10(iii), p. 124, Entry 12 (ii)], we may write (3.13) (with  $q$  replaced by  $\sqrt{q}$ ) in the form

$$(3.14) \quad \begin{aligned} &2\sqrt{(1-\alpha)(1-\beta)}(\alpha\beta)^{1/8} \\ &= \sqrt{(1-\sqrt{\alpha})(1-\sqrt{\beta})} \left( (\alpha\beta)^{1/4} + 1 - (1-\sqrt{\beta})(1-\sqrt{\alpha}) \right)^2, \end{aligned}$$

where  $\alpha = \alpha(q)$  and  $\beta = \alpha(q^7)$ ,  $\alpha(q)$  being given as in (1.6). Now, from [1, pp. 316-317], it is known that, with  $t = (\alpha\beta)^{1/8}$ ,

$$\sqrt{\alpha} := (1-t+t^2)(t(2-3t+2t^2))^{1/2} + (1-t)(t(1-t+2t^2)(2-t+t^2))^{1/2}$$

and

$$\sqrt{\beta} := (1-t+t^2)(t(2-3t+2t^2))^{1/2} - (1-t)(t(1-t+2t^2)(2-t+t^2))^{1/2}.$$

Note that

$$\begin{aligned} (1-\sqrt{\alpha})(1-\sqrt{\beta}) &= 1 + 2(t(2-3t+2t^2))^{1/2}t - 2(t(2-3t+2t^2))^{1/2} \\ &\quad - 2(t(2-3t+2t^2))^{1/2}t^2 + t^4)^{1/2}. \end{aligned}$$

Hence,

$$\delta := \sqrt{(1-\sqrt{\alpha})(1-\sqrt{\beta})} = 1-t+t^2 - \sqrt{t(2-3t+2t^2)}.$$

Hence the proof of (3.13) is reduced to showing the algebraic expression

$$2t(1-t)^4 = \delta(\delta-1-t^2),$$

which is straightforward.

#### 4. CASE $d = 1$ AND A QUADRATIC ITERATION TO $\pi$

In this section, we construct an iteration to  $\pi$  arising from the case  $d = 1$ , or in other words, the cubic base. Set  $\mathcal{A}(q) = A_1(q)$ ,  $\mathcal{B}(q) = B_1(q)$  and  $\mathcal{C}(q) = C_1(q)$ . From Lemma 2.2, we have

**Corollary 4.1.**

- (i)  $\mathcal{A}^2(q) = \mathcal{B}^2(q) + \mathcal{C}^2(q)$ .
- (ii)  $\mathcal{B}(q) = \mathcal{A}(q^2) - \mathcal{C}(q^2)$ .

We need a third relation which plays the role of Corollary 4.1(ii). The third relation, unlike that of the case  $d = 2$ , turns out to be another quadratic relation.

**Lemma 4.2.**

$$\mathcal{A}^2(q) = 4\mathcal{A}^2(q^2) - 3\mathcal{B}^2(q^2).$$

*Proof.* To prove Lemma 4.2, we first replace  $q^2$  by  $\sqrt{q}$  and observe that by [1, p. 122, Entry 10(i),(vi), p. 123, Entry 11(i), (iii)], it suffices to verify that

$$(4.1) \quad \left( \sqrt{(1+\sqrt{\alpha})(1+\sqrt{\gamma})} + 2\sqrt{s} \right)^2 = 4(1+s)^2 - 3(1-\sqrt{\alpha})(1-\sqrt{\gamma}),$$

where  $\alpha = \alpha(q)$ ,  $\gamma = \alpha(q^3)$ , and

$$(4.2) \quad s^4 = \alpha\gamma.$$

The definition of  $s$  is motivated by the definition of  $t$  in the previous section and the following modular equation of degree 3 [1, p. 230, Entry 5(ii)]:

$$(4.3) \quad (\alpha\gamma)^{1/4} + \{(1-\alpha)(1-\gamma)\}^{1/4} = 1.$$

The modular equation (4.3) also shows that

$$\{(1 - \alpha)(1 - \gamma)\} = (1 - s)^4.$$

Solving for  $\alpha$  and  $\beta$  yields

$$(4.4) \quad \sqrt{\alpha} = \sqrt{s}(1 - s + \sqrt{s^2 - s + 1}),$$

and

$$(4.5) \quad \sqrt{\gamma} = \sqrt{s}(s - 1 + \sqrt{s^2 - s + 1}).$$

Using (4.4) and (4.5), we find that

$$(1 + \sqrt{\alpha})(1 + \sqrt{\gamma}) = (\sqrt{s} + \sqrt{s^2 - s + 1})^2,$$

and the left-hand side of (4.1) becomes

$$(4.6) \quad s^2 + 8s + 1 + 6\sqrt{s}\sqrt{s^2 - s + 1}.$$

Substituting (4.4) and (4.5) into the right-hand side of (4.1), we obtain the same expression (4.6) and complete the proof of Lemma 4.2.  $\square$

Dividing the identity in Lemma 4.2 by  $\mathcal{A}^2(q^2)$ , we find that

$$(4.7) \quad \frac{\mathcal{A}^2(q)}{\mathcal{A}^2(q^2)} = 4 - 3\mathcal{Y}(q^2),$$

where  $\mathcal{Y}(q)$  is given by (1.17). Now, substituting (1.17) into Corollary 3.1(ii), we deduce that

$$\mathcal{Y}(q) = \frac{1}{4 - 3\mathcal{Y}(q^2)} \left(1 - \sqrt{1 - \mathcal{Y}(q^2)}\right)^2,$$

and solving for  $\mathcal{Y}(q^2)$  gives the duplication formula (1.16).

We are now ready to derive the quadratic iteration. Following the argument as in the previous section, we find that if

$$\mathcal{K}_{\mathcal{A}} := \frac{1}{\pi \mathcal{A}^2(e^{-\pi\sqrt{n/3})}} - 2\sqrt{\frac{n}{3}} \frac{\tilde{\mathcal{A}}(e^{-\pi\sqrt{n/3})}}{\mathcal{A}^3(e^{-\pi\sqrt{n/3})}},$$

then

$$(4.8) \quad \mathcal{K}_{\mathcal{A}}(4r) = \mathbf{m}_2^2(e^{-\pi\sqrt{r/3}}) \left( \frac{2}{\mathcal{A}^2(e^{-\pi\sqrt{r/3}})} \sqrt{\frac{r}{3}} \frac{\tilde{\mathbf{m}}_2}{\mathbf{m}_2}(e^{-\pi\sqrt{r/3}}) + \mathcal{K}_{\mathcal{A}}(r) \right),$$

where

$$\mathbf{m}_2(q) = \frac{\mathcal{A}(q)}{\mathcal{A}(q^2)}.$$

Simplifying (4.8) using (4.7), we find that

$$(4.9) \quad \mathcal{K}_{\mathcal{A}}(4r) = -3\sqrt{\frac{r}{3}} \frac{\mathcal{F}(q)}{\mathcal{A}^2(e^{-\pi\sqrt{r/3}})} + \left(4 - 3\mathcal{Y}_2(e^{-\pi\sqrt{r/3}})\right) \mathcal{K}_{\mathcal{A}}(r),$$

where

$$\mathcal{Y}_2(q) := \mathcal{Y}(q^2) \quad \text{and} \quad \mathcal{F}(q) = q \frac{d\mathcal{Y}_2(q)}{dq}.$$

We first establish the identity

$$(4.10) \quad 16\eta^6(\tau)\eta^6(3\tau) = \mathcal{B}^2(q) (\mathcal{B}^2(q) - \mathcal{A}^2(q))^2.$$

Replacing  $q$  by  $\sqrt{q}$ , we see that (4.10) is equivalent to

$$(4.11) \quad 4s(1-s)^4 = (1-\sqrt{\alpha})(1-\sqrt{\gamma})((1-\sqrt{\alpha})(1-\sqrt{\gamma})-(1+s)^2)^2,$$

where  $s$  is given by (4.2). Using (4.4) and (4.5), we show that both sides of (4.11) are equal and complete the proof.

It is known from [9, (2.1), (2.2), (2.3)] that if  $x_3(q^2)$  is given by (1.7), where  $q = e^{\pi i \tau}$ , then

$$(4.12) \quad \chi(q) := x_3(q^2)(1-x_3(q^2)) = 27 \frac{\eta^6(\tau)\eta^6(3\tau)}{\mathcal{A}^6(q)} = \frac{27}{16} \mathcal{Y}(q) (\mathcal{Y}(q) - 1)^2,$$

where the last equality follows from (4.10). Using (1.16), we find that

$$(4.13) \quad \chi(q) = \frac{27 \mathcal{Y}_2^2(q)(1-\mathcal{Y}_2(q))}{4(4-3\mathcal{Y}_2(q))^3}.$$

Differentiating (4.13) with respect to  $q$  and using the fact that [9]

$$q \frac{dx_3}{dq} = 2\mathcal{A}^2(q)\chi(q),$$

we deduce that

$$(4.14) \quad \mathcal{F}(q) = q \frac{d\mathcal{Y}_2}{dq} = \mathcal{A}^2(q) \frac{2\mathcal{Y}_2(q)(\mathcal{Y}_2(q) - 1)}{\sqrt{4 - 3\mathcal{Y}_2(q)}}.$$

Substituting (4.14) into (4.9), and setting  $r = 1$ ,  $k_n = \mathcal{K}_{\mathcal{A}}(4^n)$  and  $y_n = \mathcal{Y}_2(e^{-2^{n-1}\pi/\sqrt{3}})$ , we deduce that

$$k_n = 2^n \sqrt{3} \frac{y_{n-1}(1-y_{n-1})}{\sqrt{4-3y_{n-1}}} + (4-3y_{n-1})k_{n-1}.$$

Now, again  $k_0 = \mathcal{K}_{\mathcal{A}}(1) = 0$ . It is also known [10, Proof of Iteration 5.1] that  $\chi(e^{-\pi/\sqrt{3}}) = 1/4$ . Solving for  $y_0$  using (4.13), we deduce that  $y_0 = 8/9$ . Together with the duplication formula (1.16), we obtain Iteration 1.5.

We end this section with two apparently new transformation formulas for the hypergeometric series

$${}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; z\right) = \sum_{n=0}^{\infty} \frac{(\frac{1}{3})_n (\frac{1}{6})_n z^n}{(1)_n n!},$$

where  $(a)_n = (a)(a+1)\cdots(a+n-1)$ . First, let  $0 < q < 1$  and recall from [2, Lemma 2.6] or [9, (4.3)], and [4, p. 178] (due to Kummer), that

$$(4.15) \quad \mathcal{A}(q) = {}_2F_1\left(\frac{1}{3}, \frac{2}{3}; 1; x_3(q^2)\right) = {}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; 4\chi(q)\right).$$

From (4.12), we find that

$$(4.16) \quad \chi(q) = \frac{27}{16} \mathbf{x}^2(q)(1-\mathbf{x}(q)),$$

where

$$\mathbf{x}(q) = 1 - \mathcal{Y}(q).$$

Using the new variable  $\mathbf{x}(q)$ , we write (4.13) as

$$(4.17) \quad \chi(q) = \frac{27(1-\mathbf{x}(q^2))^2 \mathbf{x}(q^2)}{4(1+3\mathbf{x}(q^2))^3}$$

and (4.7) as

$$(4.18) \quad \frac{\mathcal{A}(q)}{\mathcal{A}(q^2)} = \sqrt{1 + 3\mathbf{x}(q^2)}.$$

Now, by (4.15) and (4.16),

$$(4.19) \quad \mathcal{A}(q^2) = {}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; \frac{27}{4}\mathbf{x}^2(q^2)(1 - \mathbf{x}(q^2))\right).$$

On the other hand,

$$(4.20) \quad \mathcal{A}(q) = {}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; 27\frac{(1 - \mathbf{x}(q^2))^2\mathbf{x}(q^2)}{(1 + 3\mathbf{x}(q^2))^3}\right).$$

Finally, using (4.18) and replacing  $\mathbf{x}(q^2)$  by  $u$ , we deduce the transformation formula

$$(4.21) \quad {}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; 27\frac{(1-u)^2u}{(1+3u)^3}\right) = \sqrt{1+3u} {}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; \frac{27}{4}(1-u)u^2\right).$$

The formula (4.21) is only valid when  $0 < u < 1/9$ . Note that the formula is not valid beyond the given range, since  $u = 1/9$  satisfies the equation

$$27\frac{(1-u)^2u}{(1+3u)^3} = 1$$

and that  ${}_2F_1(a, b; c; z)$  is divergent at  $z = 1$ .

If we now replace  $u$  by  $\frac{u}{4-3u}$ , we immediately obtain the following companion of (4.21):

$$(4.22) \quad {}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; 27\frac{u^2(1-u)}{(4-3u)^3}\right) = \frac{\sqrt{4-3u}}{2} {}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; \frac{27}{4}u(1-u)^2\right).$$

This is valid for  $0 < u < 1/3$ .

### 5. BORCHARDT'S ITERATION REVISITED

The last transformation formula (4.22) can also be derived from one of C. W. Borchardt's mean iteration formulas [3], considered in [5]. We now describe a second proof of (4.22) (which certainly gives a second proof of (4.21)), using an explicit description of this mean iteration.

Once again, let  $0 < q < 1$ . Recall from Lemma 4.2 that if we set  $a_n := a_n(q) = \mathcal{A}^2(q^{2^n})$  and  $b_n := b_n(q) = \mathcal{B}^2(q^{2^{n+1}})$ , then

$$(5.1) \quad a_{n+1} = \frac{a_n + 3b_n}{4}.$$

From Corollary 4.1(ii), we find that

$$\sqrt{b_n} = \sqrt{a_{n+2}} - \sqrt{a_{n+2} - b_{n+1}},$$

which yields

$$(5.2) \quad \sqrt{\frac{b_n}{b_{n+1}}} = \sqrt{\frac{a_{n+2}}{b_{n+1}}} - \sqrt{\frac{a_{n+2}}{b_{n+1}} - 1}.$$

From (5.2), we deduce that

$$(5.3) \quad \sqrt{\frac{b_{n+1}}{b_n}} = \sqrt{\frac{a_{n+2}}{b_{n+1}}} + \sqrt{\frac{a_{n+2}}{b_{n+1}} - 1}.$$

Adding (5.2) and (5.3) and simplifying, we deduce that

$$(5.4) \quad b_{n+1} + b_n = 2\sqrt{a_{n+2}b_n}.$$

Using (5.1) twice and squaring, we find from (5.4) that

$$(5.5) \quad b_{n+1}^2 - b_{n+1}b_n - \frac{a_nb_n - b_n^2}{4} = 0.$$

Solving (5.5) for  $b_{n+1}$  and determining the square root using the expansions of  $\mathcal{A}(q)$  and  $\mathcal{B}(q)$ , we deduce that

$$(5.6) \quad b_{n+1} = \frac{b_n + \sqrt{a_nb_n}}{2}.$$

Note that the iteration

$$(a, b) \rightarrow \left( \frac{a + 3b}{4}, \frac{b + \sqrt{ab}}{2} \right)$$

gives the iteration considered by the Borweins [5].

It is known [5, (1.3)] that the sequences  $\{a_n\}$  and  $\{b_n\}$  have a common limit,  $\mathbf{B}(a_0, b_0)$ , when  $0 < q < 1$ . This common limit is the analogue of Gauss' AGM and satisfies (by observing that  $\mathbf{B}(a_0, b_0) = \mathbf{B}(a_1, b_1)$ ) the transformation formula [5, (1.10)]

$$(5.7) \quad \mathbf{B}(x) = \frac{1 + 3x}{4} \mathbf{B} \left( \frac{2(\sqrt{x} + x)}{1 + 3x} \right),$$

where  $\mathbf{B}(x) = \mathbf{B}(1, x)$ . On the other hand, from our parametrization above, we find that, for  $0 < q < 1$ ,

$$\mathbf{B}(\mathcal{A}^2(q), \mathcal{B}^2(q^2)) = \mathbf{B}(\mathcal{A}^2(q^2), \mathcal{B}^2(q^4)) = \cdots = \mathbf{B}(\mathcal{A}^2(0), \mathcal{B}^2(0)) = 1.$$

Hence,

$$(5.8) \quad \frac{1}{\mathbf{B} \left( 1, \frac{\mathcal{B}^2(q^2)}{\mathcal{A}^2(q)} \right)} = \mathcal{A}^2(q) = \left( {}_2F_1 \left( \frac{1}{3}, \frac{1}{6}; 1; 27 \frac{(1 - \mathbf{x}(q^2))^2 \mathbf{x}(q^2)}{(1 + 3\mathbf{x}(q^2))^3} \right) \right)^2,$$

by (4.20). Next, note that by (4.7), we may write

$$(5.9) \quad \frac{\mathcal{B}^2(q^2)}{\mathcal{A}^2(q)} = \frac{\mathcal{B}^2(q^2)}{\mathcal{A}^2(q^2)} \frac{\mathcal{A}^2(q^2)}{\mathcal{A}^2(q)} = \frac{1 - \mathbf{x}(q^2)}{1 + 3\mathbf{x}(q^2)}.$$

Substituting (5.9) into (5.8) and replacing  $\mathbf{x}(q^2)$  by  $x$ , we deduce that, for  $0 < x < 1/9$ ,

$$(5.10) \quad \frac{1}{\mathbf{B} \left( 1, \frac{1-x}{1+3x} \right)} = \left( {}_2F_1 \left( \frac{1}{3}, \frac{1}{6}; 1; 27 \frac{(1-x)^2 x}{(1+3x)^3} \right) \right)^2.$$

Replacing  $x$  by  $\frac{1-x}{1+3x}$  in (5.10), we deduce that

$$(5.11) \quad \frac{1}{\mathbf{B}(1, x)} = \left( {}_2F_1 \left( \frac{1}{3}, \frac{1}{6}; 1; \frac{27}{4}(1-x)x^2 \right) \right)^2,$$

valid for  $2/3 < x < 1$ . Using (5.11) in (5.7) and taking the positive square root, we obtain

$${}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; \frac{27x(1-x)^2}{(1+3x)^3}\right) = \sqrt{\frac{1+3x}{4}} {}_2F_1\left(\frac{1}{3}, \frac{1}{6}; 1; \frac{27}{4}(1-x)x^2\right),$$

valid for  $2/3 < x < 1$ . Replacing  $x$  by  $1-x$  yields (4.22).

The Borweins and Garvan [7, Theorem 5] obtained the identity (5.11) without stating the range for which it is valid.

## 6. CONCLUSION

We have seen that functions such as  $A_d(q)$ ,  $B_d(q)$  and  $C_d(q)$  possess many interesting properties. In the cases  $d = 1$  and  $d = 2$ , we have shown that they are good candidates for furthering the study of Ramanujan's elliptic functions to alternative bases. Using other values of  $d$ , we will probably encounter new functions analogous to those studied in this article.

## ACKNOWLEDGEMENTS

This article was completed during the first author's visit to the University of Sussex as a Commonwealth Fellow. He thanks the department and Dr. R. P. Lewis for their hospitality. The work was partially done while the third author was visiting the Institute for Mathematical Sciences, National University of Singapore, in 2001. The visit was supported by the Institute and by a grant from DSTA of Singapore.

## REFERENCES

1. B. C. Berndt, *Ramanujan's Notebooks, Part III*, Springer-Verlag, New York, 1991. MR **92j**:01069
2. B. C. Berndt, S. Bhargava and F. G. Garvan, *Ramanujan's theories of elliptic functions to alternative bases*, Trans. Amer. Math. Soc. **347** (1995), 4163–4244. MR **97h**:33034
3. C. W. Borchardt, *Ueber das Arithmetisch-geometrische Mittel aus vier Elementen*, Berl. Monatsber (1876), 611–621.
4. J. M. Borwein and P. B. Borwein, *Pi and the AGM*, John Wiley and Sons, New York, 1987. MR **89a**:11134
5. J. M. Borwein and P. B. Borwein, *On the mean iteration  $(a, b) \leftarrow \left(\frac{a+3b}{4}, \frac{\sqrt{ab+b}}{2}\right)$* , Math. Comp. **187** (1989), 311–326. MR **90a**:30075
6. J. M. Borwein and P. B. Borwein, *A cubic counterpart of Jacobi's identity and the AGM*, Trans. Amer. Math. Soc. **323** (1991), 691–701. MR **91e**:33012
7. J. M. Borwein, P. B. Borwein, and F. G. Garvan, *Hypergeometric analogues of the arithmetic-geometric mean iteration*, Constr. Approx. **9**, no. 4, (1993), 509–523. MR **95b**:33002
8. J. M. Borwein and F. G. Garvan, *Approximations to  $\pi$  via the Dedekind eta function*, CMS Conference Proceedings on Organic Mathematics (Burnaby, BC), vol. 20, Amer. Math. Soc., Providence, RI, 1997, pp. 89–115. MR **98j**:11030
9. H. H. Chan, *On Ramanujan's cubic transformation formula for  ${}_2F_1(1/3, 2/3; 1; z)$* , Math. Proc. Cambridge Philos. Soc. **124** (1998), 193–204. MR **99f**:11054
10. H. H. Chan, *Ramanujan's elliptic functions to alternative bases and approximations to  $\pi$* , Number Theory for the Millennium, Proc. Millennium Conf. Number Theory (Urbana, IL 2000) (M. A. Bennette et al., eds.), A. K. Peters, Boston, 2002, to appear.
11. H. H. Chan, K. S. Chua and P. Solé, *7-modular lattices and septic base Jacobi identity*, J. Number Theory, to appear.
12. H. H. Chan and Y. L. Ong, *On Eisenstein series and  $\sum_{m,n=-\infty}^{\infty} q^{m^2+mn+n^2}$* , Proc. Amer. Math. Soc. **127** (1999), 1735–1744. MR **99i**:11029
13. S. H. Chan, *Private Communication*.
14. B. Schoeneberg, *Elliptic modular functions*, Springer-Verlag, New York, 1974. MR **54**:236

15. G. N. Watson and E. T. Whittaker, *A course of modern analysis (fourth edition)*, Cambridge University Press, New York, 1992. MR **97k**:01072

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 117543,  
REPUBLIC OF SINGAPORE

*E-mail address:* `chanhh@math.nus.edu.sg`

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 117543,  
REPUBLIC OF SINGAPORE

*E-mail address:* `matv2@nus.edu.sg`

CNRS-I3S, ESSI, ROUTE DES COLLES, 06 903 SOPHIA ANTIPOLIS, FRANCE

*E-mail address:* `ps@essi.fr`