

HEAT KERNELS ON METRIC MEASURE SPACES AND AN APPLICATION TO SEMILINEAR ELLIPTIC EQUATIONS

ALEXANDER GRIGOR'YAN, JIAXIN HU, AND KA-SING LAU

ABSTRACT. We consider a metric measure space (M, d, μ) and a *heat kernel* $p_t(x, y)$ on M satisfying certain upper and lower estimates, which depend on two parameters α and β . We show that under additional mild assumptions, these parameters are determined by the intrinsic properties of the space (M, d, μ) . Namely, α is the *Hausdorff dimension* of this space, whereas β , called the *walk dimension*, is determined via the properties of the family of *Besov spaces* $W^{\sigma, 2}$ on M . Moreover, the parameters α and β are related by the inequalities $2 \leq \beta \leq \alpha + 1$.

We prove also the embedding theorems for the space $W^{\beta/2, 2}$, and use them to obtain the existence results for weak solutions to semilinear elliptic equations on M of the form

$$-\mathcal{L}u + f(x, u) = g(x),$$

where \mathcal{L} is the generator of the semigroup associated with p_t .

The framework in this paper is applicable for a large class of fractal domains, including the generalized Sierpiński carpet in \mathbb{R}^n .

1. INTRODUCTION

We say that a triple (M, d, μ) is a *metric measure space* if (M, d) is a non-empty metric space and μ is a Borel measure on M . Given a metric measure space (M, d, μ) , a family $\{p_t\}_{t>0}$ of non-negative measurable functions $p_t(x, y)$ on $M \times M$ is called a *heat kernel* or a *transition density* if the following conditions are satisfied, for all $x, y \in M$ and $s, t > 0$:

- (1) Symmetry: $p_t(x, y) = p_t(y, x)$.
- (2) Normalization (or stochastic completeness):

$$\int_M p_t(x, y) d\mu(y) = 1.$$

- (3) Semigroup property:

$$p_{s+t}(x, y) = \int_M p_s(x, z) p_t(z, y) d\mu(z).$$

Received by the editors July 23, 2002.

2000 *Mathematics Subject Classification*. Primary 60J35; Secondary 28A80, 35J60.

The first author was partially supported by a visiting grant of the Institute of Mathematical Sciences of CUHK (the Chinese University of Hong Kong). The second author was supported by a Postdoctoral Fellowship from CUHK. The third author was partially supported by a HKRGC grant at CUHK.

(4) Identity approximation: for any $f \in L^2(M, \mu)$,

$$\int_M p_t(x, y) f(y) d\mu(y) \xrightarrow{L^2(M, \mu)} f(x) \quad \text{as } t \rightarrow 0+.$$

For example, the classical Gauss-Weierstrass function in \mathbb{R}^n ,

$$(1.1) \quad p_t(x, y) = \frac{1}{(4\pi t)^{n/2}} \exp\left(-\frac{|x-y|^2}{4t}\right),$$

satisfies this definition. Another elementary example is the Cauchy-Poisson kernel in \mathbb{R}^n :

$$(1.2) \quad p_t(x, y) = \frac{C_n}{t^n} \left(1 + \frac{|x-y|^2}{t^2}\right)^{-\frac{n+1}{2}},$$

where $C_n = \Gamma\left(\frac{n+1}{2}\right) / \pi^{(n+1)/2}$.

Any heat kernel gives rise to the *heat semigroup* $\{T_t\}_{t>0}$ where T_t is an operator in $L^2(M, \mu)$ defined by

$$(1.3) \quad T_t u(x) = \int_M p_t(x, y) u(y) d\mu(y).$$

The above properties of p_t imply that T_t is a bounded selfadjoint operator in $L^2(M, \mu)$; moreover, $\{T_t\}_{t>0}$ is a strongly continuous, positivity-preserving, contraction semigroup in $L^2(M, \mu)$. Another way of constructing such a semigroup is to set

$$(1.4) \quad T_t = \exp(t\mathcal{L}),$$

where \mathcal{L} is a non-positive-definite selfadjoint operator in $L^2(M, \mu)$ satisfying in addition the Markov property. Typically, $-\mathcal{L}$ arises as the generator of a Dirichlet form. It is not always the case that the semigroup $\{T_t\}$ defined by (1.4) possesses an integral kernel. If it does, then the integral kernel will be a heat kernel in the above sense (although some additional restrictions are needed to ensure the normalization condition).

In this note we would like to adopt the axiomatic approach to heat kernels, which to some extent is opposite to the above scheme. Namely, we will assume that a heat kernel is defined on a metric measure space, and show that this implies many interesting consequences for analysis on such a space. A similar approach was used by M. Barlow [3] and K. Pietruska-Paħuba [23], although in their works a heat kernel was assumed to be the transition density of a diffusion process on M , and in [23] the underlying space M was a subset of \mathbb{R}^n .

Let a heat kernel p_t on (M, d, μ) satisfy the following two-sided estimate, for μ -almost all $x, y \in M$ and all $t \in (0, \infty)$:

$$(1.5) \quad \frac{1}{t^{\alpha/\beta}} \Phi_1\left(\frac{d(x, y)}{t^{1/\beta}}\right) \leq p_t(x, y) \leq \frac{1}{t^{\alpha/\beta}} \Phi_2\left(\frac{d(x, y)}{t^{1/\beta}}\right),$$

where α, β are positive constants, and Φ_1 and Φ_2 are non-negative monotone decreasing functions on $[0, +\infty)$.

For example, the Gauss-Weierstrass heat kernel (1.1) satisfies (1.5) with $\alpha = n$, $\beta = 2$, and

$$\Phi_1(s) = \Phi_2(s) = \frac{1}{(4\pi)^{n/2}} \exp\left(-\frac{s^2}{4}\right).$$

The Cauchy-Poisson heat kernel (1.2) satisfies (1.5) with $\alpha = n$, $\beta = 1$, and

$$\Phi_1(s) = \Phi_2(s) = C_n (1 + s^2)^{-\frac{n+1}{2}}.$$

Note that the Gauss-Weierstrass heat kernel is generated by the classical Laplace operator in \mathbb{R}^n ,

$$-\Delta = -\sum_{k=1}^n \frac{\partial^2}{\partial x_k^2},$$

whereas the Cauchy-Poisson heat kernel is generated by the operator $-(-\Delta)^{1/2}$. More generally, for any $0 < \beta < 2$ the operator $-(-\Delta)^{\beta/2}$ generates a *symmetric stable process* of index β whose heat kernel satisfies the estimate (1.5) with $\alpha = n$ and the following functions Φ_1 and Φ_2 :

$$(1.6) \quad \Phi_i(s) = c_i (1 + s^2)^{-\frac{\alpha+\beta}{2}},$$

where c_i is a positive constant, $i = 1, 2$.

The development of analysis on fractals has brought plenty of examples of heat kernels satisfying (1.5) with functions Φ_1 and Φ_2 of the form

$$(1.7) \quad \Phi_i(s) = c'_i \exp(-c''_i s^\gamma),$$

where $\gamma > 0$, and c'_i, c''_i are positive constants. In these examples the parameter β is typically larger than 2.

The nature of the parameters α and β is of great interest. Although originally they are defined through the heat kernel, a posteriori they happen to be the invariants of the space (M, d, μ) itself, provided the function Φ_2 decays at ∞ sufficiently fast. The parameter α happens to be the *Hausdorff dimension* of M (see also [10]). The nature of the parameter β is more complicated. We will call it the *walk dimension* of the heat kernel p_t . This terminology is motivated by the following observation: if the heat kernel p_t is the transition density of a Markov process X_t on M , then (1.5) implies (under reasonable assumptions about Φ_1 and Φ_2) that the mean time t needed for the process X_t to move away to a distance r from the origin is of the order r^β (see, for example, [3, Lemma 3.9]).

In this paper, we adopt an analytic approach, which does not depend on the existence of the process X_t . Following [18], we introduce on (M, d, μ) the family $\{W^{\sigma,2}\}_{\sigma>0}$ of *Besov spaces*, which generalizes the Sobolev space $W^{1,2}$ in \mathbb{R}^n (see Section 4; note that in the notation of [18], $W^{\sigma,2} = \text{Lip}(\sigma, 2, \infty)$). The space $W^{\sigma,2}$ decreases with σ ; we define the *critical exponent* β^* of the family $W^{\sigma,2}$ by

$$\beta^* = 2 \sup \{ \sigma : \dim W^{\sigma,2} = \infty \}.$$

Clearly, the exponent β^* is an invariant of the space (M, d, μ) . For $M = \mathbb{R}^n$ and μ the Lebesgue measure, we have $\beta^* = 2$.

To describe our results, assume that $\Phi_1(1) > 0$ and consider the following three hypotheses for Φ_2 :

$$(H_0) \quad \int_0^\infty s^\alpha \Phi_2(s) \frac{ds}{s} < \infty,$$

$$(H_1) \quad \int_0^\infty s^{\alpha+\beta} \Phi_2(s) \frac{ds}{s} < \infty,$$

$$(H_2) \quad \int_0^\infty s^{\alpha+\beta+\varepsilon} \Phi_2(s) \frac{ds}{s} < \infty,$$

for some $\varepsilon > 0$. We have proved the following.

Let (H_0) hold. Then the measure of any metric ball $B(x, r) := \{y : d(x, y) < r\}$ in M satisfies the estimates

$$(1.8) \quad C^{-1}r^\alpha \leq \mu(B(x, r)) \leq Cr^\alpha,$$

for all $x \in M$ and $r > 0$, where C is a positive constant (Theorem 3.2). In particular, this implies that α is intrinsically determined by (M, d, μ) ; moreover, α is the Hausdorff dimension of M .

The walk dimension β satisfies the inequality $\beta \leq \beta^*$ where β^* is the critical exponent of the family of Besov spaces defined above (Corollary 4.3). If, in addition, (M, d) satisfies a certain *chain condition* (Definition 3.4), then $\beta \leq \alpha + 1$ (Theorem 4.8(ii)).

Let (H_1) hold. Then the domain $\mathcal{E}(\mathcal{D})$ of the Dirichlet form \mathcal{E} naturally associated with the heat kernel coincides with the Besov space $W^{\beta/2, 2}$ (Theorem 4.2). If $\alpha > \beta$, then the following embedding takes place:

$$W^{\beta/2, 2}(M, \mu) \hookrightarrow L^{2^*}(M, \mu)$$

where

$$(1.9) \quad 2^* := \frac{2\alpha}{\alpha - \beta},$$

and the case $\alpha \leq \beta$ is also discussed (Theorem 4.11).

Let (H_2) hold. Then $\beta = \beta^*$ (Theorem 4.6) and $\beta \geq 2$ (Theorem 4.8(i)).

Hence, under the hypothesis (H_2) both parameters α and β are uniquely determined by the underlying space. Note that under the hypothesis (H_0) this is not the case, because, for example, both the Gauss-Weierstrass heat kernel (1.1) and the Cauchy-Poisson heat kernel (1.2) in \mathbb{R}^n satisfy (H_0) , whereas for the former $\beta = 2$ and for the latter $\beta = 1$ (although $\alpha = n$ in both cases).

Let us mention that the characterization of the domain of the Dirichlet form on the Sierpiński gasket in terms of Besov spaces, as well as the relation $\beta = \beta^*$, was first obtained by A. Jonsson [18]. K. Pietruska-Pałuba obtained the similar conclusion for *nested* fractals [22]. Observe that the condition (H_1) under which we prove the equivalence of $\mathcal{D}(\mathcal{E})$ and $W^{\beta/2, 2}$ is optimal. Indeed, for the heat kernel generated by the operator $-(-\Delta)^{\beta/2}$ in \mathbb{R}^n the function Φ_2 is given by (1.6), and for this function the hypothesis (H_1) breaks just on the borderline, because

$$\int_0^\infty \frac{s^{\alpha+\beta}}{(1+s^2)^{\frac{\alpha+\beta}{2}}} \frac{ds}{s} = \infty.$$

On the other hand, the domain of the energy form associated with the operator $-(-\Delta)^{\beta/2}$ is known to be another Besov space $\text{Lip}(\beta/2, 2, 2)$, which is smaller than $\text{Lip}(\beta/2, 2, \infty) = W^{\beta/2, 2}$ (see [2], [23], [25]).

M. Barlow studied in [3] heat kernels on geodesic metric spaces, assuming that a heat kernel p_t is the transition density of a diffusion process and that it satisfies (1.5) with functions Φ_1, Φ_2 of the type (1.7). Under these assumptions, he claimed that the walk dimension β of p_t is uniquely determined by the intrinsic properties of the underlying space (see [3, Theorem 3.21]). Under the additional assumption

that M is a subset of \mathbb{R}^n , this claim was proved in [23]. Our contribution is that we prove the uniqueness of β for a general metric measure space (M, d) and under a rather mild and nearly optimal condition (H_2) on the heat kernel (nor do we assume existence of a diffusion process associated with p_t).

The relationship between the parameters α and β , mentioned above, is summarized in Corollary 4.9 as follows: If (M, d) satisfies the chain condition, $\Phi_1(1) > 0$ and, for some $\varepsilon > 0$,

$$(1.10) \quad \int_0^\infty s^{2\alpha+\varepsilon} \Phi_2(s) ds < \infty,$$

then

$$(1.11) \quad 2 \leq \beta \leq \alpha + 1.$$

The inequalities (1.11) were stated by Barlow in [3, Theorem 3.20], under the same set of assumptions as [3, Theorem 3.21]. Under somewhat more restrictive hypotheses, A. Stós [25] proved that $2 \leq \beta \leq \alpha + 2$. In the setting of random walks in graphs, (1.11) was proved in [26] and [14], using techniques that are not available for general metric spaces. Our contribution is that we prove (1.11) under the hypothesis (1.10), which seems to be nearly optimal.

Barlow proved in [4] that for *every* pair α, β satisfying (1.11) there exists a random walk satisfying a discrete time version of (1.5) with these parameters. There is no doubt that the same is true for continuous time heat kernels. Hence, (1.11) is the *only* restriction on α and β .

In Section 5 we apply the embedding results to treat the following semilinear elliptic equation on M :

$$(1.12) \quad -\mathcal{L}u + f(x, u) = g(x),$$

where \mathcal{L} is the generator of the semigroup T_t (the equation (1.12) arises for example when investigating the potential u in porous or other irregular domains). We prove the existence and uniqueness results for weak solutions of (1.12), which in particular imply that, for all $q \geq p \geq 2^*$, the equation

$$-\mathcal{L}u + |u|^{q-2} u = g$$

has a unique weak solution $u \in \mathcal{D}(\mathcal{E}) \cap L^p \cap L^q$, for any $g \in L^{p'}$, where $p' = \frac{p}{p-1}$ (Theorem 5.4). The existence result relies on the compact embedding of the space $\mathcal{D}(\mathcal{E})$ into L^2_{loc} , which is true without any additional assumptions about Φ_1 and Φ_2 (Theorem 4.12).

Note that the classical existence results for equation (1.12) in \mathbb{R}^n , $n > 2$, depend on the critical parameter $2^* = \frac{2n}{n-2}$ (see [19]), which matches (1.9) since $\alpha = n$ and $\beta = 2$.

Notation. The letters C, c are used to denote positive constants whose values are unimportant but depend only on the hypotheses. The values of C, c may be different at different occurrences.

For two non-negative functions $f(s)$ and $g(s)$ defined on a set S , we write

$$f(s) \simeq g(s)$$

if there is a constant c such that for all $s \in S$,

$$c^{-1}g(s) \leq f(s) \leq cg(s).$$

2. SOME EXAMPLES

Let $l \geq 3$ be an integer and let $M_0 = [0, 1]^n$ ($n \geq 2$). We divide M_0 into l^n equal subcubes. Remove a symmetric pattern of subcubes from M_0 , and denote by M_1 what remains. Repeat the same procedure for each subcube in M_1 : divide each subcube into l^n equal parts and remove the same pattern from each subcube, and denote by M_2 what remains. Continuing this way infinitely, we obtain a sequence of sets $\{M_k\}$. Set

$$\widetilde{M} = \bigcap_{k=0}^{\infty} M_k$$

and define

$$M = \bigcup_{k=0}^{\infty} l^k \widetilde{M},$$

where we write $aK = \{ax : x \in K\}$ for a real number a and a set K .

The set M is called an unbounded *generalized Sierpiński carpet* (cf. [5]); see Figure 1, which corresponds to the case $n = 2$ and $l = 3$.




FIGURE 1. Generalized Sierpiński carpet

The distance d on M is set to be the Euclidean distance, and the measure μ is the Hausdorff measure of dimension α , where α is the Hausdorff dimension of M . For any generalized Sierpiński carpet, there exists a heat kernel satisfying the following estimate:

$$(2.1) \quad p_t(x, y) \simeq \frac{1}{t^{\alpha/\beta}} \exp \left(-c \left(\frac{d(x, y)^\beta}{t} \right)^{\frac{1}{\beta-1}} \right),$$

which is a particular case of (1.5) with the functions Φ_1, Φ_2 of the form

$$\Phi(s) = c \exp \left(-Cs^{\frac{\beta}{\beta-1}} \right)$$

(see [5]). There are also plenty of other fractals such that (2.1) holds; see, for example, [3], [11].

See also [13], [14], [20] for the heat kernel estimates in the setting of graphs or manifolds.

3. VOLUME OF BALLS

Definition 3.1. We say that a heat kernel p_t on a metric measure space (M, d, μ) satisfies the hypothesis (H_0) if p_t satisfies the estimate (1.5) with some positive constants α, β and non-negative decreasing functions Φ_1 and Φ_2 such that $\Phi_1(1) > 0$ and

$$(3.1) \quad \int_0^\infty s^\alpha \Phi_2(s) \frac{ds}{s} < \infty.$$

Denote by $B(x, r)$ the metric ball in M of radius r centered at the point $x \in M$; that is,

$$B(x, r) := \{y \in M : d(x, y) < r\}.$$

Theorem 3.2. *If a heat kernel p_t satisfies the hypothesis (H_0) , then for all $x \in M$ and $r > 0$,*

$$(3.2) \quad C^{-1}r^\alpha \leq \mu(B(x, r)) \leq Cr^\alpha.$$

Remark. The condition (3.1) is very mild. Indeed, if (3.2) is given, then (3.1) means that the upper bound function

$$y \mapsto t^{-\alpha/\beta} \Phi_2 \left(\frac{d(x, y)}{t^{1/\beta}} \right)$$

has a bounded integral over M uniformly in x and t , which resembles the normalization property of the heat kernel (cf. (3.7) below).

Note also that the estimate (3.2) can be true only for a single value of α , which is called the *exponent of the volume growth* and is determined by intrinsic properties of the space (M, d, μ) . Hence, under the hypothesis (H_0) the value of the parameter α in (1.5) is an invariant of the underlying space.

Proof. Fix $x \in M$ and prove first the upper bound

$$(3.3) \quad \mu(B(x, r)) \leq Cr^\alpha,$$

for all $r > 0$. Indeed, for any $t > 0$, we have

$$(3.4) \quad \int_{B(x, r)} p_t(x, y) d\mu(y) \leq \int_M p_t(x, y) d\mu(y) = 1,$$

whence

$$\mu(B(x, r)) \leq \left(\inf_{y \in B(x, r)} p_t(x, y) \right)^{-1}.$$

Taking $t = r^\beta$ and applying the lower bound in (1.5), we obtain

$$\inf_{y \in B(x, r)} p_t(x, y) \geq \frac{1}{t^{\alpha/\beta}} \Phi_1(1) = cr^{-\alpha},$$

whence (3.3) follows. An obvious modification of this argument allows us to replace the hypothesis $\Phi_1(1) > 0$ by $\Phi_1(s_0) > 0$ for some $s_0 > 0$.

Let us prove the opposite inequality

$$(3.5) \quad \mu(B(x, r)) \geq cr^\alpha.$$

We first show that the upper bounds in (1.5) and (3.3) imply that

$$(3.6) \quad \int_{M \setminus B(x,r)} p_t(x,y) d\mu(y) \leq \frac{1}{2}, \quad \text{for all } t \leq \varepsilon r^\beta,$$

provided $\varepsilon > 0$ is sufficiently small. Setting $r_k = 2^k r$ and using the monotonicity of Φ_2 and (3.3), we obtain

$$(3.7) \quad \begin{aligned} \int_{M \setminus B(x,r)} p_t(x,y) d\mu(y) &\leq \int_{M \setminus B(x,r)} t^{-\alpha/\beta} \Phi_2 \left(\frac{d(x,y)}{t^{1/\beta}} \right) d\mu(y) \\ &\leq \sum_{k=0}^{\infty} \int_{B(x,r_{k+1}) \setminus B(x,r_k)} t^{-\alpha/\beta} \Phi_2 \left(\frac{r_k}{t^{1/\beta}} \right) d\mu(y) \\ &\leq C \sum_{k=0}^{\infty} r_k^\alpha t^{-\alpha/\beta} \Phi_2 \left(\frac{r_k}{t^{1/\beta}} \right) \\ &= C \sum_{k=0}^{\infty} \left(\frac{2^k r}{t^{1/\beta}} \right)^\alpha \Phi_2 \left(\frac{2^k r}{t^{1/\beta}} \right) \\ &\leq C \int_{\frac{1}{2}r/t^{1/\beta}}^{\infty} s^\alpha \Phi_2(s) \frac{ds}{s}. \end{aligned}$$

The last line in (3.7) is proved as follows: setting $t_k = 2^k r/t^{1/\beta}$ for $k \geq -1$ and using the monotonicity of Φ_2 , we obtain

$$\begin{aligned} \int_{\frac{1}{2}r/t^{1/\beta}}^{\infty} s^\alpha \Phi_2(s) \frac{ds}{s} &= \sum_{k=0}^{\infty} \int_{t_{k-1}}^{t_k} s^{\alpha-1} \Phi_2(s) ds \\ &\geq \frac{1}{\alpha} \sum_{k=0}^{\infty} \Phi_2(t_k) (t_k^\alpha - t_{k-1}^\alpha) \\ &= c \sum_{k=0}^{\infty} \Phi_2(t_k) t_k^\alpha. \end{aligned}$$

Since by (3.1) the integral in (3.7) is convergent, its value can be made arbitrarily small provided r^β/t is large enough, whence (3.6) follows.

From the normalization property and (3.6), we conclude that for such r and t ,

$$(3.8) \quad \int_{B(x,r)} p_t(x,y) d\mu(y) \geq \frac{1}{2},$$

whence

$$\mu(B(x,r)) \geq \frac{1}{2} \left(\sup_{y \in B(x,r)} p_t(x,y) \right)^{-1}.$$

Finally, choosing $t := \varepsilon r^\beta$ and using the upper bound

$$p_t(x,y) \leq t^{-\alpha/\beta} \Phi_2(0) = Cr^{-\alpha},$$

we obtain (3.5). \square

Note that the method we have used in the proof of Theorem 3.2 is close to [14].

Corollary 3.3. *Let (M, d, μ) be a metric measure space, and p_t be a heat kernel on M satisfying (1.5). If $\Phi_1(1) > 0$, then $\text{diam}(M) = \infty$.*

Proof. Let us show that the upper bound in (1.5) implies $\mu(M) = \infty$. Indeed, fix a point $x_0 \in M$ and observe that the family of functions $u_t(x) = p_t(x, x_0)$ satisfies the following two conditions:

$$\|u_t\|_1 = 1$$

and

$$\|u_t\|_\infty \leq Ct^{-\alpha/\beta} \rightarrow 0 \quad \text{as } t \rightarrow \infty,$$

where $\|\cdot\|_q$ stands for the $L^q(M, \mu)$ norm. Hence, we obtain

$$\mu(M) \geq \frac{\|u_t\|_1}{\|u_t\|_\infty} \rightarrow \infty \quad \text{as } t \rightarrow \infty,$$

that is, $\mu(M) = \infty$.

On the other hand, the first part of the proof of Theorem 3.2, based solely on the hypothesis $\Phi_1(1) > 0$, says that the measure of any ball is finite. Hence, M is not contained in any ball, which ends the proof. \square

Under a mild additional hypothesis about the metric space (M, d) , we will obtain a self-improved lower bound for the heat kernel.

Definition 3.4. We say that a metric space (M, d) satisfies the *chain condition* if there exists a (large) constant C such that for any two points $x, y \in M$ and for any positive integer n there exists a sequence $\{x_i\}_{i=0}^n$ of points in M such that $x_0 = x$, $x_n = y$, and

$$(3.9) \quad d(x_i, x_{i+1}) \leq C \frac{d(x, y)}{n}, \quad \text{for all } i = 0, 1, \dots, n-1.$$

The sequence $\{x_i\}_{i=0}^n$ is referred to as a *chain* connecting x and y .

For example, the chain condition is satisfied if (M, d) is a *length space*, that is, if the distance $d(x, y)$ is defined as the infimum of the length of all continuous curves connecting x and y , with a proper definition of length. On the other hand, the chain condition is not satisfied if M is a locally finite graph, and d is the graph distance.

Corollary 3.5. *Under the hypotheses of Theorem 3.2, assume in addition that $\beta > 1$ and that (M, d) satisfies the chain condition. Then the following inequality holds, for all $x, y \in M$ and $t > 0$:*

$$(3.10) \quad p_t(x, y) \geq \frac{c}{t^{\alpha/\beta}} \exp\left(-C \left(\frac{d(x, y)^\beta}{t}\right)^{\frac{1}{\beta-1}}\right).$$

Remark. The assumption $\beta > 1$ is not very restrictive. In fact, as we will see in Section 4.5, (1.5) implies $\beta \geq 2$ provided Φ_2 satisfies a condition slightly stronger than (3.1). On the other hand, the condition $\beta > 1$ does not follow from the hypotheses of Theorem 3.2, because the Cauchy-Poisson heat kernel (1.2) satisfies them whereas for this heat kernel $\beta = 1$.

Finally, let us observe that (3.10) matches the lower bound in (2.1).

Proof. By iterating the semigroup identity, we obtain, for any positive integer n and real $r > 0$,

$$\begin{aligned}
 p_t(x, y) &= \int \dots \int p_{\frac{t}{n}}(x, z_1) p_{\frac{t}{n}}(z_1, z_2) \dots p_{\frac{t}{n}}(z_{n-1}, y) d\mu(z_1) \dots d\mu(z_{n-1}) \\
 (3.11) \quad &\geq \int_{B(x_1, r)} \dots \int_{B(x_{n-1}, r)} p_{\frac{t}{n}}(x, z_1) p_{\frac{t}{n}}(z_1, z_2) \dots p_{\frac{t}{n}}(z_{n-1}, y) d\mu(z_{n-1}) \dots d\mu(z_1),
 \end{aligned}$$

where $\{x_i\}_{i=0}^n$ is a chain connecting x and y .

Denote for simplicity $z_0 = x$ and $z_n = y$. Since $z_i \in B(x_i, r)$, $0 \leq i \leq n - 1$, we obtain by the triangle inequality and (3.9),

$$d(z_i, z_{i+1}) \leq d(x_i, x_{i+1}) + 2r \leq C \frac{d(x, y)}{n} + 2r.$$

Choosing

$$(3.12) \quad r = \frac{d(x, y)}{n}$$

and using the lower bound in (1.5), we obtain

$$(3.13) \quad p_{\frac{t}{n}}(z_i, z_{i+1}) \geq \frac{1}{(t/n)^{\alpha/\beta}} \Phi_1 \left(\frac{d(z_i, z_{i+1})}{(t/n)^{1/\beta}} \right) \geq \frac{n^{\alpha/\beta}}{t^{\alpha/\beta}} \Phi_1 \left(\frac{d(x, y)}{(ctn^{\beta-1})^{1/\beta}} \right),$$

where $c = (C + 2)^{-\beta}$.

If $d(x, y)^\beta \leq t$, then (3.10) follows immediately from (1.5). Assume in the sequel that $d(x, y)^\beta > t$. Choosing n to be the least positive integer satisfying the inequality

$$ctn^{\beta-1} \geq d(x, y)^\beta,$$

we obtain from (3.13),

$$p_{\frac{t}{n}}(z_i, z_{i+1}) \geq \frac{n^{\alpha/\beta}}{t^{\alpha/\beta}} \Phi_1(1).$$

Hence, (3.11), (3.2), and (3.12) imply

$$\begin{aligned}
 p_t(x, y) &\geq \left(c \frac{n^{\alpha/\beta}}{t^{\alpha/\beta}} \right)^n \prod_{i=1}^{n-1} \mu(B(x_i, r)) \\
 &\geq \left(c \frac{n^{\alpha/\beta}}{t^{\alpha/\beta}} \right)^n (cr^\alpha)^{n-1} \\
 (3.14) \quad &= c \left(\frac{n}{t} \right)^{\alpha/\beta} \left(c \frac{d(x, y)^\beta}{n^{\beta-1}t} \right)^{(n-1)\alpha/\beta}.
 \end{aligned}$$

Since $n^{\beta-1} \simeq d(x, y)^\beta / t$, we see that

$$\left(c \frac{d(x, y)^\beta}{n^{\beta-1}t} \right)^{(n-1)\alpha/\beta} \geq \varepsilon^n$$

for some (small) positive constant ε , which together with (3.14) yields

$$p_t(x, y) \geq c \left(\frac{n}{t} \right)^{\alpha/\beta} \varepsilon^n \geq \frac{c}{t^{\alpha/\beta}} \exp(-Cn) \geq \frac{c}{t^{\alpha/\beta}} \exp \left(-C \left(\frac{d(x, y)^\beta}{t} \right)^{\frac{1}{\beta-1}} \right).$$

□

4. BESOV SPACES

Let (M, d, μ) be a metric measure space. For any $q \in [1, +\infty]$, set $L^q = L^q(M, \mu)$ and

$$\|u\|_q = \|u\|_{L^q(M, \mu)}.$$

For any $\sigma > 0$, define a non-negative functional $W_\sigma(u)$ on measurable functions on M by

$$(4.1) \quad W_\sigma(u) := \sup_{0 < r < 1} r^{-2\sigma} \int_M \left[\frac{1}{\mu(B(x, r))} \int_{B(x, r)} |u(y) - u(x)|^2 d\mu(y) \right] d\mu(x).$$

In particular, if the condition (3.2) is satisfied, then for any $\beta > 0$,

$$(4.2) \quad W_{\beta/2}(u) \simeq \sup_{0 < r < 1} r^{-\alpha-\beta} \int_M \left[\int_{B(x, r)} |u(y) - u(x)|^2 d\mu(y) \right] d\mu(x).$$

Define the space $W^{\sigma, 2}$ as follows:

$$W^{\sigma, 2} = W^{\sigma, 2}(M, d, \mu) := \{u \in L^2 : W_\sigma(u) < \infty\}.$$

It is easy to see that $W_\sigma^{1/2}$ is a semi-norm in $W^{\sigma, 2}$, and $W^{\sigma, 2}$ is a Banach space with the norm

$$\|u\|_{\sigma, 2} := \|u\|_2 + W_\sigma(u)^{1/2}$$

(see for example [18]). The space $W^{\sigma, 2}$ is one of the family of *Besov spaces*, and it is similar to the space that was denoted by $\text{Lip}(\sigma, 2, \infty)$ in [18].

4.1. The Laplace operator. Define by (1.3) the semigroup $\{T_t\}_{t>0}$ in L^2 , and consider its *infinitesimal generator* \mathcal{L} defined by

$$(4.3) \quad \mathcal{L}u := \lim_{t \rightarrow 0} \frac{T_t u - u}{t},$$

where the limit is understood in the L^2 -norm. It is natural to refer to \mathcal{L} as the *Laplace operator* of the heat kernel p_t . The *domain* $\text{dom}(\mathcal{L})$ of the operator \mathcal{L} is the space of functions $u \in L^2$ for which the limit in (4.3) exists. Since $\{T_t\}$ is a strongly continuous contraction semigroup in L^2 , by the Hille–Yosida theorem $\text{dom}(\mathcal{L})$ is dense in L^2 ; furthermore, \mathcal{L} is a selfadjoint, non-positive-definite operator, which follows from the fact that T_t is selfadjoint and contractive (see for example [8], [12], [16], or [29, Theorem 1, p. 237]).

4.2. The Dirichlet form. For any $t > 0$, define the quadratic form \mathcal{E}_t on L^2 by

$$(4.4) \quad \mathcal{E}_t[u] := \left(\frac{u - T_t u}{t}, u \right),$$

where (\cdot, \cdot) is the inner product in L^2 . An easy computation shows that \mathcal{E}_t can be equivalently defined by

$$(4.5) \quad \mathcal{E}_t[u] = \frac{1}{2t} \int_M \int_M (u(x) - u(y))^2 p_t(x, y) d\mu(y) d\mu(x).$$

In terms of the spectral resolution $\{E_\lambda\}$ of the operator $-\mathcal{L}$, \mathcal{E}_t can be expressed as follows:

$$\mathcal{E}_t[u] = \int_0^\infty \frac{1 - e^{-t\lambda}}{t} d\|E_\lambda u\|_2^2,$$

which implies that $\mathcal{E}_t[u]$ is decreasing in t (see also [7]).

Let us define a quadratic form \mathcal{E} by

$$(4.6) \quad \mathcal{E}[u] := \lim_{t \rightarrow 0^+} \mathcal{E}_t[u]$$

(where the limit may be $+\infty$ since $\mathcal{E}[u] \geq \mathcal{E}_t[u]$) and its domain $\mathcal{D}(\mathcal{E})$ by

$$\mathcal{D}(\mathcal{E}) := \{u \in L^2 : \mathcal{E}[u] < \infty\}.$$

By a standard procedure, the quadratic form $\mathcal{E}[u]$ extends to the bilinear form $\mathcal{E}(u, v)$.

From (4.5) it follows that \mathcal{E} is non-negative definite. The domain $\mathcal{D}(\mathcal{E})$ is dense in L^2 because $\mathcal{D}(\mathcal{E})$ contains $\text{dom}(\mathcal{L})$. Moreover, the form \mathcal{E} is closed; that is, the space $\mathcal{D}(\mathcal{E})$ is a Hilbert space with the inner product

$$(4.7) \quad [u, v] := (u, v) + \mathcal{E}(u, v).$$

Clearly (4.3) and (4.4) imply that

$$(4.8) \quad \mathcal{E}(u, v) = (-\mathcal{L}u, v),$$

for all $u \in \text{dom}(\mathcal{L})$ and $v \in \mathcal{D}(\mathcal{E})$.

It is easy to see from (1.3) and the definition of a heat kernel that the semigroup $\{T_t\}$ is *Markovian*, that is, $0 \leq u \leq 1$ implies $0 \leq T_t u \leq 1$. By [12, Theorem 1.4.1], the form \mathcal{E} satisfies the Markov property, that is, $u \in \mathcal{D}(\mathcal{E})$ implies $v := \min(u_+, 1) \in \mathcal{D}(\mathcal{E})$ and $\mathcal{E}[v] \leq \mathcal{E}[u]$. Hence, \mathcal{E} is a *Dirichlet form*.

4.3. Besov space as the domain of the Dirichlet form.

Definition 4.1. We say that a heat kernel p_t on a metric measure space (M, d, μ) satisfies the hypothesis (H_1) if p_t satisfies the estimate (1.5) with some positive constants α, β and non-negative decreasing functions Φ_1 and Φ_2 such that $\Phi_1(1) > 0$ and

$$(4.9) \quad \int_0^\infty s^{\alpha+\beta} \Phi_2(s) \frac{ds}{s} < \infty.$$

The following theorem identifies the domain $\mathcal{D}(\mathcal{E})$ of the Dirichlet form with a Besov space.

Theorem 4.2. *Let p_t be a heat kernel on (M, d, μ) satisfying the hypothesis (H_1) , and let \mathcal{E} be the Dirichlet form defined by (4.5) and (4.6). Then*

$$\mathcal{D}(\mathcal{E}) = W^{\beta/2, 2}$$

and for any $u \in \mathcal{D}(\mathcal{E})$,

$$(4.10) \quad \mathcal{E}[u] \simeq W_{\beta/2}(u).$$

Proof. Since the expressions $\mathcal{E}[u]$ and $W_{\beta/2}(u)$ are defined for all $u \in L^2$, it suffices to show that (4.10) holds for all $u \in L^2$ (allowing infinite values for both sides). Note that the hypotheses of Theorem 3.2 hold, so that we have the estimate (3.2) of the volumes of balls.

We first prove that

$$(4.11) \quad \mathcal{E}[u] \geq c W_{\beta/2}(u),$$

using the approach of [23]. Using the lower bound in (1.5) and the monotonicity of Φ_1 , we obtain from (4.5) that for any $r > 0$ and $t = r^\beta$,

$$\begin{aligned} \mathcal{E}[u] &\geq \frac{1}{2t} \int_M \int_{B(x,r)} (u(x) - u(y))^2 p_t(x, y) d\mu(y) d\mu(x) \\ &\geq \frac{1}{2} \left(\frac{1}{t}\right)^{\alpha/\beta+1} \Phi_1\left(\frac{r}{t^{1/\beta}}\right) \int_M \int_{B(x,r)} (u(x) - u(y))^2 d\mu(y) d\mu(x) \\ &= \frac{\Phi_1(1)}{2} r^{-(\alpha+\beta)} \int_M \int_{B(x,r)} (u(x) - u(y))^2 d\mu(y) d\mu(x). \end{aligned}$$

Together with (4.2), this implies (4.11).

Let us now prove the opposite inequality, that is,

$$\mathcal{E}[u] \leq C W_{\beta/2}(u).$$

For any $t > 0$ we have

$$(4.12) \quad \mathcal{E}_t[u] = \frac{1}{2t} \int_M \int_M (u(x) - u(y))^2 p_t(x, y) d\mu(y) d\mu(x) = A(t) + B(t)$$

where

$$(4.13) \quad A(t) = \frac{1}{2t} \int_M \int_{M \setminus B(x,1)} (u(x) - u(y))^2 p_t(x, y) d\mu(y) d\mu(x),$$

$$(4.14) \quad B(t) = \frac{1}{2t} \int_M \int_{B(x,1)} (u(x) - u(y))^2 p_t(x, y) d\mu(y) d\mu(x).$$

To estimate $A(t)$ let us observe that by (3.7),

$$(4.15) \quad \int_{M \setminus B(x,1)} p_t(x, y) d\mu(y) \leq C \int_{\frac{1}{2t^{1/\beta}}}^{\infty} s^\alpha \Phi_2(s) \frac{ds}{s} \leq C t \int_{\frac{1}{2t^{1/\beta}}}^{\infty} s^{\alpha+\beta} \Phi_2(s) \frac{ds}{s},$$

whence by (4.9),

$$(4.16) \quad \int_{M \setminus B(x,1)} p_t(x, y) d\mu(y) = o(t) \quad \text{as } t \rightarrow 0+ \text{ uniformly in } x \in M.$$

Therefore, applying the elementary inequality $(a - b)^2 \leq 2(a^2 + b^2)$ in (4.13) and using (4.16), we obtain that for small enough $t > 0$,

$$\begin{aligned} A(t) &\leq \frac{1}{t} \int_M \int_{M \setminus B(x,1)} (u(x)^2 + u(y)^2) p_t(x, y) d\mu(y) d\mu(x) \\ &= \frac{2}{t} \int_M u(x)^2 \left(\int_{M \setminus B(x,1)} p_t(x, y) d\mu(y) \right) d\mu(x) \\ &= o(1) \|u\|_2^2, \end{aligned}$$

whence

$$(4.17) \quad \lim_{t \rightarrow 0+} A(t) = 0.$$

The quantity $B(t)$ is estimated as follows using (1.5), (4.2), and (4.9):

$$\begin{aligned}
 B(t) &= \frac{1}{2t} \sum_{k=1}^{\infty} \int_M \int_{B(x, 2^{-(k-1)}) \setminus B(x, 2^{-k})} (u(x) - u(y))^2 p_t(x, y) d\mu(y) d\mu(x) \\
 &\leq \frac{1}{2} \sum_{k=1}^{\infty} \left(\frac{1}{t}\right)^{\alpha/\beta+1} \Phi_2\left(\frac{2^{-k}}{t^{1/\beta}}\right) \int_M \int_{B(x, 2^{-(k-1)})} (u(x) - u(y))^2 d\mu(y) d\mu(x) \\
 &\leq C W_{\beta/2}(u) \sum_{k=1}^{\infty} \left(\frac{2^{-k}}{t^{1/\beta}}\right)^{\alpha+\beta} \Phi_2\left(\frac{2^{-k}}{t^{1/\beta}}\right) \\
 &\leq C W_{\beta/2}(u) \int_0^{\infty} s^{\alpha+\beta} \Phi_2(s) \frac{ds}{s} \\
 (4.18) \quad &\leq C W_{\beta/2}(u).
 \end{aligned}$$

It follows from (4.12), (4.17) and (4.18) that

$$(4.19) \quad \mathcal{E}[u] = \lim_{t \rightarrow 0^+} \mathcal{E}_t[u] \leq C W_{\beta/2}(u),$$

which finishes the proof. \square

4.4. Intrinsic characterization of the walk dimension. As one can see from the above proof of Theorem 4.2, the inclusion $\mathcal{D}(\mathcal{E}) \subset W^{\beta/2, 2}$ requires only the lower estimates in (1.5) and (3.2) (and the opposite inclusion requires only the upper estimates in (1.5) and (3.2)). This observation allows us to prove the following statement.

Corollary 4.3. *If a heat kernel satisfies the hypotheses (H_0) (cf. Theorem 3.2), then $\dim W^{\beta/2, 2} = \infty$.*

Proof. Indeed, by Theorem 3.2 the hypothesis (H_0) implies (3.2), and by the above remark we obtain $\mathcal{D}(\mathcal{E}) \subset W^{\beta/2, 2}$. On the other hand, $\mathcal{D}(\mathcal{E})$ is always dense in L^2 , whereas by Corollary 3.3 we have $\dim L^2 = \infty$, whence the claim follows. \square

Definition 4.4. Let us set

$$\beta^* := 2 \sup \{ \sigma : \dim W^{\sigma, 2} = \infty \}$$

and call β^* the *critical exponent* of the family $W^{\sigma, 2}$ of the Besov spaces in (M, d, μ) .

Note that the value of β^* is an intrinsic property of the space (M, d, μ) , which is defined independently of any heat kernel. For example, for \mathbb{R}^n we have $\beta^* = 2$. In terms of the critical exponent, Corollary 4.3 can be stated as follows: if a heat kernel satisfies (H_0) , then

$$\beta \leq \beta^*,$$

where β is the walk dimension of the heat kernel. In general, it may happen that $\beta < \beta^*$, as one can see from the example of the Cauchy-Poisson heat kernel in \mathbb{R}^n where $\beta = 1$ and $\beta^* = 2$. A theorem below provides conditions to ensure $\beta = \beta^*$ (see also [17]).

Definition 4.5. We say that a heat kernel p_t on a metric measure space (M, d, μ) satisfies the hypothesis (H_2) if p_t satisfies the estimate (1.5) with some positive

constants α, β and non-negative decreasing functions Φ_1 and Φ_2 such that $\Phi_1(1) > 0$ and, for some $\varepsilon > 0$,

$$(4.20) \quad \int_0^\infty s^{\alpha+\beta+\varepsilon} \Phi_2(s) \frac{ds}{s} < \infty.$$

Note that the hypothesis (H_2) is stronger than (H_1) , and (H_1) is stronger than (H_0) , but still (H_2) is satisfied in most applications. For example, every function of the form

$$(4.21) \quad \Phi_2(s) = \exp(-cs^\gamma)$$

satisfies (H_2) provided $c > 0$ and $\gamma > 0$.

Theorem 4.6. *If p_t is a heat kernel satisfying the hypothesis (H_2) , then for any $\sigma > \beta/2$ the space $W^{\sigma,2}$ contains only zero. Consequently, $\beta = \beta^*$.*

Remark. As follows from Corollary 4.3, the conclusion of Theorem 4.6 remains true if β in (4.20) is replaced by β^* , which makes the hypotheses of Theorem 4.6 independent of β .

Proof. Let us explain why $\beta = \beta^*$ follows from the first claim. Indeed, $\dim W^{\sigma,2} = 0$ implies that $\sigma \geq \beta^*/2$, and since this is true for any $\sigma > \beta/2$, we obtain $\beta \geq \beta^*$. Since the opposite inequality holds by Corollary 4.3, we conclude that $\beta = \beta^*$.

The proof of the first claim is similar to that of Theorem 4.2. Fix $\sigma > \beta/2$ and a function $u \in W^{\sigma,2}$, and show that $\mathcal{E}[u] = 0$. Note that if (4.20) holds for some $\varepsilon > 0$, it also holds for a smaller value of ε . Thus we may assume ε to be so small that

$$(4.22) \quad 2\sigma - \beta - \varepsilon > 0.$$

Again we use the decomposition $\mathcal{E}_t[u] = A(t) + B(t)$, where $A(t)$ and $B(t)$ are defined in (4.13) and (4.14). Similar to (4.18) but using W_σ instead of $W_{\beta/2}$, we estimate $B(t)$ and obtain by (4.1), (4.22), and (4.20),

$$\begin{aligned} & t^{-\varepsilon/\beta} B(t) \\ & \leq \frac{1}{2} \sum_{k=1}^{\infty} \left(\frac{1}{t}\right)^{(\alpha+\varepsilon)/\beta+1} \Phi_2\left(\frac{2^{-k}}{t^{1/\beta}}\right) \int_M \int_{B(x, 2^{-(k-1)})} (u(x) - u(y))^2 d\mu(y) d\mu(x) \\ & \leq C W_\sigma(u) \sum_{k=1}^{\infty} 2^{-k(2\sigma+\alpha)} \left(\frac{1}{t}\right)^{(\alpha+\varepsilon)/\beta+1} \Phi_2\left(\frac{2^{-k}}{t^{1/\beta}}\right) \\ & = C W_\sigma(u) \sum_{k=1}^{\infty} 2^{-k(2\sigma-\beta-\varepsilon)} \left(\frac{2^{-k}}{t^{1/\beta}}\right)^{\alpha+\beta+\varepsilon} \Phi_2\left(\frac{2^{-k}}{t^{1/\beta}}\right) \\ & \leq C W_\sigma(u) \int_0^\infty s^{\alpha+\beta+\varepsilon} \Phi_2(s) \frac{ds}{s} \\ & \leq C W_\sigma(u). \end{aligned}$$

Together with (4.12) and (4.17) this yields

$$\mathcal{E}_t[u] \leq A(t) + C t^{\varepsilon/\beta} W_\sigma(u) \rightarrow 0 \quad \text{as } t \rightarrow 0,$$

whence

$$\mathcal{E}[u] = \lim_{t \rightarrow 0} \mathcal{E}_t[u] = 0.$$

Since $\mathcal{E}_t[u] \leq \mathcal{E}[u]$, this implies in return that $\mathcal{E}_t[u] \equiv 0$ for all $t > 0$. On the other hand, it follows from (4.5) and the lower bound in (1.5) that

$$\mathcal{E}_t[u] \geq \frac{1}{2t^{\alpha/\beta+1}} \Phi_1(1) \iint_{\{x,y: d(x,y) \leq t^{1/\beta}\}} (u(x) - u(y))^2 d\mu(y) d\mu(x),$$

which yields $u(x) = u(y)$ for μ -almost all x, y such that $d(x, y) \leq t^{1/\beta}$. Since t is arbitrary, we conclude that u is a constant function. Since $\mu(M) = \infty$ (see Corollary 3.3), we obtain $u \equiv 0$. \square

Corollary 4.7. *If a heat kernel p_t satisfies (H_2) , then the values of the parameters α and β in (1.5) are invariants of the space (M, d, μ) . In particular, if there are two heat kernels $p_t^{(1)}$ and $p_t^{(2)}$ on (M, d, μ) both satisfying (H_2) with parameters α_1, β_1 and α_2, β_2 respectively, then $\alpha_1 = \alpha_2$ and $\beta_1 = \beta_2$.*

Proof. By Theorem 3.2, $\mu(B(x, r))$ satisfies (3.2), which uniquely determines α as the exponent of the volume growth of (M, d, μ) . By Theorem 4.6, β is uniquely determined as the critical exponent of the family of Besov spaces of (M, d, μ) . \square

Remark. Although $\alpha_1 = \alpha_2$ is true already under the hypothesis (H_0) , this hypothesis is not enough to ensure $\beta_1 = \beta_2$. Indeed, \mathbb{R}^n carries the Gauss-Weierstrass heat kernel $p_t^{(1)}$ given by (1.1) and the Cauchy-Poisson heat kernel $p_t^{(2)}$ given by (1.2), and for these heat kernels we have $\alpha_1 = \alpha_2 = n$ whereas $\beta_1 = 2 \neq 1 = \beta_2$.

4.5. Inequalities for the walk dimension.

Theorem 4.8. *Let p_t be a heat kernel on a metric measure space (M, d, μ) .*

(i) *If p_t satisfies the hypothesis (H_2) (cf. Theorem 4.6), then*

$$(4.23) \quad \beta \geq 2.$$

(ii) *If p_t satisfies the hypothesis (H_0) (cf. Theorem 3.2) and the space (M, d) satisfies the chain condition (cf. Definition 3.4), then*

$$(4.24) \quad \beta \leq \alpha + 1.$$

Remark. Observe that the chain condition is essential for the inequality $\beta \leq \alpha + 1$ to be true. Indeed, assume for a moment that the claim of Theorem 4.8(ii) holds without the chain condition, and consider a new metric d_γ on M given by $d_\gamma = d^{1/\gamma}$ where $\gamma > 1$. It is easy to see that the heat kernel p_t considered on the space (M, d_γ, μ) will satisfy (1.5) with parameters $\alpha\gamma$ and $\beta\gamma$ instead of α and β , respectively. Hence, Theorem 4.8(ii) would yield $\beta\gamma \leq \alpha\gamma + 1$, which implies $\beta \leq \alpha$ because γ may be taken arbitrarily large. However, there are spaces with $\beta > \alpha$, for example the classical Sierpiński gasket. Clearly, the metric d_γ does not satisfy the chain condition; indeed, the inequality (3.9) implies

$$(4.25) \quad d_\gamma(x_i, x_{i+1}) \leq C \frac{d_\gamma(x, y)}{n^{1/\gamma}},$$

which is not good enough. If in the inequality (3.9) we replace n by $n^{1/\gamma}$, then the proof will give $\beta \leq \alpha + \gamma$ instead of $\beta \leq \alpha + 1$.

Corollary 4.9. *Let a metric space (M, d) satisfy the chain condition. Assume that a heat kernel p_t on (M, d, μ) satisfies the estimate (1.5) with positive parameters*

α, β and non-negative decreasing functions Φ_1 and Φ_2 such that $\Phi_1(1) > 0$ and, for some $\varepsilon > 0$,

$$(4.26) \quad \int_0^\infty s^{2\alpha+1+\varepsilon} \Phi_2(s) \frac{ds}{s} < \infty.$$

Then

$$(4.27) \quad 2 \leq \beta \leq \alpha + 1.$$

Proof of Corollary 4.9. Clearly, (4.26) implies (H_0) , and by Theorem 4.8(ii) we obtain $\beta \leq \alpha + 1$. Therefore, (4.26) implies (H_2) , and by Theorem 4.8(i) we obtain $\beta \geq 2$. \square

Proof of Theorem 4.8(i). Let us show that the space $W^{1,2}$ is non-trivial, which would imply by Theorem 4.6 that $\beta \geq 2$. Fix a ball $B(x_0, R)$ in M and let $u(x)$ be the test function of this ball, that is,

$$u(x) = (R - d(x, x_0))_+.$$

Since $\text{diam}(M) = \infty$ (see Corollary 3.3), this function is non-constant for large enough R . Let us show that $W_1(u) < \infty$. Indeed, by (4.2), it suffices to prove that for some constant C and for all $r \in (0, 1)$,

$$(4.28) \quad U(r) := r^{-2-\alpha} \int_M \left[\int_{B(x,r)} |u(x) - u(y)|^2 d\mu(y) \right] d\mu(x) < C.$$

Since the function u vanishes outside the ball $B(x_0, R)$ and $r < 1$, the exterior integration in (4.28) can be reduced to $B(x_0, R + 1)$. Hence, using the obvious inequality

$$|u(x) - u(y)| \leq d(x, y) \leq r,$$

we obtain

$$\begin{aligned} U(r) &= r^{-2-\alpha} \int_{B(x_0, R+1)} \left[\int_{B(x,r)} |u(y) - u(x)|^2 d\mu(y) \right] d\mu(x) \\ &\leq r^{-2-\alpha} \int_{B(x_0, R+1)} r^2 \mu(B(x, r)) d\mu(x) \\ &\leq C r^{-2-\alpha} \int_{B(x_0, R+1)} r^{2+\alpha} d\mu(x) \\ &= C \mu(B(x_0, R + 1)), \end{aligned}$$

whence (4.28) follows. \square

We precede the proof of the second part of Theorem 4.8 by two lemmas.

Lemma 4.10. Let $\{x_i\}_{i=0}^n$ be a sequence of points in a metric space (M, d) such that for some $\rho > 0$ we have $d(x_0, x_n) > 2\rho$ and

$$(4.29) \quad d(x_i, x_{i+1}) < \rho \quad \text{for all } i = 0, 1, \dots, n-1.$$

Then there exists a subsequence $\{x_{i_k}\}_{k=0}^l$ such that

- (a) $0 = i_0 < i_1 < \dots < i_l = n$;
- (b) $d(x_{i_k}, x_{i_{k+1}}) < 5\rho$ for all $k = 0, 1, \dots, l-1$;
- (c) $d(x_{i_k}, x_{i_j}) \geq 2\rho$ for all distinct $k, j = 0, 1, \dots, l$.

The significance of conditions (a), (b), (c) is that a sequence $\{x_{i_k}\}_{k=0}^l$ satisfying them gives rise to a chain of balls $B(x_{i_k}, 5\rho)$ connecting the points x_0 and x_n in such a way that each ball contains the center of the next one whereas the balls $B(x_{i_k}, \rho)$ are disjoint. This is similar to the classical ball-covering argument, but additional difficulties arise from the necessity to maintain a proper order in the set of balls.

Proof. Let us say that a sequence of indices $\{i_k\}_{k=0}^l$ is *good* if the following conditions are satisfied:

- (a') $0 = i_0 < i_1 < \dots < i_l$;
- (b') $d(x_{i_k}, x_{i_{k+1}}) < 3\rho$ for all $k = 0, 1, \dots, l-1$;
- (c') $d(x_{i_k}, x_{i_j}) \geq 2\rho$ for all distinct $k, j = 0, 1, \dots, l$.

Note that a good sequence does not necessarily have $i_l = n$ as required in condition (a). We start with a good sequence that consists of a single index $i_0 = 0$, and will successively redefine it to increase at each step the value of i_l . A terminal good sequence will be used to construct a sequence satisfying (a), (b), (c).

Assuming that $\{i_k\}_{k=0}^l$ is a good sequence, define the set of indices

$$S := \{s : i_l < s \leq n \text{ and } d(x_s, x_{i_k}) \geq 2\rho \text{ for all } k \leq l\},$$

and consider two cases.

Set S is non-empty. In this case we will redefine $\{i_k\}$ to increase i_l . Let m be the least index in S . By (4.29) and the definition of S , $m > i_l + 1$. Since $m - 1$ is not in S , we have

$$(4.30) \quad d(x_{m-1}, x_{i_k}) < 2\rho \quad \text{for some } k \leq l$$

(see Figure 2).




FIGURE 2.

By (4.30) and (b') we obtain, for the same k as in (4.30),

$$d(x_m, x_{i_k}) \leq d(x_m, x_{m-1}) + d(x_{m-1}, x_{i_k}) < 3\rho.$$

Now we modify the sequence $\{i_j\}$ as follows: keep i_0, i_1, \dots, i_k as before, forget the previously selected indices i_{k+1}, \dots, i_l , and set $i_{k+1} := m$ and $l := k + 1$.

Clearly, the new sequence $\{i_k\}_{k=0}^l$ is also good, and the value of i_l has increased (although l may have decreased). Therefore, this construction can be repeated only a finite number of times.

Set S is empty. In this case, we will use the existing good sequence to construct a sequence satisfying conditions (a), (b), (c). The set S can be empty for two reasons:

- either $i_l = n$,

- or $i_l < n$ and for any index s such that $i_l < s \leq n$ there exists $k \leq l$ such that $d(x_s, x_{i_k}) < 2\rho$.

In the first case the sequence $\{i_k\}_{k=0}^l$ already satisfies (a), (b), (c), and the proof is finished. In the second case, choose the minimal $k \leq l$ such that $d(x_n, x_{i_k}) < 2\rho$ (see Figure 3).




FIGURE 3.

The hypothesis $d(x_n, x_0) \geq 2\rho$ implies $k \geq 1$, and from (b') we obtain

$$d(x_n, x_{i_{k-1}}) \leq d(x_n, x_{i_k}) + d(x_{i_k}, x_{i_{k-1}}) < 5\rho.$$

By the minimality of k , we also have $d(x_n, x_{i_j}) \geq 2\rho$ for all $j < k$. Hence, we define the final sequence $\{i_j\}$ as follows: keep i_0, i_1, \dots, i_{k-1} as before, forget i_k, \dots, i_l , and set $i_k := n$ and $l := k$. Then this sequence satisfies (a), (b), (c). \square

Let A be a subset of M of finite measure; that is, $\mu(A) < \infty$. Then any function $u \in L^2(M, \mu)$ is integrable on A , and we can set

$$u_A := \frac{1}{\mu(A)} \int_A u \, d\mu.$$

For any two measurable sets $A, B \subset M$ of finite measure, we have the following identity:

$$\begin{aligned} \int_A \int_B |u(x) - u(y)|^2 \, d\mu(x) d\mu(y) &= \mu(A) \int_B |u - u_B|^2 \, d\mu + \mu(B) \int_A |u - u_A|^2 \, d\mu \\ (4.31) \qquad \qquad \qquad &+ \mu(A)\mu(B) |u_A - u_B|^2, \end{aligned}$$

which is proved by a straightforward computation.

Proof of Theorem 4.8(ii). The inequality $\beta \leq \alpha + 1$ will follow from Corollary 4.3 if we show that for any $\sigma > \frac{\alpha+1}{2}$ the space $W^{\sigma,2}$ is trivial; that is, $W_\sigma(u) < \infty$ implies $u \equiv 0$. To see this, let x, y be two distinct Lebesgue points in M . By the chain condition, for any positive integer n there exists a sequence of points $\{x_i\}_{i=0}^n$ such that $x_0 = x$, $x_n = y$, and

$$d(x_i, x_{i+1}) < C \frac{d(x, y)}{n} := \rho, \quad \text{for all } 0 \leq i < n.$$

Assuming that n is large enough so that $d(x, y) > 2\rho$, we obtain by Lemma 4.10 that there exists a subsequence $\{x_{i_k}\}_{k=0}^l$ (of course $l \leq n$) such that $x_{i_0} = x$, $x_{i_l} = y$, the balls $\{B(x_{i_k}, \rho)\}$ are disjoint, and

$$(4.32) \qquad \qquad \qquad d(x_{i_k}, x_{i_{k+1}}) < 5\rho,$$

for all $k = 0, 1, \dots, l-1$.

For simplicity, let $B_k := B(x_{i_k}, \rho)$, $0 \leq k \leq l - 1$. Let $r := 7\rho < 1$. We see that the domain $\{(\xi, \eta) : \xi \in B_k, \eta \in B_{k+1}\}$ is contained in the domain $\{(\xi, \eta) : \xi \in B_k, \eta \in B(\xi, r)\}$ for all $0 \leq k \leq l - 1$, by (4.32).

By (4.2), (4.31) and (3.2), we have

$$\begin{aligned} W_\sigma(u) &\geq cr^{-2\sigma-\alpha} \int_M \int_{B(\xi,r)} |u(\xi) - u(\eta)|^2 d\mu(\eta) d\mu(\xi) \\ &\geq cr^{-2\sigma-\alpha} \sum_{k=0}^{l-1} \int_{B_k} \int_{B_{k+1}} |u(\xi) - u(\eta)|^2 d\mu(\eta) d\mu(\xi) \\ &\geq cr^{-2\sigma-\alpha} \rho^{2\alpha} \sum_{k=0}^{l-1} |u_{B_k} - u_{B_{k+1}}|^2 \\ &\geq c\rho^{-2\sigma+\alpha} \frac{1}{l} \left(\sum_{k=0}^{l-1} |u_{B_k} - u_{B_{k+1}}| \right)^2 \\ &\geq c\rho^{-2\sigma+\alpha} \frac{1}{n} |u_{B_0} - u_{B_l}|^2 \\ &\geq c\rho^{-2\sigma+\alpha+1} \frac{|u_{B(x,\rho)} - u_{B(y,\rho)}|^2}{d(x,y)}. \end{aligned}$$

Therefore, using $W_\sigma(u) < \infty$ and $2\sigma > \alpha + 1$,

$$\begin{aligned} |u(x) - u(y)|^2 &= \lim_{\rho \rightarrow 0} |u_{B(x,\rho)} - u_{B(y,\rho)}|^2 \\ &\leq C d(x,y) W_\sigma(u) \lim_{\rho \rightarrow 0} \rho^{2\sigma-\alpha-1} = 0. \end{aligned}$$

Note that (3.2) implies that μ -almost all points in M are Lebesgue points, and so $u \equiv \text{const}$. Finally, $\mu(M) = \infty$ implies $u \equiv 0$. □

4.6. Embedding theorems. In addition to the spaces L^p and $W^{\sigma,2}$ defined above, let us define a Hölder space $C^\lambda = C^\lambda(M, d, \mu)$ as follows: we say $u \in C^\lambda$ if

$$\|u\|_{C^\lambda} := \|u\|_\infty + \mu\text{-ess sup}_{\substack{x,y \in M \\ 0 < d(x,y) < 1/3}} \frac{|u(x) - u(y)|}{d(x,y)^\lambda} < \infty.$$

The restriction $d(x, y) < 1/3$ here is related to the restriction $r < 1$ in the definition (4.1) of the functional $W_\sigma(u)$. Note that the $1/3$ may be replaced by any other positive constant, giving an equivalent space.

Theorem 4.11. *Assume that a metric measure space (M, d, μ) admits a heat kernel p_t satisfying the hypothesis (H_1) (cf. Theorem 4.2). Then the following are true.*

(i) *If $\alpha > \beta$, then*

$$(4.33) \quad W^{\beta/2,2} \hookrightarrow L^q \quad \text{for any } 2 \leq q \leq 2^* := \frac{2\alpha}{\alpha - \beta}.$$

That is, $u \in W^{\beta/2,2}$ implies $u \in L^q$ and

$$(4.34) \quad \|u\|_q \leq C \|u\|_{\beta/2,2}.$$

(ii) *If $\alpha = \beta$, then*

$$W^{\beta/2,2} \hookrightarrow L^q \quad \text{for any } 2 \leq q < \infty.$$

(iii) If $\alpha < \beta$, then

$$(4.35) \quad W^{\beta/2,2} \hookrightarrow C^\lambda \quad \text{for } \lambda = \frac{\beta - \alpha}{2}.$$

That is, $u \in W^{\beta/2,2}$ implies $u \in C^\lambda$ and

$$(4.36) \quad \|u\|_{C^\lambda} \leq C \|u\|_{\beta/2,2}.$$

Remark. Observe that the definitions of the function spaces $W^{\beta/2,2}$, L^q , and C^λ involved in the embedding theorems do not depend on a heat kernel. The existence of a heat kernel is used in the proof in three different ways:

- First, in (i) and (ii) we actually prove the embedding for $\mathcal{D}(\mathcal{E})$ and then use $W^{\beta/2,2} = \mathcal{D}(\mathcal{E})$, which holds by Theorem 4.2.
- Second, the embedding results for $\mathcal{D}(\mathcal{E})$ in (i) and (ii) explicitly use the estimate

$$(4.37) \quad p_t(x, y) \leq Ct^{-\alpha/\beta}.$$

- Third, the proof of (iii) uses the estimate (3.2) of the volumes of balls, which holds by Theorem 3.2. For this part, the hypothesis (H_1) can be replaced by (H_0) . Furthermore, (iii) follows from (3.2) alone, in which case β may be any number larger than α . With this understanding, the statement of (iii) is not new (see, for example, [6], [15], [21]) and is included for completeness. However, if the hypothesis (H_1) does hold, then it follows from Theorem 4.2 that any function in $\mathcal{D}(\mathcal{E})$ is Hölder continuous.

Remark. If (M, d) satisfies the chain condition (see Definition 3.4), then for any $\lambda > 1$ the space C^λ consists only of constants. On the other hand, by Corollary 4.3 $\dim W^{\beta/2,2} = \infty$, and the embedding (4.35) cannot be true if $\lambda > 1$. Hence, by Theorem 4.11(iii) we conclude that $\beta \leq \alpha + 2$ (cf. [25]). Of course, by Theorem 4.8 we have the stronger inequality $\beta \leq \alpha + 1$ under weaker hypotheses.

Proof. By Theorem 4.2, we have $W^{\beta/2,2} = \mathcal{D}(\mathcal{E})$ and

$$\|u\|_{\beta/2,2} \simeq \|u\|_2 + \mathcal{E}[u]^{1/2},$$

for any $u \in W^{\beta/2,2}$. Hence, in parts (i) and (ii), it suffices to prove that

$$\mathcal{D}(\mathcal{E}) \hookrightarrow L^q$$

(where $2 \leq q \leq 2^*$ in case (i) and $2 \leq q < \infty$ in case (ii)), and

$$(4.38) \quad \|u\|_q \leq C \left(\|u\|_2 + \mathcal{E}[u]^{1/2} \right)$$

for any $u \in \mathcal{D}(\mathcal{E})$.

Proof for case (i), $\alpha > \beta$. The upper bound in (1.5) implies that

$$(4.39) \quad \mu\text{-ess sup}_{x,y \in M} p_t(x, y) \leq Ct^{-\alpha/\beta}.$$

Using the definition (1.3) of the semigroup T_t , the Cauchy-Schwarz inequality, and the normalization property of the heat kernel, we obtain that for any $u \in L^2$,

$x \in M$, and $t > 0$,

$$\begin{aligned}
 |T_t u(x)| &\leq \int_M p_t(x, y) |u(y)| d\mu(y) \\
 &\leq \left\{ \int_M p_t(x, y) u(y)^2 d\mu(y) \right\}^{1/2} \left\{ \int_M p_t(x, y) d\mu(y) \right\}^{1/2} \\
 (4.40) \quad &\leq C t^{-\frac{\alpha}{2\beta}} \|u\|_2.
 \end{aligned}$$

In particular, for any $\nu \geq \frac{2\alpha}{\beta}$, we have

$$(4.41) \quad \|T_t u\|_\infty \leq C t^{-\frac{\nu}{4}} \|u\|_2 \quad \text{for all } 0 < t < 1.$$

Hence, the heat semigroup $\{T_t\}$ is $L^2 \rightarrow L^\infty$ ultracontractive for $0 < t < 1$. Since $\alpha > \beta$, we see that $\nu \geq 2\alpha/\beta > 2$, and so, by [9, Theorem 2.4.2, p.75] (or [7], [28]), $\mathcal{D}(\mathcal{E}) \hookrightarrow L^{\frac{2\nu}{\nu-2}}$; that is, for any $u \in \mathcal{D}(\mathcal{E})$,

$$(4.42) \quad \|u\|_{\frac{2\nu}{\nu-2}}^2 \leq C\mathcal{E}[u] + C_0 \|u\|_2^2.$$

When ν varies in $[\frac{2\alpha}{\beta}, +\infty)$, the exponent $q = \frac{2\nu}{\nu-2}$ varies in $(2, 2^*]$. Therefore, (4.42) implies (4.38) for all $2 < q \leq 2^*$. For the remaining case $q = 2$, (4.38) is trivial.

Note that if $\nu = \frac{2\alpha}{\beta}$, then (4.41) holds for all $t > 0$, which implies (4.42) with $C_0 = 0$ (see [9, Corollary 2.4.3]); that is,

$$(4.43) \quad \|u\|_{2^*}^2 \leq C\mathcal{E}[u].$$

We will need (4.43) in Section 5.

Proof for case (ii), $\alpha = \beta$. The proof is the same as above, with the following modification. Since (4.41) holds for all $\nu \geq 2\alpha/\beta = 2$, we see that (4.42) holds for all $\nu > 2$. Therefore, $q = \frac{2\nu}{\nu-2}$ takes all the values in $(2, +\infty)$, whence the claim follows.

Proof for case (iii), $\alpha < \beta$. For any $x \in M$ and $r > 0$, set

$$(4.44) \quad u_r(x) := \frac{1}{\mu(B(x, r))} \int_{B(x, r)} u(\xi) d\mu(\xi).$$

We claim that for any $u \in W^{\beta/2, 2}$, any $0 < r < 1/3$, and all $x, y \in M$ such that $d(x, y) \leq r$, the following inequality holds:

$$(4.45) \quad |u_r(x) - u_r(y)| \leq C r^\lambda W_{\beta/2}(u)^{1/2}.$$

Indeed, setting $B_1 = B(x, r)$, $B_2 = B(y, r)$, we have

$$u_r(x) = \frac{1}{\mu(B_1)} \int_{B_1} u(\xi) d\mu(\xi) = \frac{1}{\mu(B_1)\mu(B_2)} \int_{B_1} \int_{B_2} u(\xi) d\mu(\eta) d\mu(\xi),$$

and similarly

$$u_r(y) = \frac{1}{\mu(B_1)\mu(B_2)} \int_{B_1} \int_{B_2} u(\eta) d\mu(\eta) d\mu(\xi).$$

Applying the Cauchy-Schwarz inequality, (3.2) and (4.1), we obtain

$$\begin{aligned} |u_r(x) - u_r(y)|^2 &= \left\{ \frac{1}{\mu(B_1)\mu(B_2)} \int_{B_1} \int_{B_2} (u(\xi) - u(\eta)) d\mu(\eta) d\mu(\xi) \right\}^2 \\ &\leq \frac{1}{\mu(B_1)\mu(B_2)} \int_{B_1} \int_{B_2} |u(\xi) - u(\eta)|^2 d\mu(\eta) d\mu(\xi) \\ &\leq C r^{-\alpha} \int_M \left[\frac{1}{\mu(B(\xi, 3r))} \int_{B(\xi, 3r)} |u(\xi) - u(\eta)|^2 d\mu(\eta) \right] d\mu(\xi) \\ &\leq C r^{-\alpha+\beta} W_{\beta/2}(u), \end{aligned}$$

thus proving (4.45).

Similarly, one proves that for any $0 < r < 1/3$ and $x \in M$,

$$(4.46) \quad |u_{2r}(x) - u_r(x)| \leq C r^\lambda W_{\beta/2}(u)^{1/2}.$$

Let x be a Lebesgue point of u . Setting $r_k = 2^{-k}r$ for any $k = 0, 1, 2, \dots$, we obtain from (4.46) that

$$\begin{aligned} |u(x) - u_r(x)| &\leq \sum_{k=0}^{\infty} |u_{r_k}(x) - u_{r_{k+1}}(x)| \\ &\leq C \left(\sum_{k=0}^{\infty} r_k^\lambda \right) W_{\beta/2}(u)^{1/2} \\ (4.47) \quad &\leq C r^\lambda W_{\beta/2}(u)^{1/2}. \end{aligned}$$

Applying the Cauchy-Schwarz inequality

$$|u_r(x)| \leq C r^{-\alpha/2} \|u\|_2$$

and using (4.47) to some fixed value of r , say $r = 1/4$, we obtain

$$|u(x)| \leq |u(x) - u_r(x)| + |u_r(x)| \leq C \left(\|u\|_2 + W_{\beta/2}(u)^{1/2} \right),$$

whence

$$(4.48) \quad \|u\|_\infty \leq C \|u\|_{\beta/2,2}.$$

If y is another Lebesgue point of u such that $r := d(x, y) < 1/3$, then we obtain from (4.45), (4.47), and a similar inequality for y ,

$$|u(x) - u(y)| \leq |u(x) - u_r(x)| + |u_r(x) - u_r(y)| + |u_r(y) - u(y)| \leq C r^\lambda W_{\beta/2}(u)^{1/2}.$$

Hence,

$$\frac{|u(x) - u(y)|}{d(x, y)^\lambda} \leq C W_{\beta/2}(u)^{1/2},$$

which together with (4.48) yields (4.36). □

Theorem 4.12 (Compact embedding theorem). *Let (M, d, μ) be a metric measure space, and p_t be a heat kernel in M satisfying (1.5). Then for any bounded sequence $\{u_k\}$ in $\mathcal{D}(\mathcal{E})$ in the norm (4.7), there exists a subsequence $\{u_{k_i}\}$ that converges to a function $u \in L^2(M, \mu)$ in the following sense:*

$$\|u_{k_i} - u\|_{L^2(B, \mu)} \rightarrow 0,$$

for any set $B \subset M$ of finite measure.

Remark. Without any additional assumptions about the functions Φ_1 and Φ_2 , the estimate (1.5) is equivalent to

$$p_t(x, y) \leq Ct^{-\alpha/\beta},$$

and this is exactly what is used in the proof.

Proof. Let $\{u_k\}$ be a bounded sequence in $\mathcal{D}(\mathcal{E})$. Since $\{u_k\}$ is also bounded in L^2 , there exists a subsequence, still denoted by $\{u_k\}$, such that $\{u_k\}$ converges weakly to some function $u \in L^2$. Let us show that in fact $\{u_k\}$ converges to u in $L^2(B) = L^2(B, \mu)$ for any set $B \subset M$ of finite measure.

For any $t > 0$, by the triangle inequality we have

$$(4.49) \quad \|u_k - u\|_{L^2(B)} \leq \|u_k - T_t u_k\|_{L^2(M)} + \|T_t u_k - T_t u\|_{L^2(B)} + \|T_t u - u\|_{L^2(M)}.$$

For any function $v \in L^2$ we have

$$\begin{aligned} \|v - T_t v\|_2^2 &= \int_M \left(\int_M (v(x) - v(y)) p_t(x, y) d\mu(y) \right)^2 d\mu(x) \\ &\leq \int_M \left\{ \int_M p_t(x, y) d\mu(y) \int_M |v(x) - v(y)|^2 p_t(x, y) d\mu(y) \right\} d\mu(x) \\ &= 2t \mathcal{E}_t[v] \\ &\leq 2t \mathcal{E}[v]. \end{aligned}$$

Since $\mathcal{E}[u_k]$ is uniformly bounded in k by the hypothesis, we obtain that for all k and $t > 0$,

$$(4.50) \quad \|u_k - T_t u_k\|_2 \leq C\sqrt{t}.$$

Since $\{u_k\}$ converges to u weakly in L^2 and $p_t(x, \cdot) \in L^2$, we see that for μ -almost all $x \in M$,

$$T_t u_k(x) = \int_M p_t(x, y) u_k(y) d\mu(y) \xrightarrow{k \rightarrow \infty} \int_M p_t(x, y) u(y) d\mu(y) = T_t u(x).$$

Also, by (4.40) we have

$$\|T_t u_k\|_\infty \leq Ct^{-\frac{\alpha}{2\beta}} \|u_k\|_2,$$

so that the sequence $\{T_t u_k\}$ is uniformly bounded in k for any $t > 0$. Since $\{T_t u_k\}$ converges to $T_t u$ almost everywhere, the dominated convergence theorem yields

$$(4.51) \quad T_t u_k \xrightarrow{k \rightarrow \infty} T_t u \quad \text{in } L^2(B),$$

because $\mu(B) < \infty$. Hence, we obtain from (4.49), (4.50), and (4.51) that for any $t > 0$,

$$\limsup_{k \rightarrow \infty} \|u_k - u\|_{L^2(B)} \leq C\sqrt{t} + \|T_t u - u\|_{L^2(M)}.$$

Since $T_t u \rightarrow u$ in $L^2(M)$ as $t \rightarrow 0$, we finish the proof by letting $t \rightarrow 0$. □

Corollary 4.13. *Let (M, d, μ) be a metric measure space, and let p_t be a heat kernel in M satisfying (1.5) with $\Phi_1(1) > 0$. Then for any bounded sequence $\{u_k\}$ in $\mathcal{D}(\mathcal{E})$ in the norm (4.7), there exists a subsequence $\{u_{k_i}\}$ that converges to a function $u \in L^2(M, \mu)$ almost everywhere.*

Proof. By the first part of the proof of Theorem 3.2, the hypothesis $\Phi_1(1) > 0$ implies finiteness of the measure of any ball. Fix a point $x \in M$ and consider the sequence of balls $B_N = B(x, N)$, where $N = 1, 2, \dots$. By Theorem 4.12 we can assume that the sequence $\{u_k\}$ converges to $u \in L^2(M)$ in the norm of $L^2(B_N)$ for any N . Therefore, there exists a subsequence that converges almost everywhere in B_1 . From this sequence, let us select a subsequence that converges to u almost everywhere in B_2 , and so on. Using the diagonal principle, we obtain a subsequence that converges to u almost everywhere in M . \square

5. SEMILINEAR ELLIPTIC EQUATIONS

As above, let (M, d, μ) be a metric measure space that possesses a heat kernel satisfying (1.5). In this section we show the existence of generalized solutions of the equation

$$(5.1) \quad -\mathcal{L}u + f(x, u) = g(x),$$

where \mathcal{L} is the Laplace operator in M defined by (4.3) or (4.8). More precisely, we say that $u \in \mathcal{D}(\mathcal{E})$ is a generalized solution of (5.1) if the following identity holds:

$$(5.2) \quad \mathcal{E}(u, v) + \int_M f(x, u(x))v(x)d\mu(x) - \int_M g(x)v(x)d\mu(x) = 0,$$

for any test function v from a certain class to be defined below.

Fix a couple $p, q \in (1, \infty)$, set

$$E^{p,q} := \mathcal{D}(\mathcal{E}) \cap L^p \cap L^q,$$

and define the norm in $E^{p,q}$ by

$$\|u\| := \|u\|_p + \|u\|_q + \mathcal{E}[u]^{1/2}.$$

Clearly $E^{p,q}$ is a Banach space, and its dual is

$$(E^{p,q})^* = E^{p',q'},$$

where p' and q' are the Hölder conjugates to p and q , respectively.

We assume throughout this section that

$$(5.3) \quad g \in L^{p'}$$

and

$$(5.4) \quad |f(x, u)| \leq C|u|^{q-1} + f_0(x), \quad \text{for all } x \in M \text{ and } u \in \mathbb{R},$$

where f_0 is a non-negative function in $L^{q'}$.

Let us show that all the terms in (5.2) make sense if $u \in \mathcal{D}(\mathcal{E}) \cap L^q$ and $v \in E^{p,q}$. Indeed, $\mathcal{E}(u, v)$ is defined as $u, v \in \mathcal{D}(\mathcal{E})$, and the other two terms are finite by the Hölder inequality:

$$(5.5) \quad \left| \int_M gv d\mu \right| \leq \|g\|_{p'} \|v\|_p < \infty$$

and

$$(5.6) \quad \left| \int_M f(\cdot, u)v d\mu \right| \leq \|f(\cdot, u)\|_{q'} \|v\|_q \leq \left(C\|u\|_q^{q/q'} + \|f_0\|_{q'} \right) \|v\|_q < \infty.$$

Now we can give a precise definition of a generalized solution of (5.1).

Definition 5.1. Assuming that f and g satisfy (5.3) and (5.4), we say that $u \in E^{p,q}$ is a *generalized solution* of (5.1) if the identity (5.2) holds for all $v \in E^{p,q}$.

Let E be a Banach space and $I : E \rightarrow \mathbb{R}$ be a functional on E . Recall that I is *Fréchet differentiable* at $u \in E$ if there exists an element in the dual space E^* of E , denoted by $I'(u)$, such that for all $v \in E$,

$$I(u + tv) = I(u) + tI'(u)v + o(t) \quad \text{as } t \rightarrow 0.$$

The functional $I'(u)$ is called the *Fréchet derivative* of I at point u . We say that I is *continuously Fréchet differentiable* if I is Fréchet differentiable at any $u \in E$, and the mapping $u \mapsto I'(u)$ is a continuous mapping from E to E^* (see, for example, [19], [24]). Finally, we say that u is a *critical point* of I if $I'(u) = 0$.

We will show that a generalized solution of (5.1) may be obtained as a *critical point* of a functional $I(u)$, where $I(u)$ is defined by

$$(5.7) \quad I(u) := \frac{1}{2}\mathcal{E}[u] + \int_M F(x, u(x))d\mu(x) - \int_M g(x)u(x)d\mu(x),$$

and

$$(5.8) \quad F(x, u) := \int_0^u f(x, s) ds.$$

Let us show that the functional I defined by (5.7) is continuously Fréchet differentiable for suitable f and g .

Proposition 5.2. *Assume that $f(x, u)$ is continuous in $u \in \mathbb{R}$ for all $x \in M$ and satisfies (5.4), and g satisfies (5.3). Then I defined as in (5.7) is continuously Fréchet differentiable in $E^{p,q}$. Moreover, we have*

$$(5.9) \quad I'(u)v = \mathcal{E}(u, v) + \int_M f(x, u(x))v(x)d\mu(x) - \int_M g(x)v(x)d\mu(x),$$

for all $u, v \in E^{p,q}$.

Thus, u is a generalized solution of (5.1) if and only if u is a critical point of I .

Proof. The proof follows the same line as in [19]. For completeness, we sketch the proof. It is easy to see directly from the definition that the functional

$$I_0(u) = \frac{1}{2}\mathcal{E}[u] - \int_M g(x)u(x)d\mu(x)$$

is continuously Fréchet differentiable at any point $u \in E^{p,q}$, and

$$I'_0(u)v = \mathcal{E}(u, v) - \int_M g(x)v(x)d\mu(x).$$

Let us show that the remaining part of I , namely the functional

$$J(u) := \int_M F(x, u(x))d\mu(x),$$

is also continuously Fréchet differentiable, and

$$(5.10) \quad J'(u)v = \int_M f(x, u(x))v(x)d\mu(x).$$

Indeed, taking (5.10) as the definition of J' , we have, for all $u, v \in E^{p,q}$ and $-1 < t < 1$,

$$\begin{aligned} J(u + tv) - J(u) - tJ'(u)v &= \int_M \left[\int_u^{u+tv} f(x, s)ds - tf(x, u(x))v(x) \right] d\mu(x) \\ &= t \int_M (f(x, u + \theta v) - f(x, u))v(x)d\mu(x), \end{aligned}$$

where

$$\theta = \theta(x, t) \in [0, t] \subset (-1, 1).$$

By (5.6) we have $f(\cdot, u)v \in L^1$. By (5.4) and (5.6) we obtain, similarly,

$$|f(\cdot, u + \theta v)| \leq C |u + \theta v|^{q-1} + f_0 \leq C (|u| + |v|)^{q-1} + f_0 \in L^1.$$

Since $f(x, u + \theta v) \rightarrow f(x, u)$ as $t \rightarrow 0$, we conclude by the dominated convergence theorem that

$$J(u + tv) - J(u) - tJ'(u)v = o(t) \quad \text{as } t \rightarrow 0,$$

proving that J is Fréchet differentiable.

It remains to show that $J'(u)$ is continuous. For any $u_1, u_2, v \in E^{p,q}$, we have

$$\begin{aligned} |J'(u_1)v - J'(u_2)v| &= \left| \int_M (f(x, u_1(x)) - f(x, u_2(x)))v(x) d\mu(x) \right| \\ &\leq \|f(\cdot, u_1) - f(\cdot, u_2)\|_{q'} \|v\|_q \\ &\leq \|f(\cdot, u_1) - f(\cdot, u_2)\|_{q'} \|v\|_{E^{p,q}}, \end{aligned}$$

whence

$$\|J'(u_1) - J'(u_2)\|_{(E^{p,q})^*} \leq \|f(\cdot, u_1) - f(\cdot, u_2)\|_{q'}.$$

Note that the *Nemytsky operator* $\mathcal{F}u := f(x, u(x))$ is continuous from L^q to $L^{q'}$, provided that f satisfies (5.4) (see [27, Theorem 19.1]; for a bounded domain, see [1, Theorem 2.2]). Indeed, if $\mathcal{F}u$ is not continuous, then there exists a sequence $\{u_k\}$ such that $\|u_k - u\|_q \rightarrow 0$ but

$$(5.11) \quad \|\mathcal{F}u_k - \mathcal{F}u\|_{q'} \geq \varepsilon$$

for all k and some $\varepsilon > 0$. Since $\{u_k\}$ converges to u in L^q , there is a subsequence $\{u_{k_j}\}$ of $\{u_k\}$ such that $\{u_{k_j}\}$ converges to u almost everywhere in M . Fix some $R > 0$ and set $B_R = B(x_0, R)$ for a fixed point $x_0 \in M$. By (5.4) and the dominated convergence theorem, we have

$$\begin{aligned} &\lim_{j \rightarrow \infty} \|\mathcal{F}u_{k_j} - \mathcal{F}u\|_{q'}^{q'} \\ &= \lim_{j \rightarrow \infty} \left\{ \int_{B_R} |f(\cdot, u_{k_j}) - f(\cdot, u)|^{q'} d\mu + \int_{M \setminus B_R} |f(\cdot, u_{k_j}) - f(\cdot, u)|^{q'} d\mu \right\} \\ &\leq C \lim_{j \rightarrow \infty} \int_{M \setminus B_R} (|u_{k_j}|^q + |u|^q + |f_0|^{q'}) d\mu \\ &= C \lim_{j \rightarrow \infty} \int_{M \setminus B_R} (|u_{k_j} - u + u|^q + |u|^q + |f_0|^{q'}) d\mu \\ &\leq C \int_{M \setminus B_R} (|u|^q + |f_0|^{q'}) d\mu. \end{aligned}$$

Choosing R large enough, we can make the right-hand side arbitrarily small, which contradicts (5.11). Hence, $\mathcal{F}u$ is continuous from L^q to $L^{q'}$. Therefore, $J'(u)$ is continuous. \square

By Proposition 5.2, in order to prove the existence of a generalized solution of (5.1), it is enough to show that the functional I defined by (5.7) has a critical point in $E^{p,q}$; this in turn will follow if I has a minimum point in $E^{p,q}$. The following

statement provides the conditions that ensure that a functional on a Banach space has a minimum point.

Proposition 5.3 ([19, Theorem 2.5, p.14]). *Let I be a real-valued functional in a reflexive Banach space E satisfying the following conditions:*

(i) *I is bounded below; that is,*

$$\inf_{u \in E} I(u) > -\infty.$$

(ii) *I is coercive; that is, for any real a there exists b such that $I(u) \leq a$ implies $\|u\| \leq b$.*

(iii) *Any sequence $\{u_k\}$ that converges to u weakly in E has a subsequence $\{u_{k_i}\}$ such that*

$$\liminf_{i \rightarrow \infty} I(u_{k_i}) \geq I(u).$$

Then I has a minimum point in E .

Now we prove the main result of this section.

Theorem 5.4. *Assume that a metric measure space (M, d, μ) admits a heat kernel p_t satisfying the hypothesis (H_1) (cf. Theorems 4.2, 4.11), and assume that $\alpha > \beta$. Fix two real numbers p, q such that*

$$(5.12) \quad q \geq p \geq 2^* := \frac{2\alpha}{\alpha - \beta}.$$

Let $g \in L^{p'}$, and let $f(x, u)$ be a measurable function on $M \times \mathbb{R}$ that is continuous in u for any $x \in M$ and satisfies (5.4). Moreover, assume that, for all $x \in M$ and $u \in \mathbb{R}$,

$$(5.13) \quad F(x, u) := \int_0^u f(x, s) ds \geq c|u|^q + F_0(x),$$

where $F_0 \in L^1$. Then (5.1) has a generalized solution $u \in E^{p,q}$.

Remark. Here is an example of a function f that satisfies all the hypotheses of Theorem 5.4:

$$f(x, u) = |u|^{q-2} u.$$

Hence, the equation

$$(5.14) \quad -\mathcal{L}u + |u|^{q-2} u = g$$

has a generalized solution $u \in E^{p,q}$ for any $g \in L^{p'}$, provided p and q satisfy (5.12). We will see below that this solution is unique.

Proof. It suffices to show that the functional I defined by (5.7) satisfies conditions (i) – (iii) of Proposition 5.3.

Condition (i): I is bounded below. By (5.12) there exists $\theta \in [0, 1]$ such that

$$\frac{1}{p} = \frac{\theta}{2^*} + \frac{1-\theta}{q},$$

whence, for any $u \in E^{p,q}$,

$$\|u\|_p \leq \|u\|_{2^*}^\theta \|u\|_q^{1-\theta}$$

by the Hölder inequality. By (4.43), we see that

$$\|u\|_{2^*} \leq C \mathcal{E}[u]^{1/2},$$

which implies

$$(5.15) \quad \begin{aligned} \|u\|_p &\leq C \mathcal{E}[u]^{\theta/2} \|u\|_q^{1-\theta} \\ &\leq C \left(\|u\|_q + \mathcal{E}[u]^{1/2} \right). \end{aligned}$$

From (5.7), (5.13), (5.5) and (5.15), we obtain that, for any $u \in E^{p,q}$,

$$(5.16) \quad \begin{aligned} I(u) &\geq \frac{1}{2} \mathcal{E}[u] + (c\|u\|_q^q - \|F_0\|_1) - \|g\|_{p'} \|u\|_p \\ &\geq \frac{1}{2} \mathcal{E}[u] + c\|u\|_q^q - \|F_0\|_1 - C\|g\|_{p'} (\|u\|_q + \mathcal{E}[u]^{1/2}) \\ &\geq \left[\frac{1}{2} \mathcal{E}[u] - C\mathcal{E}[u]^{1/2} \right] + [c\|u\|_q^q - C\|u\|_q] - C \\ &\geq \left[\frac{1}{2} s^2 - Cs \right] + [ct^q - Ct] - C, \end{aligned}$$

where $s := \mathcal{E}(u, u)^{1/2}$ and $t := \|u\|_q$. Since $q > 1$, the value of each square bracket is bounded below, whence we conclude that I is bounded below.

Condition (ii): I is coercive. If $I(u) \leq a$ for some a , then by (5.16),

$$a \geq \left[\frac{1}{2} s^2 - Cs \right] + [ct^q - Ct] - C,$$

which implies that s and t must be bounded. Together with (5.15), this implies that u is bounded in $E^{p,q}$.

Condition (iii). Let a sequence $\{u_k\}$ converge to u weakly in $E^{p,q}$. Since $g \in L^{p'} \subset (E^{p,q})^*$, we have

$$(5.17) \quad \lim_{k \rightarrow \infty} \int_M g u_k d\mu = \int_M g u d\mu.$$

Since $u \in \mathcal{D}(\mathcal{E}) \subset (E^{p,q})^*$, we have

$$\lim_{k \rightarrow \infty} \mathcal{E}(u_k, u) = \mathcal{E}[u].$$

Applying the inequality

$$\mathcal{E}[u_k] \geq 2\mathcal{E}(u_k, u) - \mathcal{E}[u],$$

we obtain

$$(5.18) \quad \liminf_{k \rightarrow \infty} \mathcal{E}[u_k] \geq \mathcal{E}[u].$$

We are left to verify that there exists a subsequence $\{u_{k_i}\}$ such that

$$(5.19) \quad \liminf_{i \rightarrow \infty} \int_M F(\cdot, u_{k_i}) d\mu \geq \int_M F(\cdot, u) d\mu.$$

The sequence $\{u_k\}$ is bounded in $\mathcal{D}(\mathcal{E})$. Therefore, by Corollary 4.13 there exists a subsequence $\{u_{k_i}\}$ that converges to u almost everywhere in M . Therefore, we have also $F(\cdot, u_{k_i}) \rightarrow F(\cdot, u)$ almost everywhere in M . By (5.13), we have $F(\cdot, u_{k_i}) \geq F_0 \in L^1$, and (5.19) follows by Fatou's lemma. Combining (5.17)-(5.19), we complete the proof. \square

Finally, we complement Theorem 5.4 by a uniqueness result.

Proposition 5.5. *Let the function $f(x, u)$ be strictly monotone increasing in u for every $x \in M$. Then the equation (5.1) has at most one generalized solution $u \in E^{p,q}$.*

In particular, the equation (5.14) has exactly one generalized solution in $E^{p,q}$.

Proof. Let u_1 and u_2 be two generalized solutions of (5.1). Then for any $v \in E^{p,q}$ we have from (5.2),

$$\mathcal{E}(u_1 - u_2, v) + \int_M (f(\cdot, u_1) - f(\cdot, u_2)) v \, d\mu = 0.$$

Substituting $v = u_1 - u_2$, we obtain

$$\mathcal{E}[u_1 - u_2] + \int (f(\cdot, u_1) - f(\cdot, u_2))(u_1 - u_2) \, d\mu = 0.$$

By the monotonicity of $f(x, u)$ in u , both terms here are non-negative, and so each of them must vanish. In particular, we obtain

$$(f(\cdot, u_1) - f(\cdot, u_2))(u_1 - u_2) = 0$$

almost everywhere, which by the strict monotonicity of f implies $u_1 = u_2$. \square

REFERENCES

1. A. Ambrosetti and G. Prodi, *A primer of nonlinear analysis*, Cambridge University Press, 1993. MR **94f**:58016
2. N. Aronszajn and K. T. Smith, *Theory of Bessel potentials. part 1*, Ann. Inst. Fourier, Grenoble **11** (1961), 385–475. MR **26**:1485
3. M. T. Barlow, *Diffusions on fractals*, Lectures on Probability Theory and Statistics, vol. 1690, Lecture Notes Math., Springer-Verlag, 1998, pp. 1–121. MR **2000a**:60148
4. ———, *Which values of the volume growth and escape time exponents are possible for graphs?*, preprint (2001).
5. M. T. Barlow and R. F. Bass, *Brownian motion and harmonic analysis on Sierpiński carpets*, Canadian J. Math. **51** (1999), 673–744. MR **2000i**:60083
6. M. Biroli and U. Mosco, *Sobolev and isoperimetric inequalities for Dirichlet forms on homogeneous spaces*, Rend. Mat. Acc. Lincei **VI** (1) (1995), 37–44. MR **96i**:46034a
7. E. A. Carlen, S. Kusuoka, and D. W. Stroock, *Upper bounds for symmetric Markov transition functions*, Ann. Inst. Henri Poincaré-Probab. Statist. **23** (1987), 245–287. MR **88i**:35066
8. E. B. Davies, *One-parameter semigroups*, Academic Press, 1980. MR **82i**:47060
9. ———, *Heat kernels and spectral theory*, Cambridge University Press, 1990. MR **92a**:35035
10. K. J. Falconer, *Fractal geometry—mathematical foundation and applications*, John Wiley, 1992. MR **92j**:28008
11. P. J. Fitzsimmons, B. M. Hambly, and T. Kumagai, *Transition density estimates for Brownian motion on affine nested fractals*, Commun. Math. Phys. **165** (1994), 595–620. MR **95j**:60122
12. M. Fukushima, Y. Oshima Y., and M. Takeda, *Dirichlet forms and symmetric Markov processes*, Studies in Mathematics, 19, De Gruyter, 1994. MR **96f**:60126
13. A. Grigor'yan, *Estimates of heat kernels on Riemannian manifolds*, London Math. Soc. Lecture Note Ser., vol. 273, 1999, pp. 140–225. MR **2001b**:58040
14. A. Grigor'yan and A. Telcs, *Sub-Gaussian estimates of heat kernels on infinite graphs*, Duke Math. J. **109** (2001), 451–510.
15. P. Hajlasz and P. Koskela, *Sobolev met Poincaré*, Mem. Amer. Math. Soc. **145** (2000), no. 688. MR **2000j**:46063
16. E. Hille and R. S. Phillips, *Functional analysis and semi-groups*, Amer. Math. Soc., 1957. MR **19**:664d
17. J. Hu and K. S. Lau, *Riesz potentials and Laplacians on fractals*, preprint (2001).
18. A. Jonsson, *Brownian motion on fractals and function spaces*, Math. Zeit. **222** (1996), 495–504. MR **97e**:60137

19. I. Kuzin and S. Pohozaev, *Entire solutions of semilinear elliptic equations*, Birkhäuser, 1997. MR **99d**:35050
20. P. Li and S. T. Yau, *On the parabolic kernel of the Schrödinger operator*, Acta Math. **156** (1986), 153–201. MR **87f**:58156
21. R. A. Macías and C. Segovia, *Lipschitz functions on spaces of homogeneous type*, Adv. Math. **33** (1979), 257–270. MR **81c**:32017a
22. K. Pietruska-Pałuba, *Some function spaces related to the Brownian motion on simple nested fractals*, Stochastics and Stochastics Reports **67** (1999), 267–285. MR **2000h**:60078
23. ———, *On function spaces related to fractional diffusions on d -sets*, Stochastics and Stochastics Reports **70** (2000), 153–164. MR **2001i**:60123
24. P. R. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, Expository Lectures from CBMS Regional Conference held at University of Miami, Amer. Math. Soc., Providence, RI, 1986. MR **87j**:58024
25. A. Stós, *Symmetric α -stable processes on d -sets*, Bull. Pol. Acad. Sci. Math. **48** (2000), 237–245. MR **2002f**:60152
26. A. Telcs, *Random walks on graphs, electric networks and fractals*, J. Probability Theory and Related Fields **82** (1989), 435–449. MR **90h**:60065
27. M. M. Vainberg, *Variational methods for the study of nonlinear operators*, San Francisco: Holden-Day, Inc., 1964. MR **31**:638
28. N. Th. Varopoulos, *Hardy-Littlewood theory for semigroups*, J. Funct. Anal. **63** (1985), 240–260. MR **87a**:31011
29. K. Yosida, *Functional analysis*, Springer-Verlag, 1980. MR **82i**:46002

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE, LONDON, SW7 2BZ, UNITED KINGDOM
AND THE INSTITUTE OF MATHEMATICAL SCIENCES, THE CHINESE UNIVERSITY OF HONG KONG,
SHATIN, N.T., HONG KONG

E-mail address: a.grigoryan@ic.ac.uk

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084 CHINA
AND DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, N.T.,
HONG KONG

E-mail address: jxhu@math.tsinghua.edu.cn

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, N.T.,
HONG KONG

E-mail address: kslau@math.cuhk.edu.hk