

RIGIDITY IN HOLOMORPHIC AND QUASIREGULAR DYNAMICS

GAVEN J. MARTIN AND VOLKER MAYER

ABSTRACT. We consider rigidity phenomena for holomorphic functions and then more generally for uniformly quasiregular maps.

1. INTRODUCTION

1.1. Rigidity of rational functions. Fatou's problem of the density of hyperbolic maps in the space of rational functions is one of the major problems in the field of holomorphic dynamics. It has been reduced by Mañé, Sad and Sullivan [13] to the no invariant lines field (NILF) conjecture. A rational map f has an *invariant line field* if the Julia set has positive Lebesgue measure, $|\mathcal{J}_f| > 0$, and if there is a measurable map $z \in \mathcal{J}_f \mapsto \mathcal{L}_z \in [0, \pi)$ with

$$\mathcal{L}_{f(z)} = \mathcal{L}_z + \arg f'(z) \bmod \pi.$$

It is well known that the map f has an invariant line field precisely when f admits a quasiconformal deformation that is non-trivial on the Julia set. Now, the NILF-conjecture suggests that a rational function does not admit an invariant line field unless it is a map of a very special kind, that is, a Lattès map. An affirmative answer to this conjecture implies the density of hyperbolic maps [13]. We recall that a *Lattès map* is a rational function f that is obtained from semi-conjugating an expanding similarity $A(z) = \lambda z$, $|\lambda| > 1$, by a certain automorphic meromorphic function $h : \mathbb{C} \rightarrow \hat{\mathbb{C}}$. Such maps f are then obtained as solutions of the functional equation

$$(1.1) \quad f \circ h(z) = h \circ A(z), \quad z \in \mathbb{C}.$$

Next, from Zalcman's normality criteria [30, 27] we know that each point z_0 of the Julia set \mathcal{J}_f of f admits a renormalization of the following form. There are a sequence of points $z_j \rightarrow x_0$ and positive real numbers $\rho_j \searrow 0$ and a subsequence of iterates f^{k_j} such that

$$(1.2) \quad f^{k_j}(\rho_j z + z_j) \rightarrow \Psi(z), \quad z \in \mathbb{C},$$

the convergence is locally uniform in \mathbb{C} and $\Psi : \mathbb{C} \rightarrow \hat{\mathbb{C}}$ is a non-constant meromorphic limit function.

Received by the editors October 19, 1999.

2000 *Mathematics Subject Classification.* Primary 30C65; Secondary 37F45.

This research was partially supported by a grant from the Marsden Fund (NZ).

Following an analogy with the limit sets of Kleinian groups, we call a point $z_0 \in \mathcal{J}_f$ a *conical point* if the sequence above in (2) can be chosen to have the additional property that there is a constant $M < \infty$ such that

$$(1.3) \quad \frac{|z_0 - z_j|}{\rho_j} < M,$$

and we require only that the convergence be uniform on \mathbb{D} .

The set of conical points is denoted Λ_c . A typical conical point is a repelling fixed point. In general Λ_c is a large subset of the Julia set, and in many cases, for example the Lattès maps, we even have $\Lambda_c = \mathcal{J}_f$. Reinforcing the analogy with Kleinian groups, we shall see that examples of points in the Julia set which are not conical are the parabolic fixed points (see Lemma 3.5).

Several other notions of conical points appear in the literature, and we discuss these later in §3.

The following lemma gives a characterisation of a conical limit point which is easier to work with.

Lemma 1.1. *Let $z_0 \in \mathcal{J}_f$ be a conical point for the mapping f . Then there are a sequence $\rho_j \searrow 0$, an increasing sequence of natural numbers $\{k_j\}_{j=1}^\infty$ and a non-constant meromorphic function Ψ such that*

$$(1.4) \quad f^{k_j}(z_0 + \rho_j z) \rightarrow \Psi$$

uniformly in \mathbb{D} .

Proof. Let ρ_j, z_j, k_j, Ψ and M be given as in the definition of a conical limit point. Set $\phi_j(z) = \rho_j z + z_j$ and $\varphi_j(z) = \rho_j z + z_0$. Then

$$(1.5) \quad f^{k_j} \circ \varphi_j(z) = f^{k_j} \circ \phi_j(z) \circ \phi_j^{-1}(z) \circ \varphi_j(z).$$

The result now follows after passing to a subsequence, since $f^{k_j} \circ \phi_j(z) \rightarrow \Psi$ and the sequence $\phi_j^{-1}(z) \circ \varphi_j(z) = z + (z_0 - z_j)/\rho_j$ is precompact. \square

We may reformulate the NILF-conjecture, replacing role of the Julia set by Λ_c . We shall prove the following result:

Theorem 1.2. *Suppose the conical set of the rational function f has positive Lebesgue measure, $|\Lambda_c| > 0$. Then either f is a Lattès map or f has no invariant line field.*

Note that the conical set has either zero or full measure. Therefore the condition $|\Lambda_c| > 0$ implies in particular that $\mathcal{J}_f = \hat{\mathbb{C}}$. In view of the above result it would be interesting to know an example of a rational function for which the Julia set has positive measure and the conical set has zero measure. Such a phenomena appears in the space of entire functions. McMullen [19] showed that $f(z) = \sin z$ has a Julia set of positive measure and, since this function has a non-empty Fatou set, its conical set has zero measure.

Theorem 1.2 is not really new (compare [20], [12], [6]). However, we give a new and elementary proof of this fact that in particular does not involve any orbifold theory. We use the possibility of renormalization at conical points to construct explicitly the necessary automorphic function of the equation (1.1). A similar idea has been used by Berteloot and Loeb in [1], where they gave a geometric description of the Lattès maps. The proof we give also has the advantage that it does not use that f is a global holomorphic map of $\hat{\mathbb{C}}$. Therefore, Theorem 1.2 is also true for

entire functions. One of our primary motivations in looking for such a proof is its applicability to higher-dimensional analogues of holomorphic dynamics, uniformly quasiregular maps, which we now present.

1.2. Rigidity in higher dimensions. The maps we shall consider in the sequel are self-maps of the n -sphere $\overline{\mathbb{R}}^n = \mathbb{R}^n \cup \{\infty\}$, $n \geq 2$, that preserve a *conformal structure*. Such a structure is a bounded measurable map $\mu : \overline{\mathbb{R}}^n \rightarrow S(n)$, the space of symmetric positive definite $n \times n$ matrices of determinant 1. More precisely, let f be a map that is in the Sobolev class $W^{1,n}(\overline{\mathbb{R}}^n, \overline{\mathbb{R}}^n)$ whose Jacobian determinant $\det f' > 0$ almost everywhere. The pull-back action of f on a conformal structure μ is given by

$$(1.6) \quad f^* \mu(x) = \det(f'(x)^t)^{-\frac{2}{n}} f'(x) \mu(f(x)) f'(x) \quad \text{for a.e. } x \in \overline{\mathbb{R}}^n .$$

We call f μ -rational if μ is f -invariant, meaning

$$(1.7) \quad f^* \mu = \mu \quad \text{a.e.}$$

This equation has the following geometric interpretation: Consider the field of μ -spheres $\Sigma_{x,r}^\mu = \{v \in \mathbb{R}^n; {}^t v \mu(x) v = r\}$, which is in fact a measurable Euclidean ellipsoid field on $\overline{\mathbb{R}}^n$ such that the eccentricity of the ellipsoids is uniformly bounded. Now, a μ -rational map is conformal in this structure in the sense that it preserves this field of μ -spheres.

A μ -rational map is *quasiregular* (see §2 or [26] for appropriate definitions), and the dilatation of any μ -rational map only depends on the eccentricity of the ellipsoid field $\Sigma_{x,r}^\mu$. Therefore, for a given conformal structure μ there is a fixed constant $K \geq 1$ such that any μ -rational map is K -quasiregular. In particular, if f is μ -rational, then the family of all the iterates f^ℓ has a uniform dilatation bound. Such maps are called *uniformly quasiregular*, or *uqr* for brevity. Conversely, it is known that any uqr map is μ -rational with respect to some conformal structure μ [9].

In two dimensions the study of uqr maps reduces to the classical case of (holomorphic) rational functions. In fact, when $\mu \equiv Id$ is the standard Euclidean structure, then (1.7) is nothing other than the usual Cauchy-Riemann system. If $n = 2$, the Id -rational maps are precisely the holomorphic functions of $\hat{\mathbb{C}}$, and hence they are rational maps. Moreover, from the Ahlfors-Bers measurable Riemannian mapping theorem (the existence theorem for quasiconformal mappings) it follows that any uqr map in two dimensions is quasiconformally conjugate to a rational map. Hence we restrict our investigation in two dimensions to the rational functions. The picture is completely different for $n \geq 3$. From Liouville's theorem we find that the only Id -rational mappings are Möbius transformations. The only way to produce non-injective uqr maps is to allow some distortion and very little regularity of the conformal structure. Thus we consider only measurable conformal structures. Nevertheless, it is a difficult task to write down explicit examples of uqr mappings having degree at least two. This was first done by Iwaniec and Martin [9], and subsequently other examples were given in [14], [17], [18]. In particular, Mayer's paper [17] exhibits the higher-dimensional Lattès maps. They are, as before, defined by the functional equation (1.1), where this time $h : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}^n$ is a quasimeromorphic map which is automorphic with respect to some crystallographic group.

The rigidity result we present below shows that there are in fact far fewer uqr maps with a large Julia set than there are rational functions. We first explain how

this is related to Theorem 1.2. A *quasiconformal deformation* of a rational map f is again a rational function g with $g = \varphi^{-1} \circ f \circ \varphi$ and φ a quasiconformal mapping that is not a Möbius transformation. Let us say that such a deformation is *non-trivial on the Julia set* if the Lebesgue measure of the Julia set $|\mathcal{J}_f| > 0$ and if the restriction of φ to \mathcal{J}_f is not the restriction of a holomorphic map. Note that such a conjugacy induces an f -invariant conformal structure $\mu = \varphi^* Id$, and, conversely, it follows from the Ahlfors-Bers theorem that a rational map that preserves a conformal structure can be quasiconformally deformed. Since we are deforming holomorphic functions, one can replace in this discussion the conformal structures by simpler objects such as the invariant line fields (for instance the line field generated by the major axis of the ellipse field that is induced by an invariant conformal structure). Thus we can reformulate Theorem 1.2 in the following way:

If f is a rational function with $|\Lambda_c| > 0$, then either f is a Lattès map, or the standard structure Id is the unique f -invariant conformal structure.

When the dimension $n \geq 3$, this second alternative cannot occur, since an Id -rational map is a Möbius transformation. Hence the following result is the precise higher-dimensional analogue of Theorem 1.2. One might compare this with the classical Mostow rigidity theorem.

Theorem 1.3. *Either the set of conical points Λ_c of a map $f \in UQR(\overline{\mathbb{R}^n})$, $n \geq 3$, has zero measure, or f is a chaotic Lattès map (and then $\Lambda_c = \overline{\mathbb{R}^n}$).*

This higher-dimensional rigidity result suggests that the NILF-conjecture has the following generalization:

Conjecture 1.4. Suppose $f \in UQR(\overline{\mathbb{R}^n})$, $n \geq 3$, with a Julia set of positive measure (or possibly $\mathcal{J}_f = \overline{\mathbb{R}^n}$). Then f is a Lattès map.

Note that the Lattès examples are quite rare in the space of all rational functions and even in the subspace of those functions that have a conical set of positive measure. Here are two examples.

Example 1.5. The rational functions

$$f_p(z) = \left(1 - \frac{2}{z}\right)^{2p}, \quad p = 1, 2, \dots,$$

are all semi-hyperbolic with $\mathcal{J}_{f_p} = \Lambda_c = \hat{\mathbb{C}}$. But the only Lattès map of this family corresponds to the value $p = 1$. Therefore, only f_1 has a higher-dimensional uqr analogue.

The degree of the above functions does increase with p . But we may fix this degree and see the same phenomenon from Lyubich's family

$$g_\lambda(z) = 1 - \frac{\lambda}{z^2}, \quad \lambda \in \mathbb{C}^*.$$

The parameter space \mathbb{C}^* contains a subset *Irr* of unstable values (see [11] for details). There is only one parameter $\lambda = 2 \in Irr$ that corresponds to a Lattès map. On the other hand, a dense subset of *Irr* gives rise to strictly critically finite maps which have $\Lambda_c = \hat{\mathbb{C}}$. So, all these values of the big space *Irr* (which is in fact a set of positive measure, as Rees [25] has shown) reduce to one point $\lambda = 2$ in higher dimensions.

However, the aforementioned uqr map in [9] is not obtained by a Lattès construction, and the space $UQR(\overline{\mathbb{R}^n})$ may well contain many other examples with Cantor or other small Julia sets.




FIGURE 1. The set Irr

2. QUASIREGULARITY AND CONFORMAL STRUCTURES

In this section we shall recall some of the necessary facts concerning uqr maps and conformal structures. The notation we use is quite standard. \mathbb{B}^n denotes the unit ball of \mathbb{R}^n and \mathbb{S}^{n-1} its boundary. In two dimensions we often use complex notation and write for example $\mathbb{D} = \mathbb{B}^2$.

2.1. Uniformly quasiregular mappings. Let $D \subset \overline{\mathbb{R}^n}$ be a domain and $f : D \rightarrow \overline{\mathbb{R}^n}$ a mapping of Sobolev class $W_{loc}^{1,n}(D, \overline{\mathbb{R}^n})$. We shall only consider orientation-preserving mappings; thus $\det f'(x) > 0$ for a.e. $x \in D$. Such a mapping is said to be K -quasiregular, where $1 \leq K < \infty$, if

$$\max_{|h|=1} |f'(x)h| \leq K \min_{|h|=1} |f'(x)h| \quad \text{for a.e. } x \in D.$$

The smallest number K for which the above inequality holds is called the *dilatation* of f . A non-constant quasiregular mapping can be redefined on a set of measure zero so as to make it continuous, open and discrete, and we shall always assume this to be the case. If D is a domain of the compactification $\overline{\mathbb{R}^n}$ (equipped with the spherical metric; thus $\overline{\mathbb{R}^n}$ is isometric via stereographic projection with the n -sphere \mathbb{S}^n), then we use the chart at infinity $x \mapsto x/|x|^2$ to extend the notion of quasiregularity to mappings $f : D \rightarrow \overline{\mathbb{R}^n}$ in the obvious manner. Such mappings are also said to be *quasimeromorphic*. A mapping f of a domain D into itself is called *uniformly quasiregular (uqr)* if there is some $1 \leq K < \infty$ such that all the iterates f^k are K -quasiregular. We abbreviate this to $f \in UQR(D)$.

The branch set B_f is the set of points $x \in D$ for which f is not locally homeomorphic at x . In the setting of uqr mappings there are analogues of the Picard and Montel theorems due to Rickman [26]. An easy application is that the *exceptional set* \mathcal{E}_f of a K -uqr map f , the set of points that have only a finite number of images and preimages, contains at most a finite number $q = q(K, n)$ of points. Also we have the usual property

$$(2.1) \quad \bigcup_{k \geq 0} f^k(\Omega) \supset \overline{\mathbb{R}^n} \setminus \mathcal{E}_f$$

for every open set Ω that intersects the Julia set (see [15]).

The other important normality criterion is the following quasiregular version of Zalcman's lemma. It is due to Miniowitz [23].

Zalcman's Lemma. *A family \mathcal{F} of K -quasiregular mappings $f : \mathbb{B}^n \rightarrow \overline{\mathbb{R}^n}$ is not normal at x_0 if and only if there are positive numbers $\rho_j \rightarrow 0$, points $x_j \rightarrow x_0$ and mappings $f_j \in \mathcal{F}$ such that*

$$(2.2) \quad f_j(x_j + \rho_j x) \rightarrow \Psi(x)$$

spherically uniformly on compact subsets of \mathbb{R}^n , where $\Psi : \mathbb{R}^n \rightarrow \overline{\mathbb{R}^n}$ is a non-constant quasimeromorphic map.

In fact Miniowitz's proof does not include the condition $x_j \rightarrow x_0$. However, it can easily be achieved with slight modifications.

2.2. Conformal structures. Let $\mu : \overline{\mathbb{R}^n} \rightarrow S(n)$ be a measurable conformal structure as defined in §1. If we forget for a moment the boundedness and determinant 1 condition, then μ can be viewed as a standard Riemannian metric with measurable coefficients. Since the notion of conformality does not change if we rescale the metric, this explains the canonical normalisation to require that $\det \mu(x) = 1$ for a.e. $x \in \overline{\mathbb{R}^n}$. The boundedness condition is necessary in order to retain the structures that are adapted to quasiregular mappings. It implies that there is a constant $K \geq 1$ such that

$$\frac{1}{K} |X|^2 \leq {}^t X \mu(x) X \leq K |X|^2 \quad \text{for all } X \in \mathbb{R}^n \text{ and a.e. } x \in \overline{\mathbb{R}^n}.$$

As we explained in §1, a conformal structure should really be viewed as a measurable field of ellipsoids of bounded eccentricity.

Recall the pull-back of a conformal structure μ by a quasiregular mapping f :

$$f^* \mu(x) = f'(x)[\mu \circ f(x)] := \frac{{}^t f'(x) \mu(f(x)) f'(x)}{\det f'(x)^{\frac{2}{n}}} \quad \text{for a.e. } x \in \overline{\mathbb{R}^n},$$

and note that this pull-back behaves nicely under composition:

$$(f_1 \circ f_2)^* \mu = f_2^*(f_1^* \mu).$$

Consequently, if f is μ -rational ($\mu(x) = f^* \mu(x)$, a.e.), then it is automatically uniformly quasiregular. The converse is

Theorem ([9]). *A mapping of the Sobolev class $W^{1,n}(\overline{\mathbb{R}^n}, \overline{\mathbb{R}^n})$ is uniformly quasiregular if and only if it is μ -rational for some measurable conformal structure μ .*

For completeness we recall the following theorem in the form we shall use it.

Liouville’s Theorem. *Let $f : U \rightarrow V$ be quasiregular with U, V domains of \mathbb{R}^n and $n \geq 3$. If $f^*Id = Id$, then f is the restriction of a Möbius transformation.*

We have seen that in the context of quasiregular mappings there are powerful normality criteria. Recall too that if a sequence $\{f_j\}$ of K -quasiregular mappings converges to a mapping f , then f is either constant or a quasiregular mapping. It will be important in our applications to know whether or not the conformal structures $\mu_{f_j} = f_j^*Id$ also converge. In general this is not the case (see [10, IV, 5.4]).

In order to make clear the notion of convergence in the space of matrices $S(n)$ we recall now how the metric δ is defined in that space (see Helgason [7] for details). First of all, the general linear group $Gl(n, \mathbb{R})$ acts on $S(n)$ via the rule

$$A \mapsto X[A] = |\det X|^{-\frac{2}{n}} {}^t XAX, \quad X \in Gl(n, \mathbb{R}), \quad A \in S(n),$$

and the metric δ is invariant under this action. In particular,

$$\delta(A, B) = \delta(\sqrt{B}^{-1}[A], Id) = \delta(\sqrt{B}^{-1}A\sqrt{B}^{-1}, Id), \quad A, B \in S(n),$$

where \sqrt{B} is the symmetric positive defined square root of B . The distance between $A \in S(n)$ and the Id is given by

$$\delta(A, Id) = \|\log A\| = \left(\sum_{i=1}^n (\log \lambda_i)^2 \right)^{\frac{1}{2}},$$

where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of the matrix A .

The following positive answer to the problem of convergence of conformal structures is due to Tukia [29]:

Good Approximation Lemma. *Suppose the K -quasiregular mappings f_j converge to the non-constant mapping f . Suppose further that the conformal structures $\mu_{f_j} = f_j^*Id$ converge in measure to μ . Then $\mu = \mu_f := f^*Id$.*

Proof. It suffices to verify the lemma for the restriction of the mappings to an arbitrary small domain D on which f is injective. Then the f_j are also injective provided j is big enough. But for quasiconformal mappings the good approximation lemma is precisely Corollary D of [29]. □

3. CONICAL POINTS

3.1. Definition of the conical points. The concept of conical limit points has many important applications in the setting of Kleinian groups. Recall that a point $x_0 \in \mathbb{S}^{n-1}$ is called a conical point of the Kleinian group $\Gamma \subset Möb(\mathbb{B}^n)$ if there are elements $\gamma_j \in \Gamma$ and a geodesic ray $\sigma \subset \mathbb{B}^n$ ending at x_0 so that the hyperbolic distance $\delta(\gamma_j(0), \sigma)$ stays bounded and $\gamma_j(0) \rightarrow x_0$. These conical points are precisely the points that allow a nice renormalization: there is $\rho_j \rightarrow 0^+$ so that

$$\gamma_j^{-1} \circ \alpha_j(x) = \gamma_j^{-1}(\rho_j x + x_0) \rightarrow \Psi(x),$$

a non-constant conformal map. If we compare this with Zalcman’s lemma, then we see that it is natural to consider the points x_0 of the Julia set of a given map f for which we have a renormalization (2.2) with a constant translation vector $x_j \equiv x_0$. We use a local version of this to define the set of conical points as in Lemma 1.1.

Definition 3.1. A point x_0 is called a conical point for the uqr mapping f if there are a sequence of positive numbers $\rho_j \searrow 0$ and a sequence of iterates f^{k_j} so that

$$(3.1) \quad f^{k_j}(x_0 + \rho_j x) \rightarrow \Psi(x)$$

uniformly on \mathbb{B}^n , where $\Psi : \mathbb{B}^n \rightarrow \overline{\mathbb{R}}^n$ is a non-constant quasimeromorphic mapping. The set of conical points is denoted Λ_c .

From the above discussion (see also the preamble to Lemma 1.1) we see that this definition is universal in the sense that it applies equally to Kleinian groups, rational functions, and also to uqr maps.

3.2. The case of rational maps. A nice feature of the conical set is that it is always a large subset of the Julia set, and in many cases we even have the equality $\Lambda_c = \mathcal{J}_f$. In particular, this is true for all known uqr maps in dimensions $n \geq 3$. To illustrate what happens in two dimensions, we now discuss in greater detail the notion of conical points for rational functions. First of all we formulate an equivalent definition of Λ_c :

Lemma 3.2. *Let f be a rational function, and denote by $U(x_0, f^k, \delta)$ the component of $f^{-k}(\mathbb{D}(f^k(x_0), \delta))$ that contains x_0 . The point x_0 is conical if and only if there are constants $\delta > 0$, $d \in \mathbb{N}$ and a sequence $k_j \rightarrow \infty$ so that*

$$(3.2) \quad f^{k_j} : U(x_0, f^{k_j}, \delta) \rightarrow \mathbb{D}(f^{k_j}(x_0), \delta)$$

has degree no more than d .

Proof. Clearly, if $x_0 \in \Lambda_c$, then (3.2) holds with d being the degree of the non-constant limit map at x_0 . Conversely, if (3.2) is true, then we see from the maximum principle that the components $U_j = U(x_0, f^{k_j}, \delta)$ are simply connected domains. Now choose $\alpha_j(x) = \rho_j x + x_0$ with $\rho_j > 0$ maximal so that $\alpha_j(\mathbb{D}) \subset U_j$. Then $\Psi_j = f^{k_j} \circ \alpha_j$ is a normal family, and since the degree of these maps is uniformly bounded it easily follows from the distortion lemma [2, Lemma 2.2] that any limit function of a convergent subsequence of $\{\Psi_j\}$ cannot be a constant map. \square

Several other notions of conical points appear in the literature. From (3.2) we see that the above definition appears in Appendix 3 of [24], where Przytycki compares different notions of conical points. We also see that Λ_c is somehow in the spirit of the conical set of Lyubich and Minsky [12]. McMullen [21] and independently Urbanski [3] call a point conical if the mappings in (3.2) can be chosen to be conformal. Points satisfying the condition (3.1) also have been used and studied in the recent paper [6] of Haïssinsky.

Recall that a rational map f is

- *hyperbolic* if there is $\delta > 0$ so that for any $x \in \mathcal{J}_f$ and any $k \in \mathbb{N}$ the map $f^k : U(x, f^k, \delta) \rightarrow \mathbb{D}(f^k(x), \delta)$ is injective,
- *semi-hyperbolic* if there is $\delta > 0$ so that for any $x \in \mathcal{J}_f$ and any $k \in \mathbb{N}$ the degree of $f^k : U(x, f^k, \delta) \rightarrow \mathbb{D}(f^k(x), \delta)$ is at most d_0 , for some fixed d_0 , and
- *topological Collet-Eckmann* if there are $\delta > 0$ and $N \in \mathbb{N}$ so that for any $x \in \mathcal{J}_f$ there is a subsequence k_j so that first $k_{j+1} \leq Nk_j$ and second the degree of $f^{k_j} : U(x, f^{k_j}, \delta) \rightarrow \mathbb{D}(f^{k_j}(x), \delta)$ is at most d_0 , with again d_0 some fixed maximal degree.

From (3.2) it immediately follows that $\Lambda_c = \mathcal{J}_f$ in the case of a hyperbolic, semi-hyperbolic or topological Collet-Eckmann map f . Here is another fact illustrating that Λ_c is a big subset of the Julia set for general maps. It is a consequence of work of Ledrappier (see [4, p.611]).

Theorem 3.3. *Let f be a rational function and μ any f -conformal probability measure with positive Lyapunov exponent (for example, $\mu = \mu_f$, the maximal entropy measure). Then the conical set has full μ -measure, i.e. $\mu(\Lambda_c) = 1$.*

3.3. General properties of Λ_c . We finally discuss some elementary properties of the conical set Λ_c . A nice property is the following:

Lemma 3.4. *The set of conical points is completely invariant, i.e.*

$$\Lambda_c = f(\Lambda_c) = f^{-1}(\Lambda_c).$$

Proof. This follows immediately from the standard distortion estimates (see for example [26, p.37]). □

From the renormalization description we see that conical points dynamically behave like repelling points. It is no surprise, though, that neutral fixed points cannot be in Λ_c .

Lemma 3.5. *Conical points that are also fixed points are always repelling.*

Proof. Suppose x_0 is a fixed point of f . If $x_0 \in \Lambda_c$, then there are constants $\delta > 0$, $d \in \mathbb{N}$ and a sequence of components $U_j = U(x_0, f^{n_j}, \delta)$ with $\overline{U_{j+1}} \subset U_j$ and so that $f^{n_j} : U_j \rightarrow \mathbb{B}(x_0, \delta)$ has degree less or equal to d . Observe that $f^{n_{j+1}-n_j} : U_{j+1} \rightarrow U_j$ and that all but a finite number of these mappings have to be injective. It follows that x_0 is a repelling fixed point. □

In our theorems the condition $|\Lambda_c| > 0$ appears. Let us clarify that in fact we have the following dichotomy:

Lemma 3.6. *The conical set of any uqr map f has either zero or full Lebesgue measure.*

Proof. Suppose that $|\Lambda_c| > 0$. It suffices then to renormalize a density point $x_0 \in \Lambda_c$ to find an open set Ω such that Λ_c has full measure in Ω . The conclusion follows then from (2.1) since $\mathcal{J}_f \cap \Omega \neq \emptyset$. □

4. LATTÈS EXAMPLES

Since they play an essential role in this paper, we recall here the construction and some properties of the (chaotic) Lattès examples of [17]. We refer to this paper for more details.

The construction is based on the existence of quasimeromorphic versions of the Weierstrass \mathcal{P} -function (compare [16]). This is a quasimeromorphic map $h : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}^n$ that is automorphic with respect to a crystallographic group $\Gamma \subset Isom(\mathbb{R}^n)$, i.e., h has the following two properties:

- 1) $h \circ \gamma = h$ for all $\gamma \in \Gamma$, and
- 2) Γ acts transitively on fibers: Whenever $h(x_1) = h(x_2)$, there exists an element $\gamma \in \Gamma$ such that $\gamma(x_1) = x_2$.

Due to these properties, the image $\overline{\mathbb{R}}^n = h(\mathbb{R}^n)$ can be identified with the orbifold \mathbb{R}^n/Γ . Now, every expanding similarity $A(x) = \lambda Ux$, $\lambda > 1$ and $U \in O(n)$, can be projected by h to a uqr map f provided A satisfies the condition $A \circ \Gamma \circ A^{-1} \subset \Gamma$. Then we have the following diagram:

$$(4.1) \quad \begin{array}{ccc} \mathbb{R}^n & \xrightarrow{A} & \mathbb{R}^n \\ h \downarrow & & \downarrow h \\ \overline{\mathbb{R}}^n & \xrightarrow{f} & \overline{\mathbb{R}}^n \end{array}$$

In this way we get a family of uqr maps that all have a chaotic dynamical behavior: Their Julia set is the whole space $\mathcal{J}_f = \overline{\mathbb{R}}^n$. In two dimensions, with h the holomorphic Weierstrass \mathcal{P} -function and with $A(z) = 2z$, the function f_{Lat} is precisely the function Lattès constructed in order to get the first example of a rational map with $\mathcal{J}_f = \hat{\mathbb{C}}$.

It is easy to see that Lattès maps are semi-hyperbolic. Therefore $\Lambda_c = \mathcal{J}_f = \overline{\mathbb{R}}^n$.

For these Lattès maps, one knows precisely all the possible invariant conformal structures (see [17, Thm. 3]). The canonical way to associate an invariant conformal structure μ to a map f that is defined by (4.1) is to push forward the standard Euclidean structure: $Id = h^*\mu$. The special property of Lattès maps is that they may have several invariant conformal structures. In fact, several crystallographic groups Γ preserve other (constant) structures M than just the Euclidean Id structure. If A also preserves it, then M induces an f -invariant structure ν which is different from the previous one and given by $M = h^*\nu$. In the case of Lattès' original example f_{Lat} it is easy to check that any constant matrix $M \in S(2)$ gives rise to an invariant conformal structure.

In the case where h is automorphic with respect to a group that is not cocompact, one obtains two other types of examples: Maps that behave like power or Tchebyshev polynomials. This larger class of maps is called *Lattès-type* maps.

5. RIGIDITY ON THE CONICAL SET OF RATIONAL FUNCTIONS

We prove here a pointwise rigidity result which implies Theorem 1.2.

Theorem 5.1 (Pointwise rigidity for rational maps). *Let f be a rational function and μ an f -invariant nontrivial dilatation. If μ is continuous in measure at a conical point $x_0 \in \Lambda_c$, then f is a Lattès map.*

Recall that a measurable function μ is *continuous in measure* at a point x_0 if for every $\varepsilon > 0$

$$\lim_{r \rightarrow 0} \frac{1}{|\mathbb{D}(x_0, r)|} |\{x \in \mathbb{D}(x_0, r) ; \delta(\mu(x), \mu(x_0)) > \varepsilon\}| = 0,$$

and that a measurable function is continuous in measure almost everywhere (see [5, Thm. 2.9.13], where this notion is called approximate continuity).

Theorem 1.2 is a direct consequence of the above result:

Proof of Theorem 1.2. Suppose that the rational map f has an invariant line field. Then it also has a (nontrivial) f -invariant conformal structure μ . This function μ is measurable and hence continuous in measure almost everywhere. Since $|\Lambda_c| > 0$, there is $x_0 \in \Lambda_c$ with μ continuous in measure at x_0 . The result follows now from Theorem 5.1. \square

A further application is that a quasiconformal deformation cannot be smooth at a conical point.

Corollary 5.2. *Suppose f and g are rational functions and φ a quasiconformal map (but not a Möbius transformation) such that $g \circ \varphi = \varphi \circ f$. If the derivative of φ is continuous in measure at a conical point of f , then f and g are Lattès maps.*

Proof. The dilatation $\mu = \varphi^* Id$ is g -invariant and continuous in measure at a conical point. The corollary now follows from Theorem 5.1. □

There is an analogous result in the context of Kleinian groups due to Ivanov [8]. He also has a rigidity result at so-called horospherical points involving some additional smoothness of the conjugating map φ . One might introduce the notion of horospherical point in holomorphic dynamics and generalize his results to holomorphic dynamics. We shall address this elsewhere.

Proof of Theorem 5.1. Let $x_0 \in \Lambda_c$ with μ continuous in measure at x_0 . We first renormalize at x_0 : There are $n_j \rightarrow \infty$ and $\alpha_j(x) = r_j x + x_0$ so that

$$\Psi = f^{n_j} \circ \alpha_j \rightarrow \Psi : \mathbb{D} \rightarrow \Omega = \Psi(\mathbb{D})$$

uniformly with Ψ a non-constant holomorphic map. Consider the pull-back dilatations $\nu = \Psi^* \mu$ and $\nu_j = \Psi_j^* \mu$. From the f -invariance of μ it follows that

$$\nu_j = \alpha_j^* [(f^{n_j})^* \mu] = \alpha_j^* \mu = \mu \circ \alpha_j.$$

Let $\varepsilon > 0$ and consider the set

$$\mathcal{E}_{\varepsilon,j} = \{x \in \mathbb{D} ; \delta(\nu_j(x), \mu(x_0)) > \varepsilon\} = \{w \in D_j = \alpha_j(\mathbb{D}) ; \delta(\mu(w), \mu(x_0)) > \varepsilon\}.$$

Then, from the continuity in measure of μ at x_0 , we see that

$$|\mathcal{E}_{\varepsilon,j}| = \frac{1}{|D_j|} |\{w \in D_j = \alpha_j(\mathbb{D}) ; \delta(\mu(w), \mu(x_0)) > \varepsilon\}| \rightarrow 0$$

when $j \rightarrow \infty$, meaning that ν_j converges in measure to the constant dilatation $\nu \equiv \mu(x_0)$.

We claim that this constant dilatation $\nu \equiv \mu(x_0)$ is non-trivial. In fact, suppose that $\nu \equiv Id$ on \mathbb{D} . The map Ψ being holomorphic, this would imply $\mu \equiv Id$ on the domain Ω . Since $\Omega \cap \mathcal{J}_f \neq \emptyset$, there is for any $\varepsilon > 0$ an integer n so that $f^n(\Omega) \supset \hat{\mathbb{C}} \setminus \mathcal{N}_\varepsilon(\mathcal{E}_f)$, with $\mathcal{N}_\varepsilon(\mathcal{E}_f)$ the ε -neighborhood of the exceptional set. It follows that $\mu \equiv Id$, contradicting our standard hypothesis that μ is nontrivial.

Now choose a point $w \in \Omega$ of a repelling cycle and suppose that this is a fixed point of $g = f^k$. Up to composition of Ψ by a translation we may suppose that $\Psi(0) = w$. In a neighborhood U of this origin we can define φ by

$$(5.1) \quad \Psi \circ \varphi(x) = g \circ \Psi(x) \text{ for } x \in U.$$

For this holomorphic map φ , the dilatation ν is invariant, i.e. $\varphi^* \nu = \nu$. Therefore, if θ_x is the direction of the major axis of the ellipse field generated by μ , we have

$$\theta_{\varphi(x)} = \theta_x + \arg \varphi'(x) \text{ mod } \pi.$$

In fact, we shall consider the associated invariant line field. Remember that ν is constant, and so $\theta_{\varphi(x)} = \theta_x$. It follows that

$$(5.2) \quad \varphi(x) = \lambda x \text{ for some real } \lambda.$$

From (5.1) we see that Ψ linearizes g at w , and this map can be extended in the usual way using (5.1) to a global map $\Psi : \mathbb{C} \rightarrow \hat{\mathbb{C}} \setminus \mathcal{E}_f$ such that $g \circ \Psi(x) = \Psi\lambda(x)$ for every $x \in \mathbb{C}$.

We finally verify that Ψ is in fact an elliptic function. Consider two points x_1 and x_2 (neither one critical) with $\Psi(x_1) = \Psi(x_2)$. Then one can define near x_1 a map γ with $\Psi \circ \gamma = \Psi$. This holomorphic map γ also preserves ν , and, as before, it follows that $\gamma(x) = \alpha x + \beta$ (with α real). The set of all possible such mappings

$$\Gamma = \{\gamma \text{ affine with } \Psi \circ \gamma = \Psi\}$$

is a discrete group that acts transitively on fibers of Ψ . Consequently, the elements of Γ have to be isometries, and g is a power, Tchebyshev or Lattès map. Moreover, since power and Tchebyshev maps are (globally) rigid, g has to be a Lattès map.

It then follows that f itself is a Lattès map. For example, using again the invariant line field argument (5.2), one sees that Ψ linearizes f at any (necessarily repelling) fixed point, and hence the automorphic function Ψ semiconjugates a scalar multiplication and f . \square

6. RIGIDITY OF UNIFORMLY QUASIREGULAR MAPPINGS

In this section we show that the chaotic Lattès maps are the only uqr mappings with a set of conical points of positive measure.

Theorem 6.1 (Pointwise rigidity for uqr maps). *Suppose f is a μ -rational map of $\overline{\mathbb{R}}^n$, $n \geq 3$, and that μ is continuous in measure at a conical point $x_0 \in \Lambda_c$. Then f is a Lattès-type map.*

As a first consequence we have the following.

Proof of Theorem 1.3. Since a measurable function is almost everywhere continuous in measure, Theorem 1.3 follows immediately from Theorem 6.1. \square

There is another rigidity phenomenon which follows from Theorem 6.1: For a hyperbolic Blaschke product B of the unit disc, one knows that the induced circle action $B : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ is quasimetrically conjugate to $z \mapsto z^d$, where d is the degree of B . But this conjugacy is always singular, except when B is actually the power map $z \mapsto z^d$ (see [28]). In higher dimensions we also have natural uqr power maps [17]. Their Julia (and conical) set is the sphere \mathbb{S}^{n-1} , and their restrictions to this sphere are Lattès maps. Theorem 6.1 implies that all higher-dimensional “hyperbolic uqr-Blaschke products” share this property:

Corollary 6.2. *Let f be a uqr map of $\overline{\mathbb{R}}^n$, let $n \geq 4$ and suppose that $\Lambda_c = \mathcal{J}_f = \mathbb{S}^{n-1}$. Then the restriction of f to \mathbb{S}^{n-1} is a Lattès map.*

There is one dimension left, and it is natural to ask what happens when $n = 3$. This problem leads to the question of which chaotic rational maps do admit a uqr extension to the 3-ball. Lattès maps have such an extension, namely the power mappings of [17]. The above corollary suggests that these are the only exceptions.

Proof of Theorem 6.1. We start with the same renormalization argument as in the Proof of Theorem 5.1: Since $x_0 \in \Lambda_c$, there are $\rho_j \rightarrow 0^+$ and $k_j \rightarrow \infty$ so that

$$\Psi_j(x) = f^{k_j} \circ \alpha_j(x) = f^{k_j}(\rho_j x + x_0) \rightarrow \Psi(x), \quad x \in \mathbb{B}^n,$$

uniformly on \mathbb{B}^n , where Ψ is non-constant and quasimeromorphic. Consider again the pull-back structures $\nu = \Psi^*\mu$ and $\nu_j = \Psi_j^*\mu$. From the f -invariance of μ and the fact that μ is continuous in measure at x_0 , it follows, precisely as in the proof of Theorem 5.1, that ν_j converges to the constant structure $x \mapsto \mu(x_0)$. The good approximation lemma asserts that

$$\nu = \lim_{j \rightarrow \infty} \nu_j \equiv \mu(x_0).$$

Now choose $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ affine with $L^*\mu(x_0) = Id$, and denote

$$h = \Psi \circ L : \mathbb{B}^n \rightarrow \Omega = h(\mathbb{B}^n).$$

Then $h^*\mu := (\Psi \circ L)^*\mu = Id$.

The next step is to lift some iterate of f by h . Let $V = \mathbb{B}(h(0), \delta) \subset \Omega$, and U the connected component of $h^{-1}(V)$ that contains 0. If δ is sufficiently small, then $h : U \rightarrow V$ is a proper map (see [26]). Since

$$\Psi_{j+N}(x) = f^{k_{j+N}-k_j} \circ \Psi_j\left(\frac{\rho_{j+N}}{\rho_j}x\right),$$

we can choose $l = k_{j+N} - k_j \gg 0$ so that $\overline{V_0} \subset V$, where V_0 is the component of $f^{-l}(V)$ that contains $\Psi_j(0)$. Denote finally by $U_0 \subset U$ a component of $h^{-1}(V_0)$. We therefore have the following diagram of proper mappings:

$$(6.1) \quad \begin{array}{ccc} U_0 & & U \\ h \downarrow & & \downarrow h \\ V_0 & \xrightarrow{f^l} & V \end{array}$$

In this picture there is one map missing, say $A : U_0 \rightarrow U$. We obtain it in lifting f^l . In order to do this, let $W \subset V$ be a domain such that a branch $H : W \rightarrow W' = H(W) \subset U$ of h^{-1} exists, and denote by $W'_0 \subset U_0$ a component of $(f^l \circ h)^{-1}(W)$. Then we can define

$$A := H \circ f^l \circ h : W'_0 \rightarrow W'.$$

Since the h -pull-back of μ is the standard structure and μ is f -invariant, the lifted map A has the property $A^*Id = Id$.

From Liouville's theorem it follows that A is (the restriction of) a Möbius transformation. Using analytic continuation (cf. the proof of Proposition 6.3 in [18]), we see that

$$(6.2) \quad h \circ A(x) = f^l \circ h(x) \quad \text{for every } x \in U_0.$$

Since $\overline{U_0} \subset U$, the map A is a loxodromic Möbius transformation with a repelling fixed point $a_0 \in U_0$. We may suppose that the repelling fixed point is ∞ .

The map h linearizes f^l at the repelling fixed point $b = h(a)$ and, using (6.2), we can extend this map to $h : \mathbb{R}^n \rightarrow \overline{\mathbb{R}^n}$ so that the conjugation (6.2) is true globally.

At this stage we can conclude using [18, §6]. Let us just briefly outline how to proceed. Suppose that x_1, x_2 are two points that are not branch points of h and so that $h(x_1) = h(x_2) = y$. Then we can define locally an inverse branch H of h with $H(y) = x_2$. This defines locally near x_1 a map $\gamma = H \circ h$ sending x_1 onto x_2 . Clearly γ is an Id -rational map, and hence the restriction of a Möbius transformation. Using analytic continuation again, we see that $h \circ \gamma = h$ is true in the whole of \mathbb{R}^n . It suffices now to consider the set

$$\Gamma = \{\gamma \in \text{Möb}(\mathbb{R}^n) ; h \circ \gamma = h\}.$$

It turns out that Γ is a discrete group of isometries and that h is automorphic with respect to this group.

We have shown that f^l is of Lattès-type, and therefore so is f (see [18, Prop. 5.1]). \square

REFERENCES

- [1] F. Berteloot and J.-J. Loeb, *Spherical Hypersurfaces and Lattès Rational Maps*, J. Math. Pures Appl. 77 (1998), 655-666. MR **99e**:32043
- [2] L. Carleson, P.W. Jones, and J.-Ch. Yoccoz, *Julia and John*, Bol. Soc. Bras. Mat. 25 (1994), 1-30. MR **95d**:30040
- [3] M. Denker, R.D. Mauldin, Z. Nitecki, and M. Urbanski, *Conformal measures for rational functions revisited*, Fund. Math. 157 (1998), 161-173. MR **99j**:58122
- [4] A.E. Eremenko and M. Yu. Lyubich, *The dynamics of analytic transformations*, Leningrad Math. J. 1 (1990), 563-634. MR **91b**:58109
- [5] H. Federer, *Geometric Measure Theory*, Springer Verlag (1969). MR **41**:1976
- [6] P. Haïssinsky, *Rigidity and expansion for rational maps*, J. London Math. Soc. (2) 63 (2001), 128-140. MR **2001m**:37085
- [7] S. Helgason, *Differential Geometry, Lie Groups and Symmetric Spaces*, Academic Press, 1978. MR **80k**:53081
- [8] N. I. Ivanov, *Action of Möbius Transformations on Homeomorphisms: Stability and Rigidity*, Geom. Funct. Anal. 6 (1996), 79-119. MR **97e**:30075
- [9] T. Iwaniec and G.J. Martin, *Quasiregular Semigroups*, Ann. Acad. Sci. Fenn. Math. 21 (1996), 241-254. MR **97i**:30032
- [10] O. Lehto and K.I. Virtanen, *Quasiconformal mappings in the plane*, Springer-Verlag, 1973. MR **49**:9202
- [11] M. Lyubich, *The dynamics of rational transforms: the topological picture*, Russian Math. Surveys 41 : 4 (1986), no. 4, 43-117. MR **88g**:58094
- [12] M. Lyubich and Y. Minsky, *Laminations in holomorphic dynamics*, J. Diff. Geometry 47 (1997), 17-94. MR **98k**:58191
- [13] R. Mañé, P. Sad and D. Sullivan, *On the dynamics of rational maps*, Ann. Scient. Éc. Norm. Sup. (4) 16 (1983), 193-217. MR **85j**:58089
- [14] G.J. Martin, *Branch sets of uniformly quasiregular maps*, Conformal Geometry and Dynamics 1 (1997), 24-27. MR **98d**:30032
- [15] G.J. Martin and V. Mayer, *Local Dynamics of Uniformly Quasiregular mappings*, preprint.
- [16] O. Martio and U. Srebro, *Periodic Quasimeromorphic Mappings in \mathbb{R}^n* , J. d'Analyse Math. 28 (1975), 20-40.
- [17] V. Mayer, *Uniformly Quasiregular mappings of Lattès' type*, Conformal Geometry and Dynamics 1 (1997), 104-111. MR **98j**:30017
- [18] V. Mayer, *Quasiregular analogues of critically finite rational functions with parabolic orbifold*, J. d'Analyse Math. 75 (1998), 105-119. MR **2000a**:30043
- [19] C. McMullen, *Area and Hausdorff dimension of Julia sets of entire functions*, Trans. Amer. Math. Soc. 300 (1987), 329-342. MR **88a**:30057
- [20] C. McMullen, *Complex Dynamics and Renormalization*, Ann. Math. Studies, Princeton University Press (1994). MR **96b**:58097
- [21] C. McMullen, *Hausdorff dimensions and conformal dynamics II: Geometrically finite rational maps*, Comment. Math. Helv. 75 (2000), 535-593. MR **2001m**:37089
- [22] C. McMullen and D. Sullivan, *Quasiconformal Homeomorphisms and Dynamics III. The Teichmüller Space of a Holomorphic Dynamical System*, Adv. in Math. 135 (1998), 351-395. MR **99e**:58145
- [23] R. Miniowitz, *Normal families of quasimeromorphic mappings*, Proc. AMS 84 (1982), 35-43. MR **83c**:30026
- [24] F. Przytycki, *Conical limit set and Poincaré exponent for iterations of rational functions*, Trans. Amer. Math. Soc. 351 (1999), 2081-2099. MR **99h**:58110
- [25] M. Rees, *Positive measure sets of ergodic rational maps*, Ann. Scient. Éc. Norm. Sup. 19 (1986), 383-407. MR **88g**:58100
- [26] S. Rickman, *Quasiregular Mappings*, Springer-Verlag (1993). MR **95g**:30026
- [27] J.L. Schiff, *Normal families*, Universitext Springer-Verlag (1993). MR **94f**:30046

- [28] M. Shub and D. Sullivan, *Expanding endomorphisms of the circle revisited*, Ergodic Theory and Dynamical Systems 5 (1985), 285-289. MR **87g**:58104
- [29] P. Tukia, *On quasiconformal groups*, J. d'Analyse Math. 46 (1986), 318-346. MR **87m**:30043
- [30] L. Zalcman, *A heuristic principle in complex function theory*, Amer. Math. Monthly 82 (1975), 813-817. MR **52**:757

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AUCKLAND, AUCKLAND, NEW ZEALAND
E-mail address: martin@math.auckland.ac.nz

UMR 8524 DU CNRS - UFR DE MATHÉMATIQUES, UNIVERSITÉ DES SCIENCES ET TECHNOLOGIES DE LILLE, 59655 VILLENEUVE D'ASCQ CEDEX, FRANCE
E-mail address: volker.mayer@univ-lille1.fr