

CLOSED PRODUCT FORMULAS FOR EXTENSIONS OF GENERALIZED VERMA MODULES

RICCARDO BIAGIOLI

ABSTRACT. We give explicit combinatorial product formulas for the polynomials encoding the dimensions of the spaces of extensions of (g, p) -generalized Verma modules, in the cases when (g, p) corresponds to an indecomposable classic Hermitian symmetric pair. The formulas imply that these dimensions are combinatorial invariants. We also discuss how these polynomials, defined by Shelton, are related to the parabolic R -polynomials introduced by Deodhar.

1. INTRODUCTION

Let g be a complex, semisimple Lie algebra with Cartan subalgebra h and let b be a Borel subalgebra of g with nilradical n and Levi component h . Let p be a parabolic subalgebra of g containing b with nilradical $u \subseteq n$ and Levi component $m \supseteq h$. So we have Levi decompositions $b = h \oplus n$ and $p = m \oplus u$. The problem of computing the u -cohomology of irreducible highest weight (g, p) -modules is completely solved by the Kazhdan-Lusztig conjectures ([7]). On the other hand, the “Extension Problem”, namely the problem of computing the u -cohomology of the (g, p) -generalized Verma modules, is as yet unsolved in general, and there are not even conjectures for the form of the answer. If $p = b$, the modules in question are the Verma modules (see [9]). There was a conjecture that solved the problem, known as the Gabber-Joseph conjecture (see, e.g., [14] or [8]), but this was disproved by Boe in [2]. In the cases when (g, p) corresponds to an indecomposable Hermitian symmetric pair (see, e.g., [9, §4]), Shelton gives a solution to this problem. His answer is in terms of a recursion relations for the polynomials encoding the dimensions of the spaces of extensions between generalized Verma modules [17].

In this paper we solve these recursion relations, in the cases when (g, p) is an indecomposable classic Hermitian symmetric pair (see Table 1 at the end of §2). Our main results are explicit combinatorial product formulas for these polynomials. These formulas can be stated in two different ways, one in terms of Weyl group elements and one in terms of partitions or shifted partitions. Moreover, these formulas imply that these polynomials are combinatorial invariants.

The organization of the paper is as follows. In the next section we collect some definitions and results that are needed in the rest of the work. In §3 we study the case of the symmetric pair $(A_{n-1}, A_{i-1} \times A_{n-i-1})$, in §4 the case of (C_n, A_{n-1}) , in §5 the case of (D_n, A_{n-1}) , and in §6 the remaining cases (B_n, B_{n-1}) and (D_n, D_{n-1}) . In the first three cases we give two statements of these results, one in terms of

Received by the editors September 24, 2001 and, in revised form, February 11, 2002.
2000 *Mathematics Subject Classification*. Primary 17B10, 05E99; Secondary 22E47, 20F55.

permutations or signed permutations and one in terms of partitions or shifted partitions. In §7 we derive some consequences of our results and in particular we prove that these polynomials are combinatorial invariants. We also discuss how they are related to the parabolic R -polynomials introduced in [12], and the pleasant “symmetry” that follows with the Gabber-Joseph conjecture [14].

2. NOTATION, DEFINITIONS, AND PRELIMINARIES

In this section we give some definitions, notation and results that will be used in the rest of this work. We let $\mathbf{P} := \{1, 2, 3, \dots\}$, $\mathbf{N} := \mathbf{P} \cup \{0\}$, and \mathbf{Z} be the set of integers; for $a \in \mathbf{N}$ we let $[a] := \{1, 2, \dots, a\}$ (where $[0] := \emptyset$) and $[\pm n] := [-n, n] \setminus \{0\}$. The cardinality of a set A will be denoted by $|A|$. For any sequence $(a_1, \dots, a_n) \in \mathbf{Z}^n$, we let $N_1(a_1, \dots, a_n) := |\{i \in [n] : a_i < 0\}|$. Let S_n be the set of all bijections $\sigma : [n] \rightarrow [n]$. If $\sigma \in S_n$, then we write $\sigma = \sigma_1 \dots \sigma_n$ to mean that $\sigma(i) = \sigma_i$, for $i = 1, \dots, n$. If $\sigma \in S_n$, then we will also write σ in *disjoint cycle form* (see, e.g., [18], p.17) and we will usually omit writing the 1-cycles of σ . For example, if $\sigma = 64175823$, then we also write $\sigma = (2, 4, 7)(1, 6, 8, 3)$. Given $\sigma, \tau \in S_n$ we let $\sigma\tau := \sigma \circ \tau$ (composition of functions) so that, for example, $(1, 2)(2, 3) = (1, 2, 3)$. We follow Chapter 3 of [18] for poset notation and terminology. In particular, given a poset (P, \leq) and $u, v \in P$ we let $[u, v] := \{z \in P : u \leq z \leq v\}$ and call this an *interval* of P . If P has a minimal element, denoted $\hat{0}$, then we call a subset of the form $[\hat{0}, u]$, for $u \in P$, a *lower interval* of P . Given any $Q \subseteq P$ we will always consider Q as a poset with the partial ordering induced by P and call Q a *subposet* of P . We say that $z \in P$ is *join-irreducible* if it covers at most one element of P . Given two posets P and Q we write $P \cong Q$ to mean that they are isomorphic as posets.

We follow §7.2 of [19] for any undefined notation and terminology concerning partitions. By an (integer) *partition* we mean a sequence of positive integers $\lambda = (\lambda_1, \dots, \lambda_k)$ such that $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$ and we write $|\lambda| = \sum \lambda_i$. The number of parts of λ is the *length* of λ , denoted $l(\lambda)$. We identify a partition λ with its *diagram*,

$$\{(i, j) \in \mathbf{P}^2 : 1 \leq i \leq k, 1 \leq j \leq \lambda_i\}.$$

If we replace the dots (i, j) by juxtaposed squares, we obtain the *Young diagram* of λ , and we draw it rotated counterclockwise by $\frac{\pi}{4}$ radians. So, for example, the diagram of $(5, 4, 2, 1)$ is illustrated in Figure 1.

We call the elements of \mathbf{P}^2 , and hence of λ , cells. Expressions such as “to the left of”, or “directly above”, always refer to these rotated diagrams. We denote by \mathcal{P} the set of all integer partitions. For any $\mu, \lambda \in \mathcal{P}$ we define $\mu \subseteq \lambda$ if and only




FIGURE 1.

if $\mu_i \leq \lambda_i$ for all i . If we identify a partition with its (Young) diagram, then the partial order \subseteq is given simply by containment of diagrams. It is well known, and not hard to see, that this makes \mathcal{P} into a lattice, usually called *Young's lattice* (see, e.g., [19, §7.2]). Given $n \in \mathbf{P}$ and $i \in [n - 1]$ we let

$$\mathcal{P}(n, i) := \{\mu \in \mathcal{P} : \mu \subseteq (n - i)^i\}.$$

Given $\lambda = (\lambda_1, \dots, \lambda_k) \in \mathcal{P}$ we let $d(\lambda)$ be the *length of the Durfee square* of λ , so

$$(1) \quad d(\lambda) := \max \{i \in [k] : \lambda_i \geq i\}.$$

Let $\mu, \lambda \in \mathcal{P}$, $\mu \subseteq \lambda$. We then call $\lambda \setminus \mu$ a *skew partition*. Note that, in poset theoretic language, partitions (respectively, skew-partitions) are the finite order ideals (respectively, finite convex subsets) of \mathbf{P}^2 . Given two skew partitions $\rho, \nu \subseteq \mathbf{P}^2$ we write $\rho \approx \nu$ if ρ is a translate of ν . Given a skew partition $\nu \subseteq \mathbf{P}^2$ its *conjugate* is

$$\nu' := \{(j, i) \in \mathbf{P}^2 : (i, j) \in \nu\}.$$

We follow [15] for general Coxeter groups notation. In particular, given a Coxeter system (W, S) and $\sigma \in W$ we denote by $l(\sigma)$ the *length* of σ in W , with respect to S , and we let $D(\sigma) := \{s \in S : l(\sigma s) < l(\sigma)\}$. We denote by e the identity of W , and we let $T := \{\sigma s \sigma^{-1} : \sigma \in W, s \in S\}$ be the set of reflections of W . Given $J \subseteq S$ we let W_J be the *parabolic* subgroup generated by J and

$$W^J := \{\sigma \in W : l(s\sigma) > l(\sigma) \text{ for all } s \in J\}.$$

Note that $W^\emptyset = W$. The *quotient* W^J is a poset and it is partially ordered by (strong) *Bruhat order*. Recall (see, e.g., [15, §5.9]) that this means that $x \leq y$ if and only if there exist $r \in \mathbf{N}$ and $t_1, \dots, t_r \in T$ such that $t_r \dots t_1 x = y$ and $l(t_i \dots t_1 x) > l(t_{i-1} \dots t_1 x)$ for $i = 1, \dots, r$. In this order, e is the minimal element and, if W is finite, w_0^J and w_J^0 denote the unique maximal elements in W^J and W_J , respectively. Given $u, v \in W^J$, $u \leq v$, we let

$$[u, v]^J := \{z \in W^J : u \leq z \leq v\},$$

and consider $[u, v]^J$ as a poset with the partial ordering induced by W^J .

In the remainder of this section, we recall a result of Shelton, for which we need some notation from [13] and [17]. Let g be a complex, semisimple Lie algebra with Cartan subalgebra h , and let b be a Borel subalgebra of g with nilradical n and Levi component h . Let p be a parabolic subalgebra of g with nilradical u and Levi component m . Let W and W_J be the Weyl groups of g and m , respectively. Let $\Delta, \Delta(m)$ be the root systems of g and m , with positive systems Δ^+ and $\Delta^+(m)$, determined by n . Let 2ρ be the sum of the positive roots in g and fix a Δ^+ -antidominant integral weight $\lambda \in h^*$. For any $\mu \in h^*$ we denote by $M(\mu)$ the g -Verma module with highest weight $\mu - \rho$. Then $L(\mu)$ and $N(\mu)$ denote the unique irreducible quotient and the maximal m -locally finite quotient of $M(\mu)$, respectively. If μ is $\Delta^+(m)$ -dominant regular and integral, then $N(\mu)$ is the (g, p) -generalized Verma module of highest weight $\mu - \rho$. For $u \in W$ and $w \in W^J$ we denote by M_u the module $M(u\lambda)$ and by N_w the module $N(w_J^0 w \lambda)$. All Ext^* groups will be computed in \mathcal{O}_m , the usual Bernstein-Gelfand-Gelfand category of g -modules relative to the pair (g, p) (see, e.g., [1]). The following result is due to Shelton, and we refer the reader to [17] for its proof. He defines for any u and v in

W^J , a polynomial $E_{u,v}(q)$ by

$$(2) \quad E_{u,v}(q) = \sum_{k \geq 0} (-1)^{l(v)-l(u)-k} q^k \dim(\text{Ext}^k(N_u, N_v)),$$

and proves that

Theorem 2.1. *Suppose that (g, p) corresponds to an indecomposable Hermitian symmetric pair. Then for all $u, v \in W^J$:*

- i) $E_{u,v}(q) = 0$ if $u \not\leq v$;
- ii) $E_{u,u}(q) = 1$;
- iii) if $u < v$ and $s \notin D(u)$ with $us \in W^J$, then

$$E_{u,v}(q) = \begin{cases} E_{us,vs}(q) & \text{if } s \notin D(v) \text{ and } vs \in W^J; \\ (q-1)E_{us,v}(q) & \text{if } s \in D(v) \text{ and } us \not\leq vs; \\ (q-q^{-1})E_{us,v}(q) + E_{us,vs}(q) & \text{if } u \leq us \leq vs \leq v; \\ qE_{us,v}(q) & \text{if } vs \notin W^J. \end{cases}$$

In this paper we solve these recurrence relations in the cases when (g, p) is an indecomposable classic Hermitian symmetric pair (see Table 1). Our main results are explicit product formulas for these polynomials. Moreover, these formulas imply that the polynomials are combinatorial invariants.

TABLE 1. The indecomposable classic Hermitian Symmetric pairs (g, p) .

(g, p)	g	$[m, m]$
$SU(r, s)$	A_N	$A_{r-1} \times A_{s-1}$
$SO(2n-1, 2)$	B_n	B_{n-1}
$Sp(2n, \mathbf{R})$	C_n	A_{n-1}
$SO(2n-2, 2)$	D_n	D_{n-1}
$SO^*(2n)$	D_n	A_{n-1}

3. THE CASE $(A_{n-1}, A_{i-1} \times A_{n-i-1})$

In this section, we give an explicit product formula for the polynomials $E_{u,v}(q)$ in the case of the pair $(A_{n-1}, A_{i-1} \times A_{n-i-1})$. We give two different formulations of this result, one in terms of permutations and one in terms of partitions. Throughout this section we fix $n \in \mathbf{P}$ and $i \in [n-1]$, and we let $W := S_n$, $s_i := (i, i+1)$ for $i = 1, \dots, n-1$, $S := \{s_1, \dots, s_{n-1}\}$, and $J := S \setminus \{s_i\}$. It is well known that (S_n, S) is a Coxeter system of type A_{n-1} and that the following characterization holds (see, e.g., [15]).

Proposition 3.1. *Let $v \in S_n$. Then*

$$D(v) = \{s_i \in S : v(i) > v(i+1)\}.$$

For symmetric groups, the parabolic subgroups of S_n are called *Young subgroups*. In the case of maximal parabolic subgroups (i.e. $J = S \setminus \{s_i\}$) their quotients take this particularly simple form:

$$W^J = \{w \in W : w^{-1}(1) < \dots < w^{-1}(i) \text{ and } w^{-1}(i+1) < \dots < w^{-1}(n)\}.$$

Given $v \in W^J$ we associate to v the partition

$$(3) \quad \Lambda(v) := (v^{-1}(i) - i, \dots, v^{-1}(2) - 2, v^{-1}(1) - 1).$$



FIGURE 2. The lattice path corresponding to $(5, 4, 2, 1)$ if $n = 9$ and $i = 4$.




FIGURE 3. The lattice path corresponding to $(5, 4, 2, 1)$ if $n = 12$ and $i = 5$.

The following is well known (see [4, Proposition 2.8]).

Proposition 3.2. *The map Λ defined by (3) is a bijection between W^J and $\mathcal{P}(n, i)$. Furthermore $u \leq v$ in W^J if and only if $\Lambda(u) \subseteq \Lambda(v)$, and $l(v) = |\Lambda(v)|$ for all $u, v \in W^J$.*

We will find it sometimes convenient to identify a partition $\lambda \in \mathcal{P}(n, i)$ with a lattice path, with $(1, 1)$ and $(1, -1)$ steps. This path is the union of the lower boundary of the diagram of the skew-partition $(n - i)^i \setminus \lambda$ and the upper boundary of the partition λ . Note that it starts at $(0, 0)$ and ends at $(n, 2i - n)$ (equivalently, it has n steps and exactly i are $(1, 1)$ -steps). We call a $(1, 1)$ -step (respectively, $(1, -1)$ -step) an *up-step* (respectively, *down-step*). Given $j \in [n - 1]$ we say that λ has a *peak* at j if the j -th step of λ is up and its $(j + 1)$ -th step is down. For example, if we take the partition of Figure 1, $\lambda = (5, 4, 2, 1) \in \mathcal{P}(9, 4)$, then the associated path is the one shown in Figure 2 and it has peaks at 1, 3, 6, and 8. Note that this identification between partitions and paths depends on n and i . For example, the same partition $(5, 4, 2, 1)$ corresponds to the path in Figure 3 if $n = 12$ and $i = 5$. Since n and i are fixed throughout this section, this will cause no confusion.

The following elementary lemma is known (see [4, Lemma 2.9]).

Lemma 3.3. *Let $v \in W^J$, and $j \in [n - 1]$. Then $s_j \in D(v)$ if and only if $\Lambda(v)$ has a peak at $n - j$.*

Note that the k -th step of $\Lambda(v)$ is an up step if and only if

$$k \in \{n + 1 - v^{-1}(i), n + 1 - v^{-1}(i - 1), \dots, n + 1 - v^{-1}(1)\}.$$

Let $u, v \in W^J$, $u \leq v$. For $j \in [n]$ we let, following [4],

$$(4) \quad a_j(u, v) := |\{r \in u^{-1}([i]) : r < j\}| - |\{r \in v^{-1}([i]) : r < j\}|.$$

For example, if $n = 9$, $i = 4$, $u = 123564789$, and $v = 516278394$, then

$$(5) \quad (a_1(u, v), \dots, a_9(u, v)) = (0, 1, 1, 2, 1, 1, 2, 1, 1).$$

Note that it follows easily from Proposition 3.2 that $a_j(u, v) \geq 0$ for $j = 1, \dots, n$ if and only if $u \leq v$, and that $a_j(u, v) > 0$ if $j \in v^{-1}([i]) \setminus u^{-1}([i])$ and $u \leq v$. Also note that, if $u \in W^J$ and $j \in [n]$, then

$$|\{r \in u^{-1}([i]) : r < j\}| = \begin{cases} u(j) - 1, & \text{if } j \in u^{-1}([i]), \\ j + i - u(j), & \text{if } j \notin u^{-1}([i]). \end{cases}$$

This may be used to obtain a more explicit formula for $a_j(u, v)$, if desired.

We can now state and prove the main result of this section.

Theorem 3.4. *Let $u, v \in W^J$, $u \leq v$. Then*

$$(6) \quad E_{u,v}(q) = q^{l(v)-l(u)} \prod_{j \in v^{-1}([i]) \setminus u^{-1}([i])} (1 - q^{-2a_j(u,v)+1}).$$

Proof. Let, for brevity, $D_i(u, v) := v^{-1}([i]) \setminus u^{-1}([i])$. We proceed by induction on $l(w_0^J) - l(u)$. If $l(w_0^J) - l(u) = 0$, we have $w_0^J = v = u$ and the result is trivially true. So suppose that $l(w_0^J) - l(u) > 0$ and let $s = (k, k+1)$ be such that $s \notin D(u)$ and $us \in W^J$. Note that, since $u \in W^J$, this implies that $k \in u^{-1}([i])$ and $k+1 \notin u^{-1}([i])$. We have four cases to consider.

a) $s \notin D(v)$, and $vs \in W^J$.

Since $v \in W^J$, this implies that $k \in v^{-1}([i])$ and $k+1 \notin v^{-1}([i])$. Moreover, $us, vs \in W^J$ and so $D_i(us, vs) = D_i(u, v)$ and $a_j(us, vs) = a_j(u, v)$ for all $j \in [n]$, since $(us)^{-1}([i])$ is obtained from $u^{-1}([i])$ by replacing k by $k+1$, and similarly for v . Hence, by Theorem 2.1 and our induction hypothesis,

$$\begin{aligned} E_{u,v}(q) &= E_{us,vs} \\ &= q^{l(vs)-l(us)} \prod_{j \in D_i(us,vs)} (1 - q^{-2a_j(us,vs)+1}) \\ &= q^{l(v)-l(u)} \prod_{j \in D_i(u,v)} (1 - q^{-2a_j(u,v)+1}), \end{aligned}$$

as desired.

b) $s \in D(v)$, and $us \not\leq vs$.

Since $v \in W^J$ this implies that $k \notin v^{-1}([i])$, $k+1 \in v^{-1}([i])$, and $vs \in W^J$. So we have that $D_i(us, v) = D_i(u, v) \setminus \{k+1\}$,

$$(7) \quad a_j(us, v) = a_j(u, v) = a_j(us, vs)$$

for all $j \in [n] \setminus \{k+1\}$, and

$$(8) \quad a_{k+1}(us, vs) = a_{k+1}(u, v) - 2.$$

Since $us \not\leq vs$ by (8) and the note before the statement of this theorem, we conclude that $a_{k+1}(u, v) = 1$. Hence, by Theorem 2.1 and the induction hypothesis,

$$\begin{aligned} E_{u,v}(q) &= (q-1)E_{us,v}(q) \\ &= (q-1)q^{l(v)-l(u)-1} \prod_{j \in D_i(u,v) \setminus \{k+1\}} (1 - q^{-2a_j(us,v)+1}) \\ &= \frac{(q-1)q^{l(v)-l(u)-1}}{(1 - q^{-2a_{k+1}(u,v)+1})} \prod_{j \in D_i(u,v)} (1 - q^{-2a_j(u,v)+1}) \\ &= q^{l(v)-l(u)} \prod_{j \in D_i(u,v)} (1 - q^{-2a_j(u,v)+1}), \end{aligned}$$

as desired.

c) $s \in D(v)$, and $us \leq vs$.

Since $v \in W^J$ this implies that $k \notin v^{-1}([i])$, $k + 1 \in v^{-1}([i])$, and $vs \in W^J$. So we have that

$$D_i(us, v) = D_i(u, v) \setminus \{k + 1\},$$

$$D_i(us, vs) = (D_i(u, v) \setminus \{k + 1\}) \cup \{k\},$$

$a_j(us, v) = a_j(us, vs) = a_j(u, v)$ for all $j \in [n] \setminus \{k + 1\}$, $a_{k+1}(us, vs) = a_{k+1}(u, v) - 2$ and $a_{k+1}(u, v) = a_k(u, v) + 1$. Hence, by Theorem 2.1 and our induction hypothesis,

$$\begin{aligned} E_{u,v}(q) &= (q - q^{-1})E_{us,v}(q) + E_{us,vs}(q) \\ &= (q - q^{-1})q^{l(v)-l(u)-1} \prod_{j \in D_i(u,v) \setminus \{k+1\}} (1 - q^{-2a_j(us,v)+1}) \\ &\quad + q^{l(v)-l(u)-2} \prod_{j \in (D_i(u,v) \setminus \{k+1\}) \cup \{k\}} (1 - q^{-2a_j(us,vs)+1}) \\ &= (q - q^{-1})q^{l(v)-l(u)-1} \prod_{j \in D_i(u,v) \setminus \{k+1\}} (1 - q^{-2a_j(u,v)+1}) \\ &\quad + q^{l(v)-l(u)-2} (1 - q^{-2a_k(u,v)+1}) \prod_{j \in D_i(u,v) \setminus \{k+1\}} (1 - q^{-2a_j(u,v)+1}) \\ &= q^{l(v)-l(u)-2} \frac{(q^2 - q^{-2a_k(u,v)+1})}{(1 - q^{-2a_{k+1}(u,v)+1})} \prod_{j \in D_i(u,v)} (1 - q^{-2a_j(u,v)+1}) \\ &= q^{l(v)-l(u)} \prod_{j \in D_i(u,v)} (1 - q^{-2a_j(u,v)+1}), \end{aligned}$$

and the result again follows.

d) $vs \notin W^J$.

Then $s \notin D(v)$ and we have two cases. In the first one, we have $k, k + 1 \in v^{-1}([i])$, which implies that $D_i(us, v) = (D_i(u, v) \setminus \{k + 1\}) \cup \{k\}$ and $a_j(us, v) = a_j(u, v)$ for $j \in [n] \setminus \{k + 1\}$, $a_{k+1}(u, v) = a_k(us, v)$. Hence, by Theorem 2.1 and the induction hypothesis,

$$\begin{aligned} E_{u,v}(q) &= qE_{us,v}(q) \\ &= q^{l(v)-l(u)} \prod_{j \in (D_i(u,v) \setminus \{k+1\}) \cup \{k\}} (1 - q^{-2a_j(us,v)+1}) \\ &= q^{l(v)-l(u)} \frac{(1 - q^{-2a_k(us,v)+1})}{(1 - q^{-2a_{k+1}(u,v)+1})} \prod_{j \in D_i(u,v)} (1 - q^{-2a_j(u,v)+1}) \\ &= q^{l(v)-l(u)} \prod_{j \in D_i(u,v)} (1 - q^{-2a_j(u,v)+1}). \end{aligned}$$

In the second case we have, $k, k + 1 \notin v^{-1}([i])$, hence $D_i(us, v) = D_i(u, v)$ and $a_j(us, v) = a_j(u, v)$ for $j \in [n] \setminus \{k + 1\}$. Hence, by Theorem 2.1 and the induction

hypothesis,

$$\begin{aligned} E_{u,v}(q) &= qE_{us,v}(q) \\ &= q^{l(v)-l(u)} \prod_{j \in D_i(u,v)} (1 - q^{-2a_j(us,v)+1}) \\ &= q^{l(v)-l(u)} \prod_{j \in D_i(u,v)} (1 - q^{-2a_j(u,v)+1}), \end{aligned}$$

and the result again follows.

This completes the induction step and hence the proof. □

Because W^J is isomorphic, as a poset, to a lower interval in Young’s lattice (by Proposition 3.2), it is natural to rephrase Theorem 3.4 in the language of partitions rather than permutations.

Let $\mu, \lambda \in \mathcal{P}(n, i)$, with $\mu \subseteq \lambda$. Think of μ and λ as paths as explained above. Then, by Proposition 3.2, the path λ lies (weakly) above the path μ . Let $1 \leq j \leq n$ and consider the j -th step of λ (from the left). Following [4], we say that such a step is *allowable* with respect to μ if it is an up-step and the j -th step of μ is a down-step. For example, if $n = 9, i = 4, \lambda = (5, 4, 2, 1)$, and $\mu = (2, 0, 0, 0)$, then the j -th step of λ is allowable with respect to μ exactly if $j \in \{1, 3, 6\}$ (see Figure 4). Now let $\tilde{a}_j(\mu, \lambda)$ be the vertical distance (divided by two, since it is always even) between the (right end of the) j -th step of λ and the (right end of the) j -th step of μ . We then have the following result, and we refer the reader to [4, Proposition 3.3] for its proof.

Proposition 3.5. *Let $u, v \in W^J, u \leq v$. Then*

$$a_j(u, v) = \tilde{a}_{n+1-j}(\Lambda(u), \Lambda(v))$$

for $j = 1, \dots, n$. Furthermore $n + 1 - j \in v^{-1}([i]) \setminus u^{-1}([i])$ if and only if the j -th step of $\Lambda(v)$ is allowable with respect to $\Lambda(u)$.

We can now rephrase Theorem 3.4 in terms of partitions.

Corollary 3.6. *Let $u, v \in W^J, u \leq v$. Then*

$$(9) \quad E_{u,v}(q) = q^{|\lambda \setminus \mu|} \prod_j (1 - q^{-2\tilde{a}_j(\mu, \lambda)+1}),$$

where $\mu = \Lambda(u), \lambda = \Lambda(v)$ and j runs over all the allowable steps of λ with respect to μ . In particular, $E_{u,v}(q)$ depends only on $\Lambda(v) \setminus \Lambda(u)$. □

In the case of a lower interval, formula (9) takes up a particularly simple form. The proof of the next result is analogous to the one of [4, Corollary 3.5], and we leave it to the reader.

Corollary 3.7. *Let $v \in W^J$. Then*

$$E_{e,v}(q) = q^{|\mu|} \prod_{j=1}^{d(\mu)} (1 - q^{-2j+1}),$$

where $\mu = \Lambda(v)$ and $d(\mu)$ is the length of the Durfee square of μ . □




FIGURE 4.




FIGURE 5.

We close this section with an example to illustrate Theorem 3.4 and Corollary 3.6. We use the various calculations made in §2 and §3. Let $u = 123564789$ and $v = 516278394$ in $S_9^{S \setminus \{4,5\}}$; we have $D_4(u, v) = \{4, 7, 9\}$. From (5) and (6), it follows that

$$(10) \quad E_{u,v}(q) = q^{10}(1 - q^{-1})(1 - q^{-3})(1 - q^{-3}).$$

Observe that we have $\Lambda(v) = (5, 4, 2, 1) = \lambda$ and $\Lambda(u) = (2, 0, 0, 0) = \mu$. The diagram of the skew-partition $\Lambda(v) \setminus \Lambda(u)$ is drawn in Figure 4. The allowable are indicated by arrows and $\tilde{a}_1(\mu, \lambda) = 1$, $\tilde{a}_3(\mu, \lambda) = 2$, $\tilde{a}_6(\mu, \lambda) = 2$.

4. THE CASE (C_n, A_{n-1})

A partition $(\lambda_1, \lambda_2, \dots, \lambda_k)$ is *strict* if $\lambda_1 > \lambda_2 > \dots > \lambda_k$. We denote by \mathcal{P}_s the set of all (integer) strict partitions. Let

$$H := \{(i, j) \in \mathbf{P}^2 : i \leq j\}$$

with the ordering induced by the product ordering on \mathbf{P}^2 . We call the finite order ideals of H *shifted partitions*. Denote by \mathcal{I} the set of all finite order ideals of H . Note that \mathcal{I} is partially ordered by set inclusion. It is well known that this makes \mathcal{I} into a distributive lattice. We identify a shifted partition with its diagram

$$\{(i, j) \in \mathbf{P}^2 : 1 \leq i \leq k, i \leq j \leq \lambda_i - 1 + i\},$$

and as in §3 we draw it rotated counterclockwise by $\frac{\pi}{4}$ radians. So for example the diagram of $(7, 6, 5, 4, 2)$ is illustrated in Figure 5.

Let

$$\tilde{\mathcal{P}} := \{\lambda \in \mathcal{P} : \lambda \supseteq l(\lambda)^{l(\lambda)}\}.$$

Note that there are inclusion preserving bijections between strict partitions, shifted partitions and partitions in $\tilde{\mathcal{P}}$, given by

$$\begin{aligned} (\lambda_1, \lambda_2, \dots, \lambda_k) &\leftrightarrow \{(i, j) \in \mathbf{P}^2 : 1 \leq i \leq k, i \leq j \leq \lambda_i - 1 + i\} \\ &\leftrightarrow (\lambda_1, \lambda_2 + 1, \dots, \lambda_k + k - 1). \end{aligned}$$

For this reason, we will freely identify these objects. Note, however, that only the bijection between strict partitions and shifted partitions preserves size. In fact, if $\lambda, \mu, \in \tilde{\mathcal{P}}$ and $\nu \in \mathcal{P}$, with $\mu \subseteq \nu \subseteq \lambda$, then it does not necessarily follow that $\nu \in \tilde{\mathcal{P}}$. Therefore the subposet

$$\{\nu \in \mathcal{P} : \mu \subseteq \nu \subseteq \lambda\}$$

of \mathcal{P} is not isomorphic to the subposet

$$\{\nu \in \mathcal{I} : \mu \subseteq \nu \subseteq \lambda\}$$

of \mathcal{I} .

Our purpose in this section is to obtain an explicit product formula for the E -polynomials in the case of the pair (C_n, A_{n-1}) . Therefore, let S_n^B be the group of all the bijections of the set $[\pm n]$ in itself such that

$$w(-a) = -w(a)$$

for all $a \in [\pm n]$, with composition as group operation. If $w \in S_n^B$, then, following [6], we write $w = [a_1, \dots, a_n]$ to mean that $w(i) = a_i$, for $i = 1, \dots, n$, and call this the *window* notation of w . Because of this notation the group S_n^B is often called the group of all *signed permutations* on $[n]$. Since the elements of S_n^B are permutations of $[\pm n]$ we can also write them in disjoint cycle form; as in §2, we multiply elements “from the right”. We identify S_n as a subgroup of S_n^B in the natural way. For the rest of this section, we fix $n \in \mathbf{P}$ and we let $W := S_n^B$, $s_i := (i, i + 1)(-i - 1, -i)$ for $i = 1, \dots, n - 1$, $s_0 := (1, -1)$, $S := \{s_0, s_1, \dots, s_{n-1}\}$ and $J := S \setminus \{s_0\}$. It is well known that (S_n^B, S) is a Coxeter system of type B_n and that the following characterization holds (see, e.g., [6]).

Proposition 4.1. *Let $v \in S_n^B$. Then*

$$D(v) = \{s_i \in S : v(i) > v(i + 1)\},$$

where $v(0) := 0$.

From Proposition 4.1 we have that

$$W^J = \{v \in W : v^{-1}(1) < v^{-1}(2) < \dots < v^{-1}(n)\}.$$

Therefore, given $v \neq e$, $v \in W^J$, there is a unique $k \in [n]$ (in fact, $k = N_1(v)$) such that

$$(11) \quad v^{-1}(k) < 0 < v^{-1}(k + 1)$$

and we associate to v the shifted partition

$$(12) \quad \Lambda_B(v) := (-v^{-1}(1), -v^{-1}(2), \dots, -v^{-1}(k)).$$

Let

$$\tilde{\mathcal{I}}(n) := \{\lambda \in \mathcal{I} : \lambda \subseteq (n, n - 1, \dots, 2, 1)\}.$$




FIGURE 6.

The following is known:

Proposition 4.2. *The map Λ_B defined by (12) is a bijection between W^J and $\tilde{\mathcal{I}}(n)$. Furthermore $u \leq v$ in W^J if and only if $\Lambda_B(u) \subseteq \Lambda_B(v)$ and $l(v) = |\Lambda_B(v)|$ for all $u, v \in W^J$.*

As before it is convenient to identify a shifted partition $\lambda \in \tilde{\mathcal{I}}(n)$ with a lattice path with $(1, 1)$ and $(1, -1)$ steps starting at $(0, 0)$ and having n steps. We have the obvious bijection between the peaks of λ as a path and the upper peaks of λ as a partition. Note that, as in Proposition 3.2, this bijection depends on n , but for us n is fixed and so there is no confusion. For example, the partition $\lambda = (7, 6, 5, 4, 2)$ corresponds to the path in Figure 6.

Lemma 4.3. *Let $v \in W^J$ and $j \in [n - 1]$. Then $s_j \in D(v)$ if and only if $\Lambda_B(v)$ has a peak at $n - j$. Furthermore, $s_o \in D(v)$ if and only if the last step of $\Lambda_B(v)$ is up.*

This result can be proved in a way similar to Lemma 3.3 (see also Lemma 5.3) and is due to Brenti [5]. Note that i -th step of $\Lambda_B(v)$ is an up-step if and only if

$$(13) \quad i \in \{n + 1 + v^{-1}(1), n + 1 + v^{-1}(2), \dots, n + 1 + v^{-1}(k)\}.$$

Proposition 4.4. *Let $v \in W^J$ and $i \in [n]$. Then the i -th step (from the left) of $\Lambda_B(v)$ (seen as a path) is an up-step if and only if $v(n + 1 - i) < 0$.*

Proof. We know that the i -th step of $v \in W^J$ is an up-step if and only if

$$(14) \quad n + 1 - i \in \{-v^{-1}(1), -v^{-1}(2), \dots, -v^{-1}(k)\}.$$

But this, by the definition of k , happens if and only if $v(n + 1 - i) < 0$, as desired. \square

We are now ready to prove the main theorem of this section, which gives an explicit product formula for the polynomials $E_{u,v}(q)$ in the case of the symmetric pair (C_n, A_{n-1}) . As in the preceding section we give two different formulations of this result, one in terms of signed permutations and one in terms of shifted partitions.

Let $u, v \in W^J$, $u \leq v$. For $j \in [n]$ let, following [5],

$$(15) \quad b_j(u, v) := |\{r \geq j : v(r) < 0\}| - |\{r \geq j : u(r) < 0\}|.$$

For example, if $u = [4, 5, -3, -2, 6, 7, -1]$ and $v = [6, -5, 7, -4, -3, -2, -1]$, then

$$(16) \quad (b_1(u, v), \dots, b_7(u, v)) = (2, 2, 1, 2, 2, 1, 0).$$

Note that it follows from Proposition 4.2 that $b_j(u, v) \geq 0$ for $j = 1, \dots, n$ if and only if $u \leq v$. Also, if $u \leq v$, then $b_j(u, v) > 0$ when $v(j) < 0 < u(j)$. We let

$$(17) \quad N(u, v) := \{r \in [n] : u(r)v(r) < 0\}$$

and

$$(18) \quad D(u, v) := \{r \in N(u, v) : (-1)^{b_r(u, v)} < 0\}.$$

Theorem 4.5. *Let $u, v \in W^J$, $u \leq v$. Then*

$$(19) \quad E_{u, v}(q) = q^{l(v) - l(u)} \prod_{j \in D(u, v)} (1 - q^{-\tilde{b}_j(u, v)}),$$

where

$$(20) \quad \tilde{b}_j(u, v) := \begin{cases} 2b_j(u, v) - 1 & \text{if } u(j) > 0, \\ 2b_j(u, v) + 1 & \text{if } u(j) < 0. \end{cases}$$

Proof. We proceed by induction on $l(w_0^J) - l(u)$. If $l(w_0^J) - l(u) = 0$, we have $w_0^J = v = u$ and the result is trivially true. So suppose that $l(w_0^J) - l(u) > 0$ and let s be such that $s \notin D(u)$ and $us \in W^J$.

Suppose first that $s = (-i-1, -i)(i, i+1)$ for some $i \in [n-1]$; then $u(i) < u(i+1)$. Note that, since $u \in W^J$, this implies that $u(i) < 0 < u(i+1)$. We have four cases to consider.

a) $s \notin D(v)$, and $vs \in W^J$.

Since $v \in W^J$, this implies that $v(i) < 0 < v(i+1)$. Moreover $us, vs \in W^J$, and so $N(us, vs) = N(u, v)$ and $b_j(us, vs) = b_j(u, v)$ for all $j \in [n]$, so that $D(us, vs) = D(u, v)$ and $\tilde{b}_j(us, vs) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{i, i+1\}$. Hence, by Theorem 2.1 and our induction hypothesis,

$$\begin{aligned} E_{u, v}(q) &= E_{us, vs}(q) \\ &= q^{l(vs) - l(us)} \prod_{j \in D(us, vs)} (1 - q^{-\tilde{b}_j(us, vs)}) \\ &= q^{l(v) - l(u)} \prod_{j \in D(u, v)} (1 - q^{-\tilde{b}_j(u, v)}), \end{aligned}$$

as desired.

b) $s \in D(v)$, and $us \not\leq vs$.

Then $v(i) > 0 > v(i+1)$, therefore $N(us, v) = N(u, v) \setminus \{i, i+1\}$, $b_j(us, v) = b_j(u, v) = b_j(us, vs)$ for all $j \in [n] \setminus \{i+1\}$, and

$$(21) \quad b_i(u, v) = b_{i+1}(u, v) - 1.$$

Since $us \not\leq vs$, it follows from the note before the statement of the theorem that $b_{i+1}(u, v) > 0 > b_{i+1}(us, vs)$. Also $b_{i+1}(u, v) - b_{i+1}(us, vs) = 2$, so $b_{i+1}(u, v) = 1$. This and (21) imply that $i \notin D(u, v)$ and $i+1 \in D(u, v)$. It follows that $D(us, v) = D(u, v) \setminus \{i+1\}$, $\tilde{b}_j(us, v) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{i, i+1\}$, and that $\tilde{b}_{i+1}(u, v) = 2b_{i+1}(u, v) - 1 = 1$. Hence, by Theorem 2.1 and the induction

hypothesis,

$$\begin{aligned}
 E_{u,v}(q) &= (q-1)E_{us,v}(q) \\
 &= (q-1)q^{l(v)-l(u)-1} \prod_{j \in D(u,v) \setminus \{i+1\}} (1 - q^{-\tilde{b}_j(us,v)}) \\
 &= \frac{(q-1)q^{l(v)-l(u)-1}}{(1 - q^{-\tilde{b}_{i+1}(u,v)})} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}) \\
 &= q^{l(v)-l(u)} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}).
 \end{aligned}$$

c) $s \in D(v)$, and $us \leq vs$.

Then, as above, $v(i) > 0 > v(i+1)$, $N(us, v) = N(u, v) \setminus \{i, i+1\}$, $b_j(us, v) = b_j(u, v)$ for all $j \in [n] \setminus \{i+1\}$, and

$$(22) \quad b_i(u, v) = b_{i+1}(u, v) - 1.$$

It follows that $D(us, v) = D(u, v) \setminus \{i, i+1\}$ and $\tilde{b}_j(us, v) = \tilde{b}_j(u, v)$, for all $j \in [n] \setminus \{i, i+1\}$. On the other hand, we have $N(us, vs) = N(u, v)$ and $b_j(us, vs) = b_j(u, v)$ for all $j \in [n] \setminus \{i+1\}$, and $b_{i+1}(us, vs) = b_{i+1}(u, v) - 2$. It follows that $D(us, vs) = D(u, v)$ and $\tilde{b}_j(us, vs) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{i, i+1\}$. Hence, by Theorem 2.1 and the induction hypothesis,

$$\begin{aligned}
 E_{u,v}(q) &= (q - q^{-1})E_{us,v}(q) + E_{us,vs}(q) \\
 &= (q - q^{-1})q^{l(v)-l(u)-1} \prod_{j \in D(u,v) \setminus \{i, i+1\}} (1 - q^{-\tilde{b}_j(us,v)}) \\
 (23) \quad &+ q^{l(v)-l(u)-2} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(us,vs)}).
 \end{aligned}$$

From (22) we have that $i \in D(u, v)$ if and only if $i+1 \notin D(u, v)$, so we have two cases. If $i \in D(u, v)$, then $D(us, v) = D(u, v) \setminus \{i\}$ and $\tilde{b}_i(us, vs) = 2b_i(us, vs) - 1 = 2b_i(u, v) - 1 = \tilde{b}_i(u, v) - 2$ because $u(i) < 0 < u(i+1)$. Hence, by (23),

$$\begin{aligned}
 E_{u,v}(q) &= (q^2 - 1)q^{l(v)-l(u)-2} \prod_{j \in D(u,v) \setminus \{i\}} (1 - q^{-\tilde{b}_j(u,v)}) \\
 &+ q^{l(v)-l(u)-2}(1 - q^{-\tilde{b}_i(us,vs)}) \prod_{j \in D(u,v) \setminus \{i\}} (1 - q^{-\tilde{b}_j(u,v)}) \\
 &= q^{l(v)-l(u)-2} \frac{(q^2 - q^{-\tilde{b}_i(us,vs)})}{(1 - q^{-\tilde{b}_i(u,v)})} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}) \\
 &= q^{l(v)-l(u)} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}).
 \end{aligned}$$

If $i+1 \in D(u, v)$, then $D(us, v) = D(u, v) \setminus \{i+1\}$ and $\tilde{b}_{i+1}(us, vs) = 2b_{i+1}(us, vs) + 1 = (2b_{i+1}(u, v) - 1) - 2 = \tilde{b}_{i+1}(u, v) - 2$. Hence, by (23),

$$\begin{aligned} E_{u,v}(q) &= (q^2 - 1)q^{l(v)-l(u)-2} \prod_{j \in D(u,v) \setminus \{i+1\}} (1 - q^{-\tilde{b}_j(u,v)}) \\ &\quad + q^{l(v)-l(u)-2} (1 - q^{-\tilde{b}_{i+1}(us,vs)}) \prod_{j \in D(u,v) \setminus \{i+1\}} (1 - q^{-\tilde{b}_j(u,v)}) \\ &= q^{l(v)-l(u)-2} \frac{(q^2 - q^{-\tilde{b}_{i+1}(us,vs)})}{(1 - q^{-\tilde{b}_{i+1}(u,v)})} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}) \\ &= q^{l(v)-l(u)} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}), \end{aligned}$$

and the result follows also in this case.

d) $vs \notin W^J$.

Then $s \notin D(v)$ and we have two cases. In the first one, we have $v(i) < v(i+1) < 0$ and this implies that $N(us, v) = (N(u, v) \setminus \{i+1\}) \cup \{i\}$, $b_j(us, v) = b_j(u, v)$ for all $j \in [n] \setminus \{i+1\}$, $b_i(us, v) = b_{i+1}(u, v)$. It follows that

$$D(us, v) \setminus \{i\} = D(u, v) \setminus \{i+1\},$$

$i \in D(us, v)$, if and only if $i+1 \in D(u, v)$, $\tilde{b}_j(us, v) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{i, i+1\}$ and $\tilde{b}_i(us, v) = \tilde{b}_{i+1}(u, v)$. Hence, by Theorem 2.1 and the induction hypothesis, if $i \notin D(us, v)$, $D(us, v) = D(u, v)$ and the thesis easily follows. Otherwise if $i \in D(us, v)$, we have

$$\begin{aligned} E_{u,v}(q) &= qE_{us,v}(q) \\ &= qq^{l(v)-l(u)-1} \prod_{j \in D(us,v)} (1 - q^{-\tilde{b}_j(us,v)}) \\ &= q^{l(v)-l(u)} \frac{(1 - q^{-\tilde{b}_i(us,v)})}{(1 - q^{-\tilde{b}_{i+1}(u,v)})} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}) \\ &= q^{l(v)-l(u)} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}), \end{aligned}$$

because $\tilde{b}_i(us, v) = \tilde{b}_{i+1}(u, v)$.

In the second case, we have $0 < v(i) < v(i+1)$ and this implies that $N(us, v) = (N(u, v) \setminus \{i\}) \cup \{i+1\}$, $b_j(us, v) = b_j(u, v)$ for all $j \in [n] \setminus \{i+1\}$, $b_{i+1}(us, v) = b_i(u, v)$. It follows that

$$D(us, v) \setminus \{i+1\} = D(u, v) \setminus \{i\},$$

$i+1 \in D(us, v)$, if and only if $i \in D(u, v)$, $\tilde{b}_j(us, v) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{i, i+1\}$ and $\tilde{b}_{i+1}(us, v) = \tilde{b}_i(u, v)$. Hence, by Theorem 2.1 and the induction hypothesis, if $i+1 \notin D(us, v)$, $D(us, v) = D(u, v)$ and the thesis easily follows. Otherwise if

$i + 1 \in D(us, v)$, we have

$$\begin{aligned} E_{u,v}(q) &= qE_{us,v}(q) \\ &= q^{l(v)-l(u)} \prod_{j \in D(us,v)} (1 - q^{-\tilde{b}_j(us,v)}) \\ &= q^{l(v)-l(u)} \frac{(1 - q^{-\tilde{b}_{i+1}(us,v)})}{(1 - q^{-\tilde{b}_i(u,v)})} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}), \end{aligned}$$

since $\tilde{b}_{i+1}(us, v) = \tilde{b}_i(u, v)$, the result again follows.

Suppose now that $s = (-1, 1) = s_0$. Then $u(1) > 0$, and we observe that $us, vs \in W^J$. We therefore have three cases to consider.

1) $s \notin D(v)$.

Then $vs \in W^J$ and $v(1) > 0$. Hence $N(us, vs) = N(u, v)$, $b_j(us, vs) = b_j(u, v)$ for all $j \in [n]$ and so $D(us, vs) = D(u, v)$, $1 \notin D(u, v)$, $\tilde{b}_j(us, vs) = \tilde{b}_j(u, v)$, for all $j \in [n] \setminus \{1\}$ and the result follows from Theorem 2.1 and the induction hypothesis.

2) $s \in D(v)$, and $us \not\leq vs$.

Then $v(1) < 0$ so $N(us, v) = N(u, v) \setminus \{1\}$, $b_j(us, v) = b_j(u, v)$ for all $j \in [n] \setminus \{1\}$. Therefore $D(us, v) = D(u, v) \setminus \{1\}$ and $\tilde{b}_j(us, v) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{1\}$. Also, $u < v$ and $us \not\leq vs$, so by the remark before the statement of the theorem, $b_1(u, v) = 1$ and hence $\tilde{b}_1(u, v) = 1$. Hence,

$$\begin{aligned} E_{u,v}(q) &= (q - 1)E_{us,v}(q) \\ &= (q - 1)q^{l(v)-l(u)-1} \prod_{j \in D(u,v) \setminus \{1\}} (1 - q^{-\tilde{b}_j(u,v)}) \\ &= q^{l(v)-l(u)} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}). \end{aligned}$$

3) $s \in D(v)$, and $us \leq vs$.

Then $v(1) < 0$, $N(us, vs) = N(u, v)$, $N(us, v) = N(u, v) \setminus \{1\}$, $b_j(us, vs) = b_j(u, v) = b_j(us, v)$ for all $j \in [n] \setminus \{1\}$, and $b_1(us, vs) = b_1(u, v) - 2 = b_1(us, v) - 1$. It follows that

$$D(us, v) = D(u, v) \setminus \{1\} = D(us, vs) \setminus \{1\}$$

and $\tilde{b}_j(us, vs) = \tilde{b}_j(u, v) = \tilde{b}_j(us, v)$ for all $j \in [n] \setminus \{1\}$. Hence,

$$\begin{aligned} E_{u,v}(q) &= (q - q^{-1})E_{us,v}(q) + E_{us,vs}(q) \\ &= (q - q^{-1})q^{l(v)-l(u)-1} \prod_{j \in D(u,v) \setminus \{1\}} (1 - q^{-\tilde{b}_j(us,v)}) \\ (24) \quad &+ q^{l(v)-l(u)-2} \prod_{j \in D(us,vs)} (1 - q^{-\tilde{b}_j(us,vs)}). \end{aligned}$$

Now we have two cases. If $1 \in D(u, v)$, then $1 \in D(us, vs)$ and $\tilde{b}_1(us, vs) = \tilde{b}_1(u, v) - 2$, so from (24),

$$\begin{aligned} E_{u,v}(q) &= (q^2 - 1)q^{l(v)-l(u)-2} \prod_{j \in D(u,v) \setminus \{1\}} (1 - q^{-\tilde{b}_j(u,v)}) \\ &\quad + q^{l(v)-l(u)-2} (1 - q^{-\tilde{b}_1(us,vs)}) \prod_{j \in D(u,v) \setminus \{1\}} (1 - q^{-\tilde{b}_j(u,v)}) \\ &= q^{l(v)-l(u)} \frac{(1 - q^{-\tilde{b}_1(us,vs)-2})}{(1 - q^{-\tilde{b}_1(u,v)})} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}). \end{aligned}$$

If $1 \notin D(u, v)$, then $1 \notin D(us, vs)$ and from (24) we have that

$$\begin{aligned} E_{u,v}(q) &= (q^2 - 1)q^{l(v)-l(u)-2} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}) \\ &\quad + q^{l(v)-l(u)-2} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}) \\ &= q^{l(v)-l(u)} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}) \end{aligned}$$

and the result follows. This completes the induction step and hence the proof. \square

As in the previous section it is natural to rephrase Theorem 4.5 in the language of shifted partitions. Let $\mu, \lambda \in \tilde{\mathcal{I}}(n)$, with $\mu \subseteq \lambda$. We think of μ and λ as paths, as explained at the beginning of this section. Then, by Proposition 4.2, the path λ lies (weakly) above the path μ . Let $j \in [n]$ and consider the j -th step of λ (from the left). Following [5] we say that such a step is *B-allowable* with respect to μ if the j -th step of μ is not parallel to it and $\tilde{a}_j(\mu, \lambda)$ is odd. For example, if $\mu = (7, 4, 3, 0, 0, 0, 0)$ and $\lambda = (7, 6, 5, 4, 2, 0, 0)$, then the j -th step of λ is *B-allowable* with respect to μ exactly if $j \in \{2, 5\}$ (see Figure 7).

Proposition 4.6. *Let $u, v \in W^J$, $u \leq v$. Then*

$$b_j(u, v) = \tilde{a}_{n+1-j}(\Lambda_B(u), \Lambda_B(v))$$

for $i = 1, \dots, n$. Furthermore $n + 1 - j \in D(u, v)$ if and only if the j -th step of $\Lambda_B(v)$ is *B-allowable* with respect to $\Lambda_B(u)$.

This result can be proved in a way similar to Proposition 3.5 and is due Brenti [5].

We can now rephrase Theorem 4.5 in terms of shifted partitions.

Corollary 4.7. *Let $u, v \in W^J$, $u \leq v$. Then*

$$(25) \quad E_{u,v}(q) = q^{|\lambda \setminus \mu|} \prod_j (1 - q^{-\tilde{a}_j(\mu, \lambda)}),$$

where $\mu = \Lambda_B(u)$, $\lambda = \Lambda_B(v)$, j runs over all the *B-allowable* steps of λ with respect to μ , and

$$\tilde{a}_j(\mu, \lambda) := \begin{cases} 2\tilde{a}_j(\mu, \lambda) - 1 & \text{if the } j\text{-th step of } \mu \text{ is down,} \\ 2\tilde{a}_j(\mu, \lambda) + 1 & \text{if the } j\text{-th step of } \mu \text{ is up.} \end{cases}$$

In particular, $E_{u,v}(q)$ depends only on $\Lambda_B(v) \setminus \Lambda_B(u)$. \square




FIGURE 7.

In the case of a lower interval, the formulas (19) and (25) take up a particularly simple form.

Corollary 4.8. *Let $v \in W^J$. Then*

$$E_{e,v}(q) = q^{l(v)} \prod_{j=1}^{\lfloor \frac{N_1(v)+1}{2} \rfloor} (1 - q^{-4j+3}) = q^{|\lambda|} \prod_{j=1}^{\lfloor \frac{l(\lambda)+1}{2} \rfloor} (1 - q^{-4j+3}),$$

where $\lambda = \Lambda_B(v)$.

Proof. We know from Theorem 4.5 that

$$(26) \quad E_{e,v}(q) = q^{l(v)} \prod_{j \in D(e,v)} (1 - q^{-\tilde{b}_j(e,v)}).$$

Clearly, $N(e,v) = \{r \in [n] : v(r) < 0\}$, $\tilde{b}_j(e,v) = 2b_j(e,v) - 1$ for all $j \in [n]$. However, $b_j(e,v) = |\{r \geq j : v(r) < 0\}|$ for $j \in [n]$, and $D(e,v) = \{r \in N(e,v) : b_r(e,v) \text{ is odd}\}$. Hence,

$$E_{e,v}(q) = q^{l(v)} \prod_{j=1}^{\lfloor \frac{N_1(v)+1}{2} \rfloor} (1 - q^{-4j+3}),$$

as desired. From the definition we have that $l(\Lambda_B(v)) = N_1(v)$, so the second equation follows. \square

We close this section with an example. Let $n = 7$, $u = [4, 5, -3, -2, 6, 7, -1]$, and $v = [6, -5, 7, -4, -3, -2, -1] \in (S_7^B)^J$. We have $N(u,v) = \{2, 3, 5, 6\}$ and $D(u,v) = \{3, 6\}$. Since $u(3) < 0$ and $u(6) > 0$, from (20) and (19) it follows that

$$E_{u,v}(q) = q^{10}(1 - q^{-3})(1 - q^{-1}).$$

Observe that $\Lambda_B(u) = (7, 4, 3, 0, 0, 0, 0) = \mu$ and $\Lambda_B(v) = (7, 6, 5, 4, 2, 0, 0) = \lambda$. The paths corresponding to $\Lambda_B(v)$ and $\Lambda_B(u)$ are drawn in Figure 7, and the B -allowable steps are indicated by arrows. We have that $\bar{a}_2(\mu, \lambda) = 1$ and $\bar{a}_5(\mu, \lambda) = 3$.

5. THE CASE (D_n, A_{n-1})

In this section we study the E -polynomials in the case of the pair (D_n, A_{n-1}) . Hence we consider S_n^D , the subgroup of S_n^B consisting of all the signed permutations having an even number of negative entries in their window notation. More precisely,

$$S_n^D := \{w \in S_n^B : N_1(w(1), \dots, w(n)) \equiv 0 \pmod{2}\}.$$

For the rest of this section, we fix $n \in \mathbf{P}$ and we let $W := S_n^D$, $s_i := (i, i + 1) (-i - 1, -i)$ for $i = 1, \dots, n - 1$, $s_0 := (1, -2)(-1, 2)$, $S := \{s_0, s_1, \dots, s_{n-1}\}$ and



FIGURE 8.

$J := S \setminus \{s_0\}$. It is well known that (S_n^D, S) is a Coxeter system of type D_n and that the following characterization holds (see, e.g., [6]).

Proposition 5.1. *Let $v \in S_n^D$. Then*

$$D(v) = \{s_i \in S : v(i) > v(i + 1)\},$$

where $v(0) := -v(2)$ and $v(n + 1) := 0$.

As in §4 we have that $W^J = \{v \in W : v^{-1}(1) < v^{-1}(2) < \dots < v^{-1}(n)\}$ and for every $v \in W^J$ $v \neq e$, there is a unique $k \in [n]$ such that

$$(27) \quad v^{-1}(k) < 0 < v^{-1}(k + 1)$$

and we associate to v the shifted partition

$$(28) \quad \Lambda_D(v) := (-v^{-1}(1) - 1, -v^{-1}(2) - 1, \dots, -v^{-1}(k) - 1).$$

It is not so hard to see that

Proposition 5.2. *The map Λ_D defined by (28) is a bijection between W^J and $\tilde{\mathcal{I}}(n - 1)$. Furthermore $u \leq v$ in W^J if and only if $\Lambda_D(u) \subseteq \Lambda_D(v)$ and $l(v) = |\Lambda_D(v)|$ for all $u, v \in W^J$. \square*

Let \mathcal{A} be the subset of $\tilde{\mathcal{I}}(n)$ consisting of all the shifted partitions with an even number of entries different to zero. More precisely,

$$\mathcal{A} := \{\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k) \in \tilde{\mathcal{I}}(n) : k \text{ is even}\}.$$

We call this set the *even shifted partitions*. Observe that $\Lambda_B(W^J) = \mathcal{A}$. Since, for $u, v \in W^J$, $u \leq v$ in $(S_n^B)^J$ if and only if $u \leq v$ in $(S_n^D)^J$, we have an inclusion preserving bijection between the shifted partition in $\tilde{\mathcal{I}}(n - 1)$ and even shifted partitions in $\tilde{\mathcal{I}}(n)$. Thanks to this bijection we can identify $\lambda \in \tilde{\mathcal{I}}(n - 1)$ with the lattice path associated to $\lambda \in \tilde{\mathcal{I}}(n)$ as explained in §4 after Proposition 4.2. We observe that this lattice path starts at $(0, 0)$, ends after n steps, and has an even number of up-steps. For example, let $v = [-4, 5, -3, 6, -2, -1] \in (S_6^D)^J$. Then $\Lambda_D(v) = (5, 4, 2, 0)$, $\Lambda_B(v) = (6, 5, 3, 1)$ and the lattice path associate to v is drawn in Figure 8.

Lemma 5.3. *Let $v \in W^J$ and $j \in [n - 1]$. Then $s_j \in D(v)$ if and only if $\Lambda_D(v)$ has a peak at $n - j$. Furthermore, $s_o \in D(v)$ if and only if the last two steps of $\Lambda_D(v)$ are up.*

Proof. Let k be defined by (27) and $j \in [n - 1]$; we have that $s_j \in D(v)$ if and only if $v(j) > v(j + 1)$. Since $v \in W^J$, this happens if and only if $v(j) > 0 > v(j + 1)$. Equivalently, this happens if and only if $j \in v^{-1}([n])$ and $j + 1 \notin v^{-1}([n])$. However, $\Lambda_D(v)$ (as a path) has a peak at $n - j$ if and only if its $(n - j)$ -th step is up and its $(n - j + 1)$ -th step is down. But the i -th step of $\Lambda_D(v)$ is an up-step if and only if

$$(29) \quad i \in \{n + 1 + v^{-1}(1), n + 1 + v^{-1}(2), \dots, n + 1 + v^{-1}(k)\}.$$

Therefore $\Lambda_D(v)$ has a peak at $n - j$ if and only if

$$(30) \quad -j = v^{-1}(i) + 1$$

for some $i \in [k]$, and

$$(31) \quad -j \neq v^{-1}(i)$$

for all $i \in [k]$. Equivalently, if and only if $j \notin v^{-1}([-k, -1])$ and $j+1 \in v^{-1}([-k, -1])$, but by the definition of k , $i \notin v^{-1}([-k, -1])$ if and only if $i \notin v^{-1}([-n, -1])$, which is if and only if $i \in v^{-1}([n])$, for all $i \in [n]$. The result follows.

Now let $s_o \in D(v)$. This happens if and only if $v(1) + v(2) < 0$ and implies that $v(1) < v(2) < 0$. It follows that $v^{-1}(k) = -1$ and $v^{-1}(k - 1) = -2$, and that the n -th and the $n - 1$ -th steps of $\Lambda_D(v)$ are up. \square

The following is exactly the analogue of Proposition 4.4 for permutation in $(S_n^D)^J$.

Proposition 5.4. *Let $v \in W^J$ and $i \in [n]$. Then the i -th step (from the left) of $\Lambda_D(v)$ (seen as a path) is an up-step if and only if $v(n + 1 - i) < 0$. \square*

Now we are ready to prove the analogue of Theorem 4.5 for the permutations in $(S_n^D)^J$. The formula is exactly the same, but observe that the polynomials are not always the same; in fact the function length is different.

Theorem 5.5. *Let $u, v \in W^J$, $u \leq v$. Then*

$$(32) \quad E_{u,v}(q) = q^{l(v)-l(u)} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}),$$

where $\tilde{b}_j(u, v)$ is defined as in Theorem 4.5.

Proof. We proceed by induction on $l(w_0^J) - l(u)$. If $l(w_0^J) - l(u) = 0$, we have $w_0^J = v = u$ and the result is trivially true. So suppose that $l(w_0^J) - l(u) > 0$ and let s be such that $s \notin D(u)$ and $us \in W^J$. If $s = (-i - 1, -i)(i, i + 1)$ for some $i \in [n - 1]$, then the proof is exactly the same as for Theorem 4.5. So suppose that $s = (-1, 2)(1, -2) = s_0$; then we have $u(1) + u(2) > 0$. We observe that $us \in W^J$ implies that $0 < u(1) < u(2)$. Moreover we have that $b_1(u, v)$ is even for every $u, v \in W^J$, and so $1 \notin D(u, v)$. We have four cases to consider.

a) $s \notin D(v)$ and $vs \in W^J$.

We have $0 < v(1) < v(2)$. Then $N(us, vs) = N(u, v)$ and $b_j(us, vs) = b_j(u, v)$ for all $j \in [n]$ and so $D(us, vs) = D(u, v)$ and $\tilde{b}_j(us, vs) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{1, 2\}$. Hence, by Theorem 2.1 and our induction hypothesis,

$$\begin{aligned} E_{u,v}(q) &= E_{us,vs}(q) \\ &= q^{l(v)-l(u)} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}). \end{aligned}$$

b) $s \in D(v)$ and $us \not\leq vs$.

Hence $v(1) + v(2) < 0$ and $v(1) < v(2) < 0$. We have that $N(us, v) = N(u, v) \setminus \{1, 2\}$, $b_j(us, v) = b_j(u, v)$ for all $j \in [n] \setminus \{1, 2\}$. Using reasoning as in Theorem 4.5b), $b_1(u, v) > b_2(u, v) \geq 0$, so $b_1(u, v) \geq 2$; $b_2(us, vs) > b_1(us, vs)$ so $b_1(us, vs) \leq -2$, and $b_1(u, v) - b_1(us, vs) = 4$ so $b_1(u, v) = 2$ and $b_2(u, v) = 1$. So $D(us, v) =$

$D(u, v) \setminus \{2\}$ and $\tilde{b}_j(us, v) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{1, 2\}$ and $\tilde{b}_2(u, v) = 1$. Hence, by Theorem 2.1 and the induction hypothesis,

$$\begin{aligned} E_{u,v}(q) &= (q - 1)E_{us,v}(q) \\ &= (q - 1)q^{l(v)-l(u)-1} \prod_{j \in D(u,v) \setminus \{2\}} (1 - q^{-\tilde{b}_j(u,v)}) \\ &= q^{l(v)-l(u)} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}). \end{aligned}$$

c) $s \in D(v)$ and $us \leq vs$.

We have $v(1) + v(2) < 0$ and $v(1) < v(2) < 0$. So $N(us, vs) = N(u, v)$, $N(us, v) = N(u, v) \setminus \{1, 2\}$, $b_j(us, vs) = b_j(u, v) = b_j(us, v)$ for all $j \in [n] \setminus \{1, 2\}$, and $b_2(us, vs) = b_2(u, v) - 2$. It follows that

$$D(us, v) = D(u, v) \setminus \{2\}, \quad D(us, vs) = D(u, v),$$

$\tilde{b}_j(us, vs) = \tilde{b}_j(u, v) = \tilde{b}_j(us, v)$ for all $j \in [n] \setminus \{1, 2\}$, and $\tilde{b}_2(us, vs) = \tilde{b}_2(u, v) - 2$. Hence, by Theorem 2.1 and our induction hypothesis,

$$\begin{aligned} E_{u,v}(q) &= (q - q^{-1})E_{us,v}(q) + E_{us,vs}(q) \\ &= (q - q^{-1})q^{l(v)-l(u)-1} \prod_{j \in D(u,v) \setminus \{2\}} (1 - q^{-\tilde{b}_j(u,v)}) \\ &\quad + q^{l(v)-l(u)-2}(1 - q^{-\tilde{b}_2(us,vs)}) \prod_{j \in D(u,v) \setminus \{2\}} (1 - q^{-\tilde{b}_j(u,v)}) \\ &= q^{l(v)-l(u)} \frac{(1 - q^{-\tilde{b}_2(us,vs)-2})}{(1 - q^{-\tilde{b}_2(u,v)})} \prod_{j \in D(u,v)} (1 - q^{-\tilde{b}_j(u,v)}), \end{aligned}$$

and the result follows.

d) $vs \notin W^J$.

We have two cases. Suppose that $v(1) > 0$ and $v(2) < 0$. We have $N(us, v) = (N(u, v) \cup \{1\}) \setminus \{2\}$, $b_j(us, v) = b_j(us, v)$ for all $j \in [n] \setminus \{1, 2\}$, and

$$(33) \quad b_1(u, v) = b_2(u, v),$$

$$(34) \quad b_1(us, v) = b_1(u, v) - 2.$$

Since $b_1(u, v)$ is even, (33) implies that $2 \notin D(u, v)$, and (34) implies that $1 \notin D(us, v)$. It follows that $D(u, v) = D(us, v)$ and $\tilde{b}_j(us, v) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{1, 2\}$, so the thesis follows immediately by induction.

Suppose now that $v(1) < 0$ and $v(2) > 0$. Then $N(us, v) = (N(u, v) \cup \{2\}) \setminus \{1\}$, $b_j(us, v) = b_j(us, v)$ for all $j \in [n] \setminus \{1, 2\}$, and

$$(35) \quad b_1(u, v) = b_2(us, v) + 2.$$

Since $b_1(u, v)$ is even, (35) implies that $2 \notin D(us, v)$. It follows that $D(u, v) = D(us, v)$ and $\tilde{b}_j(us, v) = \tilde{b}_j(u, v)$ for all $j \in [n] \setminus \{1, 2\}$, so the thesis follows immediately by induction. This completes the proof. \square

As in the previous cases it is natural to rephrase Theorem 5.5 in the language of shifted partitions. Let $\lambda, \mu \in \mathcal{A}$, with $\lambda \subseteq \mu$. We think of μ and λ as paths as explained in §4. So using Proposition 4.6 we have that

Corollary 5.6. *Let $u, v \in W^J$, $u \leq v$. Then*

$$(36) \quad E_{u,v}(q) = q^{|\lambda \setminus \mu|} \prod_j (1 - q^{-\bar{a}_j(\mu, \lambda)})$$

where $\mu = \Lambda_D(u)$, $\lambda = \Lambda_D(v)$, j runs over all the B -allowable steps of λ with respect to μ , and $\bar{a}_j(u, v)$ is defined as in Corollary 4.7. In particular, $E_{u,v}(q)$ depends only on $\Lambda_D(v) \setminus \Lambda_D(u)$. □

For lower intervals we obtain the following

Corollary 5.7. *Let $v \in W^J$. Then*

$$E_{e,v}(q) = q^{l(v)} \prod_{j=1}^{\frac{N_1(v)}{2}} (1 - q^{-4j+3}) = q^{|\lambda|} \prod_{j=1}^{\frac{l(\lambda)}{2}} (1 - q^{-4j+3}),$$

where $\lambda = \Lambda_D(v)$.

Proof. The results follows immediately from Corollary 4.8 , observing that $N_1(v)$ is even for every $v \in W^J$. □

6. THE CASES (B_n, B_{n-1}) AND (D_n, D_{n-1})

In this section we analyze the E -polynomials in the cases (B_n, B_{n-1}) and (D_n, D_{n-1}) . We start with the first one. We use the same notations as in §4, hence $W = S_n^B$, $S = \{s_0, s_1, \dots, s_{n-1}\}$, and $s_o = (-1, 1)$, but now we let $J := S \setminus \{s_{n-1}\}$, so as to have $W_J = B_{n-1}$. It follows that the quotient W^J is a totally ordered set, more precisely, is the chain

$$W^J = \{e, s_{n-1}, \dots, s_{n-1}s_{n-2} \dots s_1s_0, s_{n-1} \dots s_1s_0s_1, \dots, s_{n-1}s_{n-2} \dots s_1s_0s_1 \dots s_{n-1}\}.$$

Proposition 6.1. *Let (W, S) be a Coxeter system, $J \subset S$ and $u, v \in W^J$ such that $u \leq v$. If $[u, v]^J$ is a chain, then*

$$E_{u,v}(q) = q^{l(v)-l(u)}(1 - q^{-1}).$$

Proof. We proceed by induction on $l(w_0^J) - l(u)$. If $l(w_0^J) - l(u) = 0$, the result is trivially true. So suppose that $l(w_0^J) - l(u) > 0$ and let $s \notin D(u)$ and $us \in W^J$. We have four cases to consider.

- a) $s \notin D(v)$ and $vs \in W^J$.

Then the result follows immediately by induction.

- b) $s \in D(v)$ and $us \not\leq vs$.

We know that $[u, v]^J$ is a chain, so $us \not\leq vs$ implies that $us = v$. It follows that

$$E_{u,v}(q) = (q - 1)E_{us,v}(q) = q(1 - q^{-1}).$$

- c) $s \in D(v)$ and $us \leq vs$.

We have

$$\begin{aligned} E_{u,v}(q) &= (q - q^{-1})q^{l(v)-l(u)-1}(1 - q^{-1}) + q^{l(v)-l(u)-2}(1 - q^{-1}) \\ &= q^{l(v)-l(u)-2}(1 - q^{-1})q^2 = q^{l(v)-l(u)}(1 - q^{-1}). \end{aligned}$$




FIGURE 9.

d) $vs \notin W^J$.

Then the result follows immediately by induction. □

By the comment preceding the proposition this settles the case (B_n, B_{n-1}) .

Let us examine the case (D_n, D_{n-1}) . Using the notations of §5 we let $W = D_n$, $S = \{s_0, \dots, s_{n-1}\}$, and $s_0 = (-1, 2)(1, -2)$. Now we let $J := S \setminus \{s_{n-1}\}$. The quotient can be written in the form

$$W^J = \{w \in W : w^{-1}(-2) < w^{-1}(1) < \dots < w^{-1}(n-1)\},$$

and its Bruhat order is drawn in Figure 9.

Moreover we know the unique reduced expression of each $w \in W^J$, in fact the $n-1$ elements in the chain on the bottom are $e, s_{n-1}, \dots, s_{n-1} \cdots s_2$, the $n-1$ elements in the chain on the top are $s_{n-1} \cdots s_2 s_0 s_1, \dots, s_{n-1} \cdots s_2 s_0 s_1 s_2 \cdots s_{n-1}$, and the remaining two elements are exactly $s_{n-1} \cdots s_2 s_0$ and $s_{n-1} \cdots s_2 s_1$. So every $i \in [2, n-1]$ identifies two elements in the quotient, $u_i := s_{n-1} \cdots s_i$ in the chain on the bottom, and $v_i := s_{n-1} \cdots s_2 s_0 s_1 \cdots s_i$ in the chain on the top. Moreover, we define $u_n := e$.

Note that for all $i \in [2, n]$ and for all $j \in [1, n-1]$ we have $l_D(u_i) = n-i$ and $l_D(v_j) = n+j-1$.

We call (u, v) a *equidistant pair* if there exists $i \in [2, n]$ such that $u = u_i$ and $v = v_{i-1}$.

The verification of the following observation is left to the reader.

Lemma 6.2. *Let $u, v \in W^J$. If (u, v) is an equidistant pair, then $u^{-1}(j) = v^{-1}(j)$ for all $j \in [2, n-1]$. □*

Proposition 6.3. *Let $u, v \in W^J$. Then*

$$E_{u,v}(q) = \begin{cases} (q^{l(v)-l(u)} - q)(1 - q^{-1}) & \text{if } (u, v) \text{ is equidistant,} \\ q^{l(v)-l(u)}(1 - q^{-1}) & \text{otherwise.} \end{cases}$$

Proof. We have three cases to consider.

1) $(u, v) = (u_i, v_j)$, with $i \in [2, n]$ and $j \in [1, n-1] \setminus \{i-1\}$.

If $j \geq i$, then $u_i = s_{n-1} \cdots s_i$ and $v_j = s_{n-1} \cdots s_2 s_0 s_1 \cdots s_j$. Hence $s_{i-1} \notin D(u_i)$ and $u_i s_{i-1} = u_{i-1} \in W^J$, while $v_j s_{i-1} \notin W^J$, so we have $E_{u_i, v_j}(q) = q E_{u_{i-1}, v_j}(q)$. We have $i-2$ steps like this to do, and one more step for $s = s_0$, so at the end we

have

$$(37) \quad E_{u_i, v_j}(q) = q^{i-2} E_{u_2, v_j}(q) = q^{i-1} E_{u_2 s_0, v_j}(q).$$

Now $[u_2 s_0, v_j]^J$ is a chain with j steps and so by Proposition 6.1 it follows that $E_{u_2 s_0, v_j}(q) = q^{j-1}(q-1)$ and so from (37) we are done.

If $j < i - 1$, after $(i - j - 2)$ steps in the diagram we have

$$(38) \quad E_{u_i, v_j}(q) = q^{i-j-2} E_{u_{j+2}, v_j}(q).$$

The next step is for $s = s_{j+1}$; $s_{j+1} \notin D(u_{j+2})$ and $s_{j+1} \notin D(v_j)$, so by the first part of the proof we have

$$E_{u_{j+2}, v_j}(q) = E_{u_{j+1}, v_{j+1}}(q) = q^{2j+1}(1 - q^{-1}),$$

and the result follows from (38).

2) (u, v) is an equidistant pair.

We proceed by induction on i . If $i = 2$, then $u_2 = s_{n-1} \cdots s_2$ and $v_1 = s_{n-1} \cdots s_2 s_0 s_1$. Hence $s_0 \notin D(u_2)$, $s_0 \in D(v_1)$ and $u_2 s_0 \not\leq v_1 s_0$, so $E_{u_2, v_1}(q) = (q-1)E_{u_2 s_0, v_1}(q)$. In the second step, $s := s_1$, we have $u_2 s_0 s_1 \not\leq v_1 s_1$, so $E_{u_2, v_1}(q) = (q-1)^2 E_{u_2 s_0 s_1, v_1}(q) = (q-1)^2$, since $v_1 = u_2 s_0 s_1$.

So suppose $i > 2$. Then $s_{i-1} \notin D(u_i)$, $u_i s_{i-1} = u_{i-1} \in W^J$, and $v_{i-1} s_{i-1} = v_{i-2} \in W^J$, so since $u_{i-1} \leq v_{i-2}$, we have

$$E_{u_i, v_{i-1}}(q) = (q - q^{-1})E_{u_{i-1}, v_{i-1}}(q) + E_{u_{i-1}, v_{i-2}}(q).$$

By case 1), $E_{u_{i-1}, v_{i-1}}(q) = q^{l(v_{i-1})-l(u_{i-1})}(1 - q^{-1})$ and by induction, $E_{u_{i-1}, v_{i-2}}(q) = q^{l(v_{i-2})-l(u_{i-1})}(1 - q^{-1})$. The result follows.

3) $[u, v]^J$ is a chain.

The result follows by Proposition 6.1. This completes the proof. □

7. CONSEQUENCES AND FURTHER REMARKS

In this section we derive some consequences of our results. We start by proving that the E -polynomials are combinatorial invariants, i.e. that they depend only on the poset $[u, v]^J$. To do this we need a purely order theoretic result on skew partitions that was first proved in [4, Lemma 5.5].

Lemma 7.1. *Let ρ, ν be two connected skew partitions that are isomorphic as posets. Then either $\rho \approx \nu$ or $\rho \approx \nu'$.*

We can now prove the main result of this section.

Corollary 7.2. *Let $J \subset S$, as in §3, §4, and §5, and $u, v \in W^J$, $x, y \in W^J$ be such that $[u, v]^J \cong [x, y]^J$. Then*

$$E_{u, v}(q) = E_{x, y}(q).$$

Proof. Now we prove this result in the case when $W = S_n$. By Proposition 3.2 we have that $[u, v]^J$ is isomorphic, as a poset, to the interval $[\Lambda(u), \Lambda(v)]$ in Young's lattice. But it follows immediately from the definitions and well-known results in the theory of partially ordered sets (see, e.g., [18, §3.4]) that the subposet of join-irreducibles of $[\Lambda(u), \Lambda(v)]$ is isomorphic to $\Lambda(v) \setminus \Lambda(u)$, where the skew partition $\Lambda(v) \setminus \Lambda(u)$ is seen as a poset. Therefore, since $[u, v]^J \cong [x, y]^J$, we conclude that $\Lambda(v) \setminus \Lambda(u) \cong \Lambda(y) \setminus \Lambda(x)$ (as poset), and the result follows from Lemma 7.1 and Corollary 3.6.

Similarly, we can prove the result for the other cases, but we need to replace Proposition 3.2, Corollary 3.6 and Λ for (C_n, A_{n-1}) with Proposition 4.2, Corollary 4.7 and Λ_B , and for (D_n, A_{n-1}) with Proposition 5.2, Corollary 5.6 and Λ_D , respectively. \square

Note that in the case when $W = S_n$ the proof of Corollary 7.2 applies whenever $[u, v]^K \cong [x, y]^H$ with $K, H \subset S$, $|K| = |H| = |S| - 1$.

We conclude this section by discussing the connections mentioned at the end of the Introduction. In [14], Gabber and Joseph define for every $u, v \in W$ a polynomial

$$R'_{u,v}(q) = \sum_{k \geq 0} (-1)^{l(v)-l(u)-k} q^k \dim(\text{Ext}^k(M_u, M_v))$$

and they conjectured (although this is not explicitly stated) that

$$R'_{u,v}(q) = R_{u,v}(q).$$

This conjecture is not true (see [2]), but the R' and the R -polynomials are not so different. In fact, Carlin shows that the R' -polynomials are monic of degree $l(v) - l(u)$ (see [8, Theorem 3.8]), as are the R -polynomials, and proves that the conjecture is true in two cases: when $l(v) - l(u) \leq 3$ (see [8, Proposition 3.13]) and when (u, v) is a Coxeter pair ([8, Proposition 3.11]). The E -polynomials play the same role as the R' -polynomials in the generalized case, so it is natural to wonder about the analogous question, i.e. if $E_{u,v}(q) = R^J_{u,v}(q)$, where $R^J_{u,v}(q)$ are the parabolic R -polynomials (see e.g., [12]). This question also has a negative answer. In fact, for example, let $v = [3, 4, 1, 2, 5] \in S_5^{\setminus \{(2,3)\}}$. Then we have that $R^J_{e,v}(q) = q^4(1 - q^{-1})(1 - q^{-2})$, while $E_{e,v}(q) = q^4(1 - q^{-1})(1 - q^{-3})$.

However, we can prove the analogue of the results of Carlin, for generalized Verma modules. The first two results are very simple, and their proofs are immediate from Theorems 3.4, 4.5 and 5.5.

Corollary 7.3. *Let $u, v \in W^J$, $u \leq v$. Then $E_{u,v}(q)$ is a monic polynomial of degree $l(v) - l(u)$.* \square

Corollary 7.4. *Let $u, v \in W^J$. If $u \leq v$, then*

$$\dim(\text{Ext}^{l(v)-l(u)}(N_u, N_v)) = 1.$$

\square

In [4] Brenti finds explicit formulas for the maximal parabolic R -polynomials of the symmetric group and, in [5] for the group of signed permutations, when $J = S \setminus \{s_0\}$. He proves the following

Theorem 7.5. *Let $u, v \in S_n^J$, $u \leq v$. Then*

$$R^J_{u,v}(q) = q^{|\Lambda(v) \setminus \Lambda(u)|} \prod_j (1 - q^{-\tilde{a}_j(\Lambda(u), \Lambda(v))})$$

where j runs over the allowable steps of $\Lambda(v)$ with respect to $\Lambda(u)$.

Theorem 7.6. *Let $u, v \in (S_n^B)^J$, $u \leq v$. Then*

$$R^J_{u,v}(q) = q^{|\Lambda_B(v) \setminus \Lambda_B(u)|} \prod_j (1 - q^{-\tilde{b}_j(\Lambda_B(u), \Lambda_B(v))})$$

where j runs over the B -allowable steps of $\Lambda_B(v)$ with respect to $\Lambda_B(u)$ and

$$\bar{b}_j(\Lambda_B(u), \Lambda_B(v)) := \begin{cases} \tilde{a}_j(\Lambda_B(u), \Lambda_B(v)) & \text{if the } j\text{-th step of } \Lambda_B(u) \text{ is down,} \\ \tilde{a}_j(\Lambda_B(u), \Lambda_B(v)) + 1 & \text{if the } j\text{-th step of } \Lambda_B(u) \text{ is up.} \end{cases}$$

The author will prove in another paper that the formula in Theorem 7.6 can be extended to the case $W = S_n^D$, but we need to replace Λ_B by Λ_D .

Our formulas are very similar to Brenti's, and it is easy to see that the following results hold. The statements and proofs are given for $W = S_n$, but the results are also true for S_n^B and S_n^D . We simply need to replace Λ by Λ_B and Λ_D , respectively.

We say that a skew partition is a *border strip* (also called a *ribbon*) if it contains no 2×2 square of cells.

Proposition 7.7. *Let $u, v \in W^J$ be such that the skew partition $\Lambda(v) \setminus \Lambda(u)$ is a border strip. Then*

$$E_{u,v}(q) = R_{u,v}^J(q).$$

□

So, in particular, we obtain the analogue of Proposition 3.13 of [8].

Corollary 7.8. *Let $u, v \in W^J$. If $l(v) - l(u) \leq 3$, then*

$$\dim(\text{Ext}^k(N_u, N_v)) = r_k^J(u, v),$$

where $r_k^J(u, v)$ is the absolute value of the coefficient of q^k in $R_{u,v}^J(q)$. □

Let $u, v \in W^J$; we call (u, v) a *generalized Coxeter pair* if $r_1^J(u, v) = l(v) - l(u)$. The next result is the analogue of Proposition 3.11 of [8].

Proposition 7.9. *Let $u, v \in W^J$. If (u, v) is a generalized Coxeter pair, then*

$$\dim(\text{Ext}^k(N_u, N_v)) = r_k^J(u, v) = \binom{n}{k},$$

for $k = 0, \dots, n$, where $n = l(v) - l(u)$.

Proof. If (u, v) is a generalized Coxeter pair, then the only possibility is that $R_{u,v}^J(q) = q^n(1 - q^{-1})^n$. This means that the skew partition $\Lambda(v) \setminus \Lambda(u)$ is a border strip, and so the result follows by Proposition 7.7. □

It seems that the situation is exactly analogous, hence we are led to think that, as for the generalized Verma modules, there should also exist a recursion formula for the ordinary ones. So, we close this paper with this proposal.

Problem. Find a recursion formula for the R' -polynomials.

ACKNOWLEDGEMENTS

I am very grateful to my advisor, Francesco Brenti, for suggesting the problem to me and especially for his generous support. I would like to thank David Vogan and George Lusztig for some useful conversations, and M.I.T. for its hospitality during the preparation of this work.

REFERENCES

- [1] I.N. Bernstein, I.M. Gelfand, S.I. Gelfand, *On a category of g -modules*, *Funct. Anal. Appl.*, **10** (1976), 1-8. MR **53**:10880
- [2] B. Boe, *A Counterexample to the Gabber-Joseph Conjecture*, *Kazhdan-Lusztig Theory and related topics*, *Contemp. Math.*, **139**, 1-2, Amer. Math. Soc., Providence, RI, 1992. MR **94a**:22032
- [3] B. Boe, T.J. Enright, B. Shelton, *Determination of the intertwining operators for holomorphically induced representations of Hermitian symmetric pairs*, *Pacific J. Math.*, **131** (1988), 39-50. MR **88m**:22038
- [4] F. Brenti, *Kazhdan-Lusztig and R -polynomials, Young's lattice, and Dyck partitions*, *Pacific J. Math.*, to appear.
- [5] F. Brenti, personal communication, December 2000.
- [6] F. Brenti, *q -Eulerian Polynomials Arising from Coxeter Groups*, *Europ. J. Combinatorics*, **15** (1994), 417-441. MR **95i**:05013
- [7] J. Brylinski, M. Kashiwara, *Kazhdan-Lusztig conjecture and holonomic systems*, *Invent. Math.*, **64** (1981), 387-410. MR **83e**:22020
- [8] K.J. Carlin, *Extensions of Verma Modules*, *Trans. Amer. Math. Soc.*, **294** (1986), 29-43. MR **87j**:17006
- [9] L. Casian, D. Collingwood, *The Kazhdan-Lusztig conjecture for generalized Verma modules*, *Math. Z.*, **195** (1987), 581-600. MR **88i**:17008
- [10] D.H. Collingwood, *The n -Homology of Harish-Chandra Modules: Generalizing a theorem of Kostant*, *Math. Ann.*, **272** (1985), 161-187. MR **87a**:22027
- [11] D.H. Collingwood, B. Shelton, *A duality theorem for extensions on induced Highest Weight modules*, *Pacific J. Math.*, **146** (1990), 227-237. MR **91m**:22029
- [12] V. Deodhar, *On some geometric aspects of Bruhat orderings. II. The parabolic analogue of Kazhdan-Lusztig polynomials*, *J. Algebra*, **111** (1987), 483-506. MR **89a**:20054
- [13] T.J. Enright, B. Shelton, *Categories of highest weight modules: applications to Hermitian symmetric spaces*, *Mem. Amer. Math. Soc.*, **367** (1987). MR **88f**:22052
- [14] O. Gabber, A. Joseph, *Towards the Kazhdan-Lusztig conjecture*, *Ann. Sci. École. Norm. Sup.*, **14** (1981), 261-302. MR **83e**:17009
- [15] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, *Cambridge Stud. Adv. Math.*, no. 29, Cambridge Univ. Press, Cambridge, 1990. MR **92h**:20002
- [16] D. Kazhdan, G. Lusztig, *Representations of Coxeter groups and Hecke algebras*, *Invent. Math.*, **53** (1979), 165-184. MR **81j**:20066
- [17] B. Shelton, *Extensions Between Generalized Verma Modules: The Hermitian Symmetric Case*, *Math. Z.*, **197** (1988), 305-318. MR **89d**:17006
- [18] R. P. Stanley, *Enumerative Combinatorics*, vol. 1, Wadsworth and Brooks/Cole, Monterey, CA, 1986. MR **87j**:05003
- [19] R. P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge Stud. Adv. Math., no. 62, Cambridge Univ. Press, Cambridge, 1999. MR **2000k**:05026

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA "LA SAPIENZA", 00185 ROMA, ITALY
Current address: LACIM, Université du Québec à Montréal, case postale 8888, succursale
 Centre-Ville, Montréal, Québec, Canada H3C 3P8
E-mail address: biagioli@math.uqam.ca