

NEWTON POLYHEDRA, UNSTABLE FACES AND THE POLES OF IGUSA’S LOCAL ZETA FUNCTION

KATHLEEN HOORNAERT

ABSTRACT. In this paper we examine when the order of a pole of Igusa’s local zeta function associated to a polynomial f is smaller than “expected”. We carry out this study in the case that f is sufficiently non-degenerate with respect to its Newton polyhedron $\Gamma(f)$, and the main result of this paper is a proof of one of the conjectures of Denef and Sargos. Our technique consists in reducing our question about the polynomial f to the same question about polynomials f_μ , where μ are faces of $\Gamma(f)$ depending on the examined pole and f_μ is obtained from f by throwing away all monomials of f whose exponents do not belong to μ . Secondly, we obtain a formula for Igusa’s local zeta function associated to a polynomial f_μ , with μ unstable, which shows that, in this case, the upperbound for the order of the examined pole is obviously smaller than “expected”.

1. INTRODUCTION

For p prime, denote the field of p -adic numbers by \mathbb{Q}_p , the ring of p -adic integers by \mathbb{Z}_p , and the finite field of p elements by \mathbb{F}_p . If R is a commutative ring with identity, we denote the set of its units by R^\times .

Definition 1.1. Let $f(x) = f(x_1, \dots, x_n) \in \mathbb{Z}_p[x_1, \dots, x_n]$ with p prime. For $z \in \mathbb{Q}_p$, $\text{ord } z \in \mathbb{Z} \cup \{\infty\}$ denotes the valuation, $|z| = p^{-\text{ord } z}$ and $\text{ac}(z) = p^{-\text{ord } z} z$ denotes the angular component. Let $\chi : \mathbb{Z}_p^\times \rightarrow \mathbb{C}^\times$ be a character of \mathbb{Z}_p^\times , i.e., a group homomorphism with finite image. We formally set $\chi(0) = 0$. To the above data we associate the following two *Igusa local zeta functions* (the global and the local functions):

$$Z_f(s, \chi) = \int_{\mathbb{Z}_p^n} \chi(\text{ac } f(x)) |f(x)|^s |dx|,$$
$$Z_{f,0}(s, \chi) = \int_{(p\mathbb{Z}_p)^n} \chi(\text{ac } f(x)) |f(x)|^s |dx|,$$

for $s \in \mathbb{C}$, $\text{Re}(s) > 0$, where $|dx|$ denotes the Haar measure on \mathbb{Q}_p^n normalized so that \mathbb{Z}_p^n has measure 1. If χ is the trivial character χ_{triv} , then we write $Z_f(s)$ (resp. $Z_{f,0}(s)$) instead of $Z_f(s, \chi_{\text{triv}})$ (resp. $Z_{f,0}(s, \chi_{\text{triv}})$).

Received by the editors March 12, 2002.

2000 *Mathematics Subject Classification.* Primary 11S40, 11D79; Secondary 14M25, 52B20, 14G10.

Key words and phrases. Igusa zeta function, Newton polyhedron, congruences, p -adic integrals.

©2003 American Mathematical Society

Igusa's local zeta function $Z_f(s)$ is directly related to the number of solutions of the congruences $f(x) \equiv 0 \pmod{p^m}$, $m = 1, 2, 3, \dots$; see, e.g., [Igu78, pp. 97–98], [Den91, Section 1.2]. Using resolutions of singularities, Igusa [Igu74] proved that $Z_f(s, \chi)$ is a rational function of p^{-s} (see also [Igu78]). An entirely different proof was obtained ten years later by Denef [Den84] using p -adic cell decomposition instead of resolutions of singularities. We denote the meromorphic continuation of $Z_f(s, \chi)$ again by $Z_f(s, \chi)$. We study the poles of Igusa's local zeta function for a special but important class of polynomials, namely the polynomials that are non-degenerate with respect to their Newton polyhedron. This study was first started by Lichtin and Meuser [LM85] for polynomials in two variables.

Definition 1.2. Let $f(x) = f(x_1, \dots, x_n) = \sum_{\omega \in \mathbb{N}^n} a_\omega x_1^{\omega_1} \cdots x_n^{\omega_n}$ be a non-zero polynomial over \mathbb{Z}_p with $f(0) = 0$. Let $\mathbb{R}^+ = \{x \in \mathbb{R} \mid x \geq 0\}$, and let $\text{supp}(f) = \{\omega \in \mathbb{N}^n \mid a_\omega \neq 0\}$ be the support of f . The *Newton polyhedron* $\Gamma(f)$ of f is defined as the convex hull in $(\mathbb{R}^+)^n$ of the set

$$\bigcup_{\omega \in \text{supp}(f)} \omega + (\mathbb{R}^+)^n.$$

The *global Newton polyhedron* $\Gamma_{\text{gl}}(f)$ of f is defined as the convex hull of $\text{supp}(f)$. It is easy to verify that $\Gamma(f) = \Gamma_{\text{gl}}(f) + (\mathbb{R}^+)^n$.

Because a Newton polyhedron is a polyhedron, every proper face is an exposed face.¹ So, every proper face of $\Gamma(f)$ is the intersection of $\Gamma(f)$ with a supporting hyperplane [Roc70, pp. 99–100]. **By the faces of $\Gamma(f)$ we mean the proper faces of $\Gamma(f)$ and the Newton polyhedron $\Gamma(f)$ itself.**

Definition 1.3. Let p be a prime number. Let f be as in Definition 1.2. For each face τ of the Newton polyhedron $\Gamma(f)$ of f , we define

$$f_\tau(x) = \sum_{\omega \in \tau} a_\omega x^\omega,$$

and we also define the polynomial $\bar{f}_\tau(x)$ with coefficients in \mathbb{F}_p by reducing each coefficient a_ω of f_τ modulo $p\mathbb{Z}_p$.

We say f is *non-degenerate over \mathbb{F}_p with respect to all the faces of its Newton polyhedron* $\Gamma(f)$ if for every² face τ of $\Gamma(f)$ the locus of the polynomial \bar{f}_τ has no singularities in $(\mathbb{F}_p^\times)^n$, or equivalently the set of congruences

$$\begin{cases} f_\tau(x) & \equiv 0 \pmod{p}, \\ \frac{\partial f_\tau}{\partial x_i}(x) & \equiv 0 \pmod{p}, \quad i = 1, \dots, n, \end{cases}$$

has no solution in $(\mathbb{Z}_p^\times)^n$.

We say f is *non-degenerate over \mathbb{F}_p with respect to the compact faces of its Newton polyhedron* if we have the same condition, but only for the compact faces of $\Gamma(f)$.

See [DH01] for some remarks about the definition of non-degenerate. In [DH01] and [Hoo02] (see Theorem 3.2) we find an explicit formula for $Z_f(s, \chi)$ that holds if f is non-degenerate over \mathbb{F}_p with respect to all the faces of its Newton polyhedron and χ is a character of \mathbb{Z}_p with conductor $c_\chi = 1$. The conductor c_χ is the smallest $c \in \mathbb{N} \setminus \{0\}$ such that χ is trivial on $1 + p^c\mathbb{Z}_p$. So, $c_{\chi_{\text{triv}}}$ is also equal to 1. It

¹Face and exposed face in the sense of [Roc70, p. 162].

²Thus also for $\Gamma(f)$.

is well known (see [Den91, Theorem 3.3]) that $Z_f(s, \chi)$ is constant as a function of s for $p \gg 0$ if the conductor $c_\chi > 1$ and f is a polynomial over \mathbb{Z} . For the exact condition of a similar result for f a polynomial over \mathbb{Z}_p and p a fixed prime number, we refer to the same reference. So, we have an explicit formula for $Z_f(s, \chi)$ for the relevant cases. The actors in the above-mentioned formula are the cones Δ_τ associated to the faces τ of $\Gamma(f)$ (see Definition 2.3) and, the number of elements N_τ of the sets $\{x \in (\mathbb{F}_p^\times)^n \mid \bar{f}_\tau(x) = 0\}$ if $\chi = \chi_{\text{triv}}$, or the sums $\sum_{a \in (\mathbb{F}_p^\times)^n} \chi(f_\tau(a))$ if $\chi \neq \chi_{\text{triv}}$. Zúñiga-Galindo [Zn99] obtained the same formula, as a special case of a more general result. For the related work of other authors we refer to [DH01] and the references therein. Using the formula from Theorem 3.2, we can get a set of possible poles for $Z_f(s, \chi)$ (which we call candidate poles) together with their expected orders, i.e., upper bounds for the orders if these candidate poles are actual poles (see Proposition 4.1, Definition 4.2, Proposition 4.5 and Proposition 4.7). This can all be expressed in terms of the faces of $\Gamma(f)$ and their dimensions.

We have special interest for the largest real candidate pole of $Z_f(s, \chi)$ different from -1 . It is well known (see Proposition 4.5) that this candidate pole equals $-1/t_0$ if $t_0 \neq 1$, and that its expected order equals κ , where (t_0, t_0, \dots, t_0) is the intersection point of the diagonal $D = \{(t, t, \dots, t) \in \mathbb{R}^n\}$ with $\Gamma(f)$, and κ is the codimension of the smallest face τ_0 of $\Gamma(f)$ that contains (t_0, t_0, \dots, t_0) . The main purpose of this article is to prove one of the conjectures of Denef and Sargos [DS92, Section 6.2] for non-degenerate polynomials in the p -adic case:

if τ_0 is an unstable face of $\Gamma(f)$ over \mathbb{F}_p with respect to f , and if $-1/t_0 \neq -1$ is a pole of $Z_f(s, \chi)$, then its order will be smaller than expected, i.e., $< \kappa$.

The notion “unstable” (see Definition 7.1) was first established by Denef and Sargos in [DS92]. The analogue of the above conjecture for the real case has been proved by Denef, Laeremans and Sargos [DLS97]. Inspired by the proof for the real case, we tried to reduce our question about the polynomial f to the same question about the polynomial f_{τ_0} . Namely, Theorem 5.3, our first main result, states that $-1/t_0$ is an actual pole of $Z_f(s, \chi)$ of the expected order if and only if the same holds for $Z_{f_{\tau_0}}(s, \chi)$. In particular, if $-1/t_0$ is not a pole of $Z_{f_{\tau_0}}(s, \chi)$ of the expected order, the same holds for $Z_f(s, \chi)$. Theorem 6.1 extends this result to arbitrary poles.

In Theorem 7.3, we will give a formula for $Z_f(s, \chi)$ that holds if $f = f_\sigma$, where $\sigma \subset \Gamma(f)$ is an unstable convex set over \mathbb{F}_p with respect to f (see Definition 7.1), and if f is non-degenerate over \mathbb{F}_p with respect to a particular set of faces of $\Gamma(f)$. The formula in this theorem, which is our second main result, is similar to the one in Theorem 3.2, but can provide a smaller set of candidate poles for $Z_f(s, \chi)$ or smaller upperbounds for their orders (see Example 8.1). Finally, Theorem 7.6 uses this formula together with Theorem 5.3 to obtain a proof of the above-mentioned conjecture of Denef and Sargos. Theorem 7.7 gives an analogous result for arbitrary poles.

I would like to thank Jan Denef for suggesting that I solve the above conjecture and for supervising this work. We also refer to the papers [PSS99] and [PS00] which treat similar questions in the archimedean setting.

2. MORE ABOUT NEWTON POLYHEDRA

In Definition 2.3, we give a partition of $(\mathbb{R}^+)^n$ in sets that are closely related to the Newton polyhedron of a polynomial f as in Definition 1.2. In order to define

these sets Δ_τ and to know more about them, we will present a selection of some well-known definitions and properties.

Definition 2.1. Let f be as in Definition 1.2. For $a \in (\mathbb{R}^+)^n$, we define

$$m(a) = \inf_{x \in \Gamma(f)} \{a \cdot x\}$$

and we define the *first meet locus* of a as

$$F(a) = \{x \in \Gamma(f) \mid a \cdot x = m(a)\},$$

where $a \cdot x$ denotes the scalar product of a and x .

Property 2.2. Let f be as in Definition 1.2 and $a \in (\mathbb{R}^+)^n$. Then $F(a)$, the first meet locus of a , is a face of $\Gamma(f)$. In particular, $F(0) = \Gamma(f)$ and $F(a)$ is a proper face of $\Gamma(f)$, if $a \neq 0$. Moreover, $F(a)$ is a compact face if and only if $a \in (\mathbb{R}^+ \setminus \{0\})^n$.

Definition 2.3. We define an equivalence relation on $(\mathbb{R}^+)^n$ by

$$a \sim a' \quad \text{if and only if} \quad F(a) = F(a').$$

If τ is a face of $\Gamma(f)$, we define the *cone associated to* τ as

$$\Delta_\tau = \{a \in (\mathbb{R}^+)^n \mid F(a) = \tau\}.$$

Note that $\Delta_{\Gamma(f)} = \{0\}$. We now study the other equivalence classes Δ_τ and their closures $\bar{\Delta}_\tau$. It is well known that Δ_τ is a relatively open subset of $(\mathbb{R}^+)^n$ and that $\bar{\Delta}_\tau$ is equal to $\{a \in (\mathbb{R}^+)^n \mid F(a) \supset \tau\}$.

Definition 2.4. A fan \mathcal{F} is a finite set of rational polyhedral cones such that

- (1) every (non-empty) face of a cone of \mathcal{F} is contained in \mathcal{F} and
- (2) the intersection of two arbitrary cones C_i and C_j in \mathcal{F} is a face of both C_i and C_j .

Lemma 2.5. Let f be as in Definition 1.2. The closures $\bar{\Delta}_\tau$ of the cones associated to the faces of $\Gamma(f)$ form a fan in $(\mathbb{R}^+)^n$. Moreover, we have the following. Let τ be a proper face of $\Gamma(f)$. Then the order-reversing map

$$\{\text{faces of } \Gamma(f) \text{ that contain } \tau\} \rightarrow \{\text{(non-empty) faces of } \bar{\Delta}_\tau\} : \sigma \mapsto \bar{\Delta}_\sigma$$

is one-to-one and onto.

Definition 2.6. A vector $a \in \mathbb{R}^n$ is called *primitive* if the components of a are integers whose greatest common divisor is 1.

Note that because a Newton polyhedron is a polyhedron, we can prove that every proper face τ of $\Gamma(f)$ is the (finite) intersection of the facets³ of $\Gamma(f)$ that contain τ . One can also prove that for every facet of $\Gamma(f)$ there exists a unique primitive vector in $\mathbb{N}^n \setminus \{0\}$ that is perpendicular to that facet.

Proposition 2.7. Let f be as in Definition 1.2. Let τ be a proper face of $\Gamma(f)$ and let $\gamma_1, \dots, \gamma_r$ be the facets of $\Gamma(f)$ that contain τ . Let a_1, \dots, a_r be the unique primitive vectors in $\mathbb{N}^n \setminus \{0\}$ that are perpendicular to $\gamma_1, \dots, \gamma_r$, respectively. Then

³Recall that a facet is an $(n - 1)$ -dimensional face.

the cones $\bar{\Delta}_\tau$ and Δ_τ are the following convex cones:

$$\bar{\Delta}_\tau = \left\{ \sum_{i=1}^r \lambda_i a_i \mid \lambda_i \in \mathbb{R}, \lambda_i \geq 0 \right\} \quad \text{and} \quad \Delta_\tau = \left\{ \sum_{i=1}^r \lambda_i a_i \mid \lambda_i \in \mathbb{R}, \lambda_i > 0 \right\}.$$

Moreover, $\dim \Delta_\tau = \dim \bar{\Delta}_\tau = \text{codim } \tau = n - \dim \tau$.

Definition 2.8. If $a_1, \dots, a_r \in \mathbb{R}^n \setminus \{0\}$, we call cone $(a_1, \dots, a_r) = \{ \sum_{i=1}^r \lambda_i a_i \mid \lambda_i \in \mathbb{R}, \lambda_i > 0 \}$ the cone strictly positively spanned by the vectors a_1, \dots, a_r . Fix a cone Δ . If there exist linearly independent vectors $a_1, \dots, a_r \in \mathbb{R}^n \setminus \{0\}$ such that $\Delta = \text{cone}(a_1, \dots, a_r)$, then Δ is called a simplicial cone. If moreover one can choose $a_1, \dots, a_r \in \mathbb{Z}^n$ (or in \mathbb{Q}^n), we say Δ is a rational simplicial cone.

Let τ be a proper face of $\Gamma(f)$ and let a_1, \dots, a_r be as in Proposition 2.7. Following from Proposition 2.7 and Lemma 2.9, one can partition the cone Δ_τ associated to the face τ into a finite number of rational simplicial cones such that each Δ_i is spanned by vectors from the set $\{a_1, \dots, a_r\}$.

Lemma 2.9. Let Δ be the cone strictly positively spanned by the vectors $a_1, \dots, a_r \in (\mathbb{R}^+)^n \setminus \{0\}$. Then there exists a finite partition of Δ into cones Δ_i , such that each Δ_i is strictly positively spanned by some vectors from the set $\{a_1, \dots, a_r\}$ that are linearly independent over \mathbb{R} .

Proof. [Den95, Lemma 2]. □

Definition 2.10. Let a_1, \dots, a_r be vectors in \mathbb{Z}^n that are linearly independent over \mathbb{Q} . We define the multiplicity of a_1, \dots, a_r as the index of the lattice $\mathbb{Z}a_1 + \dots + \mathbb{Z}a_r$ in the group of the points with integral coordinates of the vector space generated by a_1, \dots, a_r .

Remark. It is easy to verify that the multiplicity of a_1, \dots, a_r equals the number of elements in the set

$$\mathbb{Z}^n \cap \left\{ \sum_{i=1}^r \lambda_i a_i \mid 0 \leq \lambda_i < 1 \text{ for } i = 1, \dots, r \right\},$$

and also equals the greatest common divisor of the determinants of all $r \times r$ -matrices obtained by omitting columns from the matrix with rows a_1, \dots, a_r .

3. AN EXPLICIT FORMULA FOR $Z_f(s, \chi)$

In Theorem 3.2, we give a formula for $Z_f(s, \chi)$ that holds if f is non-degenerate over \mathbb{F}_p with respect to all the faces of its Newton polyhedron. In Section 7, we will give a similar formula for $Z_f(s, \chi)$ that holds if $f = f_\sigma$, where $\sigma \subset \Gamma(f)$ is an unstable convex set, and if f is non-degenerate over \mathbb{F}_p with respect to a smaller set of faces of $\Gamma(f)$.

Definition 3.1. For $k = (k_1, \dots, k_n) \in \mathbb{R}^n$, we define

$$\sigma(k) = \sum_{i=1}^n k_i.$$

Theorem 3.2. Let p be a prime number. Let f be as in Definition 1.2. Suppose that f is non-degenerate over the finite field \mathbb{F}_p with respect to all the faces of its

Newton polyhedron $\Gamma(f)$. Let χ be a character of \mathbb{Z}_p^\times with conductor $c_\chi = 1$. For each face τ of $\Gamma(f)$ denote by N_τ the number of elements in the set

$$\{a \in (\mathbb{F}_p^\times)^n \mid \bar{f}_\tau(a) = 0\}.$$

Let s be a complex variable with $\text{Re}(s) > 0$. Then

$$Z_f(s, \chi) = \sum_{\substack{\tau \text{ face} \\ \text{of } \Gamma(f)}} L_\tau S_{\Delta_\tau},$$

with

$$L_\tau = \begin{cases} p^{-n} \left((p-1)^n - pN_\tau \frac{p^s-1}{p^{s+1}-1} \right) & \text{for } \chi = \chi_{\text{triv}}, \\ p^{-n} \sum_{a \in (\mathbb{F}_p^\times)^n} \chi(f_\tau(a)) & \text{for } \chi \neq \chi_{\text{triv}}, \end{cases}$$

and

$$S_{\Delta_\tau} = \sum_{k \in \mathbb{N}^n \cap \Delta_\tau} p^{-\sigma(k)-m(k)s}$$

for each face τ of $\Gamma(f)$ (including $\tau = \Gamma(f)$).

We have $S_{\Delta_{\Gamma(f)}} = 1$ and the other S_{Δ_τ} can be calculated as follows. Take a partition of the cone Δ_τ associated to the proper face τ into rational simplicial cones Δ_i . Then clearly $S_{\Delta_\tau} = \sum_i S_{\Delta_i}$, where the summation is over the rational simplicial cones Δ_i and

$$S_{\Delta_i} = \sum_{k \in \mathbb{N}^n \cap \Delta_i} p^{-\sigma(k)-m(k)s}.$$

Let Δ_i be the cone strictly positively spanned by the linearly independent vectors $a_1, \dots, a_r \in \mathbb{N}^n \setminus \{0\}$. Then

$$S_{\Delta_i} = \frac{\sum_h p^{\sigma(h)+m(h)s}}{(p^{\sigma(a_1)+m(a_1)s} - 1) \dots (p^{\sigma(a_r)+m(a_r)s} - 1)},$$

where h runs through the elements of the set

$$\mathbb{Z}^n \cap \left\{ \sum_{j=1}^r \lambda_j a_j \mid 0 \leq \lambda_j < 1 \text{ for } j = 1, \dots, r \right\}.$$

In the theorem above, $m(\cdot)$ and $\sigma(\cdot)$ are defined as in Definitions 2.1 and 3.1.

Remarks.

- (1) Note that S_{Δ_τ} is a rational function in p^{-s} . We also denote its meromorphic continuation by S_{Δ_τ} .
- (2) We have

$$Z_{f,0}(s, \chi) = \sum_{\substack{\tau \text{ compact face} \\ \text{of } \Gamma(f)}} L_\tau S_{\Delta_\tau},$$

with L_τ and S_{Δ_τ} as in the statement of the theorem above. For this formula, we only need f to be non-degenerate over the finite field \mathbb{F}_p with respect to the compact faces of $\Gamma(f)$.

Proof. [DH01, Theorem 4.2] and [Hoo02, Theorem 3.4]. □

4. THE CANDIDATE POLES OF $Z_f(s, \chi)$ AND THEIR EXPECTED ORDERS

Proposition 4.1. *Suppose that the prime number p , the polynomial f and the character χ satisfy the conditions of Theorem 3.2. Let $\gamma_1, \dots, \gamma_r$ be all the facets of $\Gamma(f)$, and let a_1, \dots, a_r be the unique primitive vectors in $\mathbb{N}^n \setminus \{0\}$ that are perpendicular to $\gamma_1, \dots, \gamma_r$, respectively. Then the following hold:*

(i) *If s_1 is a pole of $Z_f(s)$, then*

$$s_1 = -1 + i \frac{k2\pi}{\log_e p} \quad \text{with } k \in \mathbb{Z}, \text{ or}$$

$$s_1 = -\frac{\sigma(a_j)}{m(a_j)} + i \frac{k2\pi}{m(a_j) \log_e p},$$

with $k \in \mathbb{Z}$ and $j \in \{1, \dots, r\}$ such that $m(a_j) \neq 0$.

(ii) *If s_1 is a pole of $Z_f(s)$, then $\text{Re}(s_1)$ is -1 or $\text{Re}(s_1)$ is of the form $-1/t_1$, where (t_1, t_1, \dots, t_1) is the intersection point of the diagonal $D = \{(t, t, \dots, t) \in \mathbb{R}^n\}$ with the supporting hyperplane of a facet of $\Gamma(f)$.*

(iii) *Suppose that s_1 is a pole of $Z_f(s, \chi)$ with $\chi \neq \chi_{\text{triv}}$. Then*

$$s_1 = -\frac{\sigma(a_j)}{m(a_j)} + i \frac{k2\pi}{m(a_j) \log_e p},$$

with $k \in \mathbb{Z}$ and $j \in \{1, \dots, r\}$ such that $m(a_j) \neq 0$. Moreover, $\text{Re}(s_1)$ is of the form $-1/t_1$, where (t_1, t_1, \dots, t_1) is the intersection point of the diagonal $D = \{(t, t, \dots, t) \in \mathbb{R}^n\}$ with the supporting hyperplane of a facet of $\Gamma(f)$.

Remarks.

- (1) The same is true for $Z_{f,0}(s)$ and $Z_{f,0}(s, \chi)$. Of course, then we only need f to be non-degenerate over \mathbb{F}_p with respect to all the compact faces of $\Gamma(f)$ (see Remark (2) after Theorem 3.2).
- (2) We call the complex numbers in Proposition 4.1(i) and (iii) the *candidate poles* of $Z_f(s)$ and $Z_f(s, \chi)$, respectively. The first candidate poles in (i) (with real part equal to -1) come from the L_τ , the second ones in (i) and the ones in (iii) come from the S_{Δ_τ} . In the next section, we will give more information about the largest real candidate pole coming from the S_{Δ_τ} .

Proof. [DH01, Proposition 5.1]. □

Looking at the formula from Theorem 3.2, we can give upperbounds for the orders of the poles of $Z_f(s, \chi)$.

Definition 4.2. Suppose that the prime number p , the polynomial f and the character χ satisfy the conditions of Theorem 3.2. Suppose that s_1 is a candidate pole of $Z_f(s, \chi)$, i.e., a number from the set described in Proposition 4.1. We define the *expected order of the candidate pole s_1* for $Z_f(s, \chi)$ (according to the formula for $Z_f(s, \chi)$ in Theorem 3.2) as

$$(4.1) \quad \text{Max}\{\text{order of the pole } s_1 \text{ for } L_\tau S_{\Delta_\tau} \mid \tau \text{ a face of } \Gamma(f)\},$$

where the order of the pole s_1 is defined as 0, if it is not a pole, and where L_τ and S_{Δ_τ} are defined as in Theorem 3.2. If $\text{Re}(s_1) \neq -1$ or $\chi \neq \chi_{\text{triv}}$, we omit L_τ in (4.1).

4.1. The largest real candidate pole coming from the S_{Δ_τ} .

Definition 4.3. Let $f(x_1, \dots, x_n)$ be a non-zero polynomial with $f(0) = 0$. We denote the unique intersection point of the diagonal $D = \{(t, t, \dots, t) \in \mathbb{R}^n\}$ with the boundary of the Newton polyhedron $\Gamma(f)$ of f by (t_0, t_0, \dots, t_0) . Denote the smallest face of $\Gamma(f)$ that contains (t_0, t_0, \dots, t_0) by τ_0 and its codimension in \mathbb{R}^n by κ . Put $\rho = -1/t_0$.

We search for more information about the largest real candidate pole of $Z_f(s)$ and $Z_f(s, \chi)$ coming from the S_{Δ_τ} . Lemma 4.4 implies that this is ρ and Proposition 4.5 shows that the expected order of ρ for $Z_f(s)$ and $Z_f(s, \chi)$ is either κ or $\kappa + 1$.

Lemma 4.4. Let $f(x) = f(x_1, \dots, x_n)$ be a non-zero polynomial with $f(0) = 0$. For every $a \in (\mathbb{R}^+)^n$ we have $\sigma(a) - m(a)(1/t_0) \geq 0$ with equality if and only if $\tau_0 \subset F(a)$ (or equivalently $a \in \bar{\Delta}_{\tau_0}$).

Proof. See [DH01, Lemma 5.3]. □

Proposition 4.5. Suppose that the prime number p , the polynomial f and the character χ satisfy the conditions of Theorem 3.2. Then for every pole s_1 of $Z_f(s, \chi)$, one has $\operatorname{Re}(s_1) \leq \rho$ if $\chi \neq \chi_{\text{triv}}$; otherwise $\operatorname{Re}(s_1) = -1$ or $\operatorname{Re}(s_1) \leq \rho$. Moreover, the expected order of the candidate pole ρ for $Z_f(s, \chi)$ equals $\kappa + 1$ if $\chi = \chi_{\text{triv}}$, $\rho = -1$ and there is a face $\tau \subset \tau_0$ such that $N_\tau \neq 0$; otherwise, its expected order will be κ .

Recall that the actual order is not larger than the expected order.

Proof. See [DH01, Proposition 5.5] or [Zn99]. The case of the trivial character is also a direct consequence of the material in [Den95]. □

4.2. The expected order of an arbitrary pole of $Z_f(s, \chi)$.

Definition 4.6. Suppose that the prime number p , the polynomial f and the character χ satisfy the conditions of Theorem 3.2. Suppose that s_1 is a candidate pole of $Z_f(s, \chi)$, i.e., a number from the set described in Proposition 4.1. We say that a vector $a \in (\mathbb{R}^+)^n \setminus \{0\}$ or a facet $F(a)$ contributes to the candidate pole s_1 if $p^{\sigma(a)+m(a)s_1} = 1$, or equivalently, $\operatorname{Re}(s_1) = -\sigma(a)/m(a)$ and there exists a $k \in \mathbb{Z}$ such that $\operatorname{Im}(s_1) = k2\pi/(m(a) \log_e p)$. We say that a face of $\Gamma(f)$ is a *face of pure contribution to the candidate pole s_1* if every facet that contains τ contributes to this candidate pole s_1 . We define $\kappa(s_1)$ as the maximum of the codimensions of such faces. We call such a face of codimension $\kappa(s_1)$ a *face of $\Gamma(f)$ of maximal pure contribution to the candidate pole*.

Remarks.

- (1) It can occur that not all faces of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 are disjoint. See Example 8.2.
- (2) If $s_1 = \rho$, then $\kappa(s_1) = \kappa$ and τ_0 is the only face of $\Gamma(f)$ of maximal pure contribution to the candidate pole ρ of $Z_f(s, \chi)$.

Proposition 4.7. Suppose that the prime number p , the polynomial f and the character χ satisfy the conditions of Theorem 3.2. Suppose that s_1 is a candidate pole of $Z_f(s, \chi)$, i.e., a number from the set described in Proposition 4.1.

- (i) Let μ be a face of $\Gamma(f)$ of pure contribution to the candidate pole s_1 and suppose that its codimension is k . There exists a constant $c_\mu > 0$, independent of p , such that

$$\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^k S_{\Delta_\mu} = c_\mu.$$

Define the half-space $H_1^- = \{x \in \mathbb{R}^n \mid \sum_{i=1}^n x_i \leq 1\}$ and define $\omega_{\bar{\Delta}_\mu}$ as the volume form on the vector space $\text{vct } \bar{\Delta}_\mu$ generated by $\bar{\Delta}_\mu$, normalized such that the parallelepiped spanned by a lattice basis of $\mathbb{Z}^n \cap \text{vct } \bar{\Delta}_\mu$ has volume 1. Then

$$c_\mu = k! |\text{Re}(s_1)|^k \text{Vol}_{\omega_{\bar{\Delta}_\mu}}(\bar{\Delta}_\mu \cap H_1^-),$$

where the volume is taken with respect to $\omega_{\bar{\Delta}_\mu}$.

- (ii) If $p^{s_1+1} \neq 1$ or $\chi \neq \chi_{\text{triv}}$, then the expected order of the candidate pole s_1 of $Z_f(s, \chi)$ is $\kappa(s_1)$. Otherwise, the expected order is either $\kappa(s_1)$ or $\kappa(s_1) + 1$.

Proof. [Hoo02, Lemma 4.8 and Proposition 4.9]. □

5. CHECKING IF $\rho = -1/t_0$ IS A POLE OF THE EXPECTED ORDER:
REDUCTION FROM f TO f_{τ_0}

Definition 5.1. Let $V = \{v_1, \dots, v_r\}$ be a finite subset of \mathbb{Z}^n such that its elements are linearly independent over \mathbb{Q} . Define

$$(5.1) \quad H_V = H_{\{v_1, \dots, v_r\}} = \mathbb{Z}^n \cap \left\{ \sum_{i=1}^r \lambda_i v_i \mid 0 \leq \lambda_i < 1 \text{ for } i = 1, \dots, r \right\}.$$

Note that all λ_i belong to \mathbb{Q} .

Recall that we can choose a unique representative in $H_{\{v_1, \dots, v_r\}}$ for a coset of the lattice $\mathbb{Z}v_1 + \dots + \mathbb{Z}v_r$ in the group of the points with integral coordinates of the vector space generated by v_1, \dots, v_r . So, the number of elements in $H_{\{v_1, \dots, v_r\}}$ equals the multiplicity of v_1, \dots, v_r .

In the formula for Igusa's local zeta function from Theorem 3.2, we have to calculate sums of the form $\sum_h p^{\sigma(h)+m(h)s}$, where h runs through the elements of a set of the form $H_{\{v_1, \dots, v_r\}}$. The next lemma gives a relation between those sets if some of the vectors from $\{v_1, \dots, v_r\}$ are the same and the others generate the same vector space. We use this relation in the proof of Theorem 5.3, which shows that $\rho \neq -1$ is a pole of $Z_f(s, \chi)$ of the expected order (see Proposition 4.5) if and only if the same holds for $Z_{f_{\tau_0}}(s, \chi)$. This is very surprising because the cones Δ_τ , and hence the S_{Δ_τ} for $\tau \subset \tau_0$ in the formulas for $Z_f(s, \chi)$ and $Z_{f_{\tau_0}}(s, \chi)$ from Theorem 3.2, can be different. For example, let $f(x, y, z) = xy + xz^2 + yz^2 + z^5 + y^7$. Then τ_0 is the convex hull of $(1, 1, 0)$, $(1, 0, 2)$ and $(0, 1, 2)$. So, $f_{\tau_0} = xy + xz^2 + yz^2$ and

$$\begin{aligned} \Delta_{(0,1,2)}(f) &= \text{cone}((1, 0, 0), (2, 2, 1), (3, 3, 1), (6, 1, 3)), \\ \Delta_{(0,1,2)}(f_{\tau_0}) &= \text{cone}((1, 0, 0), (2, 2, 1), (1, 1, 0), (2, 0, 1)). \end{aligned}$$

See also Figure 5.1 and Figure 8.2. In the case that the codimension of τ_0 is 1, one can easily prove Theorem 5.3. The real difficulties only occur when its codimension is larger than 1.




FIGURE 5.1. Newton polyhedron of $xy + xz^2 + yz^2 + z^5 + y^7$

Lemma 5.2. *Let A, A' and B be finite subsets of \mathbb{Z}^n such that each of the three sets A, A' and $A \cup B$ consists of linearly independent elements over \mathbb{Q} and such that A and A' generate the same vector space. Let π be the natural projection of $\text{vct } A + \text{vct } B$ onto $\text{vct } B$ (here “vct” denotes “the vector space generated by”). Then we have*

- (i) $\#(\pi^{-1}(\{b\}) \cap H_{A \cup B}) = \#H_A$ for any $b \in \pi(H_{A \cup B})$, and
- (ii) $\pi(H_{A \cup B}) = \pi(H_{A' \cup B})$.

Proof. Let $A = \{a_1, \dots, a_u\}$, $A' = \{a'_1, \dots, a'_u\}$ and $B = \{b_1, \dots, b_v\}$.

- (i) Suppose that $h = \sum_{i=1}^u \alpha_i a_i + b \in H_{A \cup B}$ with $b = \sum_{j=1}^v \beta_j b_j$. Suppose that $g = \sum_{i=1}^u \gamma_i a_i + b \in (\pi^{-1}(\{b\}) \cap H_{A \cup B})$. Then $g - h \in \mathbb{Z}^n \cap \text{vct } \{a_1, \dots, a_u\}$. So, there exists a unique $k \in H_A$ and unique elements $z_1, \dots, z_u \in \mathbb{Z}$ such that $g = h + k + \sum_{i=1}^u z_i a_i$ because $a_1, \dots, a_u \in \mathbb{Z}^n$ are linearly independent. Conversely, if $k \in H_A$, there exists a unique $g \in (\pi^{-1}(\{b\}) \cap H_{A \cup B})$ and unique elements $z_1, \dots, z_u \in \mathbb{Z}$ such that $g = h + k + \sum_{i=1}^u z_i a_i$. This shows that $\#(\pi^{-1}(\{b\}) \cap H_{A \cup B}) = \#H_A$.
- (ii) By the symmetry of the problem, it suffices to prove that $\pi(H_{A \cup B}) \subset \pi(H_{A' \cup B})$. Suppose that $h = \sum_{i=1}^u \alpha_i a_i + b \in H_{A \cup B}$ with $b = \sum_{j=1}^v \beta_j b_j$.

We now show that there also exists an element $h' = \sum_{i=1}^u \alpha'_i a'_i + b \in H_{A' \cup B}$, such that $\pi(h') = b = \pi(h)$. Because a'_1, \dots, a'_u generate the same vector space over \mathbb{Q} as a_1, \dots, a_u , there exist $\gamma_1, \dots, \gamma_u \in \mathbb{Q}$ such that $\sum_{i=1}^u \alpha_i a_i = \sum_{i=1}^u \gamma_i a'_i$. For each i , there exists a unique $0 \leq \alpha'_i < 1$ and a $z_i \in \mathbb{Z}$ such that $\gamma_i = \alpha'_i + z_i$. Because the elements a'_1, \dots, a'_u belong to \mathbb{Z}^n , it is clear that $h' = \sum_{i=1}^u \alpha'_i a'_i + b = h - \sum_{i=1}^u z_i a'_i$ is an element of \mathbb{Z}^n and hence, also of $H_{A' \cup B}$. □

Theorem 5.3. *Suppose that the prime number p , the polynomial f and the character χ satisfy the conditions of Theorem 3.2. Let τ_0, ρ and κ be defined as in Definition 4.3. Then*

$$\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^k Z_f(s, \chi) = c \lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^k Z_{f_{\tau_0}}(s, \chi),$$

where

$$k = \begin{cases} \kappa + 1 & \text{if } \rho = -1 \text{ and } \chi = \chi_{\text{triv}}, \\ \kappa & \text{otherwise,} \end{cases}$$

and $c > 0$ is a constant real number, independent of p . In fact,

$$c = \frac{\text{Vol}(\bar{\Delta}_{\tau_0}(f) \cap H_1^-)}{\text{Vol}(\bar{\Delta}_{\tau_0}(f_{\tau_0}) \cap H_1^-)},$$

where the volume is the one on the vector space generated by $\bar{\Delta}_{\tau_0}$, induced by the standard metric on \mathbb{R}^n , and H_1^- is the half-space $\{x \in \mathbb{R}^n \mid \sum_{i=1}^n x_i \leq 1\}$.

Remarks.

- (1) If it is important to know whether we examine the cone Δ_τ associated to the polynomial f or to the polynomial f_{τ_0} , we will indicate this by $\Delta_\tau(f)$ and $\Delta_\tau(f_{\tau_0})$, respectively.
- (2) The proof below also shows that a similar result holds for $Z_{f,0}(s, \chi)$, even if τ_0 is not compact. Then f has to be non-degenerate over \mathbb{F}_p with respect to all the compact faces of its Newton polyhedron.
- (3) If $\chi = \chi_{\text{triv}}$, $\rho = -1$ and the expected order of the candidate pole ρ for $Z_f(s)$ equals κ , we know by [Hoo02, Theorem 4.10] that ρ is an actual pole of $Z_f(s)$ of order κ . So, Theorem 5.3 will be useful for the other cases.

Proof. Note also that f_{τ_0} is non-degenerate over the finite field \mathbb{F}_p with respect to all the faces of its Newton polyhedron $\Gamma(f_{\tau_0})$. Indeed, suppose that τ is a face of $\Gamma(f_{\tau_0})$. Then $(f_{\tau_0})_\tau = f_{\tau_0 \cap \tau}$ and the face $\tau_0 \cap \tau$ of τ_0 is also a face of the Newton polyhedron $\Gamma(f)$, by [Roc70, p. 165]. So, we can use the formula for $Z_f(s, \chi)$ and $Z_{f_{\tau_0}}(s, \chi)$ from Theorem 3.2.

We know from [DH01, Lemma 5.4] that $\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_\tau} = 0$ for faces $\tau \not\subset \tau_0$ and that $\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_\tau} > 0$ for faces $\tau \subset \tau_0$. Moreover, $L_\tau(f) = L_\tau(f_{\tau_0})$ for faces $\tau \subset \tau_0$. So, it suffices to prove that

$$(5.2) \quad \frac{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_\tau}(f)}{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_{\tau_0}}(f)} = \frac{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_\tau}(f_{\tau_0})}{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_{\tau_0}}(f_{\tau_0})}$$

for faces $\tau \subset \tau_0$, and that

$$(5.3) \quad \frac{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_{\tau_0}}(f)}{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_{\tau_0}}(f_{\tau_0})} = \frac{\text{Vol}(\bar{\Delta}_{\tau_0}(f) \cap H_1^-)}{\text{Vol}(\bar{\Delta}_{\tau_0}(f_{\tau_0}) \cap H_1^-)}.$$

The last equality follows by Proposition 4.7 because $\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_{\tau_0}}$ equals $\kappa!|\rho|^\kappa$ times the volume of $\bar{\Delta}_{\tau_0} \cap H_1^-$ with respect to the volume form $\omega_{\bar{\Delta}_{\tau_0}}$, where $\omega_{\bar{\Delta}_{\tau_0}}$ is the volume form on $\text{vct } \bar{\Delta}_{\tau_0}$, normalized such that the parallelepiped spanned by a lattice basis of $\mathbb{Z}^n \cap \text{vct } \bar{\Delta}_{\tau_0}$ has volume 1. Because $\bar{\Delta}_{\tau_0}(f)$ and $\bar{\Delta}_{\tau_0}(f_{\tau_0})$ generate the same vector space, it is clear that we can also take the quotient of the volumes induced by the standard metric on \mathbb{R}^n .

Suppose that $\bigcup_i \Delta_i$ is a finite partition of Δ_τ into rational simplicial cones. Then $S_{\Delta_\tau} = \sum_i S_{\Delta_i}$. From the formula for S_{Δ_i} in Theorem 3.2, Lemma 4.4 and the fact that $\dim \bar{\Delta}_{\tau_0} = \kappa$, it follows that $\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_i} > 0$ if and only if (exactly) κ generators of Δ_i belong to $\bar{\Delta}_{\tau_0}$; otherwise, this limit is zero. So, to prove equality (5.2), it is only important to know which cones in the partition have κ generators in $\bar{\Delta}_{\tau_0}$. We will choose partitions of $\Delta_\tau(f)$ and $\Delta_\tau(f_{\tau_0})$ that are closely related to some chosen partitions of $\Delta_{\tau_0}(f)$ and $\Delta_{\tau_0}(f_{\tau_0})$, respectively.

Fix a finite partition of $\Delta_{\tau_0}(f)$ into rational simplicial cones and let

$$\text{cone } A_i, \text{ for } i = 1, \dots, \ell,$$

be the only cones of dimension κ (i.e., with κ generators in $\bar{\Delta}_{\tau_0}(f)$) in this partition. Here, each A_i is a set of κ vectors in \mathbb{N}^n that are linearly independent over \mathbb{Q} . Similarly, fix a finite partition of $\Delta_{\tau_0}(f_{\tau_0})$ into rational simplicial cones, let

$$\text{cone } A'_j, \text{ for } j = 1, \dots, m,$$

be the only cones of dimension κ in this partition, and let each A'_j be a set of κ vectors in \mathbb{N}^n that are linearly independent over \mathbb{Q} . Note that the sets A_i and the sets A'_j all generate the same vector space.

It is clear that $\Delta_\tau(f) \subset \Delta_\tau(f_{\tau_0})$ for every face $\tau \subset \tau_0$. Indeed, if $H_a = \{x \in \mathbb{R}^n \mid a \cdot x = b\}$ is a supporting hyperplane of $\Gamma(f)$ such that $H_a \cap \Gamma(f) = \tau$, then the same property holds if we replace $\Gamma(f)$ by $\Gamma(f_{\tau_0})$. Fix now for each face $\tau \subsetneq \tau_0$ a primitive vector

$$(5.4) \quad b_\tau \in \Delta_\tau(f) \subset \Delta_\tau(f_{\tau_0}).$$

From Lemma 2.5 and Lemma 5.5 below, it follows by induction on d that the following assertion is true for any d in $\mathbb{N} \setminus \{0\}$.

Assertion 5.4. *Suppose that the face $\tau \subsetneq \tau_0$ and that $\dim \tau = \dim \tau_0 - d$. Then there exists a partition of $\Delta_\tau(f)$ into rational simplicial cones, such that the only cones in this partition that have κ generators in $\bar{\Delta}_{\tau_0}(f)$ are precisely the cones of the form*

$$\text{cone } (A_i \cup B_\tau),$$

where $i \in \{1, \dots, \ell\}$ and B_τ is a set of the form $\{b_{\tau_1}, \dots, b_{\tau_{v-1}}, b_\tau\}$, where $v \leq d$ and $v = d$ for the cones in this partition of dimension equal to $\dim \Delta_\tau(f)$, $\tau \subsetneq \tau_{v-1} \subsetneq \dots \subsetneq \tau_1 \subsetneq \tau_0$ are faces of τ_0 and $b_{\tau_1}, \dots, b_{\tau_{v-1}}, b_\tau$ are defined in (5.4). Similarly, there exists a partition of $\Delta_\tau(f_{\tau_0})$ into rational simplicial cones such that the only cones in this partition that have κ generators in $\bar{\Delta}_{\tau_0}(f_{\tau_0})$ are precisely the cones of the form

$$\text{cone } (A'_j \cup B_\tau),$$

where $j \in \{1, \dots, m\}$ and B_τ is defined as above.

Note that one adds the same sets B_τ to the sets A'_j as one adds to the sets A_i . Also note that $\tau \subsetneq \tau_{v-1} \subsetneq \dots \subsetneq \tau_1 \subsetneq \tau_0$ for faces of τ_0 is equivalent to $\bar{\Delta}_\tau \supsetneq$

$\bar{\Delta}_{\tau_{v-1}} \supseteq \cdots \supseteq \bar{\Delta}_{\tau_1} \supseteq \bar{\Delta}_{\tau_0}$ for faces of $\bar{\Delta}_\tau$ (Lemma 2.5), and that $\dim \tau = \dim \tau_0 - d$ is equivalent to $\dim \bar{\Delta}_\tau = \dim \bar{\Delta}_{\tau_0} + d$ (Proposition 2.7).

From the formula for $S_{\text{cone}(A_i \cup B_\tau)}$ in Theorem 3.2, it follows that

$$S_{\text{cone}(A_i \cup B_\tau)} = \frac{\sum_{h \in H_{A_i \cup B_\tau}} p^{\sigma(h) + m(h)s}}{\prod_{x \in A_i \cup B_\tau} (p^{\sigma(x) + m(x)s} - 1)}.$$

We know that m is linear on $\bar{\Delta}_\tau(f)$, and so σ and m are linear on $H_{A_i \cup B_\tau}$. It is also clear that $\sigma(a) + m(a)\rho = 0$ for every $a \in \text{cone } A_i \subset \bar{\Delta}_{\tau_0}(f)$. Let π be the natural projection of $\text{vct } A_i + \text{vct } B_\tau$ onto $\text{vct } B_\tau$. Then

$$\begin{aligned} \sum_{h \in H_{A_i \cup B_\tau}} p^{\sigma(h) + m(h)\rho} &= \sum_{h \in H_{A_i \cup B_\tau}} p^{\sigma(\pi(h)) + m(\pi(h))\rho} \\ &= \#H_{A_i} \sum_{b \in \pi(H_{A_i \cup B_\tau})} p^{\sigma(b) + m(b)\rho}, \end{aligned}$$

by Lemma 5.2(i). Consequently, $\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\text{cone}(A_i \cup B_\tau)}$ equals

$$\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa \frac{\#H_{A_i}}{\prod_{x \in A_i} (p^{\sigma(x) + m(x)s} - 1)} \frac{\sum_{b \in \pi(H_{A_i \cup B_\tau})} p^{\sigma(b) + m(b)\rho}}{\prod_{x \in B_\tau} (p^{\sigma(x) + m(x)\rho} - 1)}.$$

To finish the proof, we define $S(P, V, g, s)$ for a finite subset P of \mathbb{Q}^n , a finite subset V of \mathbb{N}^n , a polynomial g and $s \in \mathbb{C}$ as

$$S(P, V, g, s) = \frac{\sum_{x \in P} p^{\sigma(x) + m(x)s}}{\prod_{x \in V} (p^{\sigma(x) + m(x)s} - 1)},$$

and where the function m is completely determined by the Newton polyhedron $\Gamma(g)$ of the polynomial g . Note that if cone V is a rational simplicial cone contained in $(\mathbb{R}^+)^n$ (i.e., V is a finite subset of \mathbb{N}^n such that its elements are linearly independent) and cone V is contained in a cone associated to a face of $\Gamma(g)$, then

$$S_{\text{cone } V} = S(H_V, V, g, s),$$

where $S_{\text{cone } V}$ is defined as in Theorem 3.2. Using this notation,

$$\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\text{cone}(A_i \cup B_\tau)}$$

equals

$$S(\pi(H_{A_i \cup B_\tau}), B_\tau, f, \rho) \lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\text{cone } A_i}.$$

Note that the sets A_i all generate the same vector space. So, Lemma 5.2(ii) implies that $\pi(H_{A_i \cup B_\tau}) = \pi(H_{A_1 \cup B_\tau})$ for every $i \in \{1, \dots, \ell\}$. Therefore,

$$\begin{aligned} &\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_\tau}(f) \\ &= \sum_{B_\tau} \sum_{i=1}^\ell \lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\text{cone}(A_i \cup B_\tau)} \\ &= \left(\sum_{B_\tau} S(\pi(H_{A_1 \cup B_\tau}), B_\tau, f, \rho) \right) \lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_{\tau_0}}(f), \end{aligned}$$

where B_τ runs through all the sets described in Assertion 5.4. Similarly,

$$\begin{aligned} & \lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_\tau}(f_{\tau_0}) \\ &= \left(\sum_{B_\tau} S(\pi(H_{A'_1 \cup B_\tau}), B_\tau, f_{\tau_0}, \rho) \right) \lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_{\tau_0}}(f_{\tau_0}). \end{aligned}$$

Note that A'_1 generates the same vector space as A_1 . So, the sets $\pi(H_{A'_1 \cup B_\tau})$ and $\pi(H_{A_1 \cup B_\tau})$ are equal, by Lemma 5.2(ii). Moreover, the $m(b)$ for $b \in \pi(H_{A_1 \cup B_\tau}) \cup B_\tau$ are also the same, because $m(b) = b \cdot x_\tau$ for every $b \in \bar{\Delta}_\tau$, where x_τ is a fixed element of $\tau \subset \tau_0$. Now, it is clear that $S(\pi(H_{A_1 \cup B_\tau}), B_\tau, f, \rho) = S(\pi(H_{A'_1 \cup B_\tau}), B_\tau, f_{\tau_0}, \rho)$ and that

$$\begin{aligned} \frac{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_\tau}(f)}{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_{\tau_0}}(f)} &= \sum_{B_\tau} S(\pi(H_{A_1 \cup B_\tau}), B_\tau, f, \rho) \\ &= \frac{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_\tau}(f_{\tau_0})}{\lim_{s \rightarrow \rho} (p^{s-\rho} - 1)^\kappa S_{\Delta_{\tau_0}}(f_{\tau_0})}, \end{aligned}$$

which proves (5.2). □

One easily proves the following lemma.

Lemma 5.5. *Suppose that $\Delta = \text{cone}\{a_1, \dots, a_r\}$ is the cone strictly positively spanned by the vectors $a_1, \dots, a_r \in (\mathbb{R}^+)^n \setminus \{0\}$. Fix an element $b \in \Delta$. Then Δ is the disjoint union of the cones of the form*

$$W = \{\beta_1 b + f \mid \beta_1 \in \mathbb{R}, \beta_1 > 0, f \in \text{ri } F\},$$

where F is a proper face of $\bar{\Delta}$ and $\text{ri } F$ is its interior relative to the affine hull of F .

6. CHECKING IF AN ARBITRARY CANDIDATE POLE IS A POLE OF THE EXPECTED ORDER: REDUCTION FROM f TO f_μ 'S

Theorem 6.1 is a generalization of Theorem 5.3 to arbitrary candidate poles s_1 of $Z_f(s, \chi)$. Because the faces of $\Gamma(f)$ of maximal pure contribution to a candidate pole s_1 with $\text{Re}(s_1) \neq \rho$ do not intersect the diagonal $D = \{(t, t, \dots, t) \in \mathbb{R}^n\}$, the situation can be more difficult.

Theorem 6.1. *Suppose that the prime number p , the polynomial f and the character χ satisfy the conditions of Theorem 3.2. Suppose that s_1 is a candidate pole of $Z_f(s, \chi)$, i.e., a number from the set described in Proposition 4.1. Suppose that, for every face μ of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 of $Z_f(s, \chi)$, the face μ is the only face of $\Gamma(f_\mu)$ of maximal pure contribution to the candidate pole s_1 of $Z_{f_\mu}(s, \chi)$. Then*

$$\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^k Z_f(s, \chi) = \sum_{\mu} c_\mu \lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^k Z_{f_\mu}(s, \chi),$$

where μ runs through the faces of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 ,

$$k = \begin{cases} \kappa(s_1) + 1 & \text{if } p^{s_1+1} = 1 \text{ and } \chi = \chi_{\text{triv}}, \\ \kappa(s_1) & \text{otherwise,} \end{cases}$$

and the $c_\mu > 0$ are constant real numbers, independent of p . In fact,

$$c_\mu = \frac{\text{Vol}(\bar{\Delta}_\mu(f) \cap H_1^-)}{\text{Vol}(\bar{\Delta}_\mu(f_\mu) \cap H_1^-)},$$

where the volume is the one on the vector space generated by $\bar{\Delta}_\mu$, induced by the standard metric on \mathbb{R}^n and H_1^- is the half-space $\{x \in \mathbb{R}^n \mid \sum_{i=1}^n x_i \leq 1\}$.

Remarks.

- (1) Suppose that μ is a face of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 . If the dimension of μ is 0 or 1, then μ is the only face of $\Gamma(f_\mu)$ of maximal pure contribution to the candidate pole s_1 of $Z_{f_\mu}(s, \chi)$. If the dimension of μ is > 1 , the condition on μ will be satisfied in most examples. Nevertheless, it is possible that μ is not the only face of $\Gamma(f_\mu)$ of maximal pure contribution to the candidate pole s_1 . For example, let $f(x, y, z) = x^7y^3z + x^4y^8 + x^8y^4 + z^7$. Then the Newton polyhedron $\Gamma(f)$ is given by the system of linear inequalities

$$\begin{cases} x \geq 0 \\ y \geq 0 \\ z \geq 0 \\ x + y + 2z \geq 12 \\ y + z \geq 4 \\ 2y + z \geq 7 \\ 27x + 25y + 44z \geq 308 \\ 7x + 4z \geq 28 \end{cases}$$

(see also Figure 6.1). We denote the facets with supporting hyperplanes given by the corresponding equalities in the inequalities above by τ_1, \dots, τ_8 . Face τ_4 is the only face of $\Gamma(f)$ of maximal pure contribution to the candidate pole $-1/3$.




FIGURE 6.1. Newton polyhedron of $x^7y^3z + x^4y^8 + x^8y^4 + z^7$




FIGURE 6.2. Newton polyhedron of $x^7y^3z + x^4y^8 + x^8y^4$

The Newton polyhedron $\Gamma(f_{\tau_4})$ of $f_{\tau_4} = x^7y^3z + x^4y^8 + x^8y^4$ is given by the system of linear inequalities

$$\begin{cases} x \geq 4 \\ y \geq 3 \\ z \geq 0 \\ x + y + 2z \geq 12 \\ y + z \geq 4 \\ 5x + 3y \geq 44 \end{cases}$$

(see also Figure 6.2). We denote the facets with supporting hyperplanes given by the corresponding equalities in the inequalities above by $\tau'_1, \tau'_2, \tau_3, \tau_4, \tau_5, \tau'_6$. The face τ_4 is not the only face of $\Gamma(f_4)$ of maximal pure contribution to the candidate pole $-1/3$: τ'_2 is also such a face.

- (2) The proof below also shows that a similar result holds for $Z_{f,0}(s, \chi)$, even if the faces of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 are not compact. Then f has to be non-degenerate over \mathbb{F}_p with respect to all the compact faces of its Newton polyhedron.

Proof. By the same argument as in the proof of Theorem 5.3, we can also show that f_μ is non-degenerate over the finite field \mathbb{F}_p with respect to all the faces of its Newton polyhedron $\Gamma(f_\mu)$ if μ is a face of $\Gamma(f)$. So, we can use the formulas for $Z_f(s, \chi)$ and $Z_{f_\mu}(s, \chi)$ from Theorem 3.2.

We know from [Hoo02, Proposition 4.9] that $\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\tau} = 0$ for faces τ not contained in any face μ of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 . It is clear that $L_\tau(f) = L_\tau(f_\mu)$ for faces τ contained in a face μ of $\Gamma(f)$. For faces τ contained in at least one face μ of $\Gamma(f)$ of maximal pure contribution to the candidate pole, we will write

$$(6.1) \quad \lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\tau}(f) = \sum_{\mu} R_\mu,$$

where μ runs through all the faces of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 and where the R_μ , which depend on τ , will be specified later. The R_μ for $\mu \not\supset \tau$ will vanish. It suffices then to show that

$$(6.2) \quad \frac{R_\mu}{\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\mu}(f)} = \frac{\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\tau}(f_\mu)}{\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\mu}(f_\mu)}$$

for faces $\tau \subset \mu$, and that

$$(6.3) \quad \frac{\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\mu(f)}}{\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\mu(f_\mu)}} = \frac{\text{Vol}(\bar{\Delta}_\mu(f) \cap H_1^-)}{\text{Vol}(\bar{\Delta}_\mu(f_\mu) \cap H_1^-)},$$

where μ is a face of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 .

The last equality follows by Proposition 4.7 and the fact that $\bar{\Delta}_\mu(f)$ and $\bar{\Delta}_\mu(f_\mu)$ generate the same vector space.

Suppose that $\bigcup_i \Delta_i$ is a finite partition of Δ_τ into rational simplicial cones such that each Δ_i has at most $\kappa(s_1)$ generators contributing to the candidate pole s_1 . This is always possible; see below. Then $S_{\Delta_\tau} = \sum_i S_{\Delta_i}$. From the formula for S_{Δ_i} in Theorem 3.2, it follows that $\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_i} = 0$ if less than $\kappa(s_1)$ generators of Δ_i contribute to the candidate pole s_1 . So, to prove equality (6.2), it is only important to know the other cones. We will choose partitions of $\Delta_\tau(f)$ and $\Delta_\tau(f_\mu)$ that are closely related to some chosen partitions of $\Delta_\mu(f)$ and $\Delta_\mu(f_\mu)$, respectively.

Fix a finite partition of $\Delta_\mu(f)$ into rational simplicial cones, and let

$$\text{cone } A_{i,\mu}, \text{ for } i = 1, \dots, \ell_\mu,$$

be the only cones of dimension $\kappa(s_1)$, i.e., with $\kappa(s_1)$ generators contributing to the candidate pole s_1 , in this partition. Here, each $A_{i,\mu}$ is a set of $\kappa(s_1)$ vectors in \mathbb{N}^n that are linearly independent over \mathbb{Q} . Similarly, fix a finite partition of $\Delta_\mu(f_\mu)$ into rational simplicial cones, and let

$$\text{cone } A'_{j,\mu}, \text{ for } j = 1, \dots, m_\mu,$$

be the only cones of dimension $\kappa(s_1)$ in this partition where each $A'_{j,\mu}$ is a set of $\kappa(s_1)$ vectors in \mathbb{N}^n that are linearly independent over \mathbb{Q} .

It is clear that $\Delta_\tau(f) \subset \Delta_\tau(f_\mu)$ for every face $\tau \subset \mu$. Now for each face $\tau \subsetneq \mu$ fix a primitive vector

$$(6.4) \quad b_\tau \in \Delta_\tau(f) \subset \Delta_\tau(f_\mu)$$

that does not contribute to the candidate pole s_1 . This is always possible because σ and m are linear on $\bar{\Delta}_\tau(f)$ and not all generators of $\Delta_\tau(f)$ contribute to the candidate pole s_1 . From Lemma 2.5 and Lemma 5.5, it follows by induction on d that the following assertion is true for any d in $\mathbb{N} \setminus \{0\}$.

Assertion 6.2. *Suppose that the face τ is a face of $\Gamma(f)$ that is contained in at least one face of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 and that $\dim \tau = n - \kappa(s_1) - d$. Then there exists a partition of $\Delta_\tau(f)$ into rational simplicial cones such that the only cones in this partition that have $\kappa(s_1)$ generators contributing to the candidate pole s_1 are precisely the cones of the form*

$$\text{cone}(A_{i,\mu} \cup B_\tau),$$

where $\mu \supset \tau$ is a face of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 , $i \in \{1, \dots, \ell_\mu\}$, and B_τ is a set of the form $\{b_{\tau_1}, \dots, b_{\tau_{v-1}}, b_\tau\}$, where $v \leq d$ and $v = d$ for the cones in this partition of dimension equal to $\dim \Delta_\tau(f)$, $\tau \subsetneq \tau_{v-1} \subsetneq \dots \subsetneq \tau_1 \subsetneq \mu$ are faces of μ and $b_{\tau_1}, \dots, b_{\tau_{v-1}}, b_\tau$ are defined in (6.4). The other cones in the partition have less than $\kappa(s_1)$ generators contributing to the candidate pole s_1 . Similarly, there exists a partition of $\Delta_\tau(f_\mu)$ into rational simplicial cones

such that the only cones in this partition that have $\kappa(s_1)$ generators contributing to the candidate pole s_1 are precisely the cones of the form

$$\text{cone}(A'_{j,\mu} \cup B_\tau),$$

where $j \in \{1, \dots, m_\mu\}$ and B_τ is defined as above. The other cones in the partition have less than $\kappa(s_1)$ generators contributing to the candidate pole s_1 .

From the formula for $S_{\text{cone}(A_{i,\mu} \cup B_\tau)}$ in Theorem 3.2, it follows that

$$S_{\text{cone}(A_{i,\mu} \cup B_\tau)} = \frac{\sum_{h \in H_{A_{i,\mu} \cup B_\tau}} p^{\sigma(h)+m(h)s}}{\prod_{x \in A_{i,\mu} \cup B_\tau} (p^{\sigma(x)+m(x)s} - 1)}.$$

We know that m is linear on $\bar{\Delta}_\tau$, and so σ and m are linear on $H_{A_{i,\mu} \cup B_\tau}$. It is also clear that $p^{\sigma(a)+m(a)s_1} = 1$ for every $a \in \text{cone } A_{i,\mu} \subset \bar{\Delta}_\mu(f)$ because all generators of $\Delta_\mu(f)$ contribute to the candidate pole s_1 . Let π be the natural projection of $\text{vct } A_{i,\mu} + \text{vct } B_\tau$ onto $\text{vct } B_\tau$. Then

$$\begin{aligned} \sum_{h \in H_{A_{i,\mu} \cup B_\tau}} p^{\sigma(h)+m(h)s_1} &= \sum_{h \in H_{A_{i,\mu} \cup B_\tau}} p^{\sigma(\pi(h))+m(\pi(h))s_1} \\ &= \#H_{A_{i,\mu}} \sum_{b \in \pi(H_{A_{i,\mu} \cup B_\tau})} p^{\sigma(b)+m(b)s_1}, \end{aligned}$$

by Lemma 5.2(i). Consequently, $\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\text{cone}(A_{i,\mu} \cup B_\tau)}$ equals

$$\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} \frac{\#H_{A_{i,\mu}}}{\prod_{x \in A_{i,\mu}} (p^{\sigma(x)+m(x)s} - 1)} \frac{\sum_{b \in \pi(H_{A_{i,\mu} \cup B_\tau})} p^{\sigma(b)+m(b)s_1}}{\prod_{x \in B_\tau} (p^{\sigma(x)+m(x)s_1} - 1)}.$$

Using the same notation as in the proof of Theorem 5.3, we have

$$\begin{aligned} &\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\text{cone}(A_{i,\mu} \cup B_\tau)} \\ &= S(\pi(H_{A_{i,\mu} \cup B_\tau}), B_\tau, f, s_1) \lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\text{cone } A_{i,\mu}}. \end{aligned}$$

Note that the sets $A_{i,\mu}$ all generate the same vector space. So, Lemma 5.2(ii) implies that $\pi(H_{A_{i,\mu} \cup B_\tau}) = \pi(H_{A_1,\mu \cup B_\tau})$ for every $i \in \{1, \dots, \ell_\mu\}$. Therefore,

$$\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\tau(f)} = \sum_{\mu} R_\mu,$$

where μ runs through the faces of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 (that contain τ) and

$$\begin{aligned} R_\mu &= \sum_{B_\tau} \sum_{i=1}^{\ell_\mu} \lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\text{cone}(A_{i,\mu} \cup B_\tau)} \\ &= \left(\sum_{B_\tau} S(\pi(H_{A_1,\mu \cup B_\tau}), B_\tau, f, s_1) \right) \lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\mu(f)}, \end{aligned}$$

where B_τ runs through all the sets described in Assertion 6.2. Similarly, for μ a fixed face of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 , we have

$$\begin{aligned} & \lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\tau}(f_\mu) \\ &= \left(\sum_{B_\tau} S(\pi(H_{A'_{1,\mu} \cup B_\tau}), B_\tau, f_\mu, s_1) \right) \lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\mu}(f_\mu), \end{aligned}$$

because μ is the only face of $\Gamma(f_\mu)$ of maximal pure contribution to the candidate pole s_1 .

Note that $A'_{1,\mu}$ generates the same vector space as $A_{1,\mu}$. So, the sets $\pi(H_{A'_{1,\mu} \cup B_\tau})$ and $\pi(H_{A_{1,\mu} \cup B_\tau})$ are equal, by Lemma 5.2(ii). Moreover, for $b \in \pi(H_{A_{1,\mu} \cup B_\tau}) \cup B_\tau$, the $m(b)$ are also the same because $m(b) = b \cdot x_\tau$ for every $b \in \bar{\Delta}_\tau$, where x_τ is a fixed element of $\tau \subset \mu$. Now, it is clear that $S(\pi(H_{A_{1,\mu} \cup B_\tau}), B_\tau, f, s_1) = S(\pi(H_{A'_{1,\mu} \cup B_\tau}), B_\tau, f_\mu, s_1)$ and that

$$\begin{aligned} \frac{R_\mu}{\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\mu}(f)} &= \sum_{B_\tau} S(\pi(H_{A_{1,\mu} \cup B_\tau}), B_\tau, f, s_1) \\ &= \frac{\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\tau}(f_\mu)}{\lim_{s \rightarrow s_1} (p^{s-s_1} - 1)^{\kappa(s_1)} S_{\Delta_\mu}(f_\mu)}, \end{aligned}$$

which proves (6.2). □

7. ANOTHER EXPLICIT FORMULA FOR $Z_f(s, \chi)$ WHEN $f = f_\sigma$,
WHERE $\sigma \subset \Gamma(f)$ IS AN UNSTABLE CONVEX SET,
AND PROOF OF THE CONJECTURE OF DENEFF AND SARGOS

We refer to [DS92] for an explanation of the choice of the name “unstable”, which is used in Definition 7.1. In contrast to [DS92], we do not only define the notion of an unstable face of $\Gamma(f)$, but also the notion of an unstable convex set. Moreover, for the following results, it is not necessary to exclude the case where $\sigma \subset \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_j = 0\}$. Of course, the face τ_0 cannot be contained in the set $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_j = 0\}$ because τ_0 must intersect the diagonal and the origin does not belong to the Newton polyhedron.

Definition 7.1. Let f be as in Definition 1.2, and let p be a prime number. A convex set σ is called *unstable over \mathbb{F}_p relative to x_j and with respect to f* , if the following two conditions are satisfied:

- (i) $\sigma \subset \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid 0 \leq x_j \leq 1\}$ and
- (ii) for every face τ of σ contained in $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_j = 1\}$, the polynomial f_τ does not vanish on $(\mathbb{F}_p^\times)^n$.

A convex set σ is called *unstable over \mathbb{F}_p with respect to f* if there exists an index j ($1 \leq j \leq n$) such that σ is unstable over \mathbb{F}_p relative to x_j and with respect to f .

We have a similar definition for a convex set σ that is unstable over \mathbb{C} with respect to f . Then f has to be a polynomial over \mathbb{C} , and in (ii) the polynomial f_τ does not vanish on $(\mathbb{C}^\times)^n$.

In Theorem 7.3, we will give a formula for $Z_f(s, \chi)$ that holds if $f = f_\sigma$ where $\sigma \subset \Gamma(f)$ is an unstable convex set over \mathbb{F}_p with respect to f and if f is non-degenerate over \mathbb{F}_p with respect to a particular set of faces of $\Gamma(f)$. This formula

will be similar to the one in Theorem 3.2. But first, we state Lemma 7.2 and give the definition of a direction of recession.

Lemma 7.2. *Let p be a prime number and $f(x) = f(x_1, \dots, x_n) \in \mathbb{Z}_p[x_1, \dots, x_n]$. Indicate by \bar{f} the polynomial over \mathbb{F}_p obtained by reducing each coefficient of f modulo $p\mathbb{Z}_p$. Let s be a complex variable with $\text{Re}(s) > 0$, and let χ be a character of \mathbb{Z}_p^\times with conductor c_χ . Let \tilde{N} be the number of elements in the set $\{a \in (\mathbb{F}_p^\times)^{n-1} \times \mathbb{F}_p \mid \bar{f}(a) = 0\}$. Suppose that the set of congruences*

$$\begin{cases} f_\tau(x) & \equiv 0 \pmod{p}, \\ \frac{\partial f_\tau}{\partial x_i}(x) & \equiv 0 \pmod{p}, \quad i = 1, \dots, n, \end{cases}$$

has no solution in $(\mathbb{Z}_p^\times)^{n-1} \times \mathbb{Z}_p$. Then one has

$$\int_{(\mathbb{Z}_p^\times)^{n-1} \times \mathbb{Z}_p} \chi(\text{ac } f(x)) |f(x)|^s |dx| = \begin{cases} p^{-n} \left((p-1)^{n-1} p - p\tilde{N} \frac{p^s-1}{p^{s+1}-1} \right) & \text{for } \chi = \chi_{\text{triv}}, \\ p^{-nc_\chi} \sum_{\substack{a \in (\mathbb{Z}_p^\times)^{n-1} \times \mathbb{Z}_p \\ a \pmod{p^{c_\chi}} \\ f(a) \not\equiv 0 \pmod{p}}} \chi(f(a)) & \text{for } \chi \neq \chi_{\text{triv}}. \end{cases}$$

Proof. The proof is similar to the proof of [DH01, Corollary 3.2] and [Hoo02, Proposition 3.2]. □

Let τ be a face of the Newton polyhedron of a polynomial f and e_i the i -th unit vector of \mathbb{R}^n . We call e_i a *direction of recession* of τ if τ recedes in the direction of e_i , i.e.,

$$\forall x \in \tau, \forall \lambda \in \mathbb{R} \text{ with } \lambda \geq 0 : x + \lambda e_i \in \tau.$$

Theorem 7.3. *Let f be as in Definition 1.2, and let p be a prime number. Suppose that f is non-degenerate over \mathbb{F}_p with respect to all the faces of $\Gamma(f)$ contained in $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 0\}$ and suppose that $f = f_\sigma$ where $\sigma \subset \Gamma(f)$ is an unstable convex set over \mathbb{F}_p relative to x_n and with respect to f . Let s be a complex variable with $\text{Re}(s) > 0$, and let χ be a character of \mathbb{Z}_p^\times with conductor $c_\chi = 1$. Then*

$$(7.1) \quad Z_f(s, \chi) = \sum_{\substack{\tau \text{ face of } \Gamma(f) \\ e_n \in \text{rec } \tau}} \tilde{L}_\tau S_{\Delta_\tau}$$

with

$$(7.2) \quad \begin{aligned} \tilde{L}_\tau &= \begin{cases} p^{-n} \left((p-1)^{n-1} p - p\tilde{N}_\tau \frac{p^s-1}{p^{s+1}-1} \right) & \text{for } \chi = \chi_{\text{triv}}, \\ p^{-n} \sum_{a \in (\mathbb{F}_p^\times)^{n-1} \times \mathbb{F}_p} \chi(f_\tau(a)) & \text{for } \chi \neq \chi_{\text{triv}}, \end{cases} \\ \tilde{N}_\tau &= \{a \in (\mathbb{F}_p^\times)^{n-1} \times \mathbb{F}_p \mid \bar{f}_\tau(a) = 0\}, \end{aligned}$$

S_{Δ_τ} as in Theorem 3.2, $e_n = (0, 0, \dots, 1)$ and $\text{rec } \tau$ the set of directions of recession of τ . Recall that $\Gamma(f)$ is also a face of $\Gamma(f)$ with e_n as direction of recession.

Moreover, suppose that s_1 is a candidate pole of $Z_f(s, \chi)$, i.e., a number from the set described in Proposition 4.1. If $e_n \notin \text{rec } \mu$ for each face μ of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 , then the following holds: If $p^{s_1+1} \neq 1$ or

$\chi \neq \chi_{\text{triv}}$ and s_1 is a pole of $Z_f(s, \chi)$, its order is $< \kappa(s_1)$; if $p^{s_1+1} = 1$ and s_1 is a pole of $Z_f(s)$, its order is $< \kappa(s_1) + 1$.

Remarks.

- (1) Note that formula (7.1) holds if f is non-degenerate over \mathbb{F}_p with respect to all the faces of $\Gamma(f)$ contained in $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 0\}$ and if $\Gamma_{\text{gl}}(f)$ is an unstable convex set over \mathbb{F}_p relative to x_n and with respect to f .

The condition “ $f = f_\sigma$ with $\sigma \subset \Gamma(f)$ an unstable convex set over \mathbb{F}_p with respect to f ” is in fact too strong. It is already enough that there exists a convex set σ such that the following holds: for each face τ of $\Gamma(f)$ with e_n a direction of recession, the polynomial $\bar{f}_{\sigma \cap \tau \cap H_1}$ does not vanish on $(\mathbb{F}_p^\times)^n$, with $H_1 = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 1\}$. Moreover, the proof below shows that formula (7.1) for $Z_f(s, \chi)$ is even true when one has, for every face τ of $\Gamma(f)$ for which e_n is a direction of recession, that the set of congruences

$$\begin{cases} f_\tau(x) & \equiv 0 \pmod p, \\ \frac{\partial f_\tau(x)}{\partial x_i} & \equiv 0 \pmod p, \quad i = 1, \dots, n, \end{cases}$$

has no solution in $(\mathbb{Z}_p^\times)^{n-1} \times \mathbb{Z}_p$.

- (2) Let $f(x)$ be a non-zero polynomial over \mathbb{Z} with $f(0) = 0$. Suppose that f is non-degenerate over \mathbb{C} with respect to all the faces of $\Gamma(f)$ contained in $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 0\}$ and suppose that $f = f_\sigma$ where $\sigma \subset \Gamma(f)$ is an unstable convex set over \mathbb{C} relative to x_n and with respect to f . Then the conditions of Theorem 7.3 are satisfied for $p \gg 0$.

Proof. Put $H_0 = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 0\}$ and $H_1 = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 1\}$. Note that the cones associated to the faces of $\Gamma(f)$ form a partition of $(\mathbb{R}^+)^n$. Moreover, for every $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$, one has $a_n = 0$ if and only if e_n is a direction of recession of $F(a_1, \dots, a_n)$. So, if $\Delta_\tau \cap H_0 \neq \emptyset$, then $\Delta_\tau = \{a \in (\mathbb{R}^+)^n \mid F(a) = \tau\}$ is contained in H_0 . Consequently, $H_0 \cap (\mathbb{R}^+)^n$ is equal to the following disjoint union:

$$\bigcup_{\tau \text{ face of } \Gamma(f)} (\Delta_\tau \cap H_0) = \bigcup_{\substack{\tau \text{ face of } \Gamma(f) \\ \Delta_\tau \subset H_0}} \Delta_\tau = \bigcup_{\substack{\tau \text{ face of } \Gamma(f) \\ e_n \in \text{rec } \tau}} \Delta_\tau.$$

Therefore, $Z_f(s, \chi)$ equals

$$\begin{aligned} & \sum_{(k_1, \dots, k_{n-1}) \in \mathbb{N}^{n-1}} \int_{\substack{(x_1, \dots, x_n) \in \mathbb{Z}_p^n \\ \text{ord}(x_1, \dots, x_{n-1}) = (k_1, \dots, k_{n-1})}} \chi(\text{ac } f(x)) |f(x)|^s |dx| \\ &= \sum_{\substack{\tau \text{ face of } \Gamma(f) \\ e_n \in \text{rec } \tau}} \sum_{(k_1, \dots, k_{n-1}, 0) \in \mathbb{N}^n \cap \Delta_\tau} \int_{\substack{(x_1, \dots, x_n) \in \mathbb{Z}_p^n \\ \text{ord}(x_1, \dots, x_{n-1}) = (k_1, \dots, k_{n-1})}} \chi(\text{ac } f(x)) |f(x)|^s |dx|. \end{aligned}$$

Suppose that τ is a face of $\Gamma(f)$ such that e_n is a direction of recession of τ (or equivalently $\Delta_\tau \subset H_0$). Suppose that $k = (k_1, \dots, k_{n-1}, 0) \in \mathbb{N}^n \cap \Delta_\tau$ and $x = (x_1, \dots, x_n) \in \mathbb{Z}_p^n$ with $\text{ord}(x_1, \dots, x_{n-1}) = (k_1, \dots, k_{n-1})$. Write $x_j = p^{k_j} u_j$ with $u_j \in \mathbb{Z}_p^\times$, for $j = 1, \dots, n - 1$. Then

$$\begin{aligned} |dx| &= p^{-\sigma(k)} |du| |dx_n| \quad \text{and} \\ x^\omega &= x_1^{\omega_1} \cdots x_{n-1}^{\omega_{n-1}} x_n^{\omega_n} \\ &= p^{(k_1, \dots, k_{n-1}) \cdot (\omega_1, \dots, \omega_{n-1})} u_1^{\omega_1} \cdots u_{n-1}^{\omega_{n-1}} x_n^{\omega_n} \\ &= p^{k \cdot \omega} u_1^{\omega_1} \cdots u_{n-1}^{\omega_{n-1}} x_n^{\omega_n}. \end{aligned}$$

We know that $F(k) = \tau$. Thus $k \cdot \omega = m(k)$ for every $\omega \in \text{supp}(f) \cap \tau$, and $k \cdot \omega \geq m(k) + 1$ for every $\omega \in \text{supp}(f) \setminus \tau$. It follows then that

$$f(x) = p^{m(k)} \left(f_\tau(u_1, \dots, u_{n-1}, x_n) + p \tilde{f}_{\tau,k}(u_1, \dots, u_{n-1}, x_n) \right),$$

where $\tilde{f}_{\tau,k}$ is a polynomial in the variables u_1, \dots, u_{n-1}, x_n over \mathbb{Z}_p (depending on τ, k and f). Hence, $Z_f(s, \chi)$ equals

$$\begin{aligned} &\sum_{\substack{\tau \text{ face of } \Gamma(f) \\ e_n \in \text{rec } \tau}} \sum_{k \in \mathbb{N}^n \cap \Delta_\tau} p^{-\sigma(k) - m(k)s} \\ &\int_{(\mathbb{Z}_p^\times)^{n-1} \times \mathbb{Z}_p} \chi \left(\text{ac} \left(f_\tau(u_1, \dots, u_{n-1}, x_n) + p \tilde{f}_{\tau,k}(u_1, \dots, u_{n-1}, x_n) \right) \right) \\ &|f_\tau(u_1, \dots, u_{n-1}, x_n) + p \tilde{f}_{\tau,k}(u_1, \dots, u_{n-1}, x_n)|^s |du| |dx|. \end{aligned}$$

Denote the last integral by \tilde{L}_τ . By the following assertion and Lemma 7.2, we obtain equality (7.2) of Theorem 7.3.

Assertion 7.4. *For every face τ of $\Gamma(f)$ with $e_n \in \text{rec } \tau$, one has that the set of congruences*

$$\begin{cases} f_\tau(x) &\equiv 0 \pmod{p}, \\ \frac{\partial f_\tau}{\partial x_i}(x) &\equiv 0 \pmod{p}, \quad i = 1, \dots, n, \end{cases}$$

has no solution in $(\mathbb{Z}_p^\times)^{n-1} \times \mathbb{Z}_p$.

Note that \tilde{L}_τ does not depend on k . Thus,

$$(7.3) \quad Z_f(s, \chi) = \sum_{\substack{\tau \text{ face of } \Gamma(f) \\ e_n \in \text{rec } \tau}} \tilde{L}_\tau S_{\Delta_\tau},$$

with

$$S_{\Delta_\tau} = \sum_{k \in \mathbb{N}^n \cap \Delta_\tau} p^{-\sigma(k) - m(k)s}.$$

Suppose moreover that $e_n \notin \text{rec } \mu$ for each face μ of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 . Then the faces τ contained in such a μ do not occur in the sum (7.3). We know that, if s_1 is a pole of S_{Δ_τ} for τ not contained in such a μ , its order is $< \kappa(s_1)$. This is shown in [Hoo02, Proposition 4.9(i)], by choosing

a partition of Δ_τ into rational simplicial cones Δ_i such that $\langle \kappa(s_1) \rangle$ generators of Δ_i contribute to s_1 . Now, the assertion about the order follows.

We now prove Assertion 7.4. Fix a face τ of $\Gamma(f)$ with $e_n \in \text{rec } \tau$. Note that because $\sigma \subset \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid 0 \leq x_n \leq 1\}$, one has $\text{supp}(f_\tau) \subset H_0 \cup H_1$ and $f_\tau = f_{\tau \cap H_0} + f_{\tau \cap H_1}$.

Case 1. $\text{supp}(f_\tau) \cap H_1 = \emptyset$. In this case $f_\tau = f_{\tau \cap H_0}$ is independent of x_n and $\tau \cap H_0$ is a face of $\Gamma(f)$. Because f is non-degenerate with respect to all the faces of $\Gamma(f)$ contained in $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 0\}$, the assertion now follows.

Case 2. $\text{supp}(f_\tau) \cap H_1 \neq \emptyset$. In this case $(\partial f_\tau / \partial x_n) = x_n^{-1} f_{\tau \cap H_1} = x_n^{-1} f_{\sigma \cap \tau \cap H_1}$ is independent of x_n and non-zero on $(\mathbb{F}_p^\times)^{n-1}$ because $\sigma \cap \tau \cap H_1$ is a face of σ (see below) contained in H_1 and σ is unstable over \mathbb{F}_p relative to x_n and with respect to f . Now the assertion follows.

We explain that $\sigma \cap \tau \cap H_1$ is a face of σ . Suppose that H is a supporting hyperplane of $\Gamma(f)$ such that $H \cap \Gamma(f) = \tau$. Then H is also a supporting hyperplane of σ such that $H \cap \sigma = \tau \cap \sigma$ because $\sigma \subset \Gamma(f)$. Consequently, $\tau \cap \sigma$ is a face of σ . Also $\sigma \cap H_1$ is a face of σ because σ is unstable. Due to the fact that the intersection of two faces is again a face, it is now clear that $\sigma \cap \tau \cap H_1$ is a face of σ . \square

A similar result holds for $Z_{f,0}(s, \chi)$.

Theorem 7.5. *Let f be as in Definition 1.2, and let p be a prime number. Suppose that f is non-degenerate over \mathbb{F}_p with respect to all the compact faces of $\Gamma(f)$ contained in $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 0\}$ and suppose that $f = f_\sigma$ where $\sigma \subset \Gamma(f)$ is an unstable convex set over \mathbb{F}_p relative to x_n and with respect to f . (In fact one only has to require condition (ii) in Definition 7.1 for the compact faces of σ contained in $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 1\}$.) Let s be a complex variable with $\text{Re}(s) > 0$, and let χ be a character of \mathbb{Z}_p^\times with conductor $c_\chi = 1$. Then*

$$Z_{f,0}(s, \chi) = \sum_{\substack{\tau \text{ face of } \Gamma(f) \\ \{e_n\} = \text{rec } \tau \cap \{e_1, \dots, e_n\}}} L'_\tau S_{\Delta_\tau}$$

with

$$(7.4) \quad \begin{aligned} L'_\tau &= \begin{cases} p^{-n} \left((p-1)^{n-1} - p N'_\tau \frac{p^s - 1}{p^{s+1} - 1} \right) & \text{for } \chi = \chi_{\text{triv}}, \\ p^{-n} \sum_{a \in (\mathbb{F}_p^\times)^{n-1} \times \{0\}} \chi(f_\tau(a)) & \text{for } \chi \neq \chi_{\text{triv}}, \end{cases} \\ N'_\tau &= \{a \in (\mathbb{F}_p^\times)^{n-1} \times \{0\} \mid \bar{f}_\tau(a) = 0\}, \end{aligned}$$

S_{Δ_τ} as in Theorem 3.2, $e_n = (0, 0, \dots, 1)$ and $\text{rec } \tau$ the set of directions of recession of τ .

Proof. The proof is almost similar to the proof of Theorem 7.3. We only give a few important steps. It is clear that $Z_{f,0}(s, \chi)$ equals

$$\begin{aligned} & \int_{(p\mathbb{Z}_p)^n} \chi(\text{ac } f(x)) |f(x)|^s |dx| \\ &= \sum_{(k_1, \dots, k_{n-1}) \in (\mathbb{N} \setminus \{0\})^{n-1}} \int_{\substack{(x_1, \dots, x_n) \in (p\mathbb{Z}_p)^n \\ \text{ord}(x_1, \dots, x_{n-1}) = (k_1, \dots, k_{n-1})}} \chi(\text{ac } f(x)) |f(x)|^s |dx| \\ &= \sum_{\substack{\tau \text{ face of } \Gamma(f) \\ \{e_n\} = \text{rec } \tau \cap \{e_1, \dots, e_n\}}} \sum_{(k_1, \dots, k_{n-1}, 0) \in \mathbb{N}^n \cap \Delta_\tau} \int_{\substack{(x_1, \dots, x_n) \in (p\mathbb{Z}_p)^n \\ \text{ord}(x_1, \dots, x_{n-1}) = (k_1, \dots, k_{n-1})}} \chi(\text{ac } f(x)) |f(x)|^s |dx| \\ &= \sum_{\substack{\tau \text{ face of } \Gamma(f) \\ \{e_n\} = \text{rec } \tau \cap \{e_1, \dots, e_n\}}} \sum_{k \in \mathbb{N}^n \cap \Delta_\tau} p^{-\sigma(k) - m(k)s} \\ & \quad \int_{(\mathbb{Z}_p^\times)^{n-1} \times p\mathbb{Z}_p} \chi \left(\text{ac} \left(f_\tau(u_1, \dots, u_{n-1}, x_n) + p\tilde{f}_{\tau,k}(u_1, \dots, u_{n-1}, x_n) \right) \right) \\ & \quad |f_\tau(u_1, \dots, u_{n-1}, x_n) + p\tilde{f}_{\tau,k}(u_1, \dots, u_{n-1}, x_n)|^s |du| |dx|, \end{aligned}$$

where $\tilde{f}_{\tau,k}$ is as in the proof of Theorem 7.3. Denote the last integral by L'_τ . By a similar argument as in Theorem 7.3, we obtain equality (7.4) of Theorem 7.5. Note that $\tau \cap H_0$ is a compact face of $\Gamma(f)$ contained in $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 0\}$ and $\sigma \cap \tau \cap H_1$ is a compact face of σ if τ is a face of $\Gamma(f)$ with $\{e_n\} = \text{rec } \tau \cap \{e_1, \dots, e_n\}$. □

We are now able to prove one of the conjectures of Denef and Sargos [DS92, Section 6.2] for the p -adic case.

Theorem 7.6. *Suppose that the prime number p , the polynomial f and the character χ satisfy the conditions of Theorem 3.2. If τ_0 is unstable over \mathbb{F}_p with respect to f , then the following hold:*

- (i) *If $\rho \neq -1$ or $\chi \neq \chi_{\text{triv}}$ and ρ is a pole of $Z_f(s, \chi)$, then its order is $< \kappa$.*
- (ii) *If $\rho = -1$ is a pole of $Z_f(s)$, then its order is $< \kappa + 1$.*

Proof. One proves by the same argument as in the proof of Theorem 5.3 that f_{τ_0} is non-degenerate over \mathbb{F}_p with respect to all the faces of its Newton polyhedron $\Gamma(f_{\tau_0})$. Then Theorem 7.6 follows immediately by Theorem 7.3 and Theorem 5.3. Indeed, if τ_0 is unstable over \mathbb{F}_p relative to x_j , the unit vector e_j does not belong to $\text{rec } \tau$ for faces $\tau \subset \tau_0$. □

A similar property holds for $Z_{f,0}(s)$ by Theorem 7.5 and Remark (2) after Theorem 5.3. In this case f has to be non-degenerate with respect to the compact faces of $\Gamma(f)$. Note that τ_0 does not have to be compact.

By using Theorem 6.1 instead of Theorem 5.3, we can prove the following result for an arbitrary candidate pole of $Z_f(s, \chi)$.

Theorem 7.7. *Suppose that the prime number p , the polynomial f and the character χ satisfy the conditions of Theorem 3.2. Suppose that s_1 is a candidate pole of $Z_f(s, \chi)$, i.e., a number from the set described in Proposition 4.1. Suppose that for every face μ of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 , the face μ is the only face of $\Gamma(f_\mu)$ of maximal pure contribution to the candidate pole s_1 , and μ is unstable over \mathbb{F}_p with respect to f . Then the following hold:*

- (i) *If $p^{s_1+1} \neq 1$ or $\chi \neq \chi_{\text{triv}}$ and s_1 is a pole of $Z_f(s, \chi)$, then its order is $< \kappa(s_1)$.*
- (ii) *If $p^{s_1+1} = 1$ and s_1 is a pole of $Z_f(s)$, then its order is $< \kappa(s_1) + 1$.*

Remark. The proof of Theorem 6.1 shows that the condition “ μ is the only face of $\Gamma(f)$ of maximal pure contribution to the candidate pole s_1 ” in Theorem 7.7 can be replaced by a weaker condition. Examples show that even this condition might be omitted. (For example, take $f(x, y, z) = x^7y^3z + x^4y^8 + x^8y^4 + z^7$ as in the example from Remark (1) after Theorem 6.1. Then the condition on μ is not satisfied for the candidate pole $-1/3$. Nevertheless, we calculated with the program Polygusa [HL00] that $-1/3$ is not an actual pole of Igusa’s local zeta function $Z_f(s)$, e.g., for $p = 3$, $p = 5$ and $p = 31$.) We would like to search for a proof or a counterexample.

A similar property holds for $Z_{f,0}(s)$. In this case f has to be non-degenerate with respect to the compact faces of $\Gamma(f)$.

Remark. In the very special case that there is only one facet of pure contribution to s_1 that is moreover an unstable simplex, J. Denef obtained the following related result.

Theorem 7.8. *Let f be a polynomial over \mathbb{Z}_p in n variables. Suppose that f is non-degenerate over \mathbb{Q}_p with respect to the compact faces of its Newton polyhedron. Suppose that $s_1 \neq 1$ is a real candidate pole of $Z_f(s)$ such that there is only one facet μ of pure contribution to s_1 . Suppose that μ is a simplex with $n - 1$ vertices contained in a coordinate hyperplane and one vertex at distance 1 from that coordinate hyperplane. Then there is no pole of*

$$\int_{p^k \mathbb{Z}_p^n} |f(x)|^s |dx|$$

with real part equal to s_1 for $k \gg 0$.

8. EXAMPLES

We use Theorem 3.2 and Theorem 7.3 to calculate Igusa’s local zeta function in the following examples. To find the cones associated to the proper faces of the Newton polyhedron, we use Proposition 2.7. To calculate the S_{Δ_r} we divide these cones into rational simplicial cones. To find the number of the solutions of the set $\mathbb{Z}^n \cap \{\sum \lambda_i a_i \mid 0 \leq \lambda_i < 1\}$, the remark after Definition 2.10 can be useful. More examples can be found in [HL00] and [DH01]. The computer program “Polygusa” [HL00] to calculate $Z_f(s)$ for a concrete polynomial f is written in Maple and is based on the formula from Theorem 3.2.




FIGURE 8.1. Newton polyhedron of $x^2y + x^8$

Example 8.1. Let $f_1(x, y) = x^2y + x^8$. It is easy to verify that the Newton polyhedron $\Gamma(f_1)$ of f_1 is defined by the system of linear inequalities

$$\begin{cases} x \geq 2 \\ y \geq 0 \\ x + 6y \geq 8. \end{cases}$$

A picture of the Newton polyhedron of f_1 and its proper faces τ_1, \dots, τ_5 is given in Figure 8.1. If p is prime, then it is easy to verify that f_1 is non-degenerate over \mathbb{F}_p with respect to all the faces of its Newton polyhedron. So, we can use the formula for $Z_{f_1}(s)$ from Theorem 3.2. More information about the cones associated to the proper faces is contained in Table 8.1. The cone Δ_{τ_4} is rational simplicial, but the multiplicity of $(1, 0), (1, 6)$ is 6 and

$$\begin{aligned} & \mathbb{Z}^2 \cap \{\lambda_1(1, 0) + \lambda_2(1, 6) \mid 0 \leq \lambda_i < 1, i = 1, 2\} \\ &= \{(0, 0), (1, 1), (1, 2), (1, 3), (1, 4), (1, 5)\}. \end{aligned}$$

Proper face τ	Generators for Δ_τ	S_{Δ_τ}
τ_1	$(1, 0)$	$\frac{1}{p^{1+2s}-1}$
τ_2	$(1, 6)$	$\frac{1}{p^{7+8s}-1}$
τ_3	$(0, 1)$	$\frac{1}{p-1}$
τ_4	$(1, 0), (1, 6)$	$\frac{1+p^{2+3s}+p^{3+4s}+p^{4+5s}+p^{5+6s}+p^{6+7s}}{(p^{1+2s}-1)(p^{7+8s}-1)}$
τ_5	$(0, 1), (1, 6)$	$\frac{1}{(p-1)(p^{7+8s}-1)}$

TABLE 8.1. The cones associated to the faces of $\Gamma(x^2y + x^8)$

The other cones Δ_τ are all simple cones. It is easy to see that $N_{\tau_2} = N_{\Gamma(f_1)} = \{(x, y) \in (\mathbb{F}_p^\times)^2 \mid x^2y + x^8\} = p - 1$ and $N_{\tau_1} = N_{\tau_3} = N_{\tau_4} = N_{\tau_5} = 0$. According to the formula for $Z_{f_1}(s)$ in Theorem 3.2, the real candidate poles of $Z_{f_1}(s)$ are $-1, -1/2$ and $-7/8$, all with expected order 1. However, $\Gamma_{\text{gl}}(f_1)$ is unstable over

\mathbb{F}_p relative to y and with respect to f_1 . So, if we use the formula for $Z_{f_1}(s)$ from Theorem 7.3, we get

$$Z_{f_1}(s) = \tilde{L}_{\Gamma(f_1)} + \tilde{L}_{\tau_1} S_{\Delta_{\tau_1}}.$$

According to this formula, it is not surprising that $-7/8$ is not an actual pole of $Z_{f_1}(s)$. Indeed,

$$Z_{f_1}(s) = \frac{p^{3s}(p-1)^2}{(p^{1+2s}-1)(p^{1+s}-1)}.$$

Let $f_2(x, y) = x^2y + x^8 + x^{10}$ and $f_3(x, y) = x^2y + x^8 + x^{10}y^9$. If p is prime and $p \neq 2$, then it is easy to verify that f_2 is non-degenerate over \mathbb{F}_p with respect to all the faces of its Newton polyhedron. If p is prime and $p \gg 0$, then f_3 is non-degenerate over \mathbb{F}_p with respect to all the faces of its Newton polyhedron. So, we can use the formulas for $Z_{f_2}(s)$ and $Z_{f_3}(s)$ from Theorem 3.2 for those primes. The Newton polyhedra of f_1 , f_2 and f_3 are all equal, and so also are the S_{Δ_τ} . Again according to the formulas for $Z_{f_2}(s)$ and $Z_{f_3}(s)$ from Theorem 3.2, the number $-7/8$ is a real candidate pole of $Z_{f_2}(s)$ and $Z_{f_3}(s)$ with expected order 1. But by the same argument as for f_1 , it is not surprising that the real number $-7/8$ is not an actual pole of $Z_{f_2}(s)$ because $\Gamma_{\text{gl}}(f_2)$ is unstable over \mathbb{F}_p relative to y and with respect to f_2 . Indeed,

$$Z_{f_2}(s) = \frac{p^{3s}(p-1)^2}{(p^{1+2s}-1)(p^{1+s}-1)}.$$

The set $\Gamma_{\text{gl}}(f_3)$ is not unstable, but

$$(8.1) \quad \lim_{s \rightarrow -7/8} (p^{s+7/8} - 1)Z_{f_3}(s) = \lim_{s \rightarrow -7/8} (p^{s+7/8} - 1)Z_{f_1}(s),$$

because $L_{\tau_2}(f_3) = L_{\tau_2}(f_1)$, $L_{\tau_4}(f_3) = L_{\tau_4}(f_1)$, $L_{\tau_5}(f_3) = L_{\tau_5}(f_1)$. One may also obtain (8.1) by applying Theorem 6.1 to f_3 and f_1 . So, $-7/8$ is not an actual pole of $Z_{f_3}(s)$ either.


Remark. The zero-sets of the polynomials f_1 , f_2 and f_3 are normal crossing divisors. By the work of Igusa [Igu74], one can also obtain that $-1/2$ and -1 are the only possible poles. One can also easily obtain the results above by the formula of Denef [Den87].

Example 8.2. Let $f_4(x, y, z) = xy + xz^2 + yz^2$. The Newton polyhedron $\Gamma(f_4)$ of f_4 is defined by the system of linear inequalities

$$\begin{cases} x \geq 0 \\ y \geq 0 \\ z \geq 0 \\ 2x + 2y + z \geq 4 \\ 2y + z \geq 2 \\ 2x + z \geq 2 \\ x + y \geq 1. \end{cases}$$

We denote the facets with supporting hyperplanes given by the corresponding equalities in the inequalities above by τ_1, \dots, τ_7 . A picture of the Newton polyhedron of f_4 is given in Figure 8.2.

There are 19 proper faces. One easily verifies that f_4 is non-degenerate over \mathbb{F}_p with respect to all the faces of its Newton polyhedron, for every prime $p \neq 2$. So, we can use the formula for $Z_{f_4}(s)$ from Theorem 3.2 for those primes. According to this formula, the candidate real poles of $Z_{f_4}(s)$ are -1 , $-5/4$, $-3/2$ and -2 ,

FIGURE 8.2. Newton polyhedron of $xy + xz^2 + yz^2$

all with expected order 1. The set of faces of $\Gamma(f_4)$ of maximal pure contribution to the candidate poles $-5/4$, $-3/2$, -2 are $\{\tau_4\}$, $\{\tau_5, \tau_6\}$, $\{\tau_7\}$, respectively (see Definition 4.2, Definition 4.6 and Proposition 4.7). Note that this example shows that two faces of $\Gamma(f_4)$ of maximal pure contribution to the same candidate pole do not have to be disjoint. However, $-3/2$ and -2 are not actual poles of $Z_{f_4}(s)$. Indeed, by using the formula for $Z_{f_4}(s)$ from Theorem 3.2 (see [HL00]), we get

$$Z_{f_4}(s) = \frac{p^{2s}(p-1)(p^{5+3s} + p^{2+s} - p^{2+2s} - p^{1+s} + p - 1)}{(p^{1+s} - 1)(p^{5+4s} - 1)}.$$

Theorem 7.7 explains why this happens. For example, $-3/2$ is not an actual pole because the faces τ_5, τ_6 of $\Gamma(f_4)$ of maximal pure contribution to the candidate pole $-3/2$ are unstable and in addition the following property is satisfied. The face τ_5 (resp. τ_6) is the only face of $\Gamma((f_4)_{\tau_5})$ (resp. $\Gamma((f_4)_{\tau_6})$) of maximal pure contribution to $-3/2$. A similar argument holds for -2 .

REFERENCES

- [Den84] J. Denef, *The rationality of the Poincaré series associated to the p -adic points on a variety*, Invent. Math. **77** (1984), 1–23. MR **86c**:11043
- [Den87] ———, *On the degree of Igusa’s local zeta function*, Amer. J. Math. **109** (1987), 991–1008. MR **89d**:11108
- [Den91] ———, *Report on Igusa’s local zeta function*, Astérisque **201–203** (1991), 359–386, Séminaire Bourbaki 1990/1991. MR **93g**:11119
- [Den95] ———, *Poles of p -adic complex powers and Newton polyhedra*, Nieuw Arch. Wisk. **13** (1995), no. 3, 289–295. MR **96m**:11106
- [DH01] J. Denef and K. Hoornaert, *Newton polyhedra and Igusa’s local zeta function*, J. Number Theory **89** (2001), 31–64. MR **2002g**:11170
- [DLS97] J. Denef, A. Laeremans, and P. Sargos, *On the largest nontrivial pole of the distribution $|f|^s$* , RIMS Kokyuroku **999** (1997), 1–9. MR **99g**:11101
- [DS92] J. Denef and P. Sargos, *Polyèdre de Newton et distribution f_+^s . II*, Math. Ann. **293** (1992), 193–211. MR **93f**:32037
- [HL00] K. Hoornaert and D. Loots, *Polygusa: a computer program for Igusa’s local zeta function*, <http://www.wis.kuleuven.ac.be/wis/algebra/kathleen.htm>, 2000.

- [Hoo02] K. Hoornaert, *Newton polyhedra and the poles of Igusa's local zeta function*, Bull. Belg. Math. Soc. - Simon Stevin **9** (2002), 589–606.
- [Igu74] J.-I. Igusa, *Complex powers and asymptotic expansions I*, J. Reine Angew. Math **268/269** (1974), 110–130; II, *ibid*, 278/279, 307–321, 1975. MR **50**:254; MR **53**:8018
- [Igu78] ———, *Lectures on forms of higher degree*, Tata Inst. Fund. Res. Lectures on Math. and Phys., vol. 59, Springer-Verlag, Heidelberg-New York-Berlin, 1978. MR **80m**:10020
- [LM85] B. Lichtin and D. Meuser, *Poles of a local zeta function and Newton polygons*, Compositio Math. **55** (1985), 313–332. MR **87a**:11120
- [PS00] D.H. Phong and J.A. Sturm, *Algebraic estimates, stability of local zeta functions, and uniform estimates for distribution functions*, Ann. of Math. **152** (2000), 277–329. MR **2002f**:11180
- [PSS99] D.H. Phong, E.M. Stein, and J.A. Sturm, *On the growth and stability of real analytic functions*, Amer. J. Math. **121** (1999), 519–554. MR **2002a**:58025
- [Roc70] R.T. Rockafellar, *Convex analysis*, Princeton University Press, Princeton, N.J., 1970. MR **43**:445
- [Zn99] W.A. Zúñiga-Galindo, *Local zeta functions and Newton polyhedra*, preprint, 1999.

DEPARTMENT OF MATHEMATICS, CATHOLIC UNIVERSITY LEUVEN, CELESTIJNENLAAN 200B,
3001 LEUVEN, BELGIUM