

MAXIMAL HOLONOMY OF INFRA-NILMANIFOLDS WITH 2-DIMENSIONAL QUATERNIONIC HEISENBERG GEOMETRY

KU YONG HA, JONG BUM LEE, AND KYUNG BAI LEE

ABSTRACT. Let $\mathbf{H}_{4n-1}(\mathbb{H})$ be the quaternionic Heisenberg group of real dimension $4n - 1$ and let I_n denote the maximal order of the holonomy groups of all infra-nilmanifolds with $\mathbf{H}_{4n-1}(\mathbb{H})$ -geometry. We prove that $I_2 = 48$. As an application, by applying Kim and Parker's result, we obtain that the minimum volume of a 2-dimensional quaternionic hyperbolic manifold with k cusps is at least $\frac{\sqrt{2}k}{720}$.

1. INTRODUCTION

The complex Heisenberg group $\mathbf{H}_{2n-1}(\mathbb{C})$ is

$$\mathbf{H}_{2n-1}(\mathbb{C}) = \mathbb{R} \tilde{\times} \mathbb{C}^{n-1}$$

with group operation given by

$$(s, \mathbf{z})(t, \mathbf{z}') = (s + t + 2\text{Im}\{\mathbf{z}\bar{\mathbf{z}}'\}, \mathbf{z} + \mathbf{z}'),$$

where $\text{Im}(\mathbf{z}\bar{\mathbf{z}}')$ is the imaginary part of the complex number $z_1\bar{z}'_1 + z_2\bar{z}'_2 + \cdots + z_{n-1}\bar{z}'_{n-1}$ for $\mathbf{z} = (z_1, z_2, \dots, z_{n-1})$, $\mathbf{z}' = (z'_1, z'_2, \dots, z'_{n-1}) \in \mathbb{C}^{n-1}$. Then $\mathbf{H}_{2n-1}(\mathbb{C})$ is a simply connected 2-step nilpotent Lie group with the center $\mathcal{Z}(\mathbf{H}_{2n-1}(\mathbb{C})) = \mathbb{R}$. Let M be an infra-nilmanifold with $\mathbf{H}_{2n-1}(\mathbb{C})$ -geometry; that is, $M = \Pi \backslash \mathbf{H}_{2n-1}(\mathbb{C})$, where Π is a torsion free, discrete, cocompact subgroup of $\mathbf{H}_{2n-1}(\mathbb{C}) \rtimes C$ for some compact subgroup C of $\text{Aut}(\mathbf{H}_{2n-1}(\mathbb{C}))$. It is well known (see for example [4]) that Π contains a cocompact lattice of $\mathbf{H}_{2n-1}(\mathbb{C})$ with index bounded above by a universal constant I . That is, I is the maximal order of the holonomy groups. Note that the analogue in the Euclidean case is a consequence of a theorem of Bieberbach which states that there are only finitely many flat manifolds in each dimension. It is shown in [8] that when $n = 3$, $I = 24$. As a consequence ([8, Corollary]), the minimum volume of a complex hyperbolic 3-manifold M of finite volume with k cusps is $k/9$, and k is at most $-24\pi^3\chi(M)$.

Received by the editors March 23, 2003 and, in revised form, August 25, 2003.

2000 *Mathematics Subject Classification*. Primary 20H15, 20F18, 20E99, 53C55.

Key words and phrases. Almost Bieberbach group, holonomy group, quaternionic Heisenberg group, quaternionic hyperbolic manifold.

This research was supported in part by grant No. R01-1999-000-00002-0(2002) from the interdisciplinary Research program, and by grant No. R14-2002-044-01002-0(2002) from ABRL of KOSEF.

This work was done while the second-named author was visiting the Department of Mathematics at the University of Oklahoma. He expresses his sincere thanks for their hospitality.

The main concern of this paper is the quaternionic Heisenberg group

$$\begin{aligned} \mathfrak{N}_n &= \mathbf{H}_{4n-1}(\mathbb{H}) = \{(s, \mathbf{x}) \mid s \in \text{Im}(\mathbb{H}), \mathbf{x} \in \mathbb{H}^{n-1}\} \\ &\approx \mathbb{R}^3 \tilde{\times} \mathbb{H}^{n-1} \end{aligned}$$

with group operation given by

$$(s, \mathbf{p})(t, \mathbf{q}) = (s + t + 2\text{Im}\{\mathbf{q}^* \cdot \mathbf{p}\}, \mathbf{p} + \mathbf{q}),$$

for the column vectors $\mathbf{p} = (p_1, p_2, \dots, p_{n-1})^t$, $\mathbf{q} = (q_1, q_2, \dots, q_{n-1})^t \in \mathbb{H}^{n-1}$, where $\text{Im}\{\mathbf{q}^* \cdot \mathbf{p}\}$ is the imaginary part of the quaternion $\bar{q}_1 p_1 + \bar{q}_2 p_2 + \dots + \bar{q}_{n-1} p_{n-1}$ seen as an element of \mathbb{R}^3 . Then \mathfrak{N}_n is a simply connected 2-step nilpotent Lie group with the center $\mathcal{Z}(\mathfrak{N}_n) = \mathbb{R}^3$.

Let M be an infra-nilmanifold with \mathfrak{N}_n -geometry; that is, $M = \Pi \backslash \mathfrak{N}_n$, where $\Pi \subset \mathfrak{N}_n \rtimes C$ is a torsion free, discrete subgroup with compact quotient, where C is a compact subgroup of $\text{Aut}(\mathfrak{N}_n)$. Such a group Π is called an *almost Bieberbach group* (=AB-group). Since $\Pi \cap \mathcal{Z}(\mathfrak{N}_n) \cong \mathbb{Z}^3$ is a lattice of $\mathcal{Z}(\mathfrak{N}_n)$, M fits $T^3 \rightarrow M \rightarrow N$, a Seifert 3-torus “bundle” over a $4(n - 1)$ -dimensional flat orbifold. When there is no singular point, it is a genuine bundle over the base space N which is a flat Riemannian $4(n - 1)$ -manifold.

It is remarkable that the group of isometries of \mathfrak{N}_2 as the boundary of the Siegel domain model of $\mathbb{H}_{\mathbb{H}}^2$ (that is, the nilpotent factor of the Iwasawa decomposition of the isometry group $P\text{Sp}(2, 1)$ of $\mathbb{H}_{\mathbb{H}}^2$) contains a maximal subgroup of automorphisms of \mathfrak{N}_2 as a real nilpotent Lie group (forgetting quaternionic structure). Therefore, the group of isometries for the natural Hermitian metric is isomorphic to the group of isometries for any left invariant metric on \mathfrak{N}_2 .

Let I_n be the maximal order of the holonomy groups of all infra-nilmanifolds with \mathfrak{N}_n -geometry. Then the main result of this paper is the following.

Main Theorem. $I_2 = 48$.

The number I_2 is significant in its own right, but here is another application. According to a recent work of Kim and Parker [5, Corollary 5.3], it is related to the minimum volume of quaternionic hyperbolic orbifolds. More precisely, let M be an n -dimensional quaternionic hyperbolic orbifold with k cusps. Then the volume of M is

$$\text{vol}(M) \geq \frac{2^{n-1}k}{3^{n-1}(2n+1)m\sqrt{2}},$$

where m is the maximal index of a lattice in any of the subgroups of $\pi_1(M)$ stabilizing a cusp. When $n = 2$, it is shown in [5, Proposition 5.8] that $m = 576$. We note that this result is still true for manifolds, and in this case $m = I_n$. Hence we have a very sharp result for manifolds; namely, the minimum volume of a 2-dimensional quaternionic hyperbolic manifold with k cusps is at least $\frac{\sqrt{2}k}{15I_2} = \frac{\sqrt{2}k}{720}$.

We are indebted to the referee who made many invaluable suggestions, as well as corrections to some mistakes of the initial version of the paper.

2. THE GROUP OF ISOMETRIES AND AFFINE STRUCTURE OF $\mathbb{R}^3 \tilde{\times} \mathbb{R}^4$

The group

$$\mathfrak{N}_n = \{(s, \mathbf{x}) \mid s \in \text{Im}(\mathbb{H}), \mathbf{x} \in \mathbb{H}^{n-1}\}$$

is the nilpotent part of the Iwasawa decomposition KAN of $P\text{Sp}(n, 1)$. It is the boundary of the Siegel domain $\mathfrak{N}_n \times \mathbb{R}^+$. The following theorem is known (see [5]).

Theorem 2.1. *With a natural Hermitian metric on $P\text{Sp}(n, 1)$, the Heisenberg isometry group is $\text{Isom}_0(\mathfrak{N}_n) = \mathfrak{N}_n \rtimes (\text{Sp}(n - 1) \times_{\mathbb{Z}_2} \text{Sp}(1))$.*

For $n = 2$, this is $\mathfrak{N}_2 \rtimes (\text{Sp}(1) \times_{\mathbb{Z}_2} \text{Sp}(1))$. We shall explain how this acts on \mathfrak{N}_2 below.

We embed \mathbb{H} into $\mathfrak{gl}(4, \mathbb{R})$ as a division algebra as follows:

$$\zeta : \mathbb{H} \longrightarrow \mathfrak{gl}(4, \mathbb{R})$$

defined by

$$(2-1) \quad \zeta(x_1 + x_2\mathbf{i} + x_3\mathbf{j} + x_4\mathbf{k}) = \begin{bmatrix} x_1 & -x_2 & -x_3 & -x_4 \\ x_2 & x_1 & -x_4 & x_3 \\ x_3 & x_4 & x_1 & -x_2 \\ x_4 & -x_3 & x_2 & x_1 \end{bmatrix}.$$

Note that if $\mathbf{q} = x_1 + x_2\mathbf{i} + x_3\mathbf{j} + x_4\mathbf{k}$ corresponds to the first column, then the second, third and fourth columns correspond to \mathbf{qi} , \mathbf{qj} , \mathbf{qk} , respectively. Then ζ is an injective homomorphism (preserving addition and multiplication).

Observe that:

- (1) $\zeta(\overline{\mathbf{q}}) = \zeta(\mathbf{q})^t$, the transpose of $\zeta(\mathbf{q})$, and
- (2) $\det(\zeta(\mathbf{q})) = |\mathbf{q}|^4$.

Using the map $\zeta : \mathbb{H} \longrightarrow \mathfrak{gl}(4, \mathbb{R})$, we can embed the group \mathfrak{N}_2 into the affine group $\text{Aff}(\mathbb{R}^7)$ as follows: For $s = s_1\mathbf{i} + s_2\mathbf{j} + s_3\mathbf{k}$, $\mathbf{x} = x_1 + x_2\mathbf{i} + x_3\mathbf{j} + x_4\mathbf{k}$,

$$(s, \mathbf{x}) \longrightarrow \begin{bmatrix} I_3 & -2\overline{\xi}(\mathbf{x}) & S \\ 0 & I_4 & X \\ 0 & 0 & 1 \end{bmatrix} \in \text{Aff}(\mathbb{R}^7) \subset \text{GL}(8, \mathbb{R}),$$

where I_3, I_4 are identity matrices of size 3 and 4 respectively,

$$S = \begin{bmatrix} s_1 \\ s_2 \\ s_3 \end{bmatrix}, \quad X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix},$$

and $\overline{\xi}(\mathbf{x})$ is obtained by removing the first row from $\zeta(\mathbf{x})^t$ (so it is a real 3×4 matrix). By abuse of notation, we shall use s and \mathbf{x} for S and X .

Then the product in the affine group is

$$\begin{bmatrix} I_3 & -2\overline{\xi}(\mathbf{x}) & s \\ 0 & I_4 & \mathbf{x} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} I_3 & -2\overline{\xi}(\mathbf{y}) & t \\ 0 & I_4 & \mathbf{y} \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} I_3 & -2\overline{\xi}(\mathbf{x} + \mathbf{y}) & s + t - 2\overline{\xi}(\mathbf{x})\mathbf{y} \\ 0 & I_4 & \mathbf{x} + \mathbf{y} \\ 0 & 0 & 1 \end{bmatrix}.$$

Since $\overline{\xi}(\mathbf{x})\mathbf{y} = \text{Im}(\overline{\mathbf{x}}\mathbf{y}) = -\text{Im}(\overline{\mathbf{y}}\mathbf{x})$,

$$s + t - 2\overline{\xi}(\mathbf{x})\mathbf{y} = s + t + 2\text{Im}(\overline{\mathbf{y}}\mathbf{x}),$$

and the above equality shows that our map $\mathfrak{N}_2 \longrightarrow \text{Aff}(\mathbb{R}^7)$ is an embedding.

Throughout the paper, σ denotes the matrix

$$(2-2) \quad \sigma = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Note that $\sigma \in O(4) - \text{SO}(4)$.

The group of isometries of \mathfrak{N}_2 fixing the identity is

$$(2-3) \quad \text{Sp}(1) \times \text{Sp}(1) = S^3 \times S^3 \subset \mathbb{H}^* \times \mathbb{H}^* \approx \text{GL}(1, \mathbb{H}) \times \mathbb{H}^*$$

which acts on the base space of the bundle $\mathbb{R}^3 \rightarrow \mathfrak{N}_2 \rightarrow \mathbb{H}$ as left and right translations:

$$(2-4) \quad (\text{GL}(1, \mathbb{H}) \times \mathbb{H}^*) \times \mathbb{H} \longrightarrow \mathbb{H},$$

$$((c, \lambda), \mathbf{x}) \mapsto c\mathbf{x}\lambda^{-1}.$$

The action of this group on \mathfrak{N}_2 can be described as follows. The left multiplication by c and right multiplication by $d = \lambda^{-1}$ (assuming the modulus of c and d are +1) correspond to conjugation by

$$\begin{bmatrix} I_3 & 0 & 0 \\ 0 & \zeta(c) & \mathbf{0} \\ 0 & 0 & 1 \end{bmatrix} \text{ and } \begin{bmatrix} \widehat{D} & 0 & 0 \\ 0 & D & \mathbf{0} \\ 0 & 0 & 1 \end{bmatrix},$$

where

$$(2-5) \quad D = \sigma\zeta(\bar{d})\sigma = \begin{bmatrix} d_1 & -d_2 & -d_3 & -d_4 \\ d_2 & d_1 & d_4 & -d_3 \\ d_3 & -d_4 & d_1 & d_2 \\ d_4 & d_3 & -d_2 & d_1 \end{bmatrix}$$

and

$$(2-6) \quad \widehat{D} = \begin{bmatrix} d_1^2 + d_2^2 - d_3^2 - d_4^2 & 2(d_2d_3 + d_1d_4) & -2(d_1d_3 - d_2d_4) \\ 2(d_2d_3 - d_1d_4) & d_1^2 - d_2^2 + d_3^2 - d_4^2 & 2(d_1d_2 + d_3d_4) \\ 2(d_1d_3 + d_2d_4) & -2(d_1d_2 - d_3d_4) & d_1^2 - d_2^2 - d_3^2 + d_4^2 \end{bmatrix}.$$

[This matrix appears in [5] already.] Notice that $D \in \text{SO}(4)$ and $\widehat{D} \in \text{SO}(3)$. Thus,

$$\begin{bmatrix} I_3 & 0 & 0 \\ 0 & \zeta(c) & \mathbf{0} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} I & -2\bar{\xi}(\mathbf{x}) & s \\ 0 & I & \mathbf{x} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} I_3 & 0 & 0 \\ 0 & \zeta(c) & \mathbf{0} \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} I & -2\bar{\xi}(\mathbf{x})\zeta(c)^{-1} & s \\ 0 & I & \zeta(c)\mathbf{x} \\ 0 & 0 & 1 \end{bmatrix}$$

and

$$\begin{bmatrix} \widehat{D} & 0 & 0 \\ 0 & D & \mathbf{0} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} I & -2\bar{\xi}(\mathbf{x}) & s \\ 0 & I & \mathbf{x} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \widehat{D} & 0 & 0 \\ 0 & D & \mathbf{0} \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} I & -2\widehat{D}\bar{\xi}(\mathbf{x})D^{-1} & s \\ 0 & I & D\mathbf{x} \\ 0 & 0 & 1 \end{bmatrix}.$$

It is easy to see that

$$\bar{\xi}(\mathbf{x})\zeta(c)^{-1} = \bar{\xi}(\zeta(c)\mathbf{x}) \quad \text{and} \quad \widehat{D}\bar{\xi}(\mathbf{x})D^{-1} = \bar{\xi}(D\mathbf{x})$$

so that both S^3 's in the group (2-3) are automorphisms of \mathfrak{N}_2 . If $\widehat{D} = I_3$, then $D = \pm I_4$. Therefore the two S^3 's intersect at $\mathbb{Z}_2 = \{\pm I_4\}$.

3. THE GROUP OF AUTOMORPHISMS OF $\mathbb{R}^3 \tilde{\times} \mathbb{R}^4$

In this section, we shall calculate $\text{Aut}(\mathfrak{N}_2)$, the group of automorphisms of the real Lie group \mathfrak{N}_2 (no quaternionic structure). From now on, we shall use $\mathfrak{N}_2 = \mathbb{R}^3 \tilde{\times} \mathbb{R}^4$ rather than $\mathbb{R}^3 \tilde{\times} \mathbb{H}$. Using the identification

$$(3-1) \quad \rho : \mathbb{H} \longrightarrow \mathbb{R}^4,$$

$$\rho(x_1 + x_2\mathbf{i} + x_3\mathbf{j} + x_4\mathbf{k}) = [x_1, x_2, x_3, x_4]^t,$$

we identify $\mathbb{R}^3 \tilde{\times} \mathbb{H}$ with $\mathbb{R}^3 \tilde{\times} \mathbb{R}^4$ by

$$(s, \mathbf{q}) \longleftrightarrow (s, \rho(\mathbf{q})).$$

Accordingly, we introduce a new notation for $\text{Im}\{\bar{\mathbf{y}}\mathbf{x}\}$. For

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}, \quad \mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix},$$

define

$$(3-2) \quad \mathcal{I}(\mathbf{x}, \mathbf{y}) = \begin{bmatrix} - \begin{vmatrix} x_1 & y_1 \\ x_2 & y_2 \end{vmatrix} + \begin{vmatrix} x_3 & y_3 \\ x_4 & y_4 \end{vmatrix} \\ - \begin{vmatrix} x_1 & y_1 \\ x_3 & y_3 \end{vmatrix} - \begin{vmatrix} x_2 & y_2 \\ x_4 & y_4 \end{vmatrix} \\ - \begin{vmatrix} x_1 & y_1 \\ x_4 & y_4 \end{vmatrix} + \begin{vmatrix} x_2 & y_2 \\ x_3 & y_3 \end{vmatrix} \end{bmatrix}.$$

We shall represent $\mathcal{I}(\mathbf{x}, \mathbf{y})$ by matrix products as follows. Using the function $\zeta : \mathbb{H} \rightarrow \mathbb{R}^4$ in (2-1), let

$$K_1 = \zeta(-\mathbf{i}), \quad K_2 = \zeta(-\mathbf{j}), \quad K_3 = \zeta(-\mathbf{k}).$$

Then

$$K_1 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{bmatrix}, \quad K_2 = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, \quad K_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix}.$$

Let σ be the matrix in (2-2) and

$$(3-3) \quad J_i = \sigma^{-1} K_i \sigma, \quad i = 1, 2, 3,$$

so that

$$J_1 = \begin{bmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{bmatrix}, \quad J_2 = \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, \quad J_3 = \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}.$$

Then, if $\mathbf{x} = \rho(\mathbf{q})$, we have

$$\begin{aligned} J_1 \mathbf{x} &= \rho(\mathbf{qi}), & J_2 \mathbf{x} &= \rho(\mathbf{qj}), & J_3 \mathbf{x} &= \rho(\mathbf{qk}), \\ -K_1 \mathbf{x} &= \rho(\mathbf{iq}), & -K_2 \mathbf{x} &= \rho(\mathbf{jq}), & -K_3 \mathbf{x} &= \rho(\mathbf{kq}). \end{aligned}$$

The definition of \mathcal{I} in (3-2) becomes

$$(3-4) \quad \mathcal{I}(\mathbf{x}, \mathbf{y}) = (\mathbf{x}^t J_1 \mathbf{y}, \mathbf{x}^t J_2 \mathbf{y}, \mathbf{x}^t J_3 \mathbf{y})^t.$$

Clearly $\mathcal{I}(\mathbf{x}, \mathbf{y}) = -\mathcal{I}(\mathbf{y}, \mathbf{x})$ and $\mathcal{I}(\mathbf{x}, \mathbf{y})$ corresponds to $\text{Im}\{\bar{\mathbf{y}}\mathbf{x}\}$. Thus the group operation in $\mathfrak{H}_2 = \mathbb{R}^3 \tilde{\times} \mathbb{R}^4$ becomes

$$(s, \mathbf{x})(t, \mathbf{y}) = (s + t + 2\mathcal{I}(\mathbf{x}, \mathbf{y}), \mathbf{x} + \mathbf{y}).$$

Since $\mathcal{I}(\mathbf{x}, \pm\mathbf{x}) = 0$, we see easily that

$$(s, \mathbf{x})^{-1} = (-s, -\mathbf{x}).$$

Thus we have

$$[(s, \mathbf{x}), (t, \mathbf{y})] = (s, \mathbf{x})^{-1}(t, \mathbf{y})^{-1}(s, \mathbf{x})(t, \mathbf{y}) = (4\mathcal{I}(\mathbf{x}, \mathbf{y}), \mathbf{0}).$$

Clearly J_1, J_2, J_3 together with K_1, K_2, K_3 form a linear basis for the vector space $\mathfrak{so}(4, \mathbb{R})$ of the skew-symmetric matrices. We denote the two subspaces of $\mathfrak{so}(4, \mathbb{R})$ as follows:

$$\begin{aligned}\mathfrak{so}(4, \mathbb{R})_\ell &= \langle J_1, J_2, J_3 \rangle, \\ \mathfrak{so}(4, \mathbb{R})_r &= \langle K_1, K_2, K_3 \rangle.\end{aligned}$$

Since $J_i K_j = K_j J_i$ for all $i, j = 1, 2, 3$, $\mathfrak{so}(4, \mathbb{R})_\ell$ and $\mathfrak{so}(4, \mathbb{R})_r$ are ideals of $\mathfrak{so}(4, \mathbb{R})$ and

$$\mathfrak{so}(4, \mathbb{R}) = \mathfrak{so}(4, \mathbb{R})_\ell \oplus \mathfrak{so}(4, \mathbb{R})_r.$$

For any $C \in \mathrm{GL}(4, \mathbb{R})$ and $V \in \mathfrak{so}(4, \mathbb{R})$,

$$J_C(V) = C^t V C$$

defines a linear isomorphism $J_C : \mathfrak{so}(4, \mathbb{R}) \rightarrow \mathfrak{so}(4, \mathbb{R})$. In fact, with respect to the basis $\{J_1, J_2, J_3, K_1, K_2, K_3\}$, it turns out that $\det(J_C) = (\det(C))^3$.

In order to calculate the group of automorphisms of \mathfrak{N}_2 , first we define

$$O(\mathbf{J}, 4) = \{C \in \mathrm{GL}(4, \mathbb{R}) \mid C^t J_i C \in \mathfrak{so}(4, \mathbb{R})_\ell \text{ for } i = 1, 2, 3\}.$$

Then $C \in O(\mathbf{J}, 4)$ if and only if the map J_C leaves the subspace $\mathfrak{so}(4, \mathbb{R})_\ell$ spanned by J_1, J_2, J_3 invariant. Therefore,

$$(3-5) \quad C^t J_i C = \lambda_{i1} J_1 + \lambda_{i2} J_2 + \lambda_{i3} J_3, \quad \lambda_{ij} \in \mathbb{R},$$

for $i = 1, 2, 3$. Then it turns out that the matrix $\lambda = (\lambda_{ij})$ is non-singular.

Now we form the column vector

$$\mathbf{J} = \begin{bmatrix} J_1 \\ J_2 \\ J_3 \end{bmatrix}$$

with entries the matrices J_1, J_2, J_3 . With some abuse of notation, we can write

$$O(\mathbf{J}, 4) = \{C \in \mathrm{GL}(4, \mathbb{R}) \mid C^t \mathbf{J} C = \lambda \mathbf{J}, \lambda \in \mathrm{GL}(3, \mathbb{R})\}.$$

Clearly $O(\mathbf{J}, 4)$ is a closed subgroup of $\mathrm{GL}(4, \mathbb{R})$. For $C \in O(\mathbf{J}, 4)$, let $\widehat{C} \in \mathrm{GL}(3, \mathbb{R})$ denote the non-singular 3×3 matrix λ which satisfies $C^t \mathbf{J} C = \lambda \mathbf{J}$. Therefore,

$$C^t \mathbf{J} C = \widehat{C} \mathbf{J}.$$

Then $C \mapsto \widehat{C}$ defines a homomorphism $\epsilon : O(\mathbf{J}, 4) \rightarrow \mathrm{GL}(3, \mathbb{R})$ by

$$\epsilon(C) = \widehat{C}$$

so that $C^t \mathbf{J} C = \epsilon(C) \mathbf{J}$.

Lemma 3.1. (1) ϵ is a homomorphism.

(2) $C \mapsto (\frac{1}{\sqrt{|\det(C)|}} \widehat{C}, \sqrt{|\det(C)|})$ defines a homomorphism $\widehat{\cdot} : O(\mathbf{J}, 4) \rightarrow O(3) \times \mathbb{R}^+$. Moreover, $\det(\widehat{C})^2 = |\det(C)|^3$.

(3) $O(\mathbf{J}, 4)$ is closed under transpose and the above homomorphism ϵ commutes with transpose.

Proof. For $C, D \in O(\mathbf{J}, 4)$,

$$\begin{aligned} (CD)^t \mathbf{J}(CD) &= D^t (C^t \mathbf{J}C) D \\ &= D^t (\widehat{C} \mathbf{J}) D \\ &= \widehat{C} \cdot D^t \mathbf{J}D \text{ (since } \widehat{C} \text{ is a "scalar" matrix)} \\ &= \widehat{C} \widehat{D} \mathbf{J} \end{aligned}$$

shows $\epsilon(CD) = \epsilon(C)\epsilon(D)$.

Now assume that $C \in O(\mathbf{J}, 4)$ and $\det(C) = \pm 1$. Let $\widehat{C} = (\lambda_{ij})$. Then the equalities (3–5) are satisfied. A direct calculation from these equalities shows that

$$|\det(C)| = \lambda_{i1}^2 + \lambda_{i2}^2 + \lambda_{i3}^2$$

for all $i = 1, 2, 3$. Let \mathbf{c}_i be the i th column of the matrix C . Then

$$\mathcal{I}(\mathbf{c}_i, \mathbf{c}_j) = \begin{bmatrix} \mathbf{c}_i^t J_1 \mathbf{c}_j \\ \mathbf{c}_i^t J_2 \mathbf{c}_j \\ \mathbf{c}_i^t J_3 \mathbf{c}_j \end{bmatrix} = \begin{bmatrix} (C^t J_1 C)_{(i,j)} \\ (C^t J_2 C)_{(i,j)} \\ (C^t J_3 C)_{(i,j)} \end{bmatrix}.$$

From the equalities (3–5), we have

$$\begin{aligned} -(C^t J_1 C)_{(1,2)} &= \lambda_{11} = (C^t J_1 C)_{(3,4)}, \\ -(C^t J_2 C)_{(1,2)} &= \lambda_{21} = (C^t J_2 C)_{(3,4)}, \\ -(C^t J_3 C)_{(1,2)} &= \lambda_{31} = (C^t J_3 C)_{(3,4)}. \end{aligned}$$

Consequently, we have

$$-\mathcal{I}(\mathbf{c}_1, \mathbf{c}_2) = \begin{bmatrix} \lambda_{11} \\ \lambda_{21} \\ \lambda_{31} \end{bmatrix} = \mathcal{I}(\mathbf{c}_3, \mathbf{c}_4).$$

Similarly, the second and third columns of $\widehat{C} = (\lambda_{ij})$ are

$$\begin{aligned} -\mathcal{I}(\mathbf{c}_1, \mathbf{c}_3) &= -\mathcal{I}(\mathbf{c}_2, \mathbf{c}_4), \\ -\mathcal{I}(\mathbf{c}_1, \mathbf{c}_4) &= \mathcal{I}(\mathbf{c}_2, \mathbf{c}_3) \end{aligned}$$

so that

$$(3-6) \quad \widehat{C} = \begin{bmatrix} \lambda_{11} & \lambda_{12} & \lambda_{13} \\ \lambda_{21} & \lambda_{22} & \lambda_{23} \\ \lambda_{31} & \lambda_{32} & \lambda_{33} \end{bmatrix} = \begin{bmatrix} -\mathcal{I}(\mathbf{c}_1, \mathbf{c}_2) & -\mathcal{I}(\mathbf{c}_1, \mathbf{c}_3) & -\mathcal{I}(\mathbf{c}_1, \mathbf{c}_4) \end{bmatrix}.$$

For the sake of calculation, we set the second matrix to (λ'_{ij}) , so

$$(3-7) \quad \widehat{C} = \begin{bmatrix} \lambda'_{11} & \lambda'_{12} & \lambda'_{13} \\ \lambda'_{21} & \lambda'_{22} & \lambda'_{23} \\ \lambda'_{31} & \lambda'_{32} & \lambda'_{33} \end{bmatrix} = \begin{bmatrix} \mathcal{I}(\mathbf{c}_3, \mathbf{c}_4) & -\mathcal{I}(\mathbf{c}_2, \mathbf{c}_4) & \mathcal{I}(\mathbf{c}_2, \mathbf{c}_3) \end{bmatrix}.$$

Of course, then $\lambda_{ij} = \lambda'_{ij}$ for all i, j . For $j \neq k$, using the equalities (3–6) and (3–7), it can be shown that

$$\begin{aligned} &2(\lambda_{j1}\lambda_{k1} + \lambda_{j2}\lambda_{k2} + \lambda_{j3}\lambda_{k3}) \\ &= (\lambda_{j1}\lambda'_{k1} + \lambda_{j2}\lambda'_{k2} + \lambda_{j3}\lambda'_{k3}) + (\lambda'_{j1}\lambda_{k1} + \lambda'_{j2}\lambda_{k2} + \lambda'_{j3}\lambda_{k3}) \\ &= 0. \end{aligned}$$

Thus any two row vectors of \widehat{C} are orthogonal. Consequently if $C \in O(\mathbf{J}, 4)$ with $\det(C) = \pm 1$, then $\widehat{C} \in O(3)$.

Suppose $C \in O(\mathbf{J}, 4)$. Let $d = \frac{1}{\sqrt[4]{|\det(C)|}}$ and $D = dI \in \text{GL}(4, \mathbb{R})$, a diagonal matrix. Then $D \in O(\mathbf{J}, 4)$ and $dC = DC \in O(\mathbf{J}, 4)$ with $\det(dC) = \pm 1$. By the above argument, $\widehat{dC} \in O(3)$ and $\widehat{dC} = \widehat{D}\widehat{C} = d^2\widehat{C}$. Hence $\frac{1}{\sqrt{|\det(C)|}}\widehat{C} \in O(3)$. This shows that

$$\widehat{\cdot} : O(\mathbf{J}, 4) \longrightarrow O(3) \times \mathbb{R}^+, \quad C \mapsto \left(\frac{1}{\sqrt{|\det(C)|}}\widehat{C}, \sqrt{|\det(C)|} \right)$$

is a homomorphism and $\det(\widehat{C})^2 = |\det(C)|^3$.

Next suppose $C \in O(\mathbf{J}, 4)$. Let $\widehat{C} = (\lambda_{ij})$. From the equalities (3-5), we have $J_1^2 = -I_4$, $J_1J_2 = -J_2J_1$, etc. Therefore,

$$(\lambda_{i1}J_1 + \lambda_{i2}J_2 + \lambda_{i3}J_3)^2 = -|\det(C)| I_4.$$

Taking inverse matrices of the equalities (3-5), we get

$$\begin{aligned} -C^{-1}J_i(C^{-1})^t &= (C^tJ_iC)^{-1} \\ &= (\lambda_{i1}J_1 + \lambda_{i2}J_2 + \lambda_{i3}J_3)^{-1} \\ &= -\frac{1}{|\det(C)|}(\lambda_{i1}J_1 + \lambda_{i2}J_2 + \lambda_{i3}J_3). \end{aligned}$$

This shows that $(C^{-1})^t$ is also an element of $O(\mathbf{J}, 4)$ with

$$((C^{-1})^t)^\wedge = \frac{1}{|\det(C)|}\widehat{C}.$$

Suppose $\det(C) = \pm 1$. Then, from the above equality, we have

$$\begin{aligned} (C^t)^\wedge &= (C^{-1})^\wedge \\ &= (C^\wedge)^{-1} \text{ (since } \epsilon \text{ is a homomorphism)} \\ &= (C^\wedge)^t \text{ (since } \widehat{C} \in O(3)). \end{aligned}$$

One can now easily see the equality $(C^t)^\wedge = (C^\wedge)^t$ for any $C \in O(\mathbf{J}, 4)$. This proves the lemma. □

Let $S^3 \subset \mathbb{H}^*$ be the group of unit quaternions

$$S^3 = \{x_1 + x_2\mathbf{i} + x_3\mathbf{j} + x_4\mathbf{k} \mid x_1^2 + x_2^2 + x_3^2 + x_4^2 = 1\}.$$

Using $\zeta : \mathbb{H} \longrightarrow \mathfrak{gl}(4, \mathbb{R})$ of (2-1), we obtain two copies of S^3 in $O(\mathbf{J}, 4)$:

$$(3-8) \quad \begin{aligned} S_\ell^3 &= \zeta(S^3), \\ S_r^3 &= \sigma\zeta(S^3)\sigma. \end{aligned}$$

Then

Proposition 3.2. (1) ϵ has image $\epsilon(S_r^3 \times \mathbb{R}) = \text{SO}(3) \times \mathbb{R}^+$.
 (2) ϵ has kernel S_ℓ^3 so that $O(\mathbf{J}, 4) = \text{SO}(4) \times \mathbb{R}^+ = (S_\ell^3 \times_{\mathbb{Z}_2} S_r^3) \times \mathbb{R}^+$.

Proof. For $\mathbf{q} = x_1 + x_2\mathbf{i} + x_3\mathbf{j} + x_4\mathbf{k}$,

$$(3-9) \quad \begin{aligned} \zeta(\mathbf{q}) &= \begin{bmatrix} x_1 & -x_2 & -x_3 & -x_4 \\ x_2 & x_1 & -x_4 & x_3 \\ x_3 & x_4 & x_1 & -x_2 \\ x_4 & -x_3 & x_2 & x_1 \end{bmatrix} \in S_\ell^3, \\ \sigma\zeta(\bar{\mathbf{q}})\sigma &= \begin{bmatrix} x_1 & -x_2 & -x_3 & -x_4 \\ x_2 & x_1 & x_4 & -x_3 \\ x_3 & -x_4 & x_1 & x_2 \\ x_4 & x_3 & -x_2 & x_1 \end{bmatrix} \in S_r^3 \end{aligned}$$

with

$$(3-10) \quad \begin{aligned} \zeta(\mathbf{q})^\wedge &= I_3, \\ (\sigma\zeta(\bar{\mathbf{q}})\sigma)^\wedge &= \begin{bmatrix} x_1^2 + x_2^2 - x_3^2 - x_4^2 & 2(x_2x_3 + x_1x_4) & -2(x_1x_3 - x_2x_4) \\ 2(x_2x_3 - x_1x_4) & x_1^2 - x_2^2 + x_3^2 - x_4^2 & 2(x_1x_2 + x_3x_4) \\ 2(x_1x_3 + x_2x_4) & -2(x_1x_2 - x_3x_4) & x_1^2 - x_2^2 - x_3^2 + x_4^2 \end{bmatrix}. \end{aligned}$$

See the equalities (2-1), (2-5) and (2-6).

Then $S_\ell^3 \times S_r^3$ corresponds to $S^3 \times S^3 \subset \mathbb{H}^* \times \mathbb{H}^* = \text{GL}(1, \mathbb{H}) \times \mathbb{H}^*$ in (2-4) acting on \mathbb{H} , by left and right multiplications. Clearly, $S_\ell^3 \cap S_r^3 = \mathbb{Z}_2 = \{\pm I_4\}$ and

$$S_\ell^3 \times_{\mathbb{Z}_2} S_r^3 = \text{SO}(4) \subset O(\mathbf{J}, 4).$$

From the equalities (3-10), the map

$$\epsilon : O(\mathbf{J}, 4) \rightarrow O(3) \times \mathbb{R}^+$$

maps S_r^3 onto $\text{SO}(3)$ (with kernel $\mathbb{Z}_2 = \{\pm I_4\}$) and S_ℓ^3 trivially.

Next we observe that the matrix $\text{diag}\{1, 1, -1\}$ is not in the image of the homomorphism. Otherwise, there would be $C \in O(\mathbf{J}, 4)$ so that

$$C^t J_1 C = J_1, \quad C^t J_2 C = J_2, \quad C^t J_3 C = -J_3.$$

Since $J_1 J_2 = -J_3$, we would have

$$\begin{aligned} J_1 J_2 &= -J_3 = C^t J_3 C = -C^t J_1 J_2 C \\ &= -C^t J_1 C \cdot C^{-1} (C^t)^{-1} \cdot C^t J_2 C \\ &= -J_1 (C^t C)^{-1} J_2, \end{aligned}$$

and $C^t C = -I$, and hence the (1,1)-entry would be $\sum_i c_{i1}^2 = -1$, which is impossible. Thus $\text{diag}\{1, 1, -1\}$ is not in the image of ϵ . Now suppose there exists $C \in O(\mathbf{J}, 4)$ so that $\widehat{C} \in O(3) \setminus \text{SO}(3)$. Then we can write $\widehat{C} = \text{diag}\{1, 1, -1\} \widehat{E}$ for some $E \in O(\mathbf{J}, 4)$. Hence $\widehat{C} E^{-1} = \text{diag}\{1, 1, -1\}$. But the right-hand side is not in the image of the homomorphism. This proves that

$$\epsilon : O(\mathbf{J}, 4) \longrightarrow \text{SO}(3) \times \mathbb{R}^+$$

is a surjective homomorphism. (Recall, for $s \neq 0$, $sI_4 \in O(\mathbf{J}, 4)$ with $(sI)^\wedge = s^2 I_3$.)

Suppose that $C \in \ker(\epsilon)$. Then $\widehat{C} = I$ and thus $C^t J_i C = J_i$ for $i = 1, 2, 3$. Since $J_1 J_2 = -J_3$, we have

$$\begin{aligned} J_1 J_2 &= -J_3 = -C^t J_3 C = C^t J_1 J_2 C = C^t J_1 C \cdot C^{-1} (C^t)^{-1} \cdot C^t J_2 C \\ &= J_1 (C^t C)^{-1} J_2, \end{aligned}$$

$C^t C = I$ and $C \in O(4)$. On the other hand, $C^t J_1 C = J_1$ and $C^t J_2 C = J_2$ immediately induce that C is of the form $\zeta(\mathbf{q})$ for some $\mathbf{q} \in S_\ell^3$. Since we know that S_ℓ^3 is already in the kernel of ϵ , we conclude that

$$\ker(\epsilon) = S_\ell^3.$$

Thus we have obtained an exact sequence

$$1 \rightarrow S_\ell^3 \rightarrow O(\mathbf{J}, 4) \rightarrow \text{SO}(3) \times \mathbb{R}^+ \rightarrow 1$$

and equalities

$$\begin{aligned} O(\mathbf{J}, 4) &= (S_\ell^3 \times_{\mathbb{Z}_2} S_r^3) \times \mathbb{R}^+ \\ &= \text{SO}(4) \times \mathbb{R}^+. \end{aligned}$$

It is clear that S_ℓ^3 and S_r^3 and \mathbb{R}^+ commute with each other. □

Corollary 3.3. *If $C \in O(\mathbf{J}, 4)$, then*

$$\det(\widehat{C}) = \det(C)^{3/2}.$$

Proof. Follows from Lemma 3.1 and Proposition 3.2. □

Lemma 3.4. *Every matrix in $S_\ell^3 \cup S_r^3$ except $\pm I$ has eigenvalues $a + bi$, $a + bi$, $a - bi$, $a - bi$ with $b \neq 0$ (two complex double roots). Conversely, suppose $A \in \text{SO}(4)$ has eigenvalues $a \pm bi$ with $b \neq 0$ of multiplicity 2. Then either $A \in S_\ell^3$ or $A \in S_r^3$.*

Proof. Let $A \in \text{SO}(4)$ have multiple eigenvalues $a \pm bi$ with $b \neq 0$. Looking at the standard maximal torus of $\text{SO}(4)$, there is $P \in \text{SO}(4)$ such that $PAP^{-1} = \begin{bmatrix} R(\theta) & 0 \\ 0 & R(\theta') \end{bmatrix}$, where $R(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$. Since $A \in \text{SO}(4)$ has multiple eigenvalues $a \pm bi$ with $b \neq 0$, either $\theta = \theta'$ or $\theta = -\theta'$. If $PAP^{-1} = \begin{bmatrix} R(\theta) & 0 \\ 0 & R(\theta) \end{bmatrix}$, then $PAP^{-1} \in S_\ell^3$; if $PAP^{-1} = \begin{bmatrix} R(\theta) & 0 \\ 0 & R(-\theta) \end{bmatrix}$, then $PAP^{-1} \in S_r^3$. Since S_ℓ^3 and S_r^3 are normal subgroups in $\text{SO}(4)$, $A \in S_r^3$ or $A \in S_\ell^3$. □

Since the center, $\mathcal{Z}(\mathfrak{N}_2) = \mathbb{R}^3$, is a characteristic subgroup of \mathfrak{N}_2 , every automorphism of \mathfrak{N}_2 restricts to an automorphism of \mathbb{R}^3 . Consequently an automorphism of \mathfrak{N}_2 induces an automorphism on the quotient group \mathbb{R}^4 . Thus there is a natural homomorphism $\text{Aut}(\mathfrak{N}_2) \rightarrow \text{Aut}(\mathbb{R}^3) \times \text{Aut}(\mathbb{R}^4)$, $\theta \mapsto (\hat{\theta}, \bar{\theta})$.

Lemma 3.5. $\text{Image}\{\text{Aut}(\mathfrak{N}_2) \rightarrow \text{Aut}(\mathbb{R}^4)\} = O(\mathbf{J}, 4)$. *Moreover, the exact sequence $\text{Aut}(\mathfrak{N}_2) \rightarrow O(\mathbf{J}, 4) \rightarrow 1$ splits.*

Proof. Let $\theta \in \text{Aut}(\mathfrak{N}_2)$. Then $(\hat{\theta}, \bar{\theta}) \in \text{Aut}(\mathbb{R}^3) \times \text{Aut}(\mathbb{R}^4)$. Since $[(s, \mathbf{x}), (t, \mathbf{y})] = (4\mathcal{I}(\mathbf{x}, \mathbf{y}), \mathbf{0})$,

$$\theta[(s, \mathbf{x}), (t, \mathbf{y})] = \theta(4\mathcal{I}(\mathbf{x}, \mathbf{y}), \mathbf{0}) = (\hat{\theta}(4\mathcal{I}(\mathbf{x}, \mathbf{y})), \mathbf{0}) = (4\hat{\theta}(\mathcal{I}(\mathbf{x}, \mathbf{y})), \mathbf{0})$$

and

$$[\theta(s, \mathbf{x}), \theta(t, \mathbf{y})] = [(*, \bar{\theta}(\mathbf{x})), (*, \bar{\theta}(\mathbf{y}))] = (4\mathcal{I}(\bar{\theta}(\mathbf{x}), \bar{\theta}(\mathbf{y})), \mathbf{0})$$

yield

$$\mathcal{I}(\bar{\theta}(\mathbf{x}), \bar{\theta}(\mathbf{y})) = \hat{\theta}(\mathcal{I}(\mathbf{x}, \mathbf{y})),$$

or, equivalently,

$$(\bar{\theta}(\mathbf{x})^t J_1 \bar{\theta}(\mathbf{y}), \bar{\theta}(\mathbf{x})^t J_2 \bar{\theta}(\mathbf{y}), \bar{\theta}(\mathbf{x})^t J_3 \bar{\theta}(\mathbf{y}))^t = \hat{\theta} \cdot (\mathbf{x}^t J_1 \mathbf{y}, \mathbf{x}^t J_2 \mathbf{y}, \mathbf{x}^t J_3 \mathbf{y})^t$$

for all \mathbf{x}, \mathbf{y} . This happens if and only if $\bar{\theta}^t \mathbf{J} \bar{\theta} = \hat{\theta} \mathbf{J}$.

Conversely, suppose that $\bar{\theta} \in O(\mathbf{J}, 4)$, i.e., $\bar{\theta}^t \mathbf{J} \bar{\theta} = \lambda \mathbf{J}$ is satisfied for some $\lambda \in \text{GL}(3, \mathbb{R})$. We define $\theta \in \text{Aut}(\mathfrak{N}_2)$ by

$$\theta(s, \mathbf{x}) = (\lambda \cdot s, \bar{\theta}(\mathbf{x})).$$

Then

$$\begin{aligned} \theta((s, \mathbf{x}) \cdot (t, \mathbf{y})) &= \theta(s + t + 2\mathcal{I}(\mathbf{x}, \mathbf{y}), \mathbf{x} + \mathbf{y}) \\ &= (\lambda \cdot (s + t + 2\mathcal{I}(\mathbf{x}, \mathbf{y})), \bar{\theta}(\mathbf{x} + \mathbf{y})) \\ &= (\lambda \cdot s + \lambda \cdot t + \lambda \cdot 2\mathcal{I}(\mathbf{x}, \mathbf{y}), \bar{\theta}(\mathbf{x} + \mathbf{y})), \\ \theta(s, \mathbf{x}) \cdot \theta(t, \mathbf{y}) &= (\lambda \cdot s, \bar{\theta}(\mathbf{x})) \cdot (\lambda \cdot t, \bar{\theta}(\mathbf{y})) \\ &= (\lambda \cdot s + \lambda \cdot t + 2\mathcal{I}(\bar{\theta}(\mathbf{x}), \bar{\theta}(\mathbf{y})), \bar{\theta}(\mathbf{x}) + \bar{\theta}(\mathbf{y})). \end{aligned}$$

Now the condition $\bar{\theta}^t \mathbf{J} \bar{\theta} = \lambda \mathbf{J}$ guarantees that

$$\lambda \cdot \mathcal{I}(\mathbf{x}, \mathbf{y}) = \mathcal{I}(\bar{\theta}(\mathbf{x}), \bar{\theta}(\mathbf{y})).$$

Thus θ is an automorphism of \mathfrak{N}_2 . Moreover, this defines a split homomorphism $O(\mathbf{J}, 4) \rightarrow \text{Aut}(\mathfrak{N}_2)$. □

Theorem 3.6 (Structure of $\text{Aut}(\mathfrak{N}_2)$).

$$\begin{aligned} \text{Aut}(\mathfrak{N}_2) &= \text{Hom}(\mathbb{R}^4, \mathbb{R}^3) \rtimes O(\mathbf{J}, 4) \\ &= \text{Hom}(\mathbb{R}^4, \mathbb{R}^3) \rtimes ((S_\ell^3 \times_{\mathbb{Z}_2} S_r^3) \times \mathbb{R}^+), \end{aligned}$$

where an element $(\eta, A) \in \text{Hom}(\mathbb{R}^4, \mathbb{R}^3) \rtimes O(\mathbf{J}, 4)$ acts by

$$(\eta, A)(s, \mathbf{x}) = (\hat{A}s + \eta(\mathbf{x}), A\mathbf{x}).$$

Proof. Let $\theta \in \text{Aut}(\mathfrak{N}_2)$. Then we have the following commutative diagram of exact sequences:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{R}^3 & \longrightarrow & \mathfrak{N}_2 & \longrightarrow & \mathbb{R}^4 \longrightarrow 1 \\ & & \downarrow \hat{\theta} & & \downarrow \theta & & \downarrow \bar{\theta} \\ 1 & \longrightarrow & \mathbb{R}^3 & \longrightarrow & \mathfrak{N}_2 & \longrightarrow & \mathbb{R}^4 \longrightarrow 1 \end{array}$$

Thus $\theta(s, \mathbf{x}) = (\hat{\theta}(s) + \eta(s, \mathbf{x}), \bar{\theta}(\mathbf{x}))$ for $(s, \mathbf{x}) \in \mathfrak{N}_2$, where $\eta : \mathfrak{N}_2 \rightarrow \mathbb{R}^3$. Since θ is a homomorphism, one can show that η is a homomorphism, i.e.,

$$\eta((s, \mathbf{x})(t, \mathbf{y})) = \eta(s, \mathbf{x}) + \eta(t, \mathbf{y}).$$

In particular, $(\hat{\theta}(s), \mathbf{0}) = \theta(s, \mathbf{0}) = (\hat{\theta}(s) + \eta(s, \mathbf{0}), \mathbf{0})$ implies that $\eta(s, \mathbf{0}) = 0$ for all $s \in \mathbb{R}^3$, and thus $\eta(s, \mathbf{x}) = \eta((s, \mathbf{0})(0, \mathbf{x})) = \eta(s, \mathbf{0}) + \eta(0, \mathbf{x}) = \eta(0, \mathbf{x})$. Hence $\eta \in \text{Hom}(\mathbb{R}^4, \mathbb{R}^3)$.

Let us find out the kernel of the surjective homomorphism of Lemma 3.5:

$$\text{Aut}(\mathfrak{N}_2) \rightarrow O(\mathbf{J}, 4), \quad \theta \mapsto \bar{\theta}.$$

Suppose that $\theta \in \text{Aut}(\mathfrak{N}_2)$ with $\bar{\theta} = \text{id}_{\mathbb{R}^4}$. Then $\hat{\theta} = \text{id}_{\mathbb{R}^3}$ and thus $\theta(s, \mathbf{x}) = (s + \eta(\mathbf{x}), \mathbf{x})$ for some $\eta \in \text{Hom}(\mathbb{R}^4, \mathbb{R}^3)$. Conversely given $\eta \in \text{Hom}(\mathbb{R}^4, \mathbb{R}^3)$, define $\theta \in \text{Aut}(\mathfrak{N}_2)$ by $\theta(s, \mathbf{x}) = (s + \eta(\mathbf{x}), \mathbf{x})$. Clearly this θ lies in the kernel of the homomorphism. Hence we have a short exact sequence

$$1 \rightarrow \text{Hom}(\mathbb{R}^4, \mathbb{R}^3) \rightarrow \text{Aut}(\mathfrak{N}_2) \rightarrow O(\mathbf{J}, 4) \rightarrow 1.$$

By Lemma 3.5, this sequence is split. □

Note that $\text{Hom}(\mathbb{R}^4, \mathbb{R}^3) \rtimes O(\mathbf{J}, 4)$ is sitting in $\text{Hom}(\mathbb{R}^4, \mathbb{R}^3) \rtimes (\text{GL}(3, \mathbb{R}) \times O(\mathbf{J}, 4))$ as $(\eta, (\hat{A}, A))$, and the action of $O(\mathbf{J}, 4)$ on $\text{Hom}(\mathbb{R}^4, \mathbb{R}^3)$ is ${}^A\eta(\mathbf{x}) = \hat{A} \cdot \eta(A^{-1}\mathbf{x})$. The group operation on $\mathfrak{N}_2 \rtimes O(\mathbf{J}, 4)$ is given by

$$\begin{aligned} ((s, \mathbf{x}), A)((t, \mathbf{y}), B) &= ((s, \mathbf{x}) \cdot {}^A(t, \mathbf{y}), AB) \\ &= ((s, \mathbf{x}) \cdot (\hat{A}t, A\mathbf{y}), AB) \\ &= ((s + \hat{A}t + 2\mathcal{I}(\mathbf{x}, A\mathbf{y}), \mathbf{x} + A\mathbf{y}), AB). \end{aligned}$$

4. THE STRUCTURE OF AB-GROUPS FOR $\mathbb{R}^3 \tilde{\times} \mathbb{R}^4$

Let $\Pi \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ be an AB-group. Then it is well known that $\Gamma = \Pi \cap \mathfrak{N}_2$, the pure translations in Π , is the maximal normal nilpotent subgroup, and $\Phi = \Pi/\Gamma$, the holonomy group of Π , is finite. Since Γ is a lattice of \mathfrak{N}_2 , $\Gamma \cap \mathcal{Z}(\mathfrak{N}_2)$ is a lattice of $\mathcal{Z}(\mathfrak{N}_2) = \mathbb{R}^3$, and $\Gamma/\Gamma \cap \mathcal{Z}(\mathfrak{N}_2)$ is a lattice of $\mathfrak{N}_2/\mathcal{Z}(\mathfrak{N}_2) = \mathbb{R}^4$. Thus $\Gamma \cap \mathcal{Z}(\mathfrak{N}_2) \cong \mathbb{Z}^3$ and $\Gamma/\Gamma \cap \mathcal{Z}(\mathfrak{N}_2) \cong \mathbb{Z}^4$.

Consider the following natural commutative diagram:

$$\begin{array}{ccccccc} & & 1 & & 1 & & \\ & & \downarrow & & \downarrow & & \\ & & \mathbb{R}^3 & \xrightarrow{=} & \mathbb{R}^3 & & \\ & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & \mathfrak{N}_2 & \longrightarrow & \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2) & \longrightarrow & \text{Aut}(\mathfrak{N}_2) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow = \\ 1 & \longrightarrow & \mathbb{R}^4 & \longrightarrow & \mathbb{R}^4 \rtimes \text{Aut}(\mathfrak{N}_2) & \longrightarrow & \text{Aut}(\mathfrak{N}_2) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \\ & & 1 & & 1 & & \end{array}$$

Recall from Theorem 3.6 that an element $(\eta, A) \in \text{Aut}(\mathfrak{N}_2) = \text{Hom}(\mathbb{R}^4, \mathbb{R}^3) \rtimes O(\mathbf{J}, 4)$ acts on $(s, \mathbf{x}) \in \mathfrak{N}_2$ by

$$(\eta, A)(s, \mathbf{x}) = (\hat{A}s + \eta(\mathbf{x}), A\mathbf{x}).$$

Thus $O(\mathbf{J}, 4)$ acts on \mathbb{R}^3 via the homomorphism $\hat{\cdot} : O(\mathbf{J}, 4) \rightarrow \text{GL}(3, \mathbb{R})$, and $O(\mathbf{J}, 4)$ acts on \mathbb{R}^4 by matrix multiplication $O(\mathbf{J}, 4) \times \mathbb{R}^4 \rightarrow \mathbb{R}^4$.

Let $Q = \Pi/\mathbb{Z}^3$. Then the previous diagram induces the following commutative diagram:

$$\begin{array}{ccccccc}
 & & 1 & & 1 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathbb{Z}^3 & \xrightarrow{\quad} & \mathbb{Z}^3 & & \\
 & & \downarrow & & \downarrow & & \\
 (4-1) & 1 & \longrightarrow & \Gamma & \longrightarrow & \Pi & \longrightarrow & \Phi & \longrightarrow & 1 \\
 & & \downarrow & & \downarrow & & \downarrow & = & & \\
 & 1 & \longrightarrow & \mathbb{Z}^4 & \longrightarrow & Q & \longrightarrow & \Phi & \longrightarrow & 1 \\
 & & \downarrow & & \downarrow & & & & & \\
 & & 1 & & 1 & & & & &
 \end{array}$$

Here $\Phi \subset O(\mathbf{J}, 4)$ acts on \mathbb{Z}^4 by matrix multiplication, and on \mathbb{Z}^3 via the homomorphism $\hat{\cdot}: O(\mathbf{J}, 4) \rightarrow GL(3, \mathbb{R})$.

Lemma 4.1. *Let $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4$ be a basis of \mathbb{R}^4 with rational entries. Then for each $i = 1, 2, 3, 4$, the set $\{\mathcal{I}(\mathbf{v}_i, \mathbf{v}_j) \mid j = 1, \dots, \hat{i}, \dots, 4\}$ forms a lattice of \mathbb{R}^3 .*

Proof. Let V be the 4×4 -matrix whose column vectors are $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4$, and let W_i be the 3×3 -matrix whose column vectors are $\mathcal{I}(\mathbf{v}_i, \mathbf{v}_j)$ with $j = 1, \dots, \hat{i}, \dots, 4$. Then we can show that

$$\det(W_1) = \det[\mathcal{I}(\mathbf{v}_1, \mathbf{v}_2), \mathcal{I}(\mathbf{v}_1, \mathbf{v}_3), \mathcal{I}(\mathbf{v}_1, \mathbf{v}_4)] = -|\mathbf{v}_1|^2 \det[\mathbf{v}_1 \mathbf{v}_2 \mathbf{v}_3 \mathbf{v}_4].$$

Similarly, $\det(W_i) = \pm |\mathbf{v}_i|^2 \det[\mathbf{v}_1 \mathbf{v}_2 \mathbf{v}_3 \mathbf{v}_4]$. Thus if $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4$ is a basis of \mathbb{R}^4 , then the set $\{\mathcal{I}(\mathbf{v}_i, \mathbf{v}_j) \mid j = 1, \dots, \hat{i}, \dots, 4\}$ spans \mathbb{R}^3 . The map $\mathcal{I}(\mathbf{x}, \mathbf{y})$ consists of polynomial functions of the entries of \mathbf{x} and \mathbf{y} . Therefore the group generated by $\{\mathcal{I}(\mathbf{v}_i, \mathbf{v}_j) \mid j = 1, \dots, \hat{i}, \dots, 4\}$ is discrete so that it forms a lattice of \mathbb{R}^3 . \square

This lemma tells us that the lattice \mathbb{Z}^4 of \mathbb{R}^4 generated by $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ yields a lattice generated by $\{4\mathcal{I}(\mathbf{v}_i, \mathbf{v}_j) \mid j = 1, \dots, \hat{i}, \dots, 4\}$, which must be contained in \mathbb{Z}^3 in the above diagram. However, the \mathbb{Z}^3 in the diagram can be finer than the lattice generated by $\{4\mathcal{I}(\mathbf{v}_i, \mathbf{v}_j) \mid j = 1, \dots, \hat{i}, \dots, 4\}$.

Recall from [6, Proposition 2] that a virtually free abelian group $1 \rightarrow \mathbb{Z}^4 \rightarrow Q \rightarrow \Phi \rightarrow 1$ is a crystallographic group if and only if the centralizer of \mathbb{Z}^4 in Q has no torsion elements. Since Φ acts effectively on \mathbb{Z}^4 , it follows that Q is naturally a 4-dimensional crystallographic group.

The finite group Φ must be in a maximal compact subgroup $SO(4)$ of $\text{Aut}(\mathfrak{N}_2) = \text{Hom}(\mathbb{R}^4, \mathbb{R}^3) \rtimes (SO(4) \times \mathbb{R}^+)$. Note that this coincides with the compact subgroup $\text{Sp}(1) \times_{\mathbb{Z}_2} \text{Sp}(1)$ in $\text{Isom}_0(\mathfrak{N}_2) = \mathfrak{N}_2 \rtimes (\text{Sp}(1) \times_{\mathbb{Z}_2} \text{Sp}(1)) = S^3 \times_{\mathbb{Z}_2} S^3 = \mathfrak{N}_2 \rtimes SO(4)$. See Theorem 2.1.

Construction of an AB-group Π from Q . For each 4-dimensional crystallographic group Q , we shall check if there exists an AB-group Π constructed from Q ; that is, a torsion free $\Pi \subset \mathfrak{N}_2 \rtimes SO(4) \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ fitting the short exact sequence

$$1 \longrightarrow \mathbb{Z}^3 \longrightarrow \Pi \longrightarrow Q \longrightarrow 1.$$

This is the key notion for our arguments and construction. We have a complete classification of 4-dimensional crystallographic groups (Q 's in the above statement). We shall use the representations of the 4-dimensional crystallographic groups given in the book [1]. Every Q has an explicit representation $Q \longrightarrow \mathbb{R}^4 \rtimes \text{GL}(4, \mathbb{Z})$ (not into $\mathbb{R}^4 \rtimes O(4)$) in this book.

5. ALGEBRAIC CRITERIA FOR EXISTENCE OF LIFTING

Corollary 5.1. *For a 4-dimensional crystallographic group Q to have an AB-group Π , its holonomy group Φ must be in $\text{SO}(4)$. Therefore, if Φ contains a matrix of determinant -1 , there is no AB-group Π from Q .*

Proof. The holonomy group Φ is a finite subgroup of $O(\mathbf{J}, 4) = \text{SO}(4) \times \mathbb{R}^+$ (by Proposition 3.2) so that $\Phi \subset \text{SO}(4)$. □

Our goal is to determine which Q will give rise to a *torsion free* Π that fits the diagram (4-1). When Q is torsion free, then Π will be automatically torsion free, but when Q contains a torsion subgroup Q_0 , we need to check whether the lift Q_0 to Π will be torsion free.

Let Π_0 be the lift of Q_0 to Π . Thus $1 \rightarrow \mathbb{Z}^3 \rightarrow \Pi_0 \rightarrow Q_0 \rightarrow 1$ is exact. The finite subgroup $Q_0 \subset Q$ injects into the holonomy group (of Q) $\Phi \subset \text{SO}(4)$ so that we can view $Q_0 \subset \text{SO}(4)$. Recall $\text{SO}(4) = S_\ell^3 \times_{\mathbb{Z}_2} S_r^3$, and S_ℓ^3 is the kernel of the homomorphism

$$\epsilon : \text{SO}(4) \longrightarrow \text{SO}(3).$$

Let $Q_1 = Q_0 \cap S_\ell^3$, and $Q_2 = Q_0/Q_1$ so that

$$1 \rightarrow Q_1 \rightarrow Q_0 \rightarrow Q_2 \rightarrow 1$$

is exact. Consider the pullbacks of the extension $1 \rightarrow \mathbb{Z}^3 \rightarrow \Pi \rightarrow Q \rightarrow 1$ by the inclusions $Q_1 \hookrightarrow Q_0 \hookrightarrow Q$. Then we have the following commutative diagram:

$$\begin{array}{ccccccc}
 & & & 1 & & 1 & \\
 & & & \downarrow & & \downarrow & \\
 & 1 & \longrightarrow & \mathbb{Z}^3 & \longrightarrow & \Pi_1 & \longrightarrow & Q_1 & \longrightarrow & 1 \\
 & & & \downarrow = & & \downarrow & & \downarrow & & \\
 (5-1) & 1 & \longrightarrow & \mathbb{Z}^3 & \longrightarrow & \Pi_0 & \longrightarrow & Q_0 & \longrightarrow & 1 \\
 & & & \downarrow & & \downarrow & & \downarrow & & \\
 & & & & & Q_2 & \xrightarrow{=} & Q_2 & & \\
 & & & \downarrow & & \downarrow & & \downarrow & & \\
 & & & 1 & & 1 & & & &
 \end{array}$$

Since Q_1 acts trivially on \mathbb{Z}^3 , $\Pi_1 = C_{\Pi_0}(\mathbb{Z}^3)$, the centralizer of \mathbb{Z}^3 in Π_0 , is torsion free. Hence by [6, Proposition 2], $\Pi_1 = C_{\Pi_0}(\mathbb{Z}^3) \cong \mathbb{Z}^3$ and $Q_1 = \Pi_1/\mathbb{Z}^3 \subset S^3$ is abelian. But every abelian subgroup of S^3 is cyclic. Hence Q_1 is cyclic. Therefore, Q_0 is a central extension of Q_1 by Q_2 .

Since Q_2 acts effectively on \mathbb{Z}^3 , Π_0 is naturally a 3-dimensional orientable Bieberbach group, and thus Q_2 must be a holonomy group, i.e, $Q_2 \cong 1, \mathbb{Z}_2, \mathbb{Z}_3, \mathbb{Z}_4, \mathbb{Z}_6$ or $\mathbb{Z}_2 \times \mathbb{Z}_2$.

In particular, we have shown the following.

Lemma 5.2. *Any finite subgroup of Q is an extension of a cyclic group by an orientable 3-dimensional holonomy group $1, \mathbb{Z}_2, \mathbb{Z}_3, \mathbb{Z}_4, \mathbb{Z}_6$ or $\mathbb{Z}_2 \times \mathbb{Z}_2$, where the cyclic group can be conjugated into S_ℓ^3 .*

Lemma 5.3. *Q cannot contain a subgroup isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$.*

Proof. Suppose $Q \subset \mathbb{R}^4 \rtimes \text{SO}(4)$ contains a finite subgroup Q_0 isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$. Then Q_0 inject into Φ . By Lemma 5.2, Q_0 is an extension of Q_1 by Q_2 , where Q_0, Q_1, Q_2 fit into the commutative diagram (5-1), $Q_1 \cong \mathbb{Z}_2$ acts trivially on \mathbb{Z}^3 and $Q_2 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ acts effectively on \mathbb{Z}^3 . By conjugation we may assume that Π_0 is the group \mathfrak{G}_6 of [9, Theorem 3.5.5]. Thus Q_0 is a finite group acting freely on the 3-torus T so that T/Q_0 is homeomorphic to $\mathbb{R}^3/\mathfrak{G}_6$. Since Q_0 is an abelian group, by [7, Theorem 6.1] or [3, Corollary 5.3], Q_0 must be $\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_4 \times \mathbb{Z}_2$ or $\mathbb{Z}_4 \times \mathbb{Z}_4$. This contradicts the assumption that $Q_0 \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$. \square

The following is crucial in eliminating the possibility of constructing a Bieberbach group Π from a crystallographic group Q .

Proposition 5.4. *Let Q be an abstract orientable crystallographic group (not yet embedded into $\mathbb{R}^4 \rtimes \text{SO}(4)$) with holonomy group Φ . Then Φ contains unique subgroups Φ_ℓ and Φ_r (up to interchanging) with the property that*

- (1) $\Phi_\ell \cap \Phi_r \subset \{\pm I\}$.
- (2) Elements from Φ_ℓ and Φ_r commute with each other.
- (3) Suppose Q_0 in the diagram (5-1) is obtained from Q . Then either $Q_1 = Q_0 \cap \Phi_\ell$ or $Q_1 = Q_0 \cap \Phi_r$.

Proof. Recall from Lemma 3.4 that $S_\ell^3 \cup S_r^3$ is exactly the subset of $\text{SO}(4)$ consisting of elements with eigenvalues $a + bi, a + bi, a - bi, a - bi$ with $b \neq 0$, together with $\pm I$. In fact, $\pm I \in S_\ell^3 \cap S_r^3$. Assume Φ is embedded in $\text{SO}(4)$ in any way. Then let

$$\begin{aligned} \Phi_\ell &= \Phi \cap S_\ell^3, \\ \Phi_r &= \Phi \cap S_r^3. \end{aligned}$$

The choice of an embedding does not change the sets Φ_ℓ and Φ_r except interchanging, because conjugation leaves the eigenvalues unchanged (this happens if $\Phi \subset \text{SO}(4)$ is conjugated by the matrix σ in (2-2)). The last claim is clear because the kernel of $\epsilon : O(\mathbf{J}, 4) \rightarrow \text{SO}(3) \times \mathbb{R}^+$ is S_ℓ^3 . \square

Procedure 5.5 (of finding Φ_ℓ and Φ_r from Q). We assume Φ lies in $\text{GL}(4, \mathbb{R})$. Look at those elements of Φ which are not $\pm I$ (since $\pm I$ must belong to both Φ_ℓ and Φ_r). Suppose $A \in \Phi$ has the eigenvalues $a \pm bi$, with $b \neq 0$, of multiplicity 2. Then put A into Φ_ℓ . For the next $B \in \Phi$ with the right eigenvalues, we look at the product AB . If this has the right eigenvalues, then put B into Φ_ℓ . Otherwise put B into Φ_r . Proceed to the next element.

Corollary 5.6. *Suppose Q contains a non-abelian torsion subgroup Q_0 of order > 12 , with all the matrices of Q_0 lying completely inside Φ_ℓ or Φ_r . Then Q does not yield a Bieberbach group Π .*

Proof. If $Q_0 \subset \Phi_\ell$, then $Q_0 = Q_1$. But this is impossible since Q_0 is not abelian. Thus we must have $Q_0 \subset \Phi_r$. Then Q_1 is at most \mathbb{Z}_2 (or trivial) so that Q_2 has order > 6 . But the order of Q_2 must be ≤ 6 by Lemma 5.2, a contradiction. \square

Corollary 5.7. *Suppose Q contains a non-abelian torsion subgroup Q_0 of order p , and let the highest order of holonomy matrices in $\Phi_\ell \cup \Phi_r$ be r . If $p/r > 6$, then Q does not yield a Bieberbach group Π .*

Proof. Since the kernel Q_1 of the map ϵ must be cyclic coming from Φ_ℓ or Φ_r , Q_2 will have order > 6 . Now apply Lemma 5.2. \square

In what follows, the bold-faced numbers associated to the 4-dimensional crystallographic groups refer to the numbering in the book [1].

Corollary 5.8. *There is no AB-group $\Pi \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ constructed from Q given in **32/12/02/004** (holonomy group of order 64).*

Proof. Let Q_0 be the subgroups of Q

$$Q_0 = \langle t_2(a, A)(b, B)(d, D), (a, A)^7(b, B)(c, C)(d, D) \rangle,$$

a non-abelian group of order 16. By looking at the eigenvalues, we find that $\Delta = Q_0 \cap (\Phi_\ell \cup \Phi_r)$ is a non-cyclic group of order 8. Further, Δ is completely in one side of $\Phi_\ell \cup \Phi_r$ (again from the eigenvalues). Since Δ is not cyclic, it cannot be in the kernel of ϵ . Therefore, $Q_1 = \{\pm I_4\}$ and Q_2 is a group of order 8, which is impossible by Lemma 5.2. \square

Recall that any almost Bieberbach group Π for \mathfrak{N}_2 is a torsion free extension of \mathbb{Z}_3 by a 4-dimensional crystallographic group Q . There are 4783 4-dimensional crystallographic groups up to isomorphism. However, not all 4-dimensional crystallographic groups are qualified here. By Corollary 5.1, we eliminate about half of those 4-dimensional crystallographic groups. By Lemmas 5.2, 5.3, and Corollaries 5.6 and 5.7 together with Lemma 3.4, we can eliminate most of the remaining 4-dimensional crystallographic groups with holonomy order ≤ 48 , except for a few cases.

6. ELIMINATION OF SPECIAL CASES

In this section, we shall prove that there is no construction of AB-groups from the crystallographic groups of type **33/08** (and **33/12**), **33/09** and **33/10**. This requires special effort.

Given a 4-dimensional crystallographic group Q , it may happen that there is no AB-group Π constructed from Q itself, but there may be one from a conjugate of Q . But, it turns out that we need to check only two cases of different representations of Q into $E_0(4) = \mathbb{R}^4 \rtimes \text{SO}(4)$ in this special case.

First, we present an example of Q , which has two distinct (non-isomorphic) constructions depending on the representations:

Example 6.1. Two affinely conjugate representations of the same group into $E_0(4)$ as crystallographic groups can result in two non-isomorphic constructions of Π .

Consider the 4-dimensional crystallographic group Q given by

$$Q = \langle (\mathbf{e}_1, I), (\mathbf{e}_2, I), (\mathbf{e}_3, I), (\mathbf{e}_4, I), (\mathbf{0}, -K_1) \rangle.$$

This is the group given in **10/01**. Conjugation by σ (see (2-2)) sends (\mathbf{e}_1, I) to $(-\mathbf{e}_1, I)$, (\mathbf{e}_i, I) to (\mathbf{e}_i, I) for $i = 2, 3, 4$. Also it sends $-K_1$ to $-J_1$. Thus, the conjugation maps Q to

$$Q' = \langle (\mathbf{e}_1, I), (\mathbf{e}_2, I), (\mathbf{e}_3, I), (\mathbf{e}_4, I), (\mathbf{0}, -J_1) \rangle.$$

Now consider the two groups

$$\begin{aligned} II &= \langle (0, \mathbf{e}_1, I), (0, \mathbf{e}_2, I), (0, \mathbf{e}_3, I), (0, \mathbf{e}_4, I), (a, \mathbf{0}, -K_1) \rangle, \\ II' &= \langle (0, \mathbf{e}_1, I), (0, \mathbf{e}_2, I), (0, \mathbf{e}_3, I), (0, \mathbf{e}_4, I), (a, \mathbf{0}, -J_1) \rangle, \end{aligned}$$

where $a = [1, 0, 0]^t$. It is quite easy to see that the subgroup Γ generated by the 4 translations is a lattice of the Heisenberg group (of rank 7), and is the Fitting subgroup (the maximal normal nilpotent subgroup) in both II and II' . Further,

$$\Gamma \cap \mathbb{R}^3 \cong \mathbb{Z}^3$$

is generated by the 3 vectors $[4, 0, 0]^t, [0, 4, 0]^t, [0, 0, 4]^t$. Now the action of the holonomy group on this characteristic subgroup \mathbb{Z}^3 is via

$$\widehat{J}_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \quad \widehat{K}_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Therefore, it is easy to see now that both II and II' are torsion free. In fact, in both cases,

$$(a, \mathbf{0}, -J_1)^4 = ([4, 0, 0]^t, \mathbf{0}, I) = (a, \mathbf{0}, -K_1)^4.$$

Thus, they are AB-groups. However, they cannot be isomorphic because the action of the holonomy groups on the center of the Fitting subgroups Γ are distinct as shown above. Recall that Q' was conjugate to Q by an element of $O(4)$.

Proposition 6.2. *Let $Q \subset \mathbb{R}^4 \rtimes \text{SO}(4)$ be a crystallographic group with holonomy group Φ . Suppose that any symmetric matrix that commutes with every element of Φ has the form λI , where $\lambda \neq 0$. Then there is an AB-group $\Pi \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ constructed from some embedding Q into $\mathbb{R}^4 \rtimes \text{SO}(4)$ if and only if there is an AB-group Π constructed from Q or $(\mathbf{0}, \sigma)^{-1}Q(\mathbf{0}, \sigma)$.*

Proof. Suppose that there is an AB-group Π constructed from some Q' , an embedding of Q into $\mathbb{R}^4 \rtimes \text{SO}(4)$. Then, by the second Bieberbach Theorem, Q' is a conjugate to Q by an element of $\mathbb{R}^4 \rtimes \text{GL}(4, \mathbb{R})$, say, (\mathbf{x}, X) . Then

$$\begin{aligned} (\mathbf{x}, X)^{-1}Q(\mathbf{x}, X) &= Q' \subset \mathbb{R}^4 \rtimes \text{SO}(4) \\ \implies X^{-1}AX &\in \text{SO}(4) \text{ for all } A \in \Phi \\ \iff (X^{-1}AX)(X^{-1}AX)^t &= I_4 \text{ for all } A \in \Phi \\ \iff ASA^t &= S \text{ for all } A \in \Phi, \end{aligned}$$

where $S = XX^t$. By the assumption, $S = kI_4$ with $k > 0$. Hence $X \in O(4) \times \mathbb{R}^+$.

If $X \in \text{SO}(4) \times \mathbb{R}^+$, then conjugation of Π by $(0, \mathbf{x}, X)^{-1}$ is an AB-group constructed from Q because $(\mathbf{x}, X)Q'(\mathbf{x}, X)^{-1} = Q$.

If $X \notin \text{SO}(4) \times \mathbb{R}^+$, then $X = \sigma Y$ for some $Y \in \text{SO}(4) \times \mathbb{R}^+$. Then

$$(0, \sigma)(\sigma\mathbf{x}, Y)Q'(\sigma\mathbf{x}, Y)^{-1}(0, \sigma)^{-1} = (\mathbf{x}, X)Q'(\mathbf{x}, X)^{-1} = Q$$

so that

$$(\sigma\mathbf{x}, Y)Q'(\sigma\mathbf{x}, Y)^{-1} = (0, \sigma)^{-1}Q(0, \sigma).$$

Therefore $(\sigma\mathbf{x}, Y)Q'(\sigma\mathbf{x}, Y)^{-1}$ is an AB-group Π constructed from $(\mathbf{0}, \sigma)^{-1}Q(\mathbf{0}, \sigma)$. The converse is trivial. \square

Proposition 6.3. *Let $Q \subset \mathbb{R}^4 \rtimes \text{SO}(4)$ be a crystallographic group generated by*

$$(\mathbf{v}_1, I), (\mathbf{v}_2, I), (\mathbf{v}_3, I), (\mathbf{v}_4, I), (\mathbf{a}_1, A_1), (\mathbf{a}_2, A_2), \dots, (\mathbf{a}_p, A_p),$$

where the subgroup $\langle (\mathbf{v}_1, I), (\mathbf{v}_2, I), (\mathbf{v}_3, I), (\mathbf{v}_4, I) \rangle$ is the maximal normal free abelian of rank 4. Then there exists an AB-group obtained from Q if and only if the group $\Pi_{(s_1, s_2, \dots, s_p)}$ generated by

$$(0, \mathbf{v}_1, I), (0, \mathbf{v}_2, I), (0, \mathbf{v}_3, I), (0, \mathbf{v}_4, I), \\ (s_1, \mathbf{a}_1, A_1), (s_2, \mathbf{a}_2, A_2), \dots, (s_p, \mathbf{a}_p, A_p)$$

(for some $s_1, s_2, \dots, s_p \in \mathbb{R}^3$) is an AB-group.

Proof. Any AB-group $\Pi \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ obtained from Q is generated by

$$(v_1, \mathbf{0}, I), (v_2, \mathbf{0}, I), (v_3, \mathbf{0}, I), \\ (w_1, \mathbf{v}_1, I), (w_2, \mathbf{v}_2, I), (w_3, \mathbf{v}_3, I), (w_4, \mathbf{v}_4, I), \\ (t_1, \mathbf{a}_1, A_1), (t_2, \mathbf{a}_2, A_2), \dots, (t_p, \mathbf{a}_p, A_p)$$

for some $v_1, v_2, v_3, w_1, w_2, w_3, w_4, t_1, t_2, \dots, t_p \in \mathbb{R}^3$. Since the group generated by

$$(w_1, \mathbf{v}_1, I), (w_2, \mathbf{v}_2, I), (w_3, \mathbf{v}_3, I), (w_4, \mathbf{v}_4, I)$$

becomes a lattice of \mathfrak{N}_2 already (see Lemma 4.1), the group generated by

$$(w_1, \mathbf{v}_1, I), (w_2, \mathbf{v}_2, I), (w_3, \mathbf{v}_3, I), (w_4, \mathbf{v}_4, I), \\ (t_1, \mathbf{a}_1, A_1), (t_2, \mathbf{a}_2, A_2), \dots, (t_p, \mathbf{a}_p, A_p)$$

(without $(v_1, \mathbf{0}, I), (v_2, \mathbf{0}, I), (v_3, \mathbf{0}, I)$) will be a subgroup of Π of finite index, and is an AB-group. Let us call this smaller group Π' .

Let $(s, \mathbf{x}, \eta, E) \in \mathbb{R}^3 \check{\times} \mathbb{R}^4 \rtimes (\text{Hom}(\mathbb{R}^4, \mathbb{R}^3) \rtimes O(\mathbf{J}, 4)) = \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$. Then for $(0, \mathbf{v}_i) \in \mathbb{R}^3 \check{\times} \mathbb{R}^4 = \mathfrak{N}_2$,

$$(s, \mathbf{x}, \eta, E)(0, \mathbf{v}_i)(s, \mathbf{x}, \eta, E)^{-1} = (\eta(\mathbf{v}_i) + 4\mathcal{I}(\mathbf{x}, E\mathbf{v}_i), E\mathbf{v}_i).$$

Thus given $w_1, w_2, w_3, w_4 \in \mathbb{R}^3$, it is obvious that there is $\eta \in \text{Hom}(\mathbb{R}^4, \mathbb{R}^3)$ such that $\eta(\mathbf{v}_i) = w_i$ for $i = 1, 2, 3, 4$. Hence $\tilde{\eta} = (0, \mathbf{0}, \eta, I) \in \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ and $\mu(\tilde{\eta})(w_i, \mathbf{v}_i, I) = (0, \mathbf{v}_i, I)$ for all $i = 1, 2, 3, 4$. Consequently, the group Π'' obtained by conjugating Π' by $\tilde{\eta}$ is another AB-group, satisfying the condition $w_1 = \dots = w_4 = 0$. [In the course of conjugation, t_i will change.]

Consequently, we have obtained an AB-group $\Pi'' = \Pi_{(s_1, s_2, \dots, s_p)} \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ generated by

$$(0, \mathbf{v}_1, I), (0, \mathbf{v}_2, I), (0, \mathbf{v}_3, I), (0, \mathbf{v}_4, I), \\ (s_1, \mathbf{a}_1, A_1), (s_2, \mathbf{a}_2, A_2), \dots, (s_p, \mathbf{a}_p, A_p)$$

for some $s_1, s_2, \dots, s_p \in \mathbb{R}^3$. □

The proposition says that if there is an AB-group Π constructed from Q , then $\Pi_{(s_1, s_2, \dots, s_p)}$ is a subgroup of Π of finite index, which is another AB-group constructed from Q . Conversely, if $\Pi_{(s_1, s_2, \dots, s_p)}$ is an AB-group (i.e., is torsion free), then we are done. Therefore, the existence/non-existence of construction is solely determined by the group $\Pi_{(s_1, s_2, \dots, s_p)}$ described above.

Let $Q \subset \mathbb{R}^4 \rtimes \text{SO}(4)$ be a crystallographic group

$$Q = \langle (\mathbf{v}_1, I), (\mathbf{v}_2, I), (\mathbf{v}_3, I), (\mathbf{v}_4, I), (\mathbf{a}_1, A_1), (\mathbf{a}_2, A_2), \dots, (\mathbf{a}_p, A_p) \rangle \\ = \langle t_1, t_2, t_3, t_4, \bar{\alpha}_1, \bar{\alpha}_2, \dots, \bar{\alpha}_p \rangle.$$

We assume that the holonomy group $\Phi = \langle A_1, A_2, \dots, A_p \rangle$ satisfies

$$\langle A_1 \rangle \triangleleft \langle A_1, A_2 \rangle \triangleleft \dots \triangleleft \langle A_1, A_2, \dots, A_p \rangle = \Phi.$$

For $s_1, s_2, \dots, s_p \in \mathbb{R}^3$, denote

$$t_1 = (0, \mathbf{v}_1, I), \quad t_2 = (0, \mathbf{v}_2, I), \quad t_3 = (0, \mathbf{v}_3, I), \quad t_4 = (0, \mathbf{v}_4, I), \\ \alpha_1 = (s_1, \mathbf{a}_1, A_1), \quad \alpha_2 = (s_2, \mathbf{a}_2, A_1), \quad \dots, \quad \alpha_p = (s_p, \mathbf{a}_p, A_p).$$

Let Π be the group generated by $t_1, t_2, t_3, t_4, \alpha_1, \alpha_2, \dots, \alpha_p$. That is,

$$\Pi = \Pi_{(s_1, s_2, \dots, s_p)} = \langle t_1, t_2, t_3, t_4, \alpha_1, \alpha_2, \dots, \alpha_p \rangle.$$

Proposition 6.4 (Description of $\Pi \cap \mathbb{R}^3$). *Every element of Π can be written as*

$$t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha_1^{q_1} \alpha_2^{q_2} \dots \alpha_p^{q_p}$$

(not uniquely). A generating set for $Z = \Pi \cap \mathbb{R}^3$ (the pure translations of the group Π in the central direction \mathbb{R}^3) is obtained by:

- (1) Take $[t_i, t_j]$ for all $i, j = 1, \dots, 4$;
- (2) Let $\omega = \alpha_1^{q_1} \alpha_2^{q_2} \dots \alpha_p^{q_p}$. If $\omega^r = (w, V, I)$ for some $r > 0$, then write V as $V = n_1 \mathbf{v}_1 + n_2 \mathbf{v}_2 + n_3 \mathbf{v}_3 + n_4 \mathbf{v}_4$, and take $t_1^{-n_1} t_2^{-n_2} t_3^{-n_3} t_4^{-n_4} \omega^r$.

Proof. Observe that

$$\langle t_1, t_2, t_3, t_4 \rangle \triangleleft \langle t_1, t_2, t_3, t_4, \bar{\alpha}_1 \rangle \triangleleft \langle t_1, t_2, t_3, t_4, \bar{\alpha}_1, \bar{\alpha}_2 \rangle \triangleleft \dots \\ \triangleleft \langle t_1, t_2, t_3, t_4, \bar{\alpha}_1, \dots, \bar{\alpha}_{p-1} \rangle \triangleleft \langle t_1, t_2, t_3, t_4, \bar{\alpha}_1, \dots, \bar{\alpha}_{p-1}, \bar{\alpha}_p \rangle,$$

which implies that every element of Π can be written as

$$\omega = t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha_1^{q_1} \alpha_2^{q_2} \dots \alpha_p^{q_p}$$

with $n_i \in \mathbb{Z}$, and $0 \leq q_i < \text{order of } A_i$, etc.

Clearly the elements in (1) and (2) represent the identity element in the quotient group Q . Conversely, an arbitrary element of Γ (the lattice of \mathfrak{N}_2 , see the diagram (4-1)) is a product of $t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4}$ and a power of $\alpha_1^{q_1} \alpha_2^{q_2} \dots \alpha_p^{q_p}$ whose matrix part is the identity. An element of Γ lies in Z if and only if the 4-dimensional vector component is zero. It may not be true that $(0, V, I) \in \Gamma$, but we know $(0, \mathbf{v}_i, I) \in \Gamma$ for $i = 1, 2, 3, 4$. We express V in terms of \mathbf{v}_i 's, and multiply such $t_i = (0, \mathbf{v}_i, I)$'s in Γ to kill V . Thus the elements in (1) and (2) form a complete set of generators for Z . □

We now can take care of the troublesome case: Let Q be an affine 4-dimensional crystallographic group given in **33/08**, **33/09** and **33/10** (holonomy group of order 96). The following matrix P will be used to conjugate the given abstract crystallographic group Q (which is embedded into $\mathbb{R}^4 \rtimes \text{GL}(4)$) into $\mathbb{R}^4 \rtimes \text{SO}(4)$:

$$(6-1) \quad P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \end{bmatrix}.$$

That is, $(\mathbf{0}, P) \in \mathbb{R}^4 \rtimes \text{GL}(4, \mathbb{R})$ conjugates Q into a genuine crystallographic group. If S is a symmetric matrix which commutes with every element of the holonomy matrices of $(\mathbf{0}, P)Q(\mathbf{0}, P)^{-1}$, it is easy to see that $S = \lambda I$ for some $\lambda \neq 0$. Therefore the condition in Proposition 6.2 is satisfied. Denote

$$(6-2) \quad R_0 = \mu(\mathbf{0}, P)(Q), \quad R_1 = \mu(\mathbf{0}, \sigma P)(Q),$$

and denote their holonomy groups by Φ_0 and Φ_1 . We need to check only *two specific embeddings* R_0 and $R_1 = (\mathbf{0}, \sigma)R_0(\mathbf{0}, \sigma)^{-1}$.

Let $R = R_0$ or R_1 . Write

$$R = \langle (\mathbf{v}_1, I), (\mathbf{v}_2, I), (\mathbf{v}_3, I), (\mathbf{v}_4, I), (\mathbf{a}, A), (\mathbf{b}, B), (\mathbf{c}, C), (\mathbf{d}, D) \rangle,$$

and denote, for some $s, t, u, v \in \mathbb{R}^3$,

$$\begin{aligned} t_1 &= (0, \mathbf{v}_1, I), & t_2 &= (0, \mathbf{v}_2, I), & t_3 &= (0, \mathbf{v}_3, I), & t_4 &= (0, \mathbf{v}_4, I), \\ \alpha &= (s, \mathbf{a}, A), & \beta &= (t, \mathbf{b}, B), & \gamma &= (u, \mathbf{c}, C), & \delta &= (v, \mathbf{d}, D). \end{aligned}$$

By Proposition 6.3, there is no AB-group from R , if and only if for any $s, t, u, v \in \mathbb{R}^3$, the group Π generated by $t_1, t_2, t_3, t_4, \alpha, \beta, \gamma, \delta$,

$$(6-3) \quad \Pi_{(s,t,u,v)} = \langle t_1, t_2, t_3, t_4, \alpha, \beta, \gamma, \delta \rangle,$$

has a non-trivial torsion element. Now let $Z = \Pi \cap \mathbb{R}^3$. Then there is no AB-group from R if and only if the following holds: For any $s, t, u, v \in \mathbb{R}^3$, there exists a non-trivial torsion element of the form

$$z t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha^p \beta^q \gamma^r \delta^o,$$

where $z \in Z$ and $n_i, p, q, r, o, k \in \mathbb{Z}$ with $0 \leq p < \text{order of } A$, etc.

Proposition 6.5. *There is no AB-group $\Pi \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ constructed from Q given in 33/08 (holonomy group of order 96).*

Proof. We work with the group $\Pi = \Pi_{(s,t,u,v)}$ in (6-3) which is to be constructed from $R = R_0$ or R_1 . We shall show the following: For any $s, t, u, v \in \mathbb{R}^3$, there exist $z \in Z$, and $n_i, p, q, r, o, k \in \mathbb{Z}$ with $0 \leq p < 2, 0 \leq q < 4, 0 \leq r < 2, 0 \leq o < 12$ such that

$$z t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha^p \beta^q \gamma^r \delta^o$$

is a non-trivial torsion element.

Case $R = R_0$. The finite subgroup $Q_0 = \langle (b, B), (d, D)^3 \rangle$ in R_0 forms a quaternion group (non-cyclic group of order 8) with $Q_1 = Q_0$. This contradicts Lemma 5.2.

Case $R = R_1$. Our R_1 has a representation

$$R_1 = \langle (\mathbf{v}_1, I), (\mathbf{v}_2, I), (\mathbf{v}_3, I), (\mathbf{v}_4, I), (\mathbf{a}, A), (\mathbf{b}, B), (\mathbf{c}, C), (\mathbf{d}, D) \rangle,$$

where

$$\mathbf{v}_1 = \begin{bmatrix} -1 \\ 0 \\ 0 \\ -1 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \quad \mathbf{v}_4 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 2 \end{bmatrix},$$

$$\begin{aligned}
 (\mathbf{a}, A) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \right), \\
 (\mathbf{b}, B) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \right), \\
 (\mathbf{c}, C) &= \left(\frac{1}{2} \begin{bmatrix} -1 \\ 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} \right), \\
 (\mathbf{d}, D) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -1 & -1 & 1 & 1 \\ -1 & 1 & -1 & 1 \\ -1 & -1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix} \right).
 \end{aligned}$$

For $s, t, u, v \in \mathbb{R}^3$, the group $\Pi = \Pi_{(s,t,u,v)}$ as in (6-3) has a generating set

$$\begin{aligned}
 t_1 &= (0, \mathbf{v}_1, I), & t_2 &= (0, \mathbf{v}_2, I), & t_3 &= (0, \mathbf{v}_3, I), & t_4 &= (0, \mathbf{v}_4, I), \\
 \alpha &= (s, \mathbf{a}, A), & \beta &= (t, \mathbf{b}, B), & \gamma &= (u, \mathbf{c}, C), & \delta &= (v, \mathbf{d}, D).
 \end{aligned}$$

Then

$$Z = \Pi \cap \mathbb{R}^3$$

is a lattice of \mathbb{R}^3 . Let

$$z = \beta^{-4} = \left(\begin{bmatrix} 0 \\ 0 \\ -4t_3 \end{bmatrix}, \mathbf{0}, I \right).$$

Then z is an element of Z , and

$$z(\beta\delta^2)^4 = \beta^{-4}(\beta\delta^2)^4 = (0, \mathbf{0}, (BD^2)^4)$$

is a torsion element of order 3 (for any choice of s, t, u, v). □

Corollary 6.6. *There is no AB-group $\Pi \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ constructed from Q given in 33/12 (holonomy group of order 192).*

Proof. Consider the subgroup S of Q generated by the following elements:

$$(\mathbf{e}_1, I), (\mathbf{e}_2, I), (\mathbf{e}_3, I), (\mathbf{e}_4, I), (\mathbf{a}, A), (\mathbf{b}, B), (\mathbf{c}, C), (\mathbf{d}, D)$$

[we removed (e, E)]. Then S will be a 4-dimensional crystallographic group, and its holonomy group is generated by A, B, C, D . Since the holonomy group of Q has order 192, the holonomy group of S must have order 96, and is normal in Q . If we prove that there is no AB-group constructed from any 4-dimensional crystallographic group with holonomy group of order 96 (in particular, there is no AB-group constructed from our S), then we will have proved that there is no AB-group constructed from Q .

In fact we can see that our S is one in the list already: The subgroup S of Q given above is actually isomorphic to the group Q' given in 33/08 with generators

$$(\mathbf{e}_1, I), (\mathbf{e}_2, I), (\mathbf{e}_3, I), (\mathbf{e}_4, I), (\mathbf{a}', A'), (\mathbf{b}', B'), (\mathbf{c}', C'), (\mathbf{d}', D').$$

Take

$$\mathbf{v} = \begin{bmatrix} 0 \\ 0 \\ -\frac{1}{2} \\ -\frac{1}{2} \end{bmatrix}.$$

Then

$$\begin{aligned} \mu(\mathbf{v}, I)(\mathbf{a}, A) &= (\mathbf{e}_3, I)(\mathbf{a}', A'), \\ \mu(\mathbf{v}, I)(\mathbf{b}, B) &= (\mathbf{e}_3, I)(\mathbf{b}', B'), \\ \mu(\mathbf{v}, I)(\mathbf{c}, C) &= (\mathbf{e}_2, I)^{-1}(\mathbf{c}', C'), \\ \mu(\mathbf{v}, I)(\mathbf{d}, D) &= (\mathbf{e}_2, I)(\mathbf{e}_4, I)(\mathbf{b}', B')(\mathbf{d}', D')^{-1}. \end{aligned}$$

This shows that $\mu(\mathbf{v}, I) : S \rightarrow Q'$ is an isomorphism. □

Proposition 6.7. *There is no AB-group $\Pi \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ constructed from Q given in **33/10** (holonomy group of order 96).*

Proof. We proceed as in the case of **33/08**. Let Q be the group given in **33/10**. Consider two embeddings R_0 and R_1 of Q into $\mathbb{R}^4 \rtimes \text{SO}(4)$ by conjugating by $(\mathbf{0}, P)$ and $(\mathbf{0}, \sigma P)$, respectively. Let

$$R_0 = \mu(\mathbf{0}, P)(Q), \quad R_1 = \mu(\mathbf{0}, \sigma P)(Q),$$

as before.

Case $R = R_0$. Our R_0 has a representation

$$R_0 = \langle (\mathbf{v}_1, I), (\mathbf{v}_2, I), (\mathbf{v}_3, I), (\mathbf{v}_4, I), (\mathbf{a}, A), (\mathbf{b}, B), (\mathbf{c}, C), (\mathbf{d}, D) \rangle,$$

where

$$\begin{aligned} \mathbf{v}_1 &= \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \quad \mathbf{v}_4 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 2 \end{bmatrix}, \\ (\mathbf{a}, A) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 1 \end{bmatrix} \right), \\ (\mathbf{b}, B) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix} \right), \\ (\mathbf{c}, C) &= \left(\frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 & -1 & -1 & -1 \\ 1 & -1 & -1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \end{bmatrix} \right), \\ (\mathbf{d}, D) &= \left(\frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ -3 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix} \right). \end{aligned}$$

For $s, t, u, v \in \mathbb{R}^3$, the group $\Pi = \Pi_{(s,t,u,v)}$ as in (6–3) has a generating set

$$\begin{aligned} t_1 &= (0, \mathbf{v}_1, I), & t_2 &= (0, \mathbf{v}_2, I), & t_3 &= (0, \mathbf{v}_3, I), & t_4 &= (0, \mathbf{v}_4, I), \\ \alpha &= (s, \mathbf{a}, A), & \beta &= (t, \mathbf{b}, B), & \gamma &= (u, \mathbf{c}, C), & \delta &= (v, \mathbf{d}, D). \end{aligned}$$

Then

$$Z = \Pi \cap \mathbb{R}^3$$

is a lattice of \mathbb{R}^3 . Let

$$z = ([t_1, t_4]^3 (\gamma\delta)^8)^{-1} = \left(\begin{bmatrix} 0 & & \\ & 0 & \\ 76 - 8u_3 & -8v_3 & \end{bmatrix}, \mathbf{0}, I \right).$$

Then z is an element of Z , and

$$z \cdot (\gamma^2\delta^2)^4 = ([t_1, t_2]^3 (\gamma\delta)^8)^{-1} \cdot (\gamma^2\delta^2)^4$$

is a torsion element of order 3 (for any choice of s, t, u, v).

Case $R = R_1$. The finite subgroup

$$Q_0 = \langle (a, A)(b, B)^3(c, C)(d, D), t_2^{-1}t_4^{-1}(b, B)^2(c, C) \rangle$$

in R_1 forms a group of order 24 with $Q_0 = \mathbb{Z}_4$, and $Q_2 \cong \mathbb{Z}_3 \times \mathbb{Z}_2$ a non-cyclic group of order 6. This contradicts Lemma 5.2. \square

Proposition 6.8. *There is no AB-group $\Pi \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ constructed from Q given in 33/09 (holonomy group of order 96).*

Proof. Let Q be the group given in 33/09. Consider two embeddings R_0 and R_1 of Q into $\mathbb{R}^4 \rtimes \text{SO}(4)$ by conjugating by $(\mathbf{0}, P)$ and $(\mathbf{0}, \sigma P)$, respectively. Let

$$(6-4) \quad R_0 = \mu(\mathbf{0}, P)(Q), \quad R_1 = \mu(\mathbf{0}, \sigma P)(Q),$$

as before.

Case $R = R_0$. Our R_0 has a representation

$$R_1 = \langle (\mathbf{v}_1, I), (\mathbf{v}_2, I), (\mathbf{v}_3, I), (\mathbf{v}_4, I), (\mathbf{a}, A), (\mathbf{b}, B), (\mathbf{c}, C), (\mathbf{d}, D) \rangle,$$

where

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \quad \mathbf{v}_4 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 2 \end{bmatrix},$$

$$\begin{aligned}
(\mathbf{a}, A) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix} \right), \\
(\mathbf{b}, B) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix} \right), \\
(\mathbf{c}, C) &= \left(\frac{1}{2} \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -1 & -1 & 1 & -1 \\ -1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & 1 & -1 \end{bmatrix} \right), \\
(\mathbf{d}, D) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} \right).
\end{aligned}$$

For $s, t, u, v \in \mathbb{R}^3$, the group $\Pi = \Pi_{(s,t,u,v)}$ as in (6-3) has a generating set

$$\begin{aligned}
t_1 = (0, \mathbf{v}_1, I), & \quad t_2 = (0, \mathbf{v}_2, I), & \quad t_3 = (0, \mathbf{v}_3, I), & \quad t_4 = (0, \mathbf{v}_4, I), \\
\alpha = (s, \mathbf{a}, A), & \quad \beta = (t, \mathbf{b}, B), & \quad \gamma = (u, \mathbf{c}, C), & \quad \delta = (v, \mathbf{d}, D).
\end{aligned}$$

Then

$$Z = \Pi \cap \mathbb{R}^3$$

is a lattice of \mathbb{R}^3 , and one can show that Z is generated by the following 12 elements (using row notation instead of column notation):

$$\begin{aligned}
w_1 &= (-1 + s_1, 1 + s_2, s_3), \\
w_2 &= (-1 + t_1, t_2, -t_3), \\
w_3 &= (v_1 + v_2, v_1 + v_2, 0), \\
w_4 &= (-t_2, 1 - t_1, -t_3), \\
w_5 &= (-1 + t_1, t_2, t_3), \\
w_6 &= (0, 0, 2s_3), \\
w_7 &= (-s_1 - s_2, -s_1 - s_2, 0), \\
w_8 &= (s_1 - s_2, -s_1 + s_2, -2), \\
w_9 &= (1 - s_1 + s_2 + t_1 + t_2, 1 + s_1 - s_2 + t_1 + t_2, 0), \\
w_{10} &= (-s_1 + s_2, s_1 - s_2, -2), \\
w_{11} &= (s_1 - s_2 + u_1 - u_2 - v_1 + v_2, -s_1 + s_2 - u_1 + u_2 + v_1 - v_2, 0), \\
w_{12} &= (0, 0, 1 + 6u_3)
\end{aligned}$$

[This, as well as other parts of calculations, was done by the program MATHEMATICA [10] and hand-checked.]

We note that

$$\begin{aligned}
(0, 0, 4) &= -w_8 - w_{10}, \\
(4, 4, 0) &= -2w_2 + 2w_4 + w_8 + 2w_9 - w_{10}, \\
(2, 2, 2) &= -w_2 + w_4 + w_9 - w_{10}
\end{aligned}$$

are elements of Z .

Let $(2x_0, 2x_0, 0) \in Z$ be a generator of the subgroup of Z consisting of the elements of the form $(x, x, 0)$. Since $(4, 4, 0) \in Z$, x_0 is a rational number. Note

also that $w_3 = (v_1 + v_2, v_1 + v_2, 0) \in Z$. Therefore, there exist integers m and p such that

$$v_1 + v_2 = m \cdot 2x_0, \quad 4 = p \cdot 2x_0.$$

(1) Suppose $p = 2\ell$, even. Then $4 = 2\ell \cdot 2x_0$ shows $2 = \ell \cdot 2x_0$, and hence $(2, 2, 0) = \ell(2x_0, 2x_0, 0) \in Z$. Since $(2, 2, 2) \in Z$, we have $(0, 0, 2) \in Z$. Consider $z = -w_{12} + 5(0, 0, 2) \in Z$. It is easy to see that $z\gamma^6 \in \Pi$ is a torsion element of order 2.

(2) Suppose $p = 2\ell + 1$, odd. Then $4 = (2\ell + 1) \cdot 2x_0$ yields

$$\begin{aligned} (2, 2, 0) &= (2\ell + 1)(x_0, x_0, 0) \\ &= \ell(2x_0, 2x_0, 0) + (x_0, x_0, 0). \end{aligned}$$

Therefore,

$$(2, 2, 2) = \ell(2x_0, 2x_0, 0) + (x_0, x_0, 2),$$

which shows $(x_0, x_0, 2) \in Z$. Let

$$z = -m(x_0, x_0, 2) \in Z.$$

Then it is easy to see that $z\delta \in \Pi$ is a torsion element of order 2.

Case $R = R_1$. The finite subgroup $Q_0 = \langle (c, C)^4, (a, A)(d, D) \rangle$ in R_1 forms a non-commutative holonomy group of order 6. This contradicts Lemma 5.2. \square

7. A CONSTRUCTION OF AN AB-GROUP WITH HOLONOMY ORDER 48

Theorem 7.1 (Existence). *There exists an almost Bieberbach group $\Pi \subset \mathfrak{N}_2 \rtimes \text{Aut}(\mathfrak{N}_2)$ whose holonomy group has order 48.*

Proof. Consider the 4-dimensional crystallographic group Q (Case **33/05/01/003**) generated by the following elements:

$$(\mathbf{e}_1, I), \quad (\mathbf{e}_2, I), \quad (\mathbf{e}_3, I), \quad (\mathbf{e}_4, I), \quad (\mathbf{a}, A), \quad (\mathbf{b}, B), \quad (\mathbf{c}, C),$$

where I is the identity matrix, \mathbf{e}_i is the unit vector with 1 in the i th coordinate, and

$$\begin{aligned} (\mathbf{a}, A) &= \left(\frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 & -1 & -1 & 2 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & -1 & 0 & 1 \end{bmatrix} \right), \\ (\mathbf{b}, B) &= \left(\frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & -1 & 0 \\ -1 & -1 & -1 & 2 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 1 \end{bmatrix} \right), \\ (\mathbf{c}, C) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 1 & 1 & -1 \\ -1 & -1 & 0 & 1 \\ 0 & 0 & 1 & -1 \end{bmatrix} \right). \end{aligned}$$

Note that

$$A^4 = I, \quad B^2 = A^2, \quad C^6 = A^2, \quad [A, B] = A^2, \quad [A, C] = BA^2, \quad [B, C] = BA^3$$

so that Q has the holonomy group of order 48.

Now we conjugate this group by the matrix

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \end{bmatrix}.$$

Then the new group has the following representation inside $\mathbb{R}^4 \rtimes \text{SO}(4)$:

$$(\mathbf{v}_1, I), \quad (\mathbf{v}_2, I), \quad (\mathbf{v}_3, I), \quad (\mathbf{v}_4, I),$$

$$\begin{aligned} (\mathbf{a}, A) &= \left(\frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix} \right), \\ (\mathbf{b}, B) &= \left(\frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ -3 \end{bmatrix}, \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix} \right), \\ (\mathbf{c}, C) &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -1 & -1 & -1 & -1 \\ -1 & 1 & 1 & -1 \\ -1 & -1 & 1 & 1 \\ 1 & -1 & 1 & -1 \end{bmatrix} \right), \end{aligned}$$

where

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \quad \mathbf{v}_4 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 2 \end{bmatrix}.$$

Next we consider the subgroup Π of $\mathbb{R}^3 \tilde{\times} \mathbb{R}^4 \rtimes \text{SO}(4)$ generated by

$$\begin{aligned} t_1 &= (0, \mathbf{v}_1, I), \quad t_2 = (0, \mathbf{v}_2, I), \quad t_3 = (0, \mathbf{v}_3, I), \quad t_4 = (0, \mathbf{v}_4, I), \\ \alpha &= \left(\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \mathbf{a}, A \right), \quad \beta = \left(\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \mathbf{b}, B \right), \quad \gamma = \left(\begin{bmatrix} -\frac{1}{6} \\ -\frac{1}{6} \\ \frac{1}{6} \end{bmatrix}, \mathbf{c}, C \right). \end{aligned}$$

Let $Z = \Pi \cap \mathbb{R}^3$. We already have a detailed description of Z in Proposition 6.4. By calculation, we see that Z is actually generated by the vectors

$$z_1 = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}, \quad z_2 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \quad z_3 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}.$$

In particular, Z is discrete and hence Π is discrete.

Next we claim that Π is torsion free. Every element of Π can be written as

$$z t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha^p \beta^q \gamma^r,$$

where $z \in Z$, and n_1, n_2, n_3, p, q, r are integers such that $0 \leq p < 4, 0 \leq q < 4$ and $0 \leq r < 12$. If $z t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha^p \beta^q \gamma^r$ is a torsion element of order g , then $(A^p B^q C^r)^g = I$. Thus it suffices to consider all the torsion elements of Φ whose

order is a prime dividing 48. The following shows the list of prime ordered elements of Φ :

Order 2 elements: $A^2, AC^3, ABC^3, A^3C^3, A^2BC^3, A^3BC^3, BC^3$.

Order 3 elements: $A^3BC^2, A^3BC^4, AC^2, A^2C^2, AC^4, BC^2, BC^4, C^4$.

It can be shown that none of the corresponding 15 equations

$$(z_1^{\ell_1} z_2^{\ell_2} z_3^{\ell_3} t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha^2 \beta^0 \gamma^0)^2 = (0, \mathbf{0}, I),$$

$$(z_1^{\ell_1} z_2^{\ell_2} z_3^{\ell_3} t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha^1 \beta^0 \gamma^3)^2 = (0, \mathbf{0}, I),$$

$$(z_1^{\ell_1} z_2^{\ell_2} z_3^{\ell_3} t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha^1 \beta^1 \gamma^3)^2 = (0, \mathbf{0}, I),$$

...

$$(z_1^{\ell_1} z_2^{\ell_2} z_3^{\ell_3} t_1^{n_1} t_2^{n_2} t_3^{n_3} t_4^{n_4} \alpha^0 \beta^0 \gamma^4)^3 = (0, \mathbf{0}, I)$$

has any integral solutions for $\ell_1, \ell_2, \ell_3, n_1, n_2, n_3, n_4$. This proves that Π is indeed torsion free, and thus is an AB-group. \square

8. NON-EXISTENCE

Now a case-by-case study of the table from [1] shows that all 4-dimensional crystallographic groups with holonomy group of order greater than 48 satisfy at least one of the disqualification criteria proved so far. Using notations from [1], we list them below. All the calculations were done using the program MATHEMATICA [10] and hand-checked.

If the holonomy group of Q contains a matrix of determinant -1 , then Q is excluded by Corollary 5.1. Also if Q contains $(\mathbb{Z}_2)^3$, then Q is excluded by Lemma 5.3.

• 1152:

33/16: $\det(G) = -1$.

• 576:

33/14: $\det(F) = -1$,

33/15: $\langle A, B^2, F \rangle \cong (\mathbb{Z}_2)^3$.

• 384:

32/21: $\det(F) = -1$.

• 288:

30/13: $\det(D) = -1$,

33/13: $\langle BD^3, C, D^6 \rangle \cong (\mathbb{Z}_2)^3$.

• 240:

31/07: $\det(C) = -1$.

• 192:

32/18: $\det(E) = -1$,

32/19: $\det(F) = -1$,

32/20/01, 03: $\langle (a, A)^2, (b, B), (c, C) \rangle \cong (\mathbb{Z}_2)^3$,

32/20/02: $\langle (a, A)^2(b, B), (a, A)(c, C), t_2(b, B) \rangle \cong (\mathbb{Z}_2)^3$,

33/12: Corollary 6.6.

- 144:
 - 23/11, 29/09, 30/11, 30/12: $\det(C) = -1$,
 - 30/10: $\langle A, C^3, D \rangle \cong (\mathbb{Z}_2)^3$,
 - 33/11: The non-abelian group $\langle AC, B \rangle$ of order 24 lies completely inside Φ_ℓ or Φ_r . Apply Corollary 5.6.
- 128:
 - 32/17: $\det(D) = -1$.
- 120:
 - 31/04, 31/05: $\det(A) = -1$,
 - 31/06: $\langle A, B, D^5 \rangle \cong (\mathbb{Z}_2)^3$.
- 96:
 - 20/22: $\det(D) = -1$,
 - 25/11: $\det(E) = -1$,
 - 32/16/01, 32/16/03: $\langle A^2, C, D \rangle \cong (\mathbb{Z}_2)^3$,
 - 32/16/02: $\langle t_2^{-1}t_4^{-1}(d, D), t_4^{-1}(c, D), t_1t_2^{-1}(a, A)^2 \rangle \cong (\mathbb{Z}_2)^3$,
 - 33/08: Proposition 6.5,
 - 33/09: Proposition 6.8,
 - 33/10: Proposition 6.7.
- 72:
 - 22/11, 23/09, 23/10: $\det(B) = -1$,
 - 23/07, 29/06, 29/08: $\det(C) = -1$,
 - 29/07: $\det(D) = -1$,
 - 23/08, 29/05: $\langle A^3, B^3, C \rangle \cong (\mathbb{Z}_2)^3$,
 - 30/07: $\langle B, BC \rangle$ generates a non-abelian group of order 72 with the highest period of the elements in $\Phi_\ell \cup \Phi_r$ 6. Apply Corollary 5.7,
 - 30/08: $\langle AB^4, BC \rangle$ induces a non-commutative holonomy of order 6, Lemma 5.2,
 - 30/09: $\langle AB^4, C \rangle$ induces a non-commutative holonomy of order 6, Lemma 5.2,
 - 33/07: $\langle A, C^2 \rangle$ generates a non-abelian group of order 72 with the highest period of the elements in $\Phi_\ell \cup \Phi_r$ 6. Apply Corollary 5.7.
- 64:
 - 19/06 (C), 32/13 (D), 32/14 (D), 32/15 (C): $\det = -1$,
 - 32/12/01/002: $\langle t_3(a, A)^4(c, C), (b, B), (c, C) \rangle \cong (\mathbb{Z}_2)^3$,
 - 32/12/01/003: $\langle t_2^{-1}(a, A)^4(b, B), (b, B), (d, D) \rangle \cong (\mathbb{Z}_2)^3$,
 - 32/12/01/004: $\langle t_1(a, A)^6(d, D), t_3^{-1}(a, A)^2(d, D), (b, B) \rangle \cong (\mathbb{Z}_2)^3$,
 - 32/12/02/002: $\langle (a, A)^4, (b, B)(c, C), t_2^{-1}(c, C) \rangle \cong (\mathbb{Z}_2)^3$,
 - 32/12/02/003: $\langle t_3^{-1}(a, A)^2(c, C), t_2^{-1}(a, A)^6(c, C), t_2t_3^{-2}(a, A)^6(b, B)(c, C) \rangle \cong (\mathbb{Z}_2)^3$,
 - 32/12/02/004: Proposition 5.8,
 - 32/12/02/005: $\langle t_2(a, A)^6(d, D), t_4^{-1}(a, A)^2(d, D), t_2(b, B) \rangle \cong (\mathbb{Z}_2)^3$,
 - 32/12/02/006: $\langle t_1^{-1}t_2^2(a, A)^6(d, D), t_2t_3(a, A)^4(b, B), t_2(b, B) \rangle \cong (\mathbb{Z}_2)^3$.

- 60:
 - 31/03/01: $\langle t_1^{-1}(d, D)^3, t_1^{-1}t_2^{-1}(b, B)(b, B) \rangle$
induces a non-commutative holonomy of order 10, Lemma 5.2,
 - 31/03/02: $\langle (b, B), (d, D) \rangle$
induces a non-commutative holonomy of order 10, Lemma 5.2.
- 48:
 - 33/05/01/003: CONSTRUCTION! (Theorem 7.1).

REFERENCES

- [1] H. Brown, R. Bülow, J. Neubüser, H. Wondratschek and H. Zassenhaus, Crystallographic Groups of Four-Dimensional Spaces, John Wiley & Sons, Inc., New York, 1978. MR58:4109
- [2] J. Cygan, A tangential convergence for bounded harmonic functions on a rank one symmetric space, *Trans. Amer. Math. Soc.*, **265** (1981), 405–418. MR83h:43010
- [3] K. Y. Ha, J. H. Jo, S. W. Kim and J. B. Lee, Classification of free actions of finite groups on the 3-torus, *Topology Appl.*, **121** (2002), 469–507. MR2003j:57056
- [4] S. Hersonsky and F. Paulin, On the Volumes of Complex Hyperbolic Manifolds, *Duke Math. J.*, **84** (1996), 719–737. MR97h:32036
- [5] I. Kim and J. R. Parker, Geometry of quaternionic hyperbolic manifolds, *Math. Proc. Cambridge Philos. Soc.* **135** (2003), 291–310.
- [6] K. B. Lee and F. Raymond, Topological, affine and isometric actions on flat Riemannian manifolds, *J. Differential Geom.*, **16** (1981), 255–269. MR84k:57027
- [7] K. B. Lee, J. Shin and S. Yokura, Free actions of finite abelian groups on the 3-torus, *Topology Appl.*, **53** (1993), 153–175. MR94j:57016
- [8] K. B. Lee and A. Szczepański, Maximal holonomy of almost Bieberbach groups for \mathbf{Heis}_5 , *Geom. Dedicata*, **87** (2001), 167–180. MR2002g:20084
- [9] J. Wolf, Spaces of Constant Curvature, 5th ed., Publish or Perish, Wilmington, 1984.
- [10] S. Wolfram, Mathematica, Wolfram Research, 1993.

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, KOREA
E-mail address: kyha@sogang.ac.kr

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, KOREA
E-mail address: jlee@sogang.ac.kr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OKLAHOMA 73019
E-mail address: kb_lee@math.ou.edu