

ON DEGREES OF IRREDUCIBLE BRAUER CHARACTERS

W. WILLEMS

ABSTRACT. Based on a large amount of examples, which we have checked so far, we conjecture that $|G|_{p'} \leq \sum_{\varphi} \varphi(1)^2$ where p is a prime and the sum runs through the set of irreducible Brauer characters in characteristic p of the finite group G . We prove the conjecture simultaneously for p -solvable groups and groups of Lie type in the defining characteristic. In non-defining characteristics we give asymptotically an affirmative answer in many cases.

1. INTRODUCTION

Let G be a finite group and let $\text{IBr}_p(G)$, resp. $\text{IBr}_p(B)$, be the set of irreducible p -Brauer characters of G , resp. of a p -block B . For $\varphi \in \text{IBr}_p(G)$ let Φ_{φ} denote the projective indecomposable character corresponding to φ . Due to a result of Brauer and Nesbitt [6] the term $|G| - \frac{|G|}{\Phi_1(1)}$, where Φ_1 denotes the projective character corresponding to the trivial character, is an upper bound for the dimension of the Jacobson radical of the p -modular group algebra of G . An obvious reformulation of this result leads to

$$(1) \quad \frac{|G|}{\Phi_1(1)} \leq \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2.$$

Moreover, equality holds if and only if G has a normal Sylow p -subgroup. This was proved by Wallace in [18] for p -solvable groups, and in full generality by Brockhaus in [7] using the classification of finite simple groups.

In case $p = 2$ the classification can be avoided to show the normality of a Sylow p -subgroup if equality holds in (1). Since Φ_{φ} is a constituent of $\varphi\Phi_1$, equality in (1) implies that $\Phi_{\varphi} = \varphi\Phi_1$. Now, if $p \mid \varphi(1)$, the trivial character has multiplicity at least 2 in $\varphi\bar{\varphi}$, and a contradiction follows by considering the scalar product relation

$$1 = (\varphi, \Phi_{\varphi}) = (\varphi, \varphi\Phi_1) = (\varphi\bar{\varphi}, \phi_1) \geq 2$$

(see [13], VII, 8.5 d)). Thus p does not divide any irreducible Brauer character degree, and a nice argument of Okuyama (see [16], Theorem 2.33) implies that for $p = 2$ the Sylow p -subgroup of G is normal.

The lower bound in (1) has been improved in [14], replacing $\Phi_1(1)$ by the spectral radius $\rho(C)$ of the Cartan matrix C , i.e. the maximum value $|\lambda|$ where λ runs through the set of complex eigenvalues of C .

However, for obvious reasons we are interested in a lower bound which only depends on terms of G . This is a weaker question than the one Brauer asked in

Received by the editors January 9, 2003 and, in revised form, October 29, 2003.
2000 *Mathematics Subject Classification*. Primary 20C20, 20G40.

Problem 15 of his famous list [5]. He actually wanted to have a characterization of the dimension of the Jacobson radical by group-theoretical properties.

In the case of p -solvable groups (the existence of a p -complement suffices) we have $\Phi_1(1) = |G|_p$ (see [11], Chap. X, 3.2). Hence the desired bound is

$$(2) \quad |G|_{p'} \leq \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2.$$

But for non- p -solvable groups, $\Phi_1(1)$ (and $\rho(C)$) may differ extremely from $|G|_p$. For instance, let $G = R(q)$ with $q = 3^{2m+1}$ ($m \geq 1$) be a Ree group and let $p = 2$. Then $\Phi_1(1) = 2(q^3 + 1) \gg 8$ and $\rho(C) \approx 16.38$ (see [15] for the Cartan matrix), but $|G|_2 = 8$ and moreover

$$\frac{|G|}{8} = |G|_{2'} \leq \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2.$$

(In order to prove the inequality it is enough to consider the $\frac{q-3m}{6}$ characters of defect zero of degree $(q^2 - 1)(q + 1 + 3m)$, see [19].) Nevertheless for the sporadic groups that we have checked so far, the bound in (2) holds true. The same bound turns out to be true if the Sylow p -subgroups of G are cyclic [14]. So for any finite group G we are led to the

Conjecture. *We always have*

$$(3) \quad |G|_{p'} \leq \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2,$$

and equality holds if and only if G has a normal Sylow p -subgroup.

Note that the *if part* is trivial, since a normal p -subgroup is always contained in the kernel of an irreducible representation in characteristic p .

Suppose that the conjecture has an affirmative answer. Thus in the extreme case $|G|_{p'} = \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2$ the group G has a normal Sylow p -subgroup, say P . By Lemma 4.26 of ([11], Chapter IV), we know that

$$\Phi_\varphi(x) = |C_P(x)|\varphi(x)$$

for p' -elements $x \in G$. On the other hand, by Lemma 3.8 of ([11], Chapter IV), we have

$$\sum_{\varphi \in \text{IBr}_p(G)} \Phi_\varphi(x)\overline{\Phi_\varphi(y)} = \begin{cases} 0 & \text{if } x \not\sim y, \\ |C_G(x)| & \text{if } x \sim y, \end{cases}$$

for p' -elements $x, y \in G$ where \sim denotes conjugation in G . Thus we get the relation

$$\sum_{\varphi \in \text{IBr}_p(G)} \varphi(x)\overline{\varphi(y)} = \begin{cases} 0 & \text{if } x \not\sim y, \\ \frac{|C_G(x)|}{|C_P(x)|} & \text{if } x \sim y. \end{cases}$$

So we may ask the

Question. Is it even true that for p' -elements $x \in G$ we always have

$$(4) \quad \frac{|C_G(x)|}{|G|_p} \leq \sum_{\varphi \in \text{IBr}_p(G)} |\varphi(x)|^2 ?$$

If so, the bound may be sharp for $x \neq 1$ even if the Sylow p -subgroup is not normal. For instance, if $p = 5$ and x is an element of order 3 in $G = L_2(16)$, then

$$3 = \frac{15}{5} = \frac{|C_G(x)|}{|G|_5} = \sum_{\varphi \in \text{IBr}_5(G)} \varphi(x)\overline{\varphi(x)} = 1^2 + 1^2 + (-1)^2 = 3.$$

Note that the inequality (4) reduces to (3) for $x = 1$.

We are furthermore tempted to ask a p -local version of (3). In this case we may replace

$$|G|_{p'} = \frac{|G|}{p^a} = \frac{\dim KG}{p^a}$$

by $\frac{\dim B}{p^d}$ where K is a splitting field of G of characteristic p and B is a p -block of defect d . Thus we ask whether

$$(5) \quad \frac{\dim B}{p^d} \leq \sum_{\varphi \in \text{IBr}_p(B)} \varphi(1)^2?$$

An affirmative answer to (5) was given by Kiyota and Wada in [14] in case G is p -solvable or B is a p -block with cyclic defect group. In the latter case equality holds if and only if the Brauer tree is a star and all irreducible Brauer characters have the same degree.

Unfortunately the principal 2-block B_0 of the alternating group A_5 shows that inequality (5) fails to be true in general, because

$$11 = \frac{\dim B_0}{2^d} > \sum_{\varphi \in \text{IBr}_p(B)} \varphi(1)^2 = 1 + 2^2 + 2^2 = 9.$$

A more advanced counterexample is the non-principal 3-block of maximal defect of $6.A_7$. This example was brought to my attention by Thomas Breuer.

2. ON THE CONJECTURE

In this section we prove the conjecture for groups which have a projective character with properties similar to those of the Steinberg character for groups of Lie type. We start with a result due to Alperin. Let Φ be any projective character of a finite group G and let $\text{IBr}(G) = \{\varphi_1, \varphi_2, \dots, \varphi_s\}$. Then Φ uniquely determines a matrix $A = (a_{ij})$ by the equations

$$\varphi_i \Phi = \sum_{j=1}^s a_{ij} \Phi_j \quad (i = 1, \dots, s).$$

If x_1, \dots, x_s are representatives of the p' -conjugacy classes of G and if C denotes the Cartan matrix of G , then we have

Lemma 2.1 ([1]).

$$\det A = \frac{\prod_{i=1}^s \Phi(x_i)}{\det C}.$$

Proof. For $C = (c_{ij})$, we have

$$\varphi_i \Phi = \sum_{j,k=1}^s a_{ij} c_{jk} \varphi_k.$$

Therefore we get the matrix equation

$$(\varphi_i(x_l)) \begin{pmatrix} \Phi(x_1) & 0 & \cdots & 0 \\ 0 & \Phi(x_2) & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \Phi(x_s) \end{pmatrix} = AC(\varphi_i(x_l)).$$

Since $(\varphi_i(x_l))$ and C are invertible ([11], Chap. IV, 3.6), the assertion follows.

Theorem 2.2. *Let G be a finite group. Suppose that G has a projective character Φ satisfying the following two conditions.*

- (a) $\Phi(1) = |G|_p$.
- (b) $\Phi(x) \neq 0$ for all p' -elements $x \in G$.

Then

$$|G|_{p'} \leq \sum_{i=1}^s \varphi_i(1)^2,$$

and equality holds if and only if G has normal Sylow p -subgroup.

Proof. Condition (b) implies that $\det A \neq 0$ in the above lemma. Thus there exists a permutation π of $\{1, \dots, s\}$ such that

$$\prod_{i=1}^s a_{\pi i, i} \neq 0.$$

This means $\varphi_{\pi i} \Phi = \Phi_i + \cdots$. If ρ denotes the regular character of G , then

$$\sum_{i=1}^s \varphi_i(1) \varphi_{\pi i} \Phi = \sum_{i=1}^s \varphi_i(1) \Phi_i + \cdots = \rho + \cdots.$$

In particular we have

$$(6) \quad \sum_{i=1}^s \varphi_i(1) \varphi_{\pi i}(1) \Phi(1) \geq \rho(1) = |G|,$$

and by condition (a)

$$(7) \quad \sum_{i=1}^s \varphi_i(1) \varphi_{\pi i}(1) \geq |G|_{p'}.$$

Applying the Cauchy-Schwarz inequality yields

$$(8) \quad \left(\sum_{i=1}^s \varphi_i(1) \varphi_{\pi i}(1) \right)^2 \leq \sum_{i=1}^s \varphi_i(1)^2 \sum_{i=1}^s \varphi_{\pi i}(1)^2,$$

and hence

$$(9) \quad \sum_{i=1}^s \varphi_i(1) \varphi_{\pi i}(1) \leq \sum_{i=1}^s \varphi_i(1)^2.$$

Putting (7) and (9) together, we obtain the desired assertion

$$|G|_{p'} \leq \sum_{i=1}^s \varphi_i(1)^2.$$

So it remains to prove that $|G|_{p'} = \sum_{i=1}^s \varphi_i(1)^2$ implies that a Sylow p -subgroup of G is normal.

By (6) we get $\varphi_{\pi_i}\Phi = \Phi_i$ for all i . In particular we have $\varphi_{\pi_1}\Phi = \Phi_1$ for the trivial character φ_1 . The equality in the Cauchy-Schwarz inequality (see (8)) forces $\varphi_{\pi_1}(1) = \varphi_1(1) = 1$. Thus we obtain $\Phi_1(1) = |G|_p$ and by a result of Brockhaus [7] the group G has a normal Sylow p -subgroup.

The properties we required on the projective character in 2.2 seem to be very strong. However there are interesting examples for which the theorem applies.

Examples 2.3. a) Let G be a finite group with a split BN -pair of characteristic p and satisfying the commutator relations (see [9]). For Φ we take the Steinberg character. Then conditions a) and b) are satisfied, since

$$\Phi(x) = |C_G(x)|_p$$

for all p' -elements x of G (see [9], 6.4.7). Note that for Chevalley groups (twisted or non-twisted) defined in characteristic p , the square of the degree of the Steinberg character already dominates the p' -part of the order of G .

b) Let G be a finite group with a p -complement. Now we may take for Φ the projective indecomposable character corresponding to the trivial character. Observe that Φ is the trivial character of a p -complement induced to G and satisfies the assumptions of 2.2.

c) More subtle is the following. Let N be a normal p -solvable subgroup of G , and suppose that $H = G/N$ possess a projective character satisfying conditions a) and b). We prove that G has a projective character with the same properties. So let $M \leq N$ be a minimal normal subgroup of G . By the inductive hypothesis the group G/M has a projective character, say Ψ , satisfying a) and b). If M is a p' -group, we may take for Φ the inflation $\Phi = \text{infl}_G(\Psi)$ of Ψ . Thus suppose that M is a p -group. Let λ be the Brauer character of the conjugation action of G on M . Now we put

$$\Phi = \lambda \text{infl}_G(\Psi).$$

That Φ is projective follows from a result of Alperin, Collins and Sibley (see [2]). Clearly,

$$\Phi(1) = \lambda(1) \text{infl}_G(\Psi)(1) = |M||H| = |G|_p$$

and

$$\Phi(x) = \lambda(x) \text{infl}_G(\Psi)(x) = |C_M(x)| \text{infl}_G(\Psi)(x) \neq 0$$

for all p' -elements $x \in G$.

3. ASYMPTOTIC RESULTS FOR GROUPS OF LIE TYPE

To give more evidence for inequality (3) we will look in this section asymptotically on groups of Lie type in non-defining characteristics. For a detailed introduction to the character theory of groups of Lie type, the reader is referred to [9].

Let G be a simple linear algebraic group over an algebraically closed field K of positive characteristic $r \neq p$, and let $G^F = G(q)$ (where $q = r^f$) denote the group of fixed points under a Frobenius map F . Furthermore let $T_w = T^F$ be the maximal torus in G^F corresponding to $w \in W = W(T)$ where W denotes the Weyl group of an F -stable maximal torus $T \subseteq G$.

Now we look at irreducible Deligne-Lusztig characters $R_{T,\theta}$ (up to a sign) where $\theta \in \widehat{T}^F = \text{Hom}(T^F, \mathbb{C}^*)$ is an irreducible character of $T_w = T^F$ and θ is in general

position. Let $\mathcal{R}(q)$ denote the set of all such $R_{T,\theta}$'s. Note that $R_{T,\theta} = R_{T,\theta'}$ if and only if $\theta = w'\theta'$ for some

$$w' \in W(T)^F = C_{w,F} = \{w' \in W \mid w'^{-1}wF(w') = w\}$$

(see [9], p. 219). Thus the number $|\mathcal{R}(q)|$ of different $R_{T,\theta}$'s is equal to the number of $W(T)^F$ -orbits of characters θ in general position. By a result of Veldkamp ([17]) this number is equal to the number of $W(T)^F$ -conjugacy classes of elements in general position in the fixed point group T^{*F^*} of the dual torus $T^* \subseteq G^*$.

For simplicity let us suppose that G^* is simply connected. In this case the regular elements in T^{*F^*} coincide with the elements in general position. (Note that $t \in G$ is regular (resp. in general position) if $C_G(t)^\circ = T$ (resp. $C_G(t) = T$).) Let l denote the Lie rank of G . Applying 3.1 and 3.2 in [10] (see also [12]), we get

$$|\mathcal{R}(q)| = \{R_{T,\theta} \mid \theta \text{ in general position}\} = \frac{q^l}{|C_{w,F}|} + O(q^{l-1}).$$

Note that $R_{T,\theta}$ is of degree

$$R_{T,\theta}(1) = \frac{|G(q)|_{r'}}{|T_w|}.$$

Thus we obtain

$$(10) \quad \sum_{R_{T,\theta} \in \mathcal{R}(q)} R_{T,\theta}(1)^2 = \left(\frac{q^l}{|C_{w,F}|} + O(q^{l-1}) \right) \frac{|G(q)|_{r'}^2}{|T_w|^2}.$$

Let N denote the number of positive roots. As polynomials in q we have

- $|T_w| = q^l + \text{lower terms in } q,$
- $|G(q)|_{r'} = q^{l+N} + \text{lower terms in } q,$
- $|G(q)| = q^{l+2N} + \text{lower terms in } q.$

Inserting these equations in (10), we obtain

$$(11) \quad \begin{aligned} \sum_{R_{T,\theta} \in \mathcal{R}(q)} R_{T,\theta}(1)^2 &= \frac{1}{|C_{w,F}|} q^{l+2N} + O(q^{l+2N-1}) \\ &= |G(q)|_{p'} + \left(\frac{1}{|C_{w,F}|} - \frac{1}{|G(q)|_p} \right) q^{l+2N} + O(q^{l+2N-1}). \end{aligned}$$

We may assume that $p \mid |G(q)|$. If we exclude the Steinberg triality ${}^3D_4(q^2)$ for a moment, then

$$|G(q)|_{r'} = \prod_{i=1}^l (q^{d_i} - \epsilon_i)$$

where $\epsilon_i \in \{1, -1\}$ and the d_i are the exponents of the underlying group (see [9], p. 75). Thus $p \mid q^{2d_i} - 1$. Note that

$$(q^{2d_i} - 1)_p < (q^{2pd_i} - 1)_p$$

and $|W|$ is independent of q (see [3], Planche II). Thus there exists an $s \in \mathbb{N}$ (for instance s a suitable power of p) such that

$$|C_{w,F}| \leq |W| < |G(q^s)|_p,$$

and therefore

$$|C_{w,F}| < |G(q^{st})|_p$$

for all $t \in \mathbb{N}$ and all $w \in W$. This is also true for ${}^3D_4(q^3)$, as a similar argument shows. By (11) we obtain

$$\sum_{R_{T,\theta} \in \mathcal{R}(q^{st})} R_{T,\theta}(1)^2 \geq |G(q^{st})|_p'$$

for all $t \geq t_0$. Now, if there exists a torus $T_w = T_w(q^{st}) \subseteq G(q^{st})$ with $p \nmid |T_w|$ for $t \geq t_0$, then all characters $R_{T,\theta}$ with θ in general position are irreducible and of p -defect zero. In particular they are irreducible Brauer characters for the prime p and we have the desired inequality

$$\sum_{\varphi \in \text{IBr}_p(G(q^{st}))} \varphi(1)^2 \geq |G(q^{st})|_p'$$

for all $t \geq t_0$. Thus we have proved

Theorem 3.1. *Let $G(q) = G^F$ be a finite group of Lie type where G is a linear simple algebraic group and G^* is simply connected. Let p be a prime with $p \nmid q$ and $p \mid |G(q)|$.*

a) *Then there exists an $s \in \mathbb{N}$ such that*

$$\sum_{R_{T,\theta} \in \mathcal{R}(q^{st})} R_{T,\theta}(1)^2 \geq |G(q^{st})|_p'$$

for all $t \geq t_0$.

b) *If $T_w = T_w(q^{st})$ is a torus in $G(q^{st})$ for some $w \in W$ and $p \nmid |T_w|$ for all $t \geq t_0$, then*

$$\sum_{\varphi \in \text{IBr}_p(G(q^{st}))} \varphi(1)^2 \geq |G(q^{st})|_p'$$

for all $t \geq t_0$.

The critical point in the whole process is the assumption $p \nmid |T_w| = |T_w(q^{st})|$ for a suitable $w \in W$ and all $t \geq t_0$. The author would like to thank an anonymous referee for pointing out the following argument.

If we replace s by $s(p - 1)$, then we have $q^s \equiv 1 \pmod p$. Note that the order of $T_w(q)$ can be written as

$$|T_w(q)| = \Phi_{n_1}(q) \cdots \Phi_{n_r}(q)$$

where Φ_n denotes the n -th cyclotomic polynomial. Now we choose $w \in W$ in such a way that $1 < n_j$ and $p \nmid n_j$ for all $j = 1, \dots, r$. Since

$$\begin{aligned} \Phi_{n_j}(q^{st}) \mid \frac{q^{st n_j} - 1}{q^{st} - 1} &= 1 + q^{st} + \dots + q^{st(n_j - 1)} \\ &\equiv 1 + 1 + \dots + 1 \pmod p \\ &\equiv n_j \pmod p, \end{aligned}$$

we get $p \nmid \Phi_{n_j}(q^{st})$ for all t ; hence $p \nmid |T_w(q^{st})|$ for all t .

The following example may illustrate the above. In particular, in the case where $G(q) = \text{PGL}(2, q)$ we are not able to find a torus as required for the prime $p = 2$. Thus our method fails for $p = 2$.

Example. Let $G^F = G(q) = \text{PGL}(l + 1, q)$ be the finite adjoint group of type A_l for $l \geq 2$ with q a power of r . Note that $G^{*F^*} = \text{SL}(l + 1, q)$ is the fixed point group of a simply connected group. Let $w \in W$ be a Coxeter element, i.e. w is a cycle of length $l + 1$, and $C_{w,F} = C_{S_{l+1}}(w) = \langle w \rangle$ where S_n is the symmetric group on

n letters. Let p be a prime with $p > l + 1$. Thus we either have $p \nmid |G(q)|$ and (3) holds with equality, or $|C_{w,F}| < |G(q)|_p$ and we have $\sum_{R_{T,\theta} \in \mathcal{R}} R_{T,\theta}(1)^2 \geq |G(q)|_{p'}$ for $q \geq q_0$ and $p \mid |G(q)|$. Let such a q be given. Now if $p \nmid |T_w| = 1 + q + \dots + q^l$, then

$$\sum_{\varphi \in \text{IBr}_p(G(q))} \varphi(1)^2 \geq |G(q)|_{p'}.$$

The condition $p \nmid |T_w|$ is satisfied for instance if $p \nmid q - 1$ and $p \mid q^s - 1$ where $2 < s < l + 1$ and $\gcd(s, l + 1) = 1$.

Let us specialize to $G(q) = \text{PGL}(2, q)$ and let $2 < p \neq r$, hence $2 = |W| < p$. Since $|G(q)| = q(q+1)(q-1)$, we have $p \nmid |T_w| = (q+1)$ or $p \nmid |T_1| = q-1$ where T_w is the Coxeter and T_1 the split torus. Thus for all q large enough and all odd primes inequality (3) holds true for $\text{PGL}(2, q)$. However in this small case we get (3) for any q . If $p = r$ we may apply 2.3 a). In case $2 \neq p \neq r$ the Sylow p -subgroups of $G(q)$ are cyclic. Thus the inequality (5) holds true for every p -block by [14], which obviously implies (3). For $p = 2$ one easily checks inequality (3) using the results of Section VIII in [8].

ADDED IN PROOF

There is an analogue of the Conjecture for p' -class lengths, namely $|G|_{p'} \leq \sum_x |G : C_G(x)|$ where the sum runs through a set of representatives of the p' -conjugacy classes, and with equality if and only if G has a normal p -complement. In contrast to characters this can easily be proved using the Frobenius Conjecture which we know to be true. We shall discuss this and similar questions in a forthcoming paper.

REFERENCES

- [1] J. ALPERIN. Projective modules and tensor products. *J. Pure and Appl. Algebra* 8 (1976), 235-241. MR53:5712
- [2] J.L. ALPERIN, M.J. COLLINS AND D.A. SIBLEY. Projective modules, filtrations and Cartan invariants. *Bull. London Math. Soc.* (3) 16 (1984), 416-420. MR85m:20009
- [3] N. BOURBAKI. *Groupes et algèbre de Lie*. Chapitres 4,5 et 6. Masson, Paris 1981. MR83g:17001
- [4] R. BRAUER. Notes on representations of finite groups. *J. London Math. Soc.* (2), 13 (1976), 162-166. MR53:3091
- [5] R. BRAUER. Representations of finite groups. *Lectures on Modern Mathematics*, Vol. 1, Wiley, New York, 1963, 133-175. MR31:2314
- [6] R. BRAUER AND C.J. NESBITT. On the modular characters of groups, *Ann. Math.*(2) 42 (1941), 556-590. MR2:309c
- [7] P. BROCKHAUS. On the radical of a group algebra. *J. Algebra* 95 (1985), 454-472. MR87h:20021
- [8] R. BURKHARDT. Die Zerlegungsmatrizen der Gruppen $\text{PSL}(2, p^f)$. *J. Algebra* 40 (1976), 75-96. MR58:864
- [9] R.W. CARTER. *Finite groups of Lie type, conjugacy classes and complex characters*. Wiley, Chichester 1985. MR87d:20060
- [10] D.I. DERIZIOTIS. On the number of conjugacy classes in finite groups of Lie type. *Comm. Algebra* 13 (1985), 1019-1045. MR86i:20067
- [11] W. FEIT. *The representation theory of finite groups*. North Holland, Amsterdam 1982. MR83g:20001
- [12] P. FLEISCHMANN AND I. JANISZCZAK. The number of regular semisimple elements for Chevalley groups of classical type. *J. Algebra* 155 (1993), 482-528. MR94f:20090
- [13] B. HUPPERT AND N. BLACKBURN. *Finite groups II*. Springer-Verlag, Berlin/Heidelberg/New York 1982. MR84i:20001a

- [14] M. KIYOTA AND T. WADA. Some remarks on the eigenvalues of the Cartan matrix in finite groups. *Commun. Algebra* 21, No.11 (1993), 3839-3860. MR94i:20021
- [15] P. LANDROCK AND G.O. MICHLER. Principal 2-blocks of simple groups of Lie type. *Trans. Am. Math. Soc.* 260 (1980), 83-111. MR81h:20013
- [16] G. NAVARRO. Characters and blocks of finite groups. London Math. Soc., Lecture Notes Series 250, Cambridge Univ. Press, Cambridge, 1998. MR2000a:20018
- [17] F.D. VELDKAMP. Regular characters and regular elements. *Comm. Algebra* 5(12) (1977), 1259-1273. MR58:16857
- [18] D.A.R. WALLACE. On the radical of a group algebra, *Proc. Am. Math. Soc.* 12 (1961), 133-137. MR22:12146
- [19] H.N. WARD. On Ree's series of simple groups. *Trans. Amer. Math. Soc.* 121 (1966), 62-89. MR33:5752

INSTITUT FÜR ALGEBRA UND GEOMETRIE, FAKULTÄT FÜR MATHEMATIK, OTTO-VON-GUERICKE-UNIVERSITÄT, 39016 MAGDEBURG, GERMANY