

k -HYPONORMALITY OF FINITE RANK PERTURBATIONS OF UNILATERAL WEIGHTED SHIFTS

RAÚL E. CURTO AND WOO YOUNG LEE

ABSTRACT. In this paper we explore finite rank perturbations of unilateral weighted shifts W_α . First, we prove that the subnormality of W_α is never stable under nonzero finite rank perturbations unless the perturbation occurs at the zeroth weight. Second, we establish that 2-hyponormality implies positive quadratic hyponormality, in the sense that the Maclaurin coefficients of $D_n(s) := \det P_n [(W_\alpha + sW_\alpha^2)^*, W_\alpha + sW_\alpha^2] P_n$ are nonnegative, for every $n \geq 0$, where P_n denotes the orthogonal projection onto the basis vectors $\{e_0, \dots, e_n\}$. Finally, for α strictly increasing and W_α 2-hyponormal, we show that for a small finite-rank perturbation α' of α , the shift $W_{\alpha'}$ remains quadratically hyponormal.

1. INTRODUCTION

Let \mathcal{H} and \mathcal{K} be complex Hilbert spaces, let $\mathcal{L}(\mathcal{H}, \mathcal{K})$ be the set of bounded linear operators from \mathcal{H} to \mathcal{K} and write $\mathcal{L}(\mathcal{H}) := \mathcal{L}(\mathcal{H}, \mathcal{H})$. An operator $T \in \mathcal{L}(\mathcal{H})$ is said to be normal if $T^*T = TT^*$, hyponormal if $T^*T \geq TT^*$, and subnormal if $T = N|_{\mathcal{H}}$, where N is normal on some Hilbert space $\mathcal{K} \supseteq \mathcal{H}$. If T is subnormal, then T is also hyponormal. Recall that given a bounded sequence of positive numbers $\alpha : \alpha_0, \alpha_1, \dots$ (called *weights*), the (*unilateral*) *weighted shift* W_α associated with α is the operator on $\ell^2(\mathbb{Z}_+)$ defined by $W_\alpha e_n := \alpha_n e_{n+1}$ for all $n \geq 0$, where $\{e_n\}_{n=0}^\infty$ is the canonical orthonormal basis for ℓ^2 . It is straightforward to check that W_α can never be *normal*, and that W_α is *hyponormal* if and only if $\alpha_n \leq \alpha_{n+1}$ for all $n \geq 0$. The Bram-Halmos criterion for subnormality states that an operator T is subnormal if and only if

$$\sum_{i,j} (T^i x_j, T^j x_i) \geq 0$$

Received by the editors December 10, 1999 and, in revised form, December 31, 2001.

2000 *Mathematics Subject Classification*. Primary 47B20, 47B35, 47B37; Secondary 47-04, 47A20, 47A57.

Key words and phrases. Weighted shifts, perturbations, subnormal, k -hyponormal, weakly k -hyponormal.

The work of the first-named author was partially supported by NSF research grants DMS-9800931 and DMS-0099357.

The work of the second-named author was partially supported by a grant (R14-2003-006-01001-0) from the Korea Science and Engineering Foundation.

for all finite collections $x_0, x_1, \dots, x_k \in \mathcal{H}$ ([2], [4, II.1.9]). It is easy to see that this is equivalent to the following positivity test:

$$(1.1) \quad \begin{pmatrix} I & T^* & \dots & T^{*k} \\ T & T^*T & \dots & T^{*k}T \\ \vdots & \vdots & \ddots & \vdots \\ T^k & T^*T^k & \dots & T^{*k}T^k \end{pmatrix} \geq 0 \quad (\text{for all } k \geq 1).$$

Condition (1.1) provides a measure of the gap between hyponormality and subnormality. In fact, the positivity condition (1.1) for $k = 1$ is equivalent to the hyponormality of T , while subnormality requires the validity of (1.1) for all k . Let $[A, B] := AB - BA$ denote the commutator of two operators A and B , and define T to be *k-hyponormal* whenever the $k \times k$ operator matrix

$$(1.2) \quad M_k(T) := ([T^{*j}, T^i]_{i,j=1}^k)$$

is positive. An application of the Choleski algorithm for operator matrices shows that the positivity of (1.2) is equivalent to the positivity of the $(k+1) \times (k+1)$ operator matrix in (1.1); the Bram-Halmos criterion can then be rephrased to say that T is subnormal if and only if T is *k-hyponormal* for every $k \geq 1$ ([16]).

Recall ([1], [16], [5]) that $T \in \mathcal{L}(\mathcal{H})$ is said to be *weakly k-hyponormal* if

$$LS(T, T^2, \dots, T^k) := \left\{ \sum_{j=1}^k \alpha_j T^j : \alpha = (\alpha_1, \dots, \alpha_k) \in \mathbb{C}^k \right\}$$

consists entirely of hyponormal operators, or equivalently, $M_k(T)$ is *weakly positive*, i.e. ([16]),

$$(1.3) \quad \left(M_k(T) \begin{pmatrix} \lambda_0 x \\ \vdots \\ \lambda_k x \end{pmatrix}, \begin{pmatrix} \lambda_0 x \\ \vdots \\ \lambda_k x \end{pmatrix} \right) \geq 0 \quad \text{for } x \in \mathcal{H} \text{ and } \lambda_0, \dots, \lambda_k \in \mathbb{C}.$$

If $k = 2$, then T is said to be *quadratically hyponormal*, and if $k = 3$, then T is said to be *cubically hyponormal*. Similarly, $T \in \mathcal{L}(\mathcal{H})$ is said to be *polynomially hyponormal* if $p(T)$ is hyponormal for every polynomial $p \in \mathbb{C}[z]$. It is known that *k-hyponormal* \Rightarrow *weakly k-hyponormal*, but the converse is not true in general.

The classes of (weakly) *k-hyponormal* operators have been studied in an attempt to bridge the gap between subnormality and hyponormality ([7], [8], [10], [11], [12], [14], [16], [19], [22]). The study of this gap has been only partially successful. For example, such a gap is not yet well described for Toeplitz operators on the Hardy space of the unit circle; in fact, even subnormality for Toeplitz operators has not been characterized (cf. [20], [6]). For weighted shifts, positive results appear in [7] and [12], although no concrete example of a weighted shift which is polynomially hyponormal and not subnormal has yet been found (the existence of such weighted shifts was established in [17] and [18]).

In the present paper we renew our efforts to help describe the above-mentioned gap between subnormality and hyponormality, with particular emphasis on polynomial hyponormality. We focus on the class of unilateral weighted shifts, and initiate a study of how the above-mentioned notions behave under finite perturbations of the weight sequence. We first obtain the following three concrete results.

(i) the subnormality of W_α is never stable under nonzero finite rank perturbations unless the perturbation is confined to the zeroth weight (Theorem 2.1);

(ii) 2-hyponormality implies *positive quadratic hyponormality*, in the sense that the Maclaurin coefficients of $D_n(s) := \det P_n [(W_\alpha + sW_\alpha^2)^*, W_\alpha + sW_\alpha^2] P_n$ are nonnegative, for every $n \geq 0$, where P_n denotes the orthogonal projection onto the basis vectors $\{e_0, \dots, e_n\}$ (Theorem 2.2); and

(iii) if α is strictly increasing and W_α is 2-hyponormal, then for α' a small perturbation of α , the shift $W_{\alpha'}$ remains positively quadratically hyponormal (Theorem 2.3).

Along the way we establish two related results, each of independent interest:

(iv) an integrality criterion for a subnormal weighted shift to have an n -step subnormal extension (Theorem 6.1); and

(v) a proof that the sets of k -hyponormal and weakly k -hyponormal operators are closed in the strong operator topology (Proposition 6.7).

2. STATEMENT OF MAIN RESULTS

C. Berger’s characterization of subnormality for unilateral weighted shifts (cf. [21], [4, III.8.16]) states that W_α is subnormal if and only if there exists a Borel probability measure μ (the so-called Berger measure of W_α) supported in $[0, \|W_\alpha\|^2]$, with $\|W_\alpha\|^2 \in \text{supp } \mu$, such that

$$\gamma_n = \int t^n d\mu(t) \quad \text{for all } n \geq 0.$$

Given an initial segment of weights $\alpha : \alpha_0, \dots, \alpha_m$, the sequence $\hat{\alpha} \in \ell^\infty(\mathbb{Z}_+)$ such that $\hat{\alpha}_i = \alpha_i$ ($i = 0, \dots, m$) is said to be *recursively generated* by α if there exist $r \geq 1$ and $\varphi_0, \dots, \varphi_{r-1} \in \mathbb{R}$ such that

$$(2.1) \quad \gamma_{n+r} = \varphi_0 \gamma_n + \dots + \varphi_{r-1} \gamma_{n+r-1} \quad (\text{for all } n \geq 0),$$

where $\gamma_0 := 1$, $\gamma_n := \alpha_0^2 \cdots \alpha_{n-1}^2$ ($n \geq 1$). In this case $W_{\hat{\alpha}}$ with weights $\hat{\alpha}$ is said to be *recursively generated*. If we let

$$(2.2) \quad g(t) := t^r - (\varphi_{r-1} t^{r-1} + \dots + \varphi_0),$$

then g has r distinct real roots $0 \leq s_0 < \dots < s_{r-1}$ ([11, Theorem 3.9]). Let

$$V := \begin{pmatrix} 1 & 1 & \dots & 1 \\ s_0 & s_1 & \dots & s_{r-1} \\ \vdots & \vdots & & \vdots \\ s_0^{r-1} & s_1^{r-1} & \dots & s_{r-1}^{r-1} \end{pmatrix}$$

and let

$$\begin{pmatrix} \rho_0 \\ \vdots \\ \rho_{r-1} \end{pmatrix} := V^{-1} \begin{pmatrix} \gamma_0 \\ \vdots \\ \gamma_{r-1} \end{pmatrix}.$$

If the associated recursively generated weighted shift $W_{\hat{\alpha}}$ is subnormal, then its Berger measure is of the form

$$\mu := \rho_0 \delta_{s_0} + \dots + \rho_{r-1} \delta_{s_{r-1}}.$$

For example, given $\alpha_0 < \alpha_1 < \alpha_2$, $W_{(\alpha_0, \alpha_1, \alpha_2)^\wedge}$ is the recursive weighted shift whose weights are calculated according to the recursive relation

$$(2.3) \quad \alpha_{n+1}^2 = \varphi_1 + \varphi_0 \frac{1}{\alpha_n^2},$$

where

$$(2.4) \quad \varphi_0 = -\frac{\alpha_0^2 \alpha_1^2 (\alpha_2^2 - \alpha_1^2)}{\alpha_1^2 - \alpha_0^2} \quad \text{and} \quad \varphi_1 = \frac{\alpha_1^2 (\alpha_2^2 - \alpha_0^2)}{\alpha_1^2 - \alpha_0^2}.$$

In this case, $W_{(\alpha_0, \alpha_1, \alpha_2)^\wedge}$ is subnormal with 2-atomic Berger measure. Let W_x denote the weighted shift whose weight sequence consists of the initial weight x followed by the weight sequence of $W_{(\alpha_0, \alpha_1, \alpha_2)^\wedge}$.

By the Density Theorem ([11, Theorem 4.2 and Corollary 4.3]), we know that if W_α is a subnormal weighted shift with weights $\alpha = \{\alpha_n\}$ and $\epsilon > 0$, then there exists a nonzero compact operator K with $\|K\| < \epsilon$ such that $W_\alpha + K$ is a recursively generated subnormal weighted shift; in fact $W_\alpha + K = W_{\widehat{\alpha^{(m)}}}$ for some $m \geq 1$, where $\alpha^{(m)} : \alpha_0, \dots, \alpha_m$. The following result shows that K cannot generally be taken to be of finite rank.

Theorem 2.1 (Finite Rank Perturbations of Subnormal Shifts). *If W_α is a subnormal weighted shift, then there exists no nonzero finite rank operator F ($\neq cP_{\{e_0\}}$) such that $W_\alpha + F$ is a subnormal weighted shift. Concretely, suppose W_α is a subnormal weighted shift with weight sequence $\alpha = \{\alpha_n\}_{n=0}^\infty$ and assume $\alpha' = \{\alpha'_n\}$ is a nonzero perturbation of α in a finite number of weights except the initial weight. Then $W_{\alpha'}$ is not subnormal.*

We next consider the self-commutator $[(W_\alpha + sW_\alpha^2)^*, W_\alpha + sW_\alpha^2]$. Let W_α be a hyponormal weighted shift. For $s \in \mathbb{C}$, we write

$$D(s) := [(W_\alpha + sW_\alpha^2)^*, W_\alpha + sW_\alpha^2]$$

and we let

$$(2.5) \quad \begin{aligned} D_n(s) &:= P_n[(W_\alpha + sW_\alpha^2)^*, W_\alpha + sW_\alpha^2]P_n \\ &= \begin{pmatrix} q_0 & \bar{r}_0 & 0 & \dots & 0 & 0 \\ r_0 & q_1 & \bar{r}_1 & \dots & 0 & 0 \\ 0 & r_1 & q_2 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & q_{n-1} & \bar{r}_{n-1} \\ 0 & 0 & 0 & \dots & r_{n-1} & q_n \end{pmatrix}, \end{aligned}$$

where P_n is the orthogonal projection onto the subspace generated by $\{e_0, \dots, e_n\}$,

$$(2.6) \quad \begin{cases} q_n := u_n + |s|^2 v_n, \\ r_n := s\sqrt{w_n}, \\ u_n := \alpha_n^2 - \alpha_{n-1}^2, \\ v_n := \alpha_n^2 \alpha_{n+1}^2 - \alpha_{n-1}^2 \alpha_{n-2}^2, \\ w_n := \alpha_n^2 (\alpha_{n+1}^2 - \alpha_{n-1}^2)^2, \end{cases}$$

and, for notational convenience, $\alpha_{-2} = \alpha_{-1} = 0$. Clearly, W_α is quadratically hyponormal if and only if $D_n(s) \geq 0$ for all $s \in \mathbb{C}$ and all $n \geq 0$. Let $d_n(\cdot) :=$

$\det(D_n(\cdot))$. Then d_n satisfies the following 2-step recursive formula:

$$(2.7) \quad d_0 = q_0, \quad d_1 = q_0q_1 - |r_0|^2, \quad d_{n+2} = q_{n+2}d_{n+1} - |r_{n+1}|^2d_n.$$

If we let $t := |s|^2$, we observe that d_n is a polynomial in t of degree $n + 1$, and if we write $d_n \equiv \sum_{i=0}^{n+1} c(n, i)t^i$, then the coefficients $c(n, i)$ satisfy a double-indexed recursive formula, namely

$$(2.8) \quad \begin{aligned} c(n + 2, i) &= u_{n+2}c(n + 1, i) + v_{n+2}c(n + 1, i - 1) - w_{n+1}c(n, i - 1), \\ c(n, 0) &= u_0 \cdots u_n, \quad c(n, n + 1) = v_0 \cdots v_n, \quad c(1, 1) = u_1v_0 + v_1u_0 - w_0 \end{aligned}$$

($n \geq 0, i \geq 1$). We say that W_α is *positively quadratically hyponormal* if $c(n, i) \geq 0$ for every $n \geq 0, 0 \leq i \leq n + 1$ (cf. [9]). Evidently, positively quadratically hyponormal \implies quadratically hyponormal. The converse, however, is not true in general (cf. [3]).

The following theorem establishes a useful relation between 2-hyponormality and positive quadratic hyponormality.

Theorem 2.2. *Let $\alpha \equiv \{\alpha_n\}_{n=0}^\infty$ be a weight sequence and assume that W_α is 2-hyponormal. Then W_α is positively quadratically hyponormal. More precisely, if W_α is 2-hyponormal, then*

$$(2.9) \quad c(n, i) \geq v_0 \cdots v_{i-1}u_i \cdots u_n \quad (n \geq 0, 0 \leq i \leq n + 1).$$

In particular, if α is strictly increasing and W_α is 2-hyponormal, then the Maclaurin coefficients of $d_n(t)$ are positive for all $n \geq 0$.

If W_α is a weighted shift with weight sequence $\alpha = \{\alpha_n\}_{n=0}^\infty$, then the *moments* of W_α are usually defined by $\beta_0 := 1, \beta_{n+1} := \alpha_n\beta_n$ ($n \geq 0$) [23]; however, we prefer to reserve this term for the sequence $\gamma_n := \beta_n^2$ ($n \geq 0$). A criterion for k -hyponormality can be given in terms of these moments ([7, Theorem 4]): if we build a $(k + 1) \times (k + 1)$ Hankel matrix $A(n; k)$ by

$$(2.10) \quad A(n; k) := \begin{pmatrix} \gamma_n & \gamma_{n+1} & \cdots & \gamma_{n+k} \\ \gamma_{n+1} & \gamma_{n+2} & \cdots & \gamma_{n+k+1} \\ \vdots & \vdots & & \vdots \\ \gamma_{n+k} & \gamma_{n+k+1} & \cdots & \gamma_{n+2k} \end{pmatrix} \quad (n \geq 0),$$

then

$$(2.11) \quad W_\alpha \text{ is } k\text{-hyponormal} \iff A(n; k) \geq 0 \quad (n \geq 0).$$

In particular, for α strictly increasing, W_α is 2-hyponormal if and only if

$$(2.12) \quad \det \begin{pmatrix} \gamma_n & \gamma_{n+1} & \gamma_{n+2} \\ \gamma_{n+1} & \gamma_{n+2} & \gamma_{n+3} \\ \gamma_{n+2} & \gamma_{n+3} & \gamma_{n+4} \end{pmatrix} \geq 0 \quad (n \geq 0).$$

One might conjecture that if W_α is a k -hyponormal weighted shift whose weight sequence is strictly increasing, then W_α remains weakly k -hyponormal under a small perturbation of the weight sequence. We will show below that this is true for $k = 2$ (Theorem 2.3).

In [12, Theorem 4.3], it was shown that the gap between 2-hyponormality and quadratic hyponormality can be detected by unilateral shifts with a weight sequence $\alpha : \sqrt{x}, (\sqrt{a}, \sqrt{b}, \sqrt{c})^\wedge$. In particular, there exists a maximum value $H_2 \equiv H_2(a, b, c)$

of x that makes $W_{\sqrt{x},(\sqrt{a},\sqrt{b},\sqrt{c})^\wedge}$ 2-hyponormal; H_2 is called the *modulus* of 2-hyponormality (cf. [12]). Any value of $x > H_2$ yields a non-2-hyponormal weighted shift. However, if $x - H_2$ is small enough, $W_{\sqrt{x},(\sqrt{a},\sqrt{b},\sqrt{c})^\wedge}$ is still quadratically hyponormal. The following theorem shows that, more generally, for finite rank perturbations of weighted shifts with strictly increasing weight sequences, there always exists a gap between 2-hyponormality and quadratic hyponormality.

Theorem 2.3 (Finite Rank Perturbations of 2-Hyponormal Shifts). *Let $\alpha = \{\alpha_n\}_{n=0}^\infty$ be a strictly increasing weight sequence. If W_α is 2-hyponormal, then W_α remains positively quadratically hyponormal under a small nonzero finite rank perturbation of α .*

3. PROOF OF THEOREM 2.1

Proof of Theorem 2.1. It suffices to show that if T is a weighted shift whose restriction to $\bigvee\{e_n, e_{n+1}, \dots\}$ ($n \geq 2$) is subnormal, then there is at most one α_{n-1} for which T is subnormal.

Let $W := T|_{\bigvee\{e_{n-1}, e_n, e_{n+1}, \dots\}}$ and $S := T|_{\bigvee\{e_n, e_{n+1}, \dots\}}$, where $n \geq 2$. Then W and S have weights $\alpha_k(W) := \alpha_{k+n-1}$ and $\alpha_k(S) := \alpha_{k+n}$ ($k \geq 0$). Thus the corresponding moments are related by the equation

$$\gamma_k(S) = \alpha_n^2 \cdots \alpha_{n+k-1}^2 = \frac{\gamma_{k+1}(W)}{\alpha_{n-1}^2}.$$

We now adapt the proof of [7, Proposition 8]. Suppose S is subnormal with associated Berger measure μ . Then $\gamma_k(S) = \int_0^{\|T\|^2} t^k d\mu$. Thus W is subnormal if and only if there exists a probability measure ν on $[0, \|T\|^2]$ such that

$$\frac{1}{\alpha_{n-1}^2} \int_0^{\|T\|^2} t^{k+1} d\nu(t) = \int_0^{\|T\|^2} t^k d\mu(t) \quad \text{for all } k \geq 0,$$

which readily implies that $t d\nu = \alpha_{n-1}^2 d\mu$. Thus W is subnormal if and only if the formula

$$(3.1) \quad d\nu := \lambda \cdot \delta_0 + \frac{\alpha_{n-1}^2}{t} d\mu$$

defines a probability measure for some $\lambda \geq 0$, where δ_0 is the point mass at the origin. In particular $\frac{1}{t} \in L^1(\mu)$ and $\mu(\{0\}) = 0$ whenever W is subnormal. If we repeat the above argument for W and $V := T|_{\bigvee\{e_{n-2}, e_{n-1}, \dots\}}$, then we should have that $\nu(\{0\}) = 0$ whenever V is subnormal. Therefore we can conclude that if V is subnormal, then $\lambda = 0$, and hence

$$(3.2) \quad d\nu = \frac{\alpha_{n-1}^2}{t} d\mu.$$

Thus we have

$$1 = \int_0^{\|T\|^2} d\nu(t) = \alpha_{n-1}^2 \int_0^{\|T\|^2} \frac{1}{t} d\mu(t),$$

so that

$$(3.3) \quad \alpha_{n-1}^2 = \left(\int_0^{\|T\|^2} \frac{1}{t} d\mu(t) \right)^{-1},$$

which implies that α_{n-1} is determined uniquely by $\{\alpha_n, \alpha_{n+1}, \dots\}$ whenever T is subnormal. This completes the proof. \square

Theorem 2.1 says that a nonzero finite rank perturbation of a subnormal shift is never subnormal unless the perturbation occurs at the initial weight. However, this is not the case for k -hyponormality. To see this we use a close relative of the Bergman shift B_+ (whose weights are given by $\alpha = \{\sqrt{\frac{n+1}{n+2}}\}_{n=0}^\infty$); it is well known that B_+ is subnormal.

Example 3.1. For $x > 0$, let T_x be the weighted shift whose weights are given by

$$\alpha_0 := \sqrt{\frac{1}{2}}, \quad \alpha_1 := \sqrt{x}, \quad \text{and} \quad \alpha_n := \sqrt{\frac{n+1}{n+2}} \quad (n \geq 2).$$

Then we have:

- (i) T_x is subnormal $\iff x = \frac{2}{3}$;
- (ii) T_x is 2-hyponormal $\iff \frac{63-\sqrt{129}}{80} \leq x \leq \frac{24}{35}$.

Proof. Assertion (i) follows from Theorem 2.1. For assertion (ii) we use (2.12): T_x is 2-hyponormal if and only if

$$\det \begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{2}x \\ \frac{1}{2} & \frac{1}{2}x & \frac{3}{8}x \\ \frac{1}{2}x & \frac{3}{8}x & \frac{3}{10}x \end{pmatrix} \geq 0 \quad \text{and} \quad \det \begin{pmatrix} \frac{1}{2} & \frac{1}{2}x & \frac{3}{8}x \\ \frac{1}{2}x & \frac{3}{8}x & \frac{3}{10}x \\ \frac{3}{8}x & \frac{3}{10}x & \frac{1}{4}x \end{pmatrix} \geq 0,$$

or equivalently, $\frac{63-\sqrt{129}}{80} \leq x \leq \frac{24}{35}$. \square

For perturbations of recursive subnormal shifts of the form $W_{(\sqrt{a}, \sqrt{b}, \sqrt{c})^\wedge}$, subnormality and 2-hyponormality coincide.

Theorem 3.2. Let $\alpha = \{\alpha_n\}_{n=0}^\infty$ be recursively generated by $\sqrt{a}, \sqrt{b}, \sqrt{c}$. If T_x is the weighted shift whose weights are given by $\alpha_x : \alpha_0, \dots, \alpha_{j-1}, \sqrt{x}, \alpha_{j+1}, \dots$, then we have

$$T_x \text{ is subnormal} \iff T_x \text{ is 2-hyponormal} \iff \begin{cases} x = \alpha_j^2 & \text{if } j \geq 1; \\ x \leq a & \text{if } j = 0. \end{cases}$$

Proof. Since α is recursively generated by $\sqrt{a}, \sqrt{b}, \sqrt{c}$, we have that $\alpha_0^2 = a$, $\alpha_1^2 = b$, $\alpha_2^2 = c$,

$$(3.4) \quad \begin{aligned} \alpha_3^2 &= \frac{b(c^2 - 2ac + ab)}{c(b - a)}, \quad \text{and} \\ \alpha_4^2 &= \frac{bc^3 - 4abc^2 + 2ab^2c + a^2bc - a^2b^2 + a^2c^2}{(b - a)(c^2 - 2ac + ab)}. \end{aligned}$$

Case 1 ($j = 0$): It is evident that T_x is subnormal if and only if $x \leq a$. For 2-hyponormality observe by (2.12) that T_x is 2-hyponormal if and only if

$$\det \begin{pmatrix} 1 & x & bx \\ x & bx & bcx \\ bx & bcx & \alpha_3^2 bcx \end{pmatrix} \geq 0,$$

or equivalently, $x \leq a$.

Case 2 ($j \geq 1$): Without loss of generality we may assume that $j = 1$ and $a = 1$. Thus $\alpha_1 = \sqrt{x}$. Then by Theorem 2.1, T_x is subnormal if and only if $x = b$. On the other hand, by (2.12), T_x is 2-hyponormal if and only if

$$\det \begin{pmatrix} 1 & 1 & x \\ 1 & x & cx \\ x & cx & \alpha_3^2 cx \end{pmatrix} \geq 0 \quad \text{and} \quad \det \begin{pmatrix} 1 & x & cx \\ x & cx & \alpha_3^2 cx \\ cx & \alpha_3^2 cx & \alpha_3^2 \alpha_4^2 cx \end{pmatrix} \geq 0.$$

Thus a direct calculation with the specific forms of α_3, α_4 given in (3.4) shows that T_x is 2-hyponormal if and only if $(x - b) \left(x - \frac{b(c^2 - 2c + b)}{b - 1} \right) \leq 0$ and $x \leq b$. Since $b \leq \frac{b(c^2 - 2c + b)}{b - 1}$, it follows that T_x is 2-hyponormal if and only if $x = b$. This completes the proof. \square

4. PROOF OF THEOREM 2.2

With the notation in (2.6), we let

$$p_n := u_n v_{n+1} - w_n \quad (n \geq 0).$$

We then have:

Lemma 4.1. *If $\alpha \equiv \{\alpha_n\}_{n=0}^\infty$ is a strictly increasing weight sequence, then the following statements are equivalent:*

- (i) W_α is 2-hyponormal;
- (ii) $\alpha_{n+1}^2 (u_{n+1} + u_{n+2})^2 \leq u_{n+1} v_{n+2} \quad (n \geq 0)$;
- (iii) $\frac{\alpha_n^2}{\alpha_{n+2}^2} \frac{u_{n+2}}{u_{n+3}} \leq \frac{u_{n+1}}{u_{n+2}} \quad (n \geq 0)$;
- (iv) $p_n \geq 0 \quad (n \geq 0)$.

Proof. This follows from a straightforward calculation. \square

Proof of Theorem 2.2. If α is not strictly increasing, then α is flat, by the argument of [7, Corollary 6], i.e., $\alpha_0 = \alpha_1 = \alpha_2 = \dots$. Then

$$(4.1) \quad D_n(s) = \begin{pmatrix} \alpha_0^2 + |s|^2 \alpha_0^4 & s \alpha_0^3 \\ s \alpha_0^3 & |s|^2 \alpha_0^4 \end{pmatrix} \oplus 0_\infty$$

(cf. (2.5)), so that (2.9) is evident. Thus we may assume that α is strictly increasing, so that $u_n > 0$, $v_n > 0$ and $w_n > 0$ for all $n \geq 0$. Recall that if we write $d_n(t) := \sum_{i=0}^{n+1} c(n, i) t^i$, then the $c(n, i)$'s satisfy the following recursive formulas (cf. (2.8)):

$$(4.2) \quad \begin{aligned} c(n+2, i) &= u_{n+2} c(n+1, i) + v_{n+2} c(n+1, i-1) \\ &\quad - w_{n+1} c(n, i-1) \quad (n \geq 0, 1 \leq i \leq n). \end{aligned}$$

Also, $c(n, n+1) = v_0 \cdots v_n$ (again by (2.8)) and $p_n := u_n v_{n+1} - w_n \geq 0$ ($n \geq 0$), by Lemma 4.1. A straightforward calculation shows that

$$(4.3) \quad \begin{aligned} d_0(t) &= u_0 + v_0 t; \\ d_1(t) &= u_0 u_1 + (v_0 u_1 + p_0) t + v_0 v_1 t^2; \\ d_2(t) &= u_0 u_1 u_2 + (v_0 u_1 u_2 + u_0 p_1 + u_2 p_0) t \\ &\quad + (v_0 v_1 u_2 + v_0 p_1 + v_2 p_0) t^2 + v_0 v_1 v_2 t^3. \end{aligned}$$

Evidently,

$$(4.4) \quad c(n, i) \geq 0 \quad (0 \leq n \leq 2, 0 \leq i \leq n+1).$$

Define

$$\beta(n, i) := c(n, i) - v_0 \cdots v_{i-1} u_i \cdots u_n \quad (n \geq 1, 1 \leq i \leq n).$$

For every $n \geq 1$, we now have

$$(4.5) \quad c(n, i) = \begin{cases} u_0 \cdots u_n \geq 0 & (i = 0), \\ v_0 \cdots v_{i-1} u_i \cdots u_n + \beta(n, i) & (1 \leq i \leq n), \\ v_0 \cdots v_n \geq 0 & (i = n + 1). \end{cases}$$

For notational convenience we let $\beta(n, 0) := 0$ for every $n \geq 0$. □

Claim 1. For $n \geq 1$,

$$(4.6) \quad c(n, n) \geq u_n c(n - 1, n) \geq 0.$$

Proof of Claim 1. We use mathematical induction. For $n = 1$,

$$c(1, 1) = v_0 u_1 + p_0 \geq u_1 c(0, 1) \geq 0,$$

and

$$\begin{aligned} c(n + 1, n + 1) &= u_{n+1} c(n, n + 1) + v_{n+1} c(n, n) - w_n c(n - 1, n) \\ &\geq u_{n+1} c(n, n + 1) + v_{n+1} u_n c(n - 1, n) - w_n c(n - 1, n) \\ &\quad \text{(by the inductive hypothesis)} \\ &= u_{n+1} c(n, n + 1) + p_n c(n - 1, n) \\ &\geq u_{n+1} c(n, n + 1), \end{aligned}$$

which proves Claim 1.

Claim 2. For $n \geq 2$,

$$(4.7) \quad \beta(n, i) \geq u_n \beta(n - 1, i) \geq 0 \quad (0 \leq i \leq n - 1).$$

Proof of Claim 2. We use mathematical induction. If $n = 2$ and $i = 0$, this is trivial. Also,

$$\beta(2, 1) = u_0 p_1 + u_2 p_0 = u_0 p_1 + u_2 \beta(1, 1) \geq u_2 \beta(1, 1) \geq 0.$$

Assume that (4.7) holds. We shall prove that

$$\beta(n + 1, i) \geq u_{n+1} \beta(n, i) \geq 0 \quad (0 \leq i \leq n).$$

For,

$$\begin{aligned} \beta(n + 1, i) + v_0 \cdots v_{i-1} u_i \cdots u_{n+1} &= c(n + 1, i) \quad \text{(by (4.2))} \\ &= u_{n+1} c(n, i) + v_{n+1} c(n, i - 1) - w_n c(n - 1, i - 1) \\ &= u_{n+1} \left(\beta(n, i) + v_0 \cdots v_{i-1} u_i \cdots u_n \right) \\ &\quad + v_{n+1} \left(\beta(n, i - 1) + v_0 \cdots v_{i-2} u_{i-1} \cdots u_n \right) \\ &\quad - w_n \left(\beta(n - 1, i - 1) + v_0 \cdots v_{i-2} u_{i-1} \cdots u_{n-1} \right), \end{aligned}$$

so that

$$\begin{aligned}
 \beta(n + 1, i) &= u_{n+1}\beta(n, i) + v_{n+1}\beta(n, i - 1) - w_n\beta(n - 1, i - 1) \\
 &\quad + v_0 \cdots v_{i-2}u_{i-1} \cdots u_{n-1} (u_n v_{n+1} - w_n) \\
 &= u_{n+1}\beta(n, i) + v_{n+1}\beta(n, i - 1) - w_n\beta(n - 1, i - 1) \\
 &\quad + (v_0 \cdots v_{i-2}u_{i-1} \cdots u_{n-1})p_n \\
 &\geq u_{n+1}\beta(n, i) + v_{n+1}u_n\beta(n - 1, i - 1) - w_n\beta(n - 1, i - 1) \\
 &\quad \text{(by the inductive hypothesis and Lemma 4.1;} \\
 &\quad \text{observe that } i - 1 \leq n - 1, \text{ so (4.7) applies)} \\
 &= u_{n+1}\beta(n, i) + p_n\beta(n - 1, i - 1) \\
 &\geq u_{n+1}\beta(n, i),
 \end{aligned}$$

which proves Claim 2.

By Claim 2 and (4.5), we can see that $c(n, i) \geq 0$ for all $n \geq 0$ and $1 \leq i \leq n - 1$. Therefore (4.4), (4.5), Claim 1 and Claim 2 imply

$$c(n, i) \geq v_0 \cdots v_{i-1}u_i \cdots u_n \quad (n \geq 0, 0 \leq i \leq n + 1).$$

This completes the proof.

5. PROOF OF THEOREM 2.3

To prove Theorem 2.3 we need:

Lemma 5.1 ([15, Lemma 2.3]). *Let $\alpha \equiv \{\alpha_n\}_{n=0}^\infty$ be a strictly increasing weight sequence. If W_α is 2-hyponormal, then the sequence of quotients*

$$(5.1) \quad \Theta_n := \frac{u_{n+1}}{u_{n+2}} \quad (n \geq 0)$$

is bounded away from 0 and from ∞ . More precisely,

$$(5.2) \quad 1 \leq \Theta_n \leq \frac{u_1}{u_2} \left(\frac{\|W_\alpha\|^2}{\alpha_0\alpha_1} \right)^2 \quad \text{for sufficiently large } n.$$

In particular, $\{u_n\}_{n=0}^\infty$ is eventually decreasing.

Proof of Theorem 2.3. By Theorem 2.2, W_α is strictly positively quadratically hyponormal, in the sense that all coefficients of $d_n(t)$ are positive for all $n \geq 0$. Note that finite rank perturbations of α affect a finite number of values of u_n , v_n and w_n . More concretely, if α' is a perturbation of α in the weights $\{\alpha_0, \dots, \alpha_N\}$, then u_n , v_n , w_n and p_n are invariant under α' for $n \geq N + 3$. In particular, $p_n \geq 0$ for $n \geq N + 3$.

Claim 1. *For $n \geq 3$, $0 \leq i \leq n + 1$,*

$$\begin{aligned}
 c(n, i) &= u_n c(n - 1, i) + p_{n-1} c(n - 2, i - 1) \\
 (5.3) \quad &+ \sum_{k=4}^n p_{k-2} \left(\prod_{j=k}^n v_j \right) c(k - 3, i - n + k - 2) + v_n \cdots v_3 \rho_{i-n+1},
 \end{aligned}$$

where

$$\rho_{i-n+1} = \begin{cases} 0 & (i < n - 1), \\ u_0 p_1 & (i = n - 1), \\ v_0 p_1 + v_2 p_0 & (i = n), \\ v_0 v_1 v_2 & (i = n + 1) \end{cases}$$

(cf. [12, Proof of Theorem 4.3]).

Proof of Claim 1. We use induction. For $n = 3$, $0 \leq i \leq 4$,

$$\begin{aligned} c(3, i) &= u_3 c(2, i) + v_3 c(2, i - 1) - w_2 c(1, i - 1) \\ &= u_3 c(2, i) + v_3 \left(u_2 c(1, i - 1) + v_2 c(1, i - 2) - w_1 c(0, i - 2) \right) \\ &\quad - w_2 c(1, i - 1) \\ &= u_3 c(2, i) + p_2 c(1, i - 1) + v_3 \left(v_2 c(1, i - 2) - w_1 c(0, i - 2) \right) \\ &= u_3 c(2, i) + p_2 c(1, i - 1) + v_3 \rho_{i-2}, \end{aligned}$$

where by (4.3),

$$\rho_{i-2} = \begin{cases} 0 & (i < 2), \\ u_0 p_1 & (i = 2), \\ v_0 p_1 + v_2 p_0 & (i = 3), \\ v_0 v_1 v_2 & (i = 4). \end{cases}$$

Now,

$$\begin{aligned} c(n + 1, i) &= u_{n+1} c(n, i) + v_{n+1} c(n, i - 1) - w_n c(n - 1, i - 1) \\ &= u_{n+1} c(n, i) + v_{n+1} \left(u_n c(n - 1, i - 1) + p_{n-1} c(n - 2, i - 2) \right. \\ &\quad \left. + \sum_{k=4}^n p_{k-2} \left(\prod_{j=k}^n v_j \right) c(k - 3, i - n + k - 3) + v_n \cdots v_3 \rho_{i-n} \right) \\ &\quad - w_n c(n - 1, i - 1) \\ &= u_{n+1} c(n, i) + p_n c(n - 1, i - 1) + v_{n+1} p_{n-1} c(n - 2, i - 2) \\ &\quad + v_{n+1} \sum_{k=4}^n p_{k-2} \left(\prod_{j=k}^n v_j \right) c(k - 3, i - n + k - 3) + v_{n+1} \cdots v_3 \rho_{i-n} \\ &\quad \text{(by the inductive hypothesis)} \\ &= u_{n+1} c(n, i) + p_n c(n - 1, i - 1) \\ &\quad + \sum_{k=4}^{n+1} p_{k-2} \left(\prod_{j=k}^{n+1} v_j \right) c(k - 3, i - n + k - 3) + v_{n+1} \cdots v_3 \rho_{i-n}, \end{aligned}$$

which proves Claim 1.

Write $u'_n, v'_n, w'_n, p'_n, \rho'_n$, and $c'(\cdot, \cdot)$ for the entities corresponding to α' . If $p_n > 0$ for every $n = 0, \dots, N + 2$, then in view of Claim 1, we can choose a small perturbation such that $p'_n > 0$ ($0 \leq n \leq N + 2$) and therefore $c'(n, i) > 0$ for all $n \geq 0$ and $0 \leq i \leq n + 1$, which implies that $W_{\alpha'}$ is also positively quadratically hyponormal. If instead $p_n = 0$ for some $n = 0, \dots, N + 2$, careful inspection of (5.3) reveals that without loss of generality we may assume $p_0 = \dots = p_{N+2} = 0$. By Theorem 2.2, we have that for a sufficiently small perturbation α' of α ,

$$(5.4) \quad c'(n, i) > 0 \quad (0 \leq n \leq N + 2, 0 \leq i \leq n + 1) \quad \text{and} \quad c'(n, n + 1) > 0 \quad (n \geq 0).$$

Write

$$k_n := \frac{v_n}{u_n} \quad (n = 2, 3, \dots).$$

Claim 2. $\{k_n\}_{n=2}^\infty$ is bounded.

Proof of Claim 2. Observe that

$$(5.5) \quad \begin{aligned} k_n &= \frac{v_n}{u_n} = \frac{\alpha_n^2 \alpha_{n+1}^2 - \alpha_{n-1}^2 \alpha_{n-2}^2}{\alpha_n^2 - \alpha_{n-1}^2} \\ &= \alpha_n^2 + \alpha_{n-1}^2 + \alpha_n^2 \frac{\alpha_{n+1}^2 - \alpha_n^2}{\alpha_n^2 - \alpha_{n-1}^2} + \alpha_{n-1}^2 \frac{\alpha_{n-1}^2 - \alpha_{n-2}^2}{\alpha_n^2 - \alpha_{n-1}^2}. \end{aligned}$$

Therefore if W_α is 2-hyponormal, then by Lemma 5.1, the sequences

$$\left\{ \frac{\alpha_{n+1}^2 - \alpha_n^2}{\alpha_n^2 - \alpha_{n-1}^2} \right\}_{n=2}^\infty \quad \text{and} \quad \left\{ \frac{\alpha_{n-1}^2 - \alpha_{n-2}^2}{\alpha_n^2 - \alpha_{n-1}^2} \right\}_{n=2}^\infty$$

are both bounded, so that $\{k_n\}_{n=2}^\infty$ is bounded. This proves Claim 2.

Write $k := \sup_n k_n$. Without loss of generality we assume $k < 1$ (this is possible from the observation that $c\alpha$ induces $\{c^2 k_n\}$). Choose a sufficiently small perturbation α' of α such that if we let

$$(5.6) \quad h := \sup_{\substack{0 \leq \ell \leq N+2 \\ 0 \leq m \leq 1}} \left| \sum_{k=4}^{N+4} p'_{k-2} \left(\prod_{j=k}^{N+3} v'_j \right) c'(k-3, \ell) + v'_{N+3} \cdots v'_3 \rho'_m \right|,$$

then

$$(5.7) \quad c'(N+3, i) - \frac{1}{1-k} h > 0 \quad (0 \leq i \leq N+3)$$

(this is always possible because, by Theorem 2.2, we can choose a sufficiently small $|p'_i|$ such that

$$c'(N+3, i) > v_0 \cdots v_{i-1} u_i \cdots u_{N+3} - \epsilon \quad \text{and} \quad |h| < (1-k)(v_0 \cdots v_{i-1} u_i \cdots u_{N+3} - \epsilon)$$

for any small $\epsilon > 0$).

Claim 3. For $j \geq 4$ and $0 \leq i \leq N + j$,

$$(5.8) \quad c'(N + j, i) \geq u_{N+j} \cdots u_{N+4} \left(c'(N + 3, i) - \sum_{n=1}^{j-3} k^n h \right).$$

Proof of Claim 3. We use induction. If $j = 4$, then by Claim 1 and (5.6),

$$\begin{aligned} c'(N + 4, i) &= u'_{N+4} c'(N + 3, i) + p'_{N+3} c'(N + 2, i - 1) \\ &\quad + v'_{N+4} \sum_{k=4}^{N+4} p'_{k-2} \left(\prod_{j=k}^{N+3} v'_j \right) c'(k - 3, i - N + k - 6) \\ &\quad + v'_{N+4} \cdots v'_3 \rho'_{i-(N+3)} \\ &\geq u'_{N+4} c'(N + 3, i) + p'_{N+3} c'(N + 2, i - 1) - v'_{N+4} h \\ &\geq u_{N+4} (c'(N + 3, i) - k_{N+4} h) \\ &\geq u_{N+4} (c'(N + 3, i) - k h), \end{aligned}$$

because $u'_{N+4} = u_{N+4}$, $v'_{N+4} = v_{N+4}$ and $p'_{N+3} = p_{N+3} \geq 0$. Now suppose (5.8) holds for some $j \geq 4$. By Claim 1, we have that for $j \geq 4$,

$$\begin{aligned} c'(N + j + 1, i) &= u'_{N+j+1} c'(N + j, i) + p'_{N+j} c(N + j - 1, i - 1) \\ &\quad + \sum_{k=4}^{N+j+1} p'_{k-2} \left(\prod_{j=k}^{N+j+1} v'_j \right) c'(k - 3, i - N + k - j - 3) \\ &\quad + v'_{N+j+1} \cdots v'_3 \rho'_{i-(N+j)} \\ &= u'_{N+j+1} c'(N + j, i) + p'_{N+j} c(N + j - 1, i - 1) \\ &\quad + \sum_{k=N+5}^{N+j+1} p'_{k-2} \left(\prod_{j=k}^{N+j+1} v'_j \right) c'(k - 3, i - N + k - j - 3) \\ &\quad + \sum_{k=4}^{N+4} p'_{k-2} \left(\prod_{j=k}^{N+j+1} v'_j \right) c'(k - 3, i - N + k - j - 3) \\ &\quad + v'_{N+j+1} \cdots v'_3 \rho'_{i-(N+j)}. \end{aligned}$$

Since $p'_n = p_n > 0$ for $n \geq N + 3$ and $c'(n, \ell) > 0$ for $0 \leq n \leq N + j$ by the inductive hypothesis, it follows that

$$(5.9) \quad p'_{N+j} c(N + j - 1, i - 1) + \sum_{k=N+5}^{N+j+1} p'_{k-2} \left(\prod_{j=k}^{N+j+1} v'_j \right) c'(k - 3, i - N + k - j - 3) \geq 0.$$

By the inductive hypothesis and (5.9),

$$\begin{aligned}
& c'(N+j+1, i) \\
& \geq u'_{N+j+1} c'(N+j, i) + \sum_{k=4}^{N+4} p'_{k-2} \left(\prod_{j=k}^{N+j+1} v'_j \right) c'(k-3, i-N+k-j-3) \\
& \quad + v'_{N+j+1} \cdots v'_3 \rho'_{i-(N+j)} \\
& \geq u_{N+j+1} u_{N+j} \cdots u_{N+4} \left(c'(N+3, i) - \sum_{n=1}^{j-3} k^n h \right) \\
& \quad + v_{N+j+1} v_{N+j} \cdots v_{N+4} \left(\sum_{k=4}^{N+4} p'_{k-2} \left(\prod_{j=k}^{N+3} v'_j \right) c'(k-3, i-N+k-j-3) \right. \\
& \qquad \qquad \qquad \left. + v'_{N+3} \cdots v'_3 \rho'_{i-(N+j)} \right) \\
& \geq u_{N+j+1} u_{N+j} \cdots u_{N+4} \left(c'(N+3, i) - \sum_{n=1}^{j-3} k^n h \right) - v_{N+j+1} v_{N+j} \cdots v_{N+4} h \\
& = u_{N+j+1} u_{N+j} \cdots u_{N+4} \left(c'(N+3, i) - \sum_{n=1}^{j-3} k^n h - k_{N+j+1} k_{N+j} \cdots k_{N+4} h \right) \\
& \geq u_{N+j+1} u_{N+j} \cdots u_{N+4} \left(c'(N+3, i) - \sum_{n=1}^{j-2} k^n h \right),
\end{aligned}$$

which proves Claim 3.

Since $\sum_{n=1}^j k^n < \frac{1}{1-k}$ for every $j > 1$, it follows from Claim 3 and (5.7) that

$$(5.10) \quad c'(N+j, i) > 0 \quad \text{for } j \geq 4 \text{ and } 0 \leq i \leq N+j.$$

It thus follows from (5.4) and (5.10) that $c'(n, i) > 0$ for every $n \geq 0$ and $0 \leq i \leq n+1$. Therefore $W_{\alpha'}$ is also positively quadratically hyponormal. This completes the proof. \square

Corollary 5.2. *Let W_{α} be a weighted shift such that $\alpha_{j-1} < \alpha_j$ for some $j \geq 1$, and let T_x be the weighted shift with weight sequence*

$$\alpha_x : \alpha_0, \dots, \alpha_{j-1}, x, \alpha_{j+1}, \dots.$$

Then $\{x : T_x \text{ is 2-hyponormal}\}$ is a proper closed subset of $\{x : T_x \text{ is quadratically hyponormal}\}$ whenever the latter set is nonempty.

Proof. Write

$$H_2 := \{x : T_x \text{ is 2-hyponormal}\}.$$

Without loss of generality, we can assume that H_2 is nonempty, and that $j = 1$. Recall that a 2-hyponormal weighted shift with two equal weights is of the form $\alpha_0 = \alpha_1 = \alpha_2 = \dots$ or $\alpha_0 < \alpha_1 = \alpha_2 = \alpha_3 = \dots$. Let $x_m := \inf H_2$. By Proposition 6.7 below, T_{x_m} is hyponormal. Then $x_m > \alpha_0$. By assumption, $x_m < \alpha_2$. Thus $\alpha_0, x_m, \alpha_2, \alpha_3, \dots$ is strictly increasing. Now we apply Theorem 2.3 to obtain x' such that $\alpha_0 < x' < x_m$ and $T_{x'}$ is quadratically hyponormal. However $T_{x'}$ is not 2-hyponormal by the definition of x_m . The proof is complete. \square

The following question arises naturally:

Question 5.3. Let α be a strictly increasing weight sequence and let $k \geq 3$. If W_α is a k -hyponormal weighted shift, does it follow that W_α is weakly k -hyponormal under a small perturbation of the weight sequence?

6. OTHER RELATED RESULTS

6.1. Subnormal extensions. Let $\alpha : \alpha_0, \alpha_1, \dots$ be a weight sequence, let $x_i > 0$ for $1 \leq i \leq n$, and let $(x_n, \dots, x_1)\alpha : x_n, \dots, x_1, \alpha_0, \alpha_1, \dots$ be the augmented weight sequence. We say that $W_{(x_n, \dots, x_1)\alpha}$ is an *extension* (or *n-step extension*) of W_α . Observe that

$$W_{(x_n, \dots, x_1)\alpha} |_{\vee\{e_n, e_{n+1}, \dots\}} \cong W_\alpha.$$

The hypothesis $F \neq cP_{\{e_0\}}$ in Theorem 2.1 is essential. Indeed, there exist infinitely many one-step subnormal extensions of a subnormal weighted shift whenever one such extension exists. Recall ([7, Proposition 8]) that if W_α is a weighted shift whose restriction to $\vee\{e_1, e_2, \dots\}$ is subnormal with associated measure μ , then W_α is subnormal if and only if

- (i) $\frac{1}{t} \in L^1(\mu)$;
- (ii) $\alpha_0^2 \leq (\|\frac{1}{t}\|_{L^1(\mu)})^{-1}$.

Also note that there may not exist any one-step subnormal extension of the subnormal weighted shift: for example, if W_α is the Bergman shift, then the corresponding Berger measure is $\mu(t) = t$, and hence $\frac{1}{t}$ is not integrable with respect to μ ; therefore W_α does not admit any subnormal extension. A similar situation arises when μ has an atom at $\{0\}$.

More generally we have:

Theorem 6.1 (Subnormal Extensions). *Let W_α be a subnormal weighted shift with weights $\alpha : \alpha_0, \alpha_1, \dots$ and let μ be the corresponding Berger measure. Then $W_{(x_n, \dots, x_1)\alpha}$ is subnormal if and only if*

- (i) $\frac{1}{t^n} \in L^1(\mu)$;
- (ii) $x_j = \left(\frac{\|\frac{1}{t^{j-1}}\|_{L^1(\mu)}}{\|\frac{1}{t^j}\|_{L^1(\mu)}} \right)^{\frac{1}{2}}$ for $1 \leq j \leq n-1$;
- (iii) $x_n \leq \left(\frac{\|\frac{1}{t^{n-1}}\|_{L^1(\mu)}}{\|\frac{1}{t^n}\|_{L^1(\mu)}} \right)^{\frac{1}{2}}$.

In particular, if we put

$$S := \{(x_1, \dots, x_n) \in \mathbb{R}^n : W_{(x_n, \dots, x_1)\alpha} \text{ is subnormal}\},$$

then either $S = \emptyset$ or S is a line segment in \mathbb{R}^n .

Proof. Write $W_j := W_{(x_n, \dots, x_1)\alpha} |_{\vee\{e_{n-j}, e_{n-j+1}, \dots\}}$ ($1 \leq j \leq n$) and hence $W_n = W_{(x_n, \dots, x_1)\alpha}$. By the argument used to establish (3.2) we have that W_1 is subnormal with associated measure ν_1 if and only if

- (i) $\frac{1}{t} \in L^1(\mu)$;
- (ii) $d\nu_1 = \frac{x_1^2}{t} d\mu$, or equivalently, $x_1^2 = \left(\int_0^{\|W_\alpha\|^2} \frac{1}{t} d\mu(t) \right)^{-1}$.

Inductively W_{n-1} is subnormal with associated measure ν_{n-1} if and only if

- (i) W_{n-2} is subnormal;
- (ii) $\frac{1}{t^{n-1}} \in L^1(\mu)$;

$$(iii) \quad d\nu_{n-1} = \frac{x_{n-1}^2}{t} d\nu_{n-2} = \dots = \frac{x_{n-1}^2 \dots x_1^2}{t^{n-1}} d\mu, \text{ or equivalently,}$$

$$x_{n-1}^2 = \frac{\int_0^{\|W_\alpha\|^2} \frac{1}{t^{n-2}} d\mu(t)}{\int_0^{\|W_\alpha\|^2} \frac{1}{t^{n-1}} d\mu(t)}.$$

Therefore W_n is subnormal if and only if

- (i) W_{n-1} is subnormal;
- (ii) $\frac{1}{t^n} \in L^1(\mu)$;
- (iii) $x_n^2 \leq \left(\int_0^{\|W_\alpha\|^2} \frac{1}{t} d\nu_{n-1}\right)^{-1} = \left(\int_0^{\|W_\alpha\|^2} \frac{x_{n-1}^2 \dots x_1^2}{t^n} d\mu(t)\right)^{-1}$
 $= \frac{\int_0^{\|W_\alpha\|^2} \frac{1}{t^{n-1}} d\mu(t)}{\int_0^{\|W_\alpha\|^2} \frac{1}{t^n} d\mu(t)}.$

□

Corollary 6.2. *If W_α is a subnormal weighted shift with associated measure μ , there exists an n -step subnormal extension of W_α if and only if $\frac{1}{t^n} \in L^1(\mu)$.*

For the next result we refer to the notation in (2.1) and (2.2).

Corollary 6.3. *A recursively generated subnormal shift with $\varphi_0 \neq 0$ admits an n -step subnormal extension for every $n \geq 1$.*

Proof. The assumption about φ_0 implies that the zeros of $g(t)$ are positive, so that $s_0 > 0$. Thus for every $n \geq 1$, $\frac{1}{t^n}$ is integrable with respect to the corresponding Berger measure $\mu = \rho_0 \delta_{s_0} + \dots + \rho_{r-1} \delta_{s_{r-1}}$. By Corollary 6.2, there exists an n -step subnormal extension. □

We need not expect that for arbitrary recursively generated shifts, 2-hyponormality and subnormality coincide as in Theorem 3.2. For example, if $\alpha : \sqrt{\frac{1}{2}}, \sqrt{x}, (\sqrt{3}, \sqrt{\frac{10}{3}}, \sqrt{\frac{17}{5}})^\wedge$, then by (2.12) and Theorem 6.1,

- (i) T_x is 2-hyponormal $\iff 4 - \sqrt{6} \leq x \leq 2$;
- (ii) T_x is subnormal $\iff x = 2$.

A straightforward calculation shows, however, that T_x is 3-hyponormal if and only if $x = 2$; for,

$$A(0; 3) := \begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{2}x & \frac{3}{2}x \\ \frac{1}{2} & \frac{1}{2}x & \frac{3}{2}x & 5x \\ \frac{1}{2}x & \frac{3}{2}x & 5x & 17x \\ \frac{3}{2}x & 5x & 17x & 58x \end{pmatrix} \geq 0 \iff x = 2.$$

This behavior is typical of general recursively generated weighted shifts: we show in [13] that subnormality is equivalent to k -hyponormality for some $k \geq 2$.

6.2. Convexity and closedness. Next, we will show that canonical rank-one perturbations of k -hyponormal weighted shifts which preserve k -hyponormality form a convex set. To see this we need an auxiliary result.

Lemma 6.4. *Let $I = \{1, \dots, n\} \times \{1, \dots, n\}$ and let J be a symmetric subset of I . Let $A = (a_{ij}) \in M_n(\mathbb{C})$ and let $C = (c_{ij}) \in M_n(\mathbb{C})$ be given by*

$$c_{ij} = \begin{cases} c a_{ij} & \text{if } (i, j) \in J \\ a_{ij} & \text{if } (i, j) \in I \setminus J \end{cases} \quad (c > 0).$$

If A and C are positive semidefinite, then $B = (b_{ij}) \in M_n(\mathbb{C})$ defined by

$$b_{ij} = \begin{cases} b a_{ij} & \text{if } (i, j) \in J \\ a_{ij} & \text{if } (i, j) \in I \setminus J \end{cases} \quad (b \in [1, c] \text{ or } [c, 1])$$

is also positive semidefinite.

Proof. Without loss of generality we may assume $c > 1$. If $b = 1$ or $b = c$, the assertion is trivial. Thus we assume $1 < b < c$. The result is now a consequence of the following observation. If $[D]_{(i,j)}$ denotes the (i, j) -entry of the matrix D , then

$$\begin{aligned} \left[\frac{c-b}{c-1} \left(A + \frac{b-1}{c-b} C \right) \right]_{(i,j)} &= \begin{cases} \frac{c-b}{c-1} \left(1 + \frac{b-1}{c-b} c \right) a_{ij} & \text{if } (i, j) \in J, \\ \frac{c-b}{c-1} \left(1 + \frac{b-1}{c-b} \right) a_{ij} & \text{if } (i, j) \in I \setminus J, \end{cases} \\ &= \begin{cases} b a_{ij} & \text{if } (i, j) \in J, \\ a_{ij} & \text{if } (i, j) \in I \setminus J, \end{cases} \\ &= [B]_{(i,j)}, \end{aligned}$$

which is positive semidefinite because positive semidefinite matrices in $M_n(\mathbb{C})$ form a cone. \square

An immediate consequence of Lemma 6.4 is that positivity of a matrix forms a convex set with respect to a fixed diagonal location; i.e., if

$$A_x = \begin{pmatrix} * & * & * \\ & x & * \\ * & * & * \end{pmatrix},$$

then $\{x : A_x \text{ is positive semidefinite}\}$ is convex.

We now have:

Theorem 6.5. *Let $\alpha = \{\alpha_n\}_{n=0}^\infty$ be a weight sequence, let $k \geq 1$, and let $j \geq 0$. Define $\alpha^{(j)}(x) : \alpha_0, \dots, \alpha_{j-1}, x, \alpha_{j+1}, \dots$. Assume W_α is k -hyponormal and define*

$$\Omega_\alpha^{k,j} := \{x : W_{\alpha^{(j)}(x)} \text{ is } k\text{-hyponormal}\}.$$

Then $\Omega_\alpha^{k,j}$ is a closed interval.

Proof. Suppose $x_1, x_2 \in \Omega_\alpha^{k,j}$ with $x_1 < x_2$. Then by (2.11), the $(k+1) \times (k+1)$ Hankel matrix

$$A_{x_i}(n; k) := \begin{pmatrix} \gamma_n & \gamma_{n+1} & \dots & \gamma_{n+k} \\ \gamma_{n+1} & \gamma_{n+2} & \dots & \gamma_{n+k+1} \\ \vdots & \vdots & & \vdots \\ \gamma_{n+k} & \gamma_{n+k+1} & \dots & \gamma_{n+2k} \end{pmatrix} \quad (n \geq 0; i = 1, 2)$$

is positive, where A_{x_i} corresponds to $\alpha^{(j)}(x_i)$. We must show that $tx_1 + (1-t)x_2 \in \Omega_\alpha^{k,j}$ ($0 < t < 1$), i.e.,

$$A_{tx_1+(1-t)x_2}(n; k) \geq 0 \quad (n \geq 0, 0 < t < 1).$$

Observe that it suffices to establish the positivity of the $2k$ Hankel matrices corresponding to $\alpha^{(j)}(tx_1 + (1-t)x_2)$ such that $tx_1 + (1-t)x_2$ appears as a factor in at least one entry but not in every entry. A moment's thought reveals that without loss of generality we may assume $j = 2k$. Observe that

$$A_{z_1}(n; k) - A_{z_2}(n; k) = (z_1^2 - z_2^2) H(n; k)$$

for some Hankel matrix $H(n; k)$. For notational convenience, we abbreviate $A_z(n; k)$ as A_z . Then

$$A_{tx_1+(1-t)x_2} = \begin{cases} t^2 A_{x_1} + (1-t)^2 A_{x_2} + 2t(1-t)A_{\sqrt{x_1x_2}} & \text{for } 0 \leq n \leq 2k, \\ \left(t + (1-t)\frac{x_2}{x_1}\right)^2 A_{x_1} & \text{for } n \geq 2k+1. \end{cases}$$

Since $A_{x_1} \geq 0$, $A_{x_2} \geq 0$ and $A_{\sqrt{x_1x_2}}$ have the form described by Lemma 6.4 and since $x_1 < \sqrt{x_1x_2} < x_2$, it follows from Lemma 6.4 that $A_{\sqrt{x_1x_2}} \geq 0$. Thus evidently, $A_{tx_1+(1-t)x_2} \geq 0$, and therefore $tx_1 + (1-t)x_2 \in \Omega_\alpha^{k,j}$. This shows that $\Omega_\alpha^{k,j}$ is an interval. The closedness of the interval follows from Proposition 6.7 below. \square

In [17] and [18], it was shown that there exists a nonsubnormal polynomially hyponormal operator. Also in [22], it was shown that there exists a nonsubnormal polynomially hyponormal operator if and only if there exists one which is also a weighted shift. However, no concrete weighted shift has yet been found. As a strategy for finding such a shift, we would like to suggest the following:

Question 6.6. Does it follow that the polynomial hyponormality of a weighted shift is stable under small perturbations of the weight sequence?

If the answer to Question 6.6 were affirmative then we would easily find a polynomially hyponormal nonsubnormal (even non-2-hyponormal) weighted shift; for example, if

$$\alpha : 1, \sqrt{x}, (\sqrt{3}, \sqrt{\frac{10}{3}}, \sqrt{\frac{17}{5}})^\wedge$$

and T_x is the weighted shift associated with α , then by Theorem 3.2, T_x is subnormal $\Leftrightarrow x = 2$, whereas T_x is polynomially hyponormal $\Leftrightarrow 2 - \delta_1 < x < 2 + \delta_2$ for some $\delta_1, \delta_2 > 0$ provided the answer to Question 6.6 is yes; therefore for sufficiently small $\epsilon > 0$,

$$\alpha_\epsilon : 1, \sqrt{2+\epsilon}, (\sqrt{3}, \sqrt{\frac{10}{3}}, \sqrt{\frac{17}{5}})^\wedge$$

would induce a non-2-hyponormal polynomially hyponormal weighted shift.

The answer to Question 6.6 for weak k -hyponormality is negative. In fact we have:

Proposition 6.7. (i) *The set of k -hyponormal operators is sot-closed.*

(ii) *The set of weakly k -hyponormal operators is sot-closed.*

Proof. Suppose $T_\eta \in \mathcal{L}(\mathcal{H})$ and $T_\eta \rightarrow T$ in *sot*. Then, by the Uniform Boundedness Principle, $\{\|T_\eta\|\}_\eta$ is bounded. Thus $T_\eta^{*i}T_\eta^j \rightarrow T^{*i}T^j$ in *sot* for every i, j , so that $M_k(T_\eta) \rightarrow M_k(T)$ in *sot* (where $M_k(T)$ is as in (1.2)).

(i) In this case $M_k(T_\eta) \geq 0$ for all η , so $M_k(T) \geq 0$, i.e., T is k -hyponormal.

(ii) Here, $M_k(T_\eta)$ is weakly positive for all η . By (1.3), $M_k(T)$ is also weakly positive, i.e., T is weakly k -hyponormal. \square

REFERENCES

1. A. Athavale, *On joint hyponormality of operators*, Proc. Amer. Math. Soc. **103** (1988), 417–423. MR0943059 (89f:47033)
2. J. Bram, *Subnormal operators*, Duke Math. J. **22** (1955), 75–94. MR0068129 (16:835a)
3. Y.B. Choi, J.K. Han and W.Y. Lee, *One-step extension of the Bergman shift*, Proc. Amer. Math. Soc. **128** (2000), 3639–3646. MR1694855 (2001b:47037)

4. J.B. Conway, *The Theory of Subnormal Operators*, Math. Surveys and Monographs, vol. 36, Amer. Math. Soc., Providence, 1991. MR1112128 (92h:47026)
5. J.B. Conway and W. Szymanski, *Linear combination of hyponormal operators*, Rocky Mountain J. Math. **18** (1988), 695–705. MR0972659 (90a:47059)
6. C. Cowen, *Hyponormal and subnormal Toeplitz operators*, Surveys of Some Recent Results in Operator Theory, I (J.B. Conway and B.B. Morrel, eds.), Pitman Research Notes in Mathematics, Vol. **171**, Longman, 1988, pp. 155–167. MR0958573 (90j:47022)
7. R.E. Curto, *Quadratically hyponormal weighted shifts*, Integral Equations Operator Theory **13** (1990), 49–66. MR1025673 (90k:47061)
8. ———, *Joint hyponormality: A bridge between hyponormality and subnormality*, Proc. Sympos. Pure Math., vol. 51, Part 2, Amer. Math. Soc., Providence, 1990, pp. 69–91. MR1077422 (91k:47049)
9. ———, *An operator theoretic approach to truncated moment problems*, in *Linear Operators*, Banach Center Publications **38** (1997), 75–104. MR1457002 (99c:47014)
10. R.E. Curto and L.A. Fialkow, *Recursiveness, positivity, and truncated moment problems*, Houston J. Math. **17** (1991), 603–635. MR1147276 (93a:47016)
11. ———, *Recursively generated weighted shifts and the subnormal completion problem*, Integral Equations Operator Theory **17** (1993), 202–246. MR1233668 (94h:47050)
12. ———, *Recursively generated weighted shifts and the subnormal completion problem, II*, Integral Equations Operator Theory **18** (1994), 369–426. MR1265443 (94m:47044)
13. R.E. Curto, I.B. Jung and W.Y. Lee, *Extensions and extremality of recursively generated weighted shifts*, Proc. Amer. Math. Soc. **130** (2002), 565–576. MR1862138 (2002i:47041)
14. R.E. Curto and W.Y. Lee, *Joint hyponormality of Toeplitz pairs*, Mem. Amer. Math. Soc. no. 712, Amer. Math. Soc., Providence, 2001. MR1810770 (2002c:47042)
15. ———, *Towards a model theory for 2-hyponormal operators*, Integral Equations Operator Theory **44** (2002), 290–315. MR1933654 (2003m:47036)
16. R.E. Curto, P.S. Muhly and J. Xia, *Hyponormal pairs of commuting operators*, Contributions to Operator Theory and Its Applications (Mesa, AZ, 1987) (I. Gohberg, J.W. Helton and L. Rodman, eds.), Operator Theory: Advances and Applications, vol. 35, Birkhäuser, Basel–Boston, 1988, pp. 1–22. MR1017663 (90m:47037)
17. R.E. Curto and M. Putinar, *Existence of non-subnormal polynomially hyponormal operators*, Bull. Amer. Math. Soc. (N.S.) **25** (1991), 373–378. MR1091568 (93e:47028)
18. ———, *Nearly subnormal operators and moment problems*, J. Funct. Anal. **115** (1993), 480–497. MR1234402 (95d:47024)
19. R.G. Douglas, V.I. Paulsen, and K. Yan, *Operator theory and algebraic geometry*, Bull. Amer. Math. Soc. (N.S.) **20** (1989), 67–71. MR0955316 (90f:47028)
20. P.R. Halmos, *Ten problems in Hilbert space*, Bull. Amer. Math. Soc. **76** (1970), 887–933. MR0270173 (42:5066)
21. ———, *A Hilbert Space Problem Book, 2nd ed.*, Springer, New York, 1982. MR0675952 (84e:47001)
22. S. McCullough and V. Paulsen, *A note on joint hyponormality*, Proc. Amer. Math. Soc. **107** (1989), 187–195. MR0972236 (90a:47062)
23. A. Shields, *Weighted shift operators and analytic function theory*, Math. Surveys **13** (1974), 49–128. MR0361899 (50:14341)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, IOWA CITY, IOWA 52242

E-mail address: rcurto@math.uiowa.edu

DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 151-742, KOREA

E-mail address: wylee@math.snu.ac.kr