

INVARIANTS, BOOLEAN ALGEBRAS AND ACA_0^+

RICHARD A. SHORE

ABSTRACT. The sentences asserting the existence of invariants for mathematical structures are usually third order ones. We develop a general approach to analyzing the strength of such statements in second order arithmetic in the spirit of reverse mathematics. We discuss a number of simple examples that are equivalent to ACA_0 . Our major results are that the existence of elementary equivalence invariants for Boolean algebras and isomorphism invariants for dense Boolean algebras are both of the same strength as ACA_0^+ . This system corresponds to the assertion that $X^{(\omega)}$ (the arithmetic jump of X) exists for every set X . These are essentially the first theorems known to be of this proof theoretic strength. The proof begins with an analogous result about these invariants on recursive (dense) Boolean algebras coding $0^{(\omega)}$.

1. INTRODUCTION

We are interested in measuring the complexity of standard mathematical theorems and constructions in various ways. Our two primary approaches are recursion theoretic and proof theoretic. The first is the domain of recursive mathematics as in Ershov et al. [1998]. The second is that of reverse mathematics as in Simpson [1999]. The typical theorem to be analyzed is an existence theorem of the form $\forall n \in \mathbb{N} \exists m \in \mathbb{N} \dots$ or $\forall f \in \mathbb{N}^{\mathbb{N}} \exists g \in \mathbb{N}^{\mathbb{N}}$. The first corresponds to a function $h : \mathbb{N} \rightarrow \mathbb{N}$ and the second to a functional $H : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$. From the recursion theoretic point of view, we want to know the complexity of h , or of H (in the sense of measuring the complexity of a witness g relative to that of the given f). In either case, we would like lower and upper bounds on the complexity. The standard measure of (relative) complexity here is that of Turing as given by Turing reducibility and the usual benchmarks along the Turing degrees (the recursive functions, $\mathbf{0}$; the halting problem, $\mathbf{0}'$; true arithmetic, $\mathbf{0}^{(\omega)}$; the hyperarithmetic hierarchy to Kleene's \mathcal{O} , etc.). The classical methods of proving high complexity include diagonalization, coding, priority arguments, index set and other hierarchy type theorems.

Reverse mathematics, on the other hand, studies the (proof theoretic) complexity of mathematical theorems by determining the axioms needed to establish them. This involves determining first the appropriate setting for such axioms. The choice is almost always second order arithmetic and one restricts one's interest to countable, or at least "separable" structures. Next, one must isolate particular systems

Received by the editors March 22, 2004.

2000 *Mathematics Subject Classification*. Primary 03B25, 03B30, 03C57, 03D28, 03D35, 03D45, 03F35, 06E05.

The author was partially supported by NSF Grant DMS-0100035.

©2005 Richard A. Shore

of axioms in which one can prove (most of) the classical theorems of countable mathematics. The intuition here is that the axiom systems correspond to a hierarchy of construction or existence principles and are usually graded by types of comprehension (i.e. set existence) assertions. Finally, one hopes to give a precise calibration of the complexity of a given theorem by showing that (relative to some weak base theory of arithmetic) it is actually equivalent to the axiom system used to prove the theorem. (This means that over the base theory one can prove the other axioms of the system from the statement of the classical theorem being studied. This reversal of proving “axioms” from “theorems” is the source of the name, reverse mathematics.)

In fact, the two approaches are closely connected in both their methods and results with each of the standard systems of reverse mathematics corresponding to one of the classical markers of complexity in recursion theory. Proofs of reversals very often correspond to proofs that any witness for particular instances of the existence theorem in question would compute (in the sense of Turing degree) some degree such as $\mathbf{0}'$ or \mathcal{O} . Proofs of the theorem in the systems often correspond to proofs that solutions can be found that are, e.g. recursive, recursive in $\mathbf{0}'$, etc. At times both proof theoretic (e.g. conservation results) and model theoretic (e.g. nonstandard models) are used as well. We will list the standard systems of reverse mathematics and explain their recursion theoretic counterparts in the next section. Our goal will then be to explore a type of mathematical classification of structures – the existence of invariants – that often provides central and fundamental theorems and is naturally related to issues of complexity. This type of theorem, however, presents unusual challenges to the subject of reverse mathematics.

Typically, such theorems begin with a class \mathcal{K} of structures which we want to classify up to some equivalence relation. The classification is embodied by an assertion of the form that there is an assignment of invariants to members of the class such that two structures have the same invariant if and only if they are equivalent. For this to have some content the invariants (and equality on them) should be somehow simpler than the structures (and the equivalence relation of interest on them). The archetypical equivalence relation is, of course, isomorphism and the invariant is generally viewed as a type of dimension or degree. Standard examples here include vector spaces (dimension) and algebraically closed fields (transcendence degree). Of course, other equivalence relations are studied as well and one looks for invariants up to homotopy, homeomorphism, elementary equivalence or other restricted types of identification of structures. Frequently, the invariants are cardinal numbers corresponding to the size of some sort of basis for the structure. If the structures are countable, then these are numbers in ω or ω itself. At other times (e.g. Ulm invariants for p -groups) the analysis of the structures proceeds by some kind of ordinal decomposition and the invariants are then ordinals (below ω_1 for countable structures). In other situations the invariants may be real numbers or other objects, although in the countable case these may be as complicated as the structures themselves and so do not provide much information.

In any case, formalizing the existence of invariants for a class \mathcal{K} of even countable mathematical structures is beyond the scope of the setting for reverse mathematics. It typically asserts the existence of a functional \mathcal{F} from a class \mathcal{K} of mathematical structures to the set of invariants such as \mathbb{N} , \aleph_1 , etc., that classifies \mathcal{A} up to \equiv , the equivalence relation of interest, in the sense that $F(\mathcal{A}) = F(\mathcal{B}) \Leftrightarrow \mathcal{A} \equiv \mathcal{B}$. Even

when the invariants are numbers and \mathcal{K} is definable in second order arithmetic this basic assertion of the existence of invariants for \mathcal{K} is a third order existential one as \mathcal{F} is a map from an uncountable class of countable structures:

$$(*) \exists (\mathcal{F} : \mathcal{K} \rightarrow \mathbb{N})(\forall A_1, A_2 \in \mathcal{K})(F(A_1) = F(A_2) \Leftrightarrow A_1 \equiv A_2).$$

(We denote this as $(*\mathcal{K}, \equiv)$ when we want to make the class and equivalence relation explicit.) Reverse mathematics only allows one to discuss numbers and sets of numbers, not sets of sets of numbers as are needed here. Thus to even begin our analysis we must first find a way to capture the assertion in some second order way.

In the standard direction of proving the existence of invariants there is usually not much to worry about. A classical proof provides an analysis that specifies a particular procedure for identifying the invariant. One can analyze first this procedure as recursive, arithmetic, hyperarithmetic, etc., and then (in terms of axiom systems) the proof that the procedure produces the same invariant for two structures if and only if they are equivalent. Thus, the assertion that this particular procedure works as desired can be proven in some appropriate subsystem Ax of second order arithmetic, and so gives us an upper bound on the (recursion and proof theoretic) complexity of the existence of invariants. Formally, we produce a formula $\Phi(A, n)$ of second order arithmetic such that $Ax \vdash \phi$ where ϕ is the sentence saying that Φ properly defines the invariant n associated with A :

$$\begin{aligned} \phi : & (\forall A \in \mathcal{K})(\exists n \in \mathbb{N})\Phi(A, n) \& \\ & (\forall A_1, A_2 \in \mathcal{K})(\forall n_1, n_2 \in \mathbb{N})(\Phi(A_1, n_1) \& \Phi(A_2, n_2) \rightarrow (n_1 = n_2 \leftrightarrow A_1 \equiv A_2)). \end{aligned}$$

Here we are assuming that our class \mathcal{K} is definable in second order arithmetic and use $A \in \mathcal{K}$ as an abbreviation for quantification over sets satisfying this definition. Similarly, we are assuming that the equivalence relation \equiv being considered is also definable in second order arithmetic. In all the cases we consider, coding the class of structures of interest in second order arithmetic will be routine and we will simply assume that we have such a coding. The notion of isomorphisms is even easier. That of elementary equivalence will, however, take some care (see Definitions 4.5–4.7).

For example, consider the case where \mathcal{K} is the set of countable (and so, without loss of generality, with domain \mathbb{N}) vector spaces over the rationals and the equivalence relation of interest is isomorphism. Here the invariant is the dimension and the required definition Φ says that n (possibly ω) is the cardinality of a maximal independent set of vectors. As each of the questions of whether there exists an independent set of size n is at the same level of complexity (Σ_2^0), arithmetic comprehension (or the iterated Turing jump) is clearly sufficient to produce the answer and so the invariant. (The proof that two vector spaces with the same dimension are isomorphic is even simpler.) Thus we have a sentence ϕ as required above for the class of vector spaces which is provable using arithmetic comprehension (ACA_0).

For the reverse direction we need some second order consequence of the existence of invariants that implies the axiom system Ax used to prove the correctness of the particular procedure used in the forward direction. Our proposal is to use the special case of the theorem for countable subclasses of our class of structures. More precisely, we consider the following sentence which will play a central role in our analysis and to which we will refer repeatedly as ψ :

$$\psi : \forall \langle A_i \mid i \in \mathbb{N} \rangle (\forall i (A_i \in \mathcal{K}) \rightarrow \exists \langle n_i \mid i \in \mathbb{N} \rangle (\forall i (n_i \in \mathbb{N}) \& \forall i, j \in \mathbb{N} (A_i \equiv A_j \leftrightarrow n_i = n_j))).$$

(We are again using membership in \mathcal{K} as a shorthand for its definition.) This assertion is obviously weaker than the existence of number invariants for the class \mathcal{K} . We can now hope to prove that it implies whatever system was used to prove the forward direction of the existence result (in the sense of providing a specific procedure defining the invariant from the structure). In the case of vector spaces over \mathbb{Q} with isomorphism, it is not difficult to show (Theorem 2.1) that, for this choice of \mathcal{K} and \equiv , ψ does imply arithmetic comprehension in RCA_0 , the standard weak base theory of reverse mathematics. Thus we are justified in asserting that the existence of number invariants for isomorphism of countable vector spaces over \mathbb{Q} is equivalent to arithmetic comprehension (ACA_0).

We believe the plan proposed here should be intuitively convincing for the claim that we capture the strength of the existence of invariants by these results (at least for the specific choices of ϕ and ψ that we present). From a recursion theoretic point of view, there will be no difficulty in that we can (and do) show that the procedure for computing the invariant of A is recursive in the recursion theoretic counterpart of Ax (the jump, the ω -jump or the hyperjump of A as the case may be). On the other hand, we can (and will) show that for any given set X there are instances of ψ for sequences $\langle A_i \rangle$ recursive in X for which any witness sequence $\langle n_i \rangle$ computes the corresponding jump of X .

For the proof theoretic point of view, however, we should say a few words about the formalities of what types of sentences should be acceptable and where the “mathematical” proofs that $\phi \rightarrow (*)$ and $(*) \rightarrow \psi$ take place. The issue here is, as Carl Jockusch pointed out to us, that in the cases of interest all these sentences are theorems of ZFC. What would then prevent us from choosing any theorems of ZFC (expressible in second order arithmetic) as ϕ and ψ ? Such choices, of course, would be nonsensical and irrelevant to any claims of capturing the complexity of some instance of $(*)$. (All the other ingredients in our claim for the equivalence of $(*)$ and Ax are assertions in second order arithmetic and can take place unproblematically in RCA_0 .)

First we suggest that any possible dependence on the mathematical content of the assertion be removed by making the implication $\phi \rightarrow (*)$ independent of the choices of \mathcal{K} and \equiv . Formally we require that

$$(1.1) \quad \text{ZFC} \vdash (\forall \text{ class of structures } \mathcal{K})(\forall \text{ equivalence relations } \equiv \text{ on } \mathcal{K})[\phi \rightarrow (*)]$$

where here, of course, we leave $A \in \mathcal{K}$ in ϕ as a formula of set theory. This seems to be sufficient to support the claim that ϕ is at least as strong as $(*)$ and is certainly true of the class of sentences that we have proposed as candidates for ϕ .

We should, of course, make the same requirement for the implication $(*) \rightarrow \psi$ but this is not sufficient. Here we are asserting that ψ contains some of the strength of $(*)$ but want to also prove that it is itself strong. We must take care not to allow some theorem of ZFC (not mentioning \mathcal{K} or \equiv say) as a candidate that itself proves Ax without any connection to $(*)$. Indeed the sentence ψ that we have proposed to use is a theorem of ZFC for every \mathcal{K} and \equiv . (As there are only countably many A_i , there are only countably many equivalence classes, and so we can number them accordingly.) Thus our worry here is that the strength asserted for ψ has no connection to its dependence on $(*)$ but is induced by the use of comprehension in ZFC. To eliminate this possibility we require that the proof that $(*) \rightarrow \psi$ take place in a version of set theory with very limited comprehension. A very strong version of this proposal is that the implication be provable in rudimentary set theory. This

is the system B_0^{set} of Simpson [1999, VII.3] whose only axioms are extensionality, infinity and that the universe is closed under the rudimentary operations which are essentially the standard ones for the primitive recursive functions on \mathbb{N} without the recursion scheme. This system corresponds to Δ_0 comprehension (see Jensen [1972]) so no application of comprehension in the metatheory can be the source of the strength of ψ . Rather, the only source of the power of comprehension is the existential assertion of $(*)$ itself. For our proposed ψ , it is clear that the implication $(*) \rightarrow \psi$ is provable with only Δ_0 comprehension. It only asserts the existence of the composition of two given functions – \mathcal{F} given by $(*)$ and the one from $\omega \rightarrow \omega$ corresponding to the given sequence $\langle A_i \rangle$ – and so is easily seen to exist by Δ_0 comprehension. A final desideratum for ψ might be that the axiom system Ax claimed to be of the same strength as $(*)$ also prove ψ (all for the same choice of \mathcal{K} and \equiv , of course).

Thus we will be willing to claim that the existence of invariants for some \mathcal{K} and \equiv has the same strength as an axiom system Ax (of second order arithmetic) if we have formulas $\phi(\mathcal{K}, \equiv)$ and $\psi(\mathcal{K}, \equiv)$ (both of third order arithmetic or its set theoretic analog but with the only nonsecond order atomic subformulas being of the form $A \in \mathcal{K}$ and $A_1 \equiv A_2$ and no third order quantification allowed) such that $RCA_0 \vdash Ax \rightarrow \phi$ and $RCA_0 \vdash \psi \rightarrow Ax$ (both with appropriate definitions replacing the occurrences of \mathcal{K} and \equiv), $Ax \vdash \psi$, $ZFC \vdash (\forall \text{class of structures } \mathcal{K})(\forall \text{equivalence relations } \equiv \text{ on } \mathcal{K})[(*) \rightarrow \phi]$ and $B_0^{set} \vdash (*) \rightarrow \psi$.

Now that we have an approach to analyzing the strength of the existence of invariants for different classes of structures we can consider particular classes of interest. As we have indicated, the case of vector spaces is easily seen to be equivalent to the standard system ACA_0 (Theorem 2.1). The situation is the same for algebraically closed fields (Theorem 2.2). Our goal here is to consider a class that presents a second challenge to the standard approach of reverse mathematics by being provably equivalent to an axiomatic system that lies strictly between two of the standard systems. The class of structures that we analyze is that of Boolean algebras. Now the isomorphism relation for arbitrary Boolean algebras is Σ_1^1 complete (see §3), so no simply defined invariants for isomorphism can exist. Indeed, the existence of invariants for isomorphisms, even for countable sequences of structures in the sense of ψ , implies $\Pi_1^1\text{-CA}_0$, the system of reverse mathematics corresponding to the existence of Kleene's \mathcal{O} (or equivalently the hyperjump). On the other hand, $\Pi_1^1\text{-CA}_0$ is enough to prove ψ for isomorphism for any class \mathcal{K} . However, it is a classical result of Tarski [1949] (see Monk [1989, 18] or Goncharov [1997, 2.3]) that there are number invariants for elementary equivalence of Boolean algebras. Our goal is to prove (Theorem 6.10) that the existence of any number invariants, in the sense of $(*)$, is equivalent (over RCA_0), in the sense of our analysis above in terms of ϕ and ψ , to an axiom system ACA_0^+ that is intermediate between two of the standard systems, ACA_0 and ATR_0 . We believe that this is the first real example of a theorem known to be equivalent to ACA_0^+ . This system corresponds to the existence of the arithmetic jump, $A^{(\omega)}$, for each set A and the recursion theoretic version of our main theorem (Theorem 5.10) is that determining the elementary invariants for recursive Boolean algebras is of the same complexity (in the sense of Turing degree) as $0^{(\omega)}$.

Some might say that the use of the logical notion of elementary equivalence is the source of the problem and that it alone might require ACA_0^+ . Indeed, one must

be careful to develop a reasonable notion of elementary equivalence that makes sense in RCA_0 . Following the approach to satisfaction from Simpson [1999], we do this in Definitions 4.5–4.7. Another objection might stem from a possible general aversion (which we do not share) to logical notions as the proper objects of study for reverse mathematics. This concern can be alleviated by noting that these invariants are, in fact, isomorphism invariants for an algebraically defined subclass of Boolean algebras. These are the dense Boolean algebras as presented in Goncharov [1997]. The required algebraic notions are given in Definitions 4.1–4.3. Intuitively these algebras play the same role for Boolean algebras that dense linear orderings play for linear orderings. The prime example is the atomless Boolean algebra but there are, of course, countably many others. (From the logical or model theoretic point of view these are just the saturated countable Boolean algebras.) We will show that the existence of isomorphism invariants for this class also implies the existence of the ω -jump, and so has the same strength as the existence of elementary equivalence invariants for all countable Boolean algebras. These results will be Theorems 5.10 and 5.19 for the recursion theoretic version and Theorems 6.6 and 6.9 for the reverse mathematics one.

2. AXIOM SYSTEMS AND EXAMPLES

All our axiom systems for reverse mathematics are phrased in terms of the language of second order arithmetic, that is the usual first order language of arithmetic augmented by set variables and the membership relation \in . Each system also contains the standard basic axioms for $+$, \cdot and $<$ (which say that \mathbf{N} is an ordered semiring). In addition, they all include one of two forms of induction. The first form is the full induction scheme **I**.

$$(\mathbf{I}): \varphi(0) \wedge \forall n(\varphi(n) \rightarrow \varphi(n+1)) \rightarrow \forall n\varphi(n),$$

for every formula. The other form of induction that we consider is an axiom that permits the application of induction only to sets (that one knows to exist):

$$(\mathbf{I}_0): 0 \in X \wedge \forall n(n \in X \rightarrow n+1 \in X) \rightarrow \forall n(n \in X).$$

We call the system consisting of **(I)** and the basic axioms of ordered semirings P . If we replace **(I)** by **(I₀)**, we call the system P_0 . In general, if we have any system S containing **(I)**, we denote by S_0 the system in which **(I)** is replaced by **(I₀)**. All the systems we shall consider will be defined by adding various types of set existence axioms to P or P_0 . We begin with a base theory strong enough to do basic arithmetic and elementary combinatorics.

(RCA): RCA, for recursive comprehension axiom, is a system just strong enough to prove the existence of the recursive sets but not of \emptyset' nor indeed of any nonrecursive set. In addition to P its axioms include the schemes of Δ_1^0 comprehension and Σ_1^0 induction:

$$(\Delta_1^0\text{-CA}_0) \forall n(\varphi(n) \leftrightarrow \psi(n)) \rightarrow \exists X \forall n(n \in X \leftrightarrow \varphi(n))$$

for all Σ_1^0 formulas φ and Π_1^0 formulas ψ in which X is not free.

$$(\Sigma_1^0\text{-I}) (\varphi(0) \wedge \forall n(\varphi(n) \rightarrow \varphi(n+1))) \rightarrow \forall n\varphi(n)$$

for all Σ_1^0 formulas φ .

The next system says that every infinite binary tree has an infinite path. It is connected to the low basis theorem of recursion theory which says that every such tree has an infinite path whose jump is recursive in that of the tree itself.

(WKL): WKL, for weak König's lemma, asserts, in addition to RCA, that every infinite subtree of $2^{<\omega}$ has an infinite path.

We next move up to arithmetic comprehension.

(ACA): ACA, for arithmetic comprehension axioms, consists of P plus the comprehension scheme for arithmetic formulas:

(ACA) $\exists X \forall n (n \in X \leftrightarrow \varphi(n))$ for every arithmetic formula φ in which X is not free.

In recursion theoretic terms, ACA₀ proves the existence of \emptyset' and by relativization it proves and, in fact is equivalent to, the existence of X' for every set X . The next system corresponds to the existence of all (relativized) H -sets, i.e. the existence of the H_e^X (and so the hyperarithmetical hierarchy up to e) for each $e \in \mathcal{O}^X$ for every set X .

(ATR): ATR, for arithmetical transfinite recursion, consists of P plus the assertion that arithmetic comprehension can be iterated along any countable well ordering.

(ATR) If X is a set coding a well ordering $<_X$ with domain D and Y is a code for a set of arithmetic formulas $\varphi_x(z, Z)$ (indexed by $x \in D$) each with one free set variable and one free number variable, then there is a sequence $\langle K_x | x \in D \rangle$ of sets such that if y is the immediate successor of x in $<_X$, then $\forall n (n \in K_y \leftrightarrow \varphi_x(n, K_x))$ and if x is a limit point in $<_X$, then K_x is $\bigoplus \{K_y | y <_X x\}$.

The systems climbing up to full second order arithmetic (i.e. comprehension for all formulas) are classified by the syntactic level of the second order formulas for which we assume a comprehension axiom.

(Π_n^1 -CA): Π_n^1 -CA, for Π_n^1 comprehension axiom, is the system P plus comprehension for Π_n^1 formulas:

(Π_n^1 -CA) $\exists X \forall k (k \in X \leftrightarrow \varphi(k))$ for every Π_n^1 formula φ in which X is not free.

The recursion theoretic equivalent of the simplest of these systems, Π_1^1 -CA, is the existence of \mathcal{O}^X , the hyperjump of X , for every set X . Together with the four systems listed above, it makes up the standard list of axiomatic stems of reverse mathematics. Almost all theorems of classical mathematics whose proof theoretic complexity has been determined have turned out to be equivalent to one of them.

As simple examples of our approach to invariants, we now analyze the existence of number invariants for isomorphism of vector spaces over a fixed field and algebraically closed fields of fixed characteristic.

Theorem 2.1. *Let F be a field and \mathcal{V} the class of countable vector spaces over F . The existence of number invariants for the isomorphism relation on \mathcal{V} , $(*\mathcal{V}, \cong)$, is of the same complexity as ACA₀. This is also true if we restrict our class to vector spaces of any fixed set of dimensions as long as the set has cardinality at least 2.*

Proof. Our formula says that V has dimension $n - 1$ for $n > 0$ and ∞ for $n = 0$:

$\phi(V, n)$:

$$[n > 0 \ \& \ \exists \langle x_2, \dots, x_n \rangle \in V^{n-1} \forall \langle c_2, \dots, c_n \rangle \in F^{n-1} (c_2 x_2 + \dots + c_n x_n = 0 \Leftrightarrow c_2 = \dots = c_n = 0) \ \& \ \neg \exists \langle x_1, \dots, x_n \rangle \in V^n \forall \langle c_1, \dots, c_n \rangle \in F^n (c_1 x_1 + \dots + c_n x_n = 0 \Leftrightarrow c_1 = \dots = c_n = 0)] \vee [n = 0 \ \& \ \forall m [\exists \langle x_2, \dots, x_m \rangle \in V^{m-1} \forall \langle c_2, \dots, c_m \rangle \in F^{m-1} (c_2 x_2 + \dots + c_m x_m = 0 \Leftrightarrow c_2 = \dots = c_m = 0)]]$$

One can easily develop enough of the usual theory of vector spaces in ACA₀ to prove that every vector space has a unique dimension (in this sense) and that any two of the same dimension are isomorphic. (See Simpson [1999, III.4] for a sketch and Friedman, Simpson and Smith [1983] for more details.) For the other direction, consider any two dimensions $n < m$ and the class \mathcal{K} of vector spaces with dimensions in A with $n, m \in A$ (where we allow ∞ as a value of m for the countable vector space of largest dimension). We need to build a sequence V_i of vector spaces

such that any witness to ψ will compute $0'$ (and so, by relativization, ψ implies the existence of X' for every X). We take V_0 and V_1 to be fixed spaces of dimension n and m , respectively. We define the rest of the V_i uniformly recursively so that we build V_i as a space of dimension n until we see that $i \in 0'$ at which point we start extending the space so that at the end it will have dimension m . Now if $\langle n_i \rangle$ is any witness for $\langle V_i \rangle$ as required by ψ , it is clear that, for $i > 1$, $i \in 0' \Leftrightarrow n_i = m$. \square

Theorem 2.2. *Let \mathcal{A} be the class of countable algebraically closed fields of characteristic p . The existence of number invariants for the isomorphism relation on \mathcal{A} , $(*\mathcal{A}, \cong)$, is of the same complexity as ACA_0 . This is also true if we restrict our class to fields of any fixed set of transcendence degrees as long as the set has cardinality at least 2.*

Proof. The proof is the same as for vector spaces except that we replace linear (in)dependence by algebraic (in)dependence (over the prime field). \square

We next introduce the system ACA^+ that captures the instances of the existence of elementary invariants for Boolean algebras that we analyze. Recursion theoretically, it corresponds to the existence of the arithmetic (or ω) jump of X for every set X .

(ACA⁺): $\forall X \exists A (A^{[0]} = X \ \& \ (\forall n) [A^{[n+1]} = (A^{[n]})']$.

As an axiomatic system in reverse mathematics ACA^+ was first studied in Blass, Hirst and Simpson [1987]. They proved in RCA_0 that Hindman's theorem implies ACA_0 and is implied by ACA_0^+ . The precise strength of this theorem is still open. We do not know of any other theorems that are even candidates for equivalence with ACA_0^+ . (We are leaving out uniform infinite versions of theorems that are each equivalent to ACA_0 such as Ramsey's theorem for n -tuples, RT_2^n . Each one for $n \geq 3$, is equivalent to ACA_0 in such a way that the assertion that for any sequence F_n of colorings of n -tuples there is a sequence H_n so that each H_n is homogeneous for F_n is equivalent to ACA_0^+ .)

3. BOOLEAN ALGEBRAS

We take the standard definition of the class \mathcal{B} of Boolean algebras B with constants 0 and 1 and operations \vee, \wedge and \neg . For convenience we also use the (Δ_0) defined relation \leq and function $-$: $x \leq y \Leftrightarrow x \wedge y = x$ and $x - y = x \wedge \neg y$. A general reference for information about Boolean algebras is Monk [1989].

As we mentioned in the introduction, the isomorphism problem for Boolean algebras is Σ_1^1 complete, and so has no reasonable notion of invariant. In fact, there is a single recursive Boolean algebra B (the Harrison Boolean algebra) such that for any Σ_1^1 formula $\Omega(n)$ there is a recursive sequence of algebras B_n such that $\Omega(n) \Leftrightarrow B_n \cong B$. (This result is presumably folklore. A proof for the analogous statement about linear orderings can be found in White [2000, 5.4]. The result for Boolean algebras follows by taking the interval algebras (Definition 5.3) of the linear orderings constructed there.) Thus the existence of invariants even for all countable sets of Boolean algebras up to isomorphism, i.e. ψ for \mathcal{B} and \cong , implies $\Pi_1^1\text{-CA}_0$. Of course, ψ is provable in $\Pi_1^1\text{-CA}_0$ for any (definable) class \mathcal{K} of structures as isomorphism is a Σ_1^1 relation. Thus there is not much to say about isomorphism invariants for arbitrary classes of Boolean algebras. We will investigate a coarser relation – elementary equivalence – on all Boolean algebras and isomorphism restricted to a subclasses of Boolean algebras – the “dense” Boolean algebras.

Tarski [1949] showed that there are only countably many completions T_i of the theory T of Boolean algebras and they are uniformly axiomatizable. Thus the theory of Boolean algebras is decidable. This, of course, supplies number invariants for elementary equivalence of countable Boolean algebras in the sense of (*): $\mathcal{F}(A) = i \Leftrightarrow A \models T_i$ and so $\mathcal{F}(A) = \mathcal{F}(B) \Leftrightarrow A \equiv B$. Our tasks are now first to analyze a proof of this theorem to see that we can define this relation $\Phi(A, i)$ and prove in ACA_0^+ that it has the properties described in ϕ . For the reversal we want to construct a uniformly recursive sequence B_i of Boolean algebras so that any witness function assigning them number invariants for elementary equivalence in the sense of ψ computes $0^{(\omega)}$. Together these will prove our main result:

Theorem 6.10. *The existence of number invariants for elementary equivalence of countable Boolean algebras or for isomorphism of dense countable Boolean algebras has the same complexity in the sense of reverse mathematics as ACA_0^+ .*

We begin with a series of definitions that are needed just to describe the theories T_i and the Tarski invariants.

Definition 3.1. Let a be an element of a Boolean algebra B . a is an *atom* if and only if $0 < a$ but $\neg\exists x(0 < x < a)$. The set of atoms of B is denoted by $At(B)$. The element a is *atomic* if and only if $\forall x \leq a \exists z \leq x (x \neq 0 \rightarrow z \text{ is an atom})$. It is *atomless* if and only if no $x \leq a$ is an atom.

Now Tarski's analysis of a Boolean algebra B proceeds by taking successive quotients by uniformly defined ideals much the way the Cantor-Bendixson analysis of a topological space or a Boolean algebra works (Monk [1989, 6]). The difference is that the Cantor-Bendixson analysis uses at each step the ideal generated by atoms. Now, Tarski's analysis uses a larger ideal. We follow the analysis as in Monk [1989, 7] which is very neat or Goncharov [1997] which is more constructive, and so often better suited to our purposes. We refer to these sources for many basic algebraic facts. For example, the ideal we want is generated by the atomic and the atomless elements. That it can also be defined via single joins is 2.2.1 of Goncharov [1997].

Definition 3.2. For B a Boolean algebra, $I(B) = \{x \in B \mid \exists y, z (y \text{ is atomic} \ \& \ z \text{ is atomless} \ \& \ x = y \vee z)\}$. We let $B' = B/I(B)$. We call B' the *Tarski derivative* of B .

We now define the sequence of Boolean algebras constructed by successively taking quotients by this ideal.

Definition 3.3. Given a Boolean algebra B we define a sequence $B^{[n]}$ of Boolean algebras and a sequence I_n of ideals of B as follows: $B^{[0]} = B$; $B^{[n+1]} = B^{[n]}/I(B^{[n]})$. The canonical embeddings are given by $\pi_{m,n} : B^{[m]} \rightarrow B^{[n]}$ for $m \leq n$. We denote $\pi_{0,n}$ by π_n . The desired sequence of ideals of B is given by $I_0(B) = \{0\}$, $I_{n+1}(B) = \pi_n^{-1}[I(B^{[n]})]$ so $B^{[n]} \cong B/I_n(B)$.

We can now define the elementary invariants for Boolean algebras as a triple of natural numbers or symbols $-1, \infty$ encoding the level at which this process stops and the type of the final nonzero algebra produced.

Definition 3.4. The invariant of the Boolean algebra B , $inv(B)$, is a triple of numbers (or symbols $-1, \infty$):

- (1) $inv(B) = \langle -1, 0, 0 \rangle$ if $B = \{0\}$;
- (2) $inv(B) = \langle \infty, 0, 0 \rangle$ if $\forall i (B^{[i]} \neq \{0\})$;

- (3) $inv(B) = \langle k, l, m \rangle$ if
- (a) $B^{[k]} \neq \{0\}, B^{[k+1]} = \{0\}$,
 - (b) $l = \min\{\infty, \|At(B^{[k]})\|\}$,
 - (c) $m = 0$ if $B^{[k]}$ is atomic and $m = 1$ otherwise.

We denote the entries of this triple by $inv_i(B)$ for $i = 1, 2, 3$.

Tarski showed that these invariants characterize all Boolean algebras up to elementary equivalence

Theorem 3.5 ([1949, Tarski]). *For all Boolean algebras B and A , $B \equiv A \Leftrightarrow inv(B) = inv(A)$.*

We now wish to argue that we can prove the existence of the map inv as a definable procedure and the fact that it characterizes elementary equivalence in ACA_0^+ . It should be clear that in ACA_0 we can show for every Boolean algebra B that $At_0(B)$, the sets of atomic and atomless elements of B , $I(B)$, $B/I(B)$ and the canonical map $\pi : B \rightarrow B/I(B)$ all exist. The sequence of these predicates and maps over all n , however, do not obviously provably exist in ACA_0 . Indeed, even the existence of all “finite” initial segments of these sequences does not seem to be provable in ACA_0 although the existence for the sequences of length n are, of course, provable for all n in ACA by induction. This will be an important issue in our consideration of reversals but for now it is sufficient to note that all these (even infinite) sequences of subsets of B provably exist in ACA_0^+ . The point is that they are all uniformly recursive in $B^{(\omega)}$. Thus the map inv is recursive in $B^{(\omega+1)}$. (We need to quantify over the sequences $B^{[n]}, I_n$ to check if $inv_1(B) = \infty$. Otherwise, we just need to check the individual $B^{[n]}$ for their cardinality and atomicity and each one is recursive in a finite (in the model) number of jumps, and so in $B^{(\omega)}$ (in the model). Thus we can define inv and prove that the definition defines a function in ACA_0^+ .)

To finish the proof of Theorem 3.5 in ACA_0^+ we have to define elementary equivalence, \equiv , and go through a suitable proof. Again in, say RCA_0 , the definition of \equiv is delicate and we will deal with it more carefully when we prove the reversals in Definitions 4.5–4.7. For now, it is straightforward to check that any reasonably constructive standard inductive definition of satisfaction for a structure A for a recursive language L can be carried out uniformly recursively in $A^{(\omega)}$, and so we can adopt the standard definition of \equiv as $A \equiv B \Leftrightarrow \forall \rho \in L(A \models \rho \leftrightarrow B \models \rho)$. The proof of Tarski’s theorem itself requires some care. The most commonly found proofs (as in Chang and Keisler [1977, 5.5.10] or Monk [1989, 18] use saturated or special models of some sort that rely on the existence of uncountable cardinals. Tarski’s original proof (according to Doner and Hodges [1988]) was a quantifier elimination proof in an expanded language and presumably the most constructive of them all. For the purposes of our argument in ACA_0^+ , the easiest route to follow is the one done in detail in Goncharov [1977, 2.3], as it can be straightforwardly carried out in ACA_0^+ . It is also the one defining the dense Boolean algebras and explicitly dealing with the isomorphism problem for that class. We now turn to these matters and the proof of Tarski’s theorem in ACA_0^+ .

4. DENSE BOOLEAN ALGEBRAS

We begin with the analog of our invariants for individual elements and formal definitions in the language of Boolean algebras of the predicates we will use to

describe the sequences of algebras and ideals of Definition 3.3. These will be the key ingredients in the proof of existence of isomorphism invariants for the dense Boolean algebras and the associated proof of Tarski’s theorem.

Definition 4.1. The invariant, $inv(a)$ (or $inv_B(a)$, if we need to make B explicit), of an element a of the Boolean algebra B is a triple of numbers (or symbols $-1, \infty$):

- (1) $inv(a) = \langle -1, 0, 0 \rangle$ if $a = 0$;
- (2) $inv(a) = \langle \infty, 0, 0 \rangle$ if $\forall i(a \notin I_n)$;
- (3) $inv(a) = \langle k, l, m \rangle$ if
 - (a) $a \notin I_k, a \in I_{k+1}$,
 - (b) $l = \min\{\infty, \|\{x \in At(B^{[k]}) \ \& \ x \leq a\}\|$
 - (c) $m = 0$ if in $B^{[k]}$ there are no nonzero atomless elements below a and $m = 1$ otherwise.

We denote the entries of this triple by $inv_i(a)$ for $i = 1, 2, 3$. Note that $inv(B) = inv_B(1)$.

Definition 4.2. We define unary predicates $\mathcal{I}_n, \mathcal{A}_n, \mathcal{B}_n$ and \mathcal{C}_n and the associated formulas in the language of Boolean algebras that characterize them by induction:

- $\mathcal{I}_0(x) \Leftrightarrow x = 0$;
- $\mathcal{A}_0(x) \Leftrightarrow \neg \mathcal{I}_0(x) \ \& \ (\forall y \leq x)(y = 0 \vee y = x)$;
- $\mathcal{B}_0(x) \Leftrightarrow (\neg \exists y \leq x)(\mathcal{A}_0(y))$;
- $\mathcal{C}_0(x) \Leftrightarrow (\neg \exists y \leq x)(\neg \mathcal{I}_0(y) \ \wedge \ \mathcal{B}_0(y))$;
- $\mathcal{I}_{n+1}(x) \Leftrightarrow \exists y, z(\mathcal{B}_n(y) \ \& \ \mathcal{C}_n(z) \ \& \ x = y \vee z)$;
- $\mathcal{A}_{n+1}(x) \Leftrightarrow \neg \mathcal{I}_{n+1}(x) \ \& \ (\forall y \leq x)(\mathcal{I}_{n+1}(y) \ \vee \ \mathcal{I}_{n+1}(x - y))$;
- $\mathcal{B}_{n+1}(x) \Leftrightarrow (\neg \exists y \leq x)(\mathcal{A}_{n+1}(y))$;
- $\mathcal{C}_{n+1}(x) \Leftrightarrow (\neg \exists y \leq x)(\neg \mathcal{I}_{n+1}(y) \ \wedge \ \mathcal{B}_{n+1}(y))$.

We note that these formulas $\mathcal{I}_n, \mathcal{A}_n, \mathcal{B}_n$ and \mathcal{C}_n are equivalent (by prenexing rules) to a recursive list of formulas which are $\Sigma_{4n}, \Pi_{4n+1}, \Pi_{4n+2}$ and Π_{4n+3} , respectively. It is proven in Goncharov [1997, 2.2] that they define the notions of being in I_n , an atom of $B^{[n]}$, atomless in $B^{[n]}$ and atomic in $B^{[n]}$, respectively. There is no problem in carrying out his proofs in ACA_0^+ . Thus in ACA_0^+ we can extend B to a structure with these predicates for all n . In particular, in ACA_0^+ we can prove that if $A \equiv B$, then $inv(A) = inv(B)$.

To prove the converse, that inv does determine elementary type, we continue to follow Goncharov [1997, 2.3] by defining dense Boolean algebras and noting some of their properties.

Definition 4.3. A Boolean algebra B is *dense* if $\forall b \in B \forall k \in \omega$:

- (1) $k < inv_1(b) \Rightarrow \exists a \leq b(inv(a) = \langle k, \infty, 0 \rangle)$, and
- (2) $inv_1(b) = \infty$ or $inv_2(b) = \infty \Rightarrow (\exists a \leq b)$
 $[inv_1(a) = inv_1(b) = inv_1(b - a) \ \& \ inv_2(a) = inv_2(b) = inv_2(b - a)]$.

We denote by \mathcal{DB} the class of dense Boolean algebras.

Goncharov [1997, 2.3] proves that every countable Boolean algebra has an elementary extension B^* which is dense and that any two countable dense Boolean algebras with the same invariant are isomorphic. This then shows that any two countable Boolean algebras with the same invariant are elementary equivalent, and so establishes Tarski’s theorem. (For uncountable Boolean algebras it follows from

the countable case by the downward Skolem-Löwenheim theorem and the fact mentioned above that the value of inv is determined by the elementary theory of a Boolean algebra.)

Given B , the construction of a dense elementary extension B^* proceeds by building an ascending chain of algebras each satisfying the complete elementary diagram of B (with constants for all its elements) and adding on the witnesses needed in the definition of density as one goes along. The existence of the required extension at each step follows from a compactness argument that depends on the existence of witnesses for any finite subset of conditions in the original algebra based only on the characteristic of the given b . The existence of these witnesses is proven using only arithmetic comprehension once one has the existence of $B^{(\omega)}$ (to supply the elementary diagram of B). Of course, the compactness theorem holds in ACA_0 as well. Thus the existence of the desired chain can always be proven in ACA_0^+ . Each element in this chain is a model of the complete diagram of B , and so an elementary extension of B as is its union B^* (again all this is provable in ACA_0^+). As all the required witnesses for density were put in along the way, B^* is dense, and so it is the required extension of B .

The proof that any two countable dense Boolean algebras A and B with the same characteristic are isomorphic proceeds by considering the relation

$$S = \{(a, b) \in A \times B \mid inv_A(a) = inv_B(b) \ \& \ inv_A(\neg a) = inv_B(\neg b)\}$$

and proving that S satisfies the Vaught Criterion [1954] (as described in Goncharov [1997, 1.5]) for carrying out the back and forth argument needed to prove isomorphism of the Boolean algebras. The existence of S follows from the existence of the function inv (as characterized in Definition 4.2) and ACA_0 and so from ACA_0^+ . That it satisfies the Vaught Criterion follows from density in ACA_0 (given the existence of the function inv on A and B). The construction of the isomorphism from S is the usual back and forth argument that can be carried out in ACA_0 . Thus the following results are provable in ACA_0^+ :

Theorem 4.4 (ACA_0^+). *For Boolean algebras A and B , $A \equiv B$ if and only if $inv(A) = inv(B)$. Moreover, if A and B are dense, then $A \cong B$ if and only if $inv(A) = inv(B)$.*

For the sake of our eventual reversal results in RCA_0 we will reconsider these arguments in ACA_0 for algebras with $inv_1 = k < \infty$ under assumptions that allow us to make sense in RCA_0 or ACA_0 of the notion of elementary equivalence for the structures we will consider. We begin with our promised treatment of satisfaction and elementary equivalence in RCA_0 (or at least ACA_0). We model our definitions on Simpson's [1999, II.8.9] notion of a weak model. The notion of a structure for a language (having a domain and functions and relations interpreting the appropriate symbols of the language) is standard. The interpretation of variable free terms in the structure and satisfaction for atomic sentences are defined as usual (using recursion) in RCA_0 .

Definition 4.5 (RCA_0). If A is a structure for a language L (with equality) and S is a set of sentences of L_A (L expanded by constants \underline{a} for all elements a of A) containing the atomic sentences of L_A which is closed under propositional combinations and substitution instances (by variable free terms of L_A) of subformulas, then $F : S \rightarrow \{0, 1\}$ is a (partial) truth assignment for A (defined on S) if the

following hold for any formulas σ, τ in S :

- (1) If σ is atomic, then $F[\sigma] = 1 \Leftrightarrow A \models \sigma$.
- (2) $F[\neg\sigma] = 1 - F[\sigma]$ for any $\sigma \in S$.
- (3) $F[\sigma \& \tau] = 1 \Leftrightarrow F[\sigma] = 1$ and $F[\tau] = 1$.
- (4) $F[\exists x \sigma(x)] = 1 \Leftrightarrow (\exists a \in A)(F[\sigma(\underline{a})] = 1)$.

Of course, given two assignments F and F' for a structure A defined on sets S and S' respectively, it is easy to prove by induction in RCA_0 that the two assignments agree on $S \cap S'$. Thus we can define the (partial) satisfaction function.

Definition 4.6. For a structure A and sentence σ of its language L we set the (partial) satisfaction function $Sat(A, \sigma) = i \in \{0, 1\}$ if there is a truth assignment F for A such that $F[\sigma] = i$ and let it be undefined otherwise.

Definition 4.7 (RCA_0). Two structures A and B for a language L are elementary equivalent, $A \equiv B$, if, for every sentence σ of L , $Sat(A, \sigma) = Sat(B, \sigma)$ in the sense that, if either is defined, they are both defined and equal.

It is now immediate that \equiv is an equivalence relation coarser than isomorphism.

Proposition 4.8 (RCA_0). *If $A \cong B$, then $A \equiv B$.*

In ACA_0^+ there is always a truth assignment on S_ω , the set of all sentences, and it and our definition of \equiv agree with the usual notions. Indeed, in ACA we can prove by induction that there is a truth assignment to the sets S_n consisting of all sentences of L_A of quantifier depth at most n . Thus the notions of satisfaction and elementary equivalence also have their usual properties in this system as well. In ACA_0 alone, however, we cannot prove that Sat is defined on all sentences for all structures. (This requires the existence of $A^{(n)}$ for every n which follows from either ACA_0^+ or ACA .) We can, however, hope to build specific structures A (e.g. decidable ones) for which we can prove that Sat is defined for all $\sigma \in L$. This is the course we will follow in our reversal argument for Boolean algebras.

The crucial notions are the predicates $\mathcal{I}_n, \mathcal{A}_n, \mathcal{B}_n$ and \mathcal{C}_n (for $n \leq \text{inv}_1(B)$), as defined above. We want to consider Boolean algebras B for which we can extend the structure to include the predicates $\mathcal{I}_n, \mathcal{A}_n, \mathcal{B}_n$ and \mathcal{C}_n (for $n \leq \text{inv}_1(B)$) so as to satisfy the formulas of Definition 4.2. Given a Boolean algebra in the extended language which satisfies these axioms, we can prove in ACA_0 that they correctly represent the associated notions, and so define the invariant of each element of the algebra. In the other direction, given a function h on B we can use these predicates and axioms to characterize the requirements for h to be the function inv_1 (or inv) as in Definition 6.4. Although we will not need it here, it is possible to prove in ACA_0 that there is a total satisfaction function (even in the extended language) for algebras with such predicates (or functions) defined on them. It is uniquely determined by the 1-quantifier theory in the extended language. (All this would follow from the appropriate quantifier elimination result underlying Tarski [1949] or the one for model completeness in Ershov [1964].) One can also prove that the algebras have dense elementary extensions (which necessarily have the same invariant). Moreover, the proof of Goncharov [1997, 2.3] discussed above, that two dense Boolean algebras of the same characteristic are isomorphic, can be carried out in ACA_0 given these predicates as they effectively determine $\text{inv}(a)$ for every element a of the algebras.

We next (§5) turn to the analysis of the recursion theoretic complexity for $\psi(\mathcal{B}, \equiv)$ and of $\psi(\mathcal{DB}, \cong)$. We will then use those constructions to establish our proof theoretic reversals in §6.

5. RECURSION THEORETIC COMPLEXITY

Our plan is first to provide a uniformly recursive set of Boolean algebras $B_{k,l,m}$ such that $\text{inv}(B_{k,l,m}) = \langle k, l, m \rangle$. Next, by induction we construct uniformly recursive Boolean algebras, $A_{i,j}$, which code $0^{(i)}$ in the following sense:

$$\begin{aligned} \text{inv}_1(A_{i,j}) &= i \Leftrightarrow j \in 0^{(i)}, \\ \text{inv}_1(A_{i,j}) &= i + 1 \Leftrightarrow j \notin 0^{(i)}. \end{aligned}$$

If we now apply the assumption of the existence of invariants for countable sequences in the sense of ψ to the sequence C_q recursively listing all the $B_{k,l,m}$ and $A_{i,j}$, we code $0^{(\omega)}$ in the sense that any sequence $\langle n_q \rangle$ witnessing ψ for $\langle C_q \rangle$ computes $0^{(\omega)}$: To see if $i \in 0^{(j)}$, find a q such that $C_q = A_{i,j}$ and then $\langle k, l, m \rangle$ and p such that $B_{k,l,m} = C_p$ and $n_p = n_q$. (Such numbers exist since the $B_{k,l,m}$ include all possible invariants and so include all possible elementary equivalence types with inv_1 finite.) Now $i \in 0^{(j)}$ if and only if $k = i$. On the other hand, our analysis in §4 shows that for every sequence $\langle A_i \rangle$ of Boolean algebras the desired witness $\langle n_i \rangle$ for ψ is given by $\langle \text{inv}(A_i) \rangle$ and this sequence is recursive in $\langle A_i \rangle^{(\omega)}$. Thus we characterize the recursion theoretic strength of $(*\mathcal{B}, \equiv)$ as that of the ω -jump.

We now need some definitions and construction methods for Boolean algebras. Our primary building blocks will be algebras generated from linear orderings. We refer the reader to Monk [1989, I.6.15] and Goncharov [1997, 1.6 and 3.2] for general information about these interval algebras. We begin with some simple linear orderings and operations on linear orderings that we will use to construct the orderings whose interval algebras will be our desired $B_{k,l,m}$ and $A_{i,j}$.

Definition 5.1. We let \mathbf{m} denote the order type of an m -element linear ordering; ω that of the first infinite ordinal and η the order type of the rationals.

Definition 5.2. We define operations $+$, $\sum_{i \in M}$, \cdot , n and $*$ on linear orderings L_i and M by describing the resulting order types. We leave the formalities of defining particular representations of the order types to the reader.

- (1) $L_1 + L_2$, the sum of the two orderings, has the order type of L_1 followed by L_2 .
- (2) $\sum_{i \in M} L_i$ has the order type produced by replacing the element i of M (assumed to have underlying set \mathbb{N}) by the ordering L_i .
- (3) $L_1 \cdot L_2$ the product of the two orderings, has the order type gotten by replacing each element of L_2 by a copy of L_1 (and so it is the ordering on pairs $\langle x_1, y_1 \rangle \in L_1 \times L_2$ given by $\langle x_1, y_1 \rangle < \langle x_2, y_2 \rangle \Leftrightarrow y_1 < y_2$ or $(y_1 = y_2$ and $x_1 < x_2)$).
- (4) L_1^n , the n^{th} power of L_1 , is defined as usual given by repeated multiplication: $L_1^1 = L_1$; $L_1^{n+1} = L_1^n \cdot L_1$.

Definition 5.3. If L is a linear ordering with a first element, then $\text{Intalg}(L)$ is the Boolean algebra of finite unions of half open intervals $[a, b)$ of \mathcal{L} where b can be ∞ . (The understanding here is that $[a, \infty) = \{x | x \geq a\}$.) We often abuse notation by using L in place of $\text{Intalg}(L)$. For example, we write $\text{inv}(L)$ for $\text{inv}(\text{Intalg}(L))$. We also carry over notions and notation. For example, an ordering L or interval

$[a, b]$ is *atomic* if every subinterval contains an *atom*, i.e. a subinterval $[x, y]$ such that $[x, y] = \{x\}$. Similarly, L or $[a, b]$ is *atomless* iff it is dense with a first element (i.e. of the form $\mathbf{1} + \eta$).

Note that, when we deal with interval algebras, the orderings generating them all have first elements. We think of ∞ as a symbol used to define half open intervals but not as an element of the ordering. It is not hard to see that every countable Boolean algebra is isomorphic to the interval algebra of a countable linear order (Monk [1989, I.6.15.10]). We will generally work directly with the linear orders. To that end, we want to point out a few facts about the correspondences between L and $Intalg(L)$ that are relevant to the analysis described by the Tarski invariants. They can be found in Monk [1989, I.6.15] or Goncharov [1997, 1.6 and 3.2].

Definition 5.4. A subset S of L is *convex* if $x, y \in S$ and $x < z < y$ implies that $z \in S$. An equivalence relation \sim on L is *convex* if every one of its equivalence classes is convex.

Proposition 5.5. *There is a one-one correspondence between ideals I of $Intalg(L)$ and convex equivalence relations \sim on L such that $Intalg(L)/I \cong Intalg(L/\sim)$. Here L/\sim is the linear ordering of equivalence classes $[x], [y]$ of \sim given by $[x] < [y] \Leftrightarrow \forall w \sim x \forall z \sim y (w < z)$. The convention here is that if a final segment of L is collapsed to a single equivalence class, then it is removed from L/\sim and its role is taken by ∞ . (That is, we identify the final segment with the “external” point ∞ .) For a given ideal I and canonical projection π , the corresponding equivalence class \sim is given by $x \sim y \Leftrightarrow \pi(x) = \pi(y)$.*

Definition 5.6. In accordance with this proposition, we define the equivalence relation \sim_T corresponding to the Tarski ideal $I(Intalg(L))$ by $x \sim_T y \Leftrightarrow [x, y]$ is a finite sum of atomic and atomless suborderings. We denote L/\sim_T by L' and so $Intalg(L') \cong Intalg(L)/I(Intalg(L)) = (Intalg(L))'$. As with the corresponding algebras we let $L^{[0]} = L$ and $L^{[n+1]} = (L^{[n]})'$.

Now for the canonical algebras $B_{k,l,m}$ with invariant $\langle k, l, m \rangle$ we take those of Morozov [1982] which are not only uniformly recursive (indeed, uniformly decidable) but dense as well.

Proposition 5.7 ([1982, Morozov]). *Let $P = (\mathbf{1} + \eta + \mathbf{1} + \omega \cdot \eta) \cdot \omega$. The following table lists linear orders whose interval algebras are uniformly decidable, dense and have the indicated invariants:*

$L_{n,0,1}$	$P^n \cdot (\mathbf{1} + \eta)$	$\langle n, m, 1 \rangle$
$L_{n,m,0}$	$P^n \cdot \mathbf{m}$	$\langle n, m, 0 \rangle$
$L_{n,m,1}$	$P^n \cdot (\mathbf{1} + \eta + \mathbf{m})$	$\langle n, 0, 1 \rangle$
$L_{n,\infty,0}$	$P^n \cdot (\mathbf{1} + \omega \cdot \eta)$	$\langle n, \infty, 0 \rangle$
$L_{n,\infty,1}$	$P^n \cdot (\mathbf{1} + \eta + \mathbf{1} + \omega \cdot \eta)$	$\langle n, \infty, 1 \rangle$

Thus we may take $B_{k,l,m}$ to be $Intalg(L_{k,l,m})$.

Next, we must describe the algebras $A_{i,j}$ that code $0^{(i)}$. Our plan is to uniformly construct recursive linear orderings $L_{n,j}^\Sigma$ and $L_{n,j}^\Pi$ such that

- $inv_1(L_{n,j}^\Sigma) = n + 1 \Leftrightarrow j \in 0^{(n)}$ (Σ_n case),
- $inv_1(L_{n,j}^\Sigma) = n \Leftrightarrow j \notin 0^{(n)}$ (Π_n case),
- $inv_1(L_{n,j}^\Pi) = n \Leftrightarrow j \in 0^{(n)}$ (Σ_n case),
- $inv_1(L_{n,j}^\Pi) = n + 1 \Leftrightarrow j \notin 0^{(n)}$ (Π_n case).

We begin with $n = 1$:

- $L_{1,j}^\Sigma = P^2 \Leftrightarrow j \in 0'$ (Σ_1 case),
- $L_{1,j}^\Sigma = P \Leftrightarrow j \notin 0'$ (Π_1 case),
- $L_{1,j}^\Pi = P \Leftrightarrow j \in 0'$ (Σ_1 case),
- $L_{1,j}^\Pi = P^2 \Leftrightarrow j \notin 0'$ (Π_1 case).

Note that the desired properties for $n = 1$ hold (in ACA_0) as $inv_1(P) = 1$ and $inv_1(P^2) = 2$. In fact, if $B = Intalg(P)$, then $B/I(B) = \{0, 1\}$ and so $inv(B) = \langle 1, 1, 0 \rangle$ while $P^2/I(P^2) = P$ and so $inv(P^2) = \langle 2, 1, 0 \rangle$.

Now each Σ_{n+1} (Π_{n+1}) fact such as $j \in 0^{(n+1)}$ ($j \notin 0^{(n+1)}$) is uniformly of the form $\exists x(f(n, j, x) \notin 0^{(n)})$ ($\forall x(f(n, j, x) \in 0^{(n)})$) for some fixed recursive f . Moreover, we may choose f such that $f(n, j, x) \notin 0^{(n)} \& y > x \rightarrow f(n, j, y) \notin 0^{(n)}$. (This is just quantifier manipulation to replace $\exists z \dots$ with $\exists x \exists z < x \dots$ etc.) We now define the rest of our orderings as follows:

- $L_{n+1,j}^\Sigma = P^{n+1} + P^{n+1} + \sum_{k \in P} L_{n,f(n,j,k)}^\Pi$,
- $L_{n+1,j}^\Pi = P^{n+1} + P^{n+1} + \sum_{k \in P} L_{n,f(n,j,k)}^\Sigma$.

We let $\mathcal{L}_n = \{L_{n,j}^\Pi | j \in \omega\} \cup \{L_{n,j}^\Sigma | j \in \omega\}$. To verify that the invariants of these orderings are as required we proceed by induction on n to prove various facts about the orderings including that the coding works. We begin with a general lemma.

Lemma 5.8. *If, for every $i \in \omega$, $inv_1(L_i) \geq 1$ for every L_i and $L = \sum_{i \in M} L_i$, then $L' = \sum_{i \in M} L'_i$.*

Proof. Consider the equivalence relation $x \sim_T y$ on L . If x and y belong to the same copy of L_k in M , then they are clearly T-equivalent in M if and only if they are in L_k as this depends only on the order type of $[x, y]$. If x and y are not in L_i and L_j for successive $i, j \in M$, then $x \not\sim_T y$ as the interval between them contains some L_k which by assumption is not a finite sum of atomic and atomless orderings. If x and y are in adjacent L_i, L_j , then they can be identified only if x is identified with all the points to its right in L_i , i.e. the final segment of L_i beginning with x is a finite sum of atomic and atomless orderings (and so disappears in L'_i) and y is identified with the first element of L_j in L'_j . \square

Theorem 5.9. *If $L \in \mathcal{L}_n$, then $inv_1(L)$ is n or $n + 1$. Moreover,*

1. $L = L_{n,j}^\Sigma \& j \in 0^{(n)} \Rightarrow inv_1(L) = n + 1$ and $K^{[n+1]} = \mathbf{1}$ for every final segment K of L ;
2. $L = L_{n,j}^\Sigma \& j \notin 0^{(n)} \Rightarrow inv_1(L) = n$ & $L^{[n]} = \mathbf{1}$ if $n = 1$ and $L^{[n]} = \mathbf{m}$ for some $m > 1$ if $n > 1$;
3. $L = L_{n,j}^\Pi \& j \notin 0^{(n)} \Rightarrow inv_1(L) = n + 1$ and $K^{[n+1]} = \mathbf{1}$ for every final segment K of L ;
4. $L = L_{n,j}^\Pi \& j \in 0^{(n)} \Rightarrow inv_1(L) = n$ & $L^{[n]} = \mathbf{1}$ if $n = 1$ and $L^{[n]} = \mathbf{m}$ for some $m > 1$ if $n > 1$.

Proof. We proceed by induction on n . We have already noted that for $n = 1$ the desired facts about invariants and Tarski derivatives of L are true. For the final segments just note that in cases 1 and 3 they all contain (and are contained in) final segments isomorphic to the whole ordering.

For the inductive step at $n + 1$ we consider the cases in order. For notational convenience, we choose L_k in each case so that $L = P^{n+1} + P^{n+1} + \sum_{k \in P} L_k$.

1. Here $\text{inv}_1(L_k)$ is n for finitely many k and $n + 1$ for the rest by induction. Recursively applying Lemma 5.8 we see that $L^{[n]} = P + P + \sum_{k \in P} L_k^{[n]}$. For the cofinitely many k for which $\text{inv}_1(L_k) = n + 1$ the entry in the k^{th} spot in the sum corresponding to $L^{[n+1]}$ is $L_k^{[n+1]}$ which is a single point (by induction). For the finitely many k for which $\text{inv}_1(L_k) = n$, $L_k^{[n]}$ is \mathbf{m}_k for some $m_k \in \mathbb{N}$ (by induction). Thus in $L^{[n+1]}$ it is identified with the least element in $L^{[n]}$ to the right of all the points in $L_k^{[n]}$ or the greatest one to its left if there is one (if both exist they are also identified) and otherwise becomes $\mathbf{1}$, i.e. a single point. Thus $L^{[n+1]}$ is $\mathbf{1} + \mathbf{1} + P$ (finitely many points have been removed from P but the result is still isomorphic to P) and so $L^{[n+2]} = \mathbf{1}$ as required. Of course, the same argument works for the sum over the L_k for k in any final segment of P .

2. Here $\text{inv}_1(L_k) = n$ for every k by induction. If $n = 1$, $L'_k = \mathbf{1}$ for every k and so $L' = P + P + P$ and $L^{[2]} = \mathbf{3}$ as required. If $n > 1$, $L^{[n]} = P + P + V$ where V is atomic, and so $L^{[n+1]} = \mathbf{2}$ again as required.

3. Here $\text{inv}_1(L_k) = n + 1$ and $L_k^{[n+1]} = \mathbf{1}$ for every k by induction. Thus $L^{[n+1]} = \mathbf{1} + \mathbf{1} + P$, and so $L^{[n+2]} = \mathbf{1}$ as required. Of course, the same is true for the sum over the L_k for k in any final segment of P .

4. If $n = 1$, then L is $P^2 + P^2$ followed by a sum over P in which finitely many points have been replaced by P^2 and the rest by P . It is clear that, in this case, $L^{[2]}$ is \mathbf{m} where m is three more than the number of points that have been replaced by P^2 . (The final segment of L is essentially P^2 and so contributes one point to $L^{[2]}$. The initial segment is $P^2 + P^2$ and contributes two points as does every other copy of P^2 in L . The points of P replaced by a copy of P which are to the left of a location where a copy of P^2 is inserted are again single points in L' , and so these intervals in L' are isomorphic to proper segments $[x, y)$ of P which are then identified with the least point of the next copy of P^2 in $L^{[2]}$.)

Finally, if $n > 1$, then for finitely many k , $\text{inv}_1(L_k) = n + 1$ and $L_k^{[n+1]} = \mathbf{1}$ while for all the rest $\text{inv}_1(L_k) = n$ and $L_k^{[n]} = \mathbf{m}_k$ for some $m_k > 1$. In this case, $L^{[n]}$ is $P + P + V$ where V is a sum over P in which finitely many of the points are replaced by copies of P and the rest are replaced by the \mathbf{m}_k which are atomic. Thus $L^{[n+1]}$ is again two more points than the number of copies of P in V . (Each copy of P contributes one point. The atomic portions caught between two copies of P are identified with the first point of the copy of P immediately to their right. The final atomic segment is identified with ∞ and so is removed.) \square

We can now prove that the recursion theoretic complexity of the existence of elementary equivalence invariants for recursive sequences of Boolean algebras is that of $0^{(\omega)}$.

Theorem 5.10. *There is a uniformly recursive sequence $\langle A_i \rangle$ of Boolean algebras such that $0^{(\omega)}$ is recursive in any witness $\langle n_i \rangle$ to the instance of $\psi(\mathcal{B}, \equiv)$ corresponding to $\langle A_i \rangle$.*

Proof. Let the members of the sequence $\langle A_i \rangle$ be the $B_{k,l,m}$ and the interval algebras of the orders L in $\bigcup \{ \mathcal{L}_n \mid n \in \omega \}$ as defined above in some recursive order of type ω . To decide if $j \in 0^{(\omega)}$ (recursively in $\langle n_i \rangle$) find numbers k, l, m, p and q such that $A_p = B_{k,l,m}$, $A_q = L_{n,j}^\Sigma$ and $n_p = n_q$. (Such numbers exist since the $B_{k,l,m}$ include all possible invariants and so all possible elementary equivalence types with inv_1

finite.) By Theorem 5.9, $inv_1(A_q)$ is $n + 1$ or n depending on whether $j \in 0^{(n)}$ or not. As we also know that $inv_1(A_q) = k$ since $A_q \equiv B_{k,l,m}$, we know that $j \in 0$ if and only if $k = n + 1$. \square

We now work towards a proof that the Boolean algebras appearing in the above proof are all dense.

Lemma 5.11 (Goncharov [1997, 2.24]). *If $a, b \in A$ a Boolean algebra and $a \wedge b = 0$, then*

- (1) $inv_1(a \vee b) = \max\{inv_1(a), inv_1(b)\}$;
- (2) $inv_1(a) = inv_1(b) \Rightarrow inv_2(a \vee b) = inv_2(a) + inv_2(b)$;
- (3) $inv_1(a) < inv_1(b) \Rightarrow inv(a \vee b) = inv(b)$.

Corollary 5.12. *If $a \in Intalg(L)$ and (as must necessarily be true for some choice of x_i, y_i and n) $a = \bigcup\{[x_i, y_i] \mid i < n\}$, then there is an $i < n$ such that $inv_1(a) = inv_1([x_i, y_i])$ and if $inv_2(a) = \infty$, then there is an $i < n$ such that $inv_1(a) = inv_1([x_i, y_i])$ and $inv_2([x_i, y_i]) = \infty$.*

Lemma 5.13. *If K, L and M are linear orderings with first elements and $x, y \in L$ (with y possibly being the ∞ of L which we identify with the first element of M), then $inv^L([x, y]) = inv^{K+L+M}([x, y])$ where we use the superscript to indicate that the evaluation of the invariant is taking place in the specified Boolean algebra (i.e. the interval algebra of the specified ordering).*

Proof. This is immediate from the fact that points x and y are identified under the Tarski equivalence relation if and only if $[x, y]$ can be divided up into a finite sum of subintervals each of which is atomic or atomless and this depends only on the order type of $[x, y]$ and not on the ambient order containing it. \square

Definition 5.14. We say that a linear ordering L is *B-dense* if $Intalg(L)$ is a dense Boolean algebra.

Remark 5.15. By Corollary 5.12 if we want to prove that a linear ordering L is B-dense we need only verify the requirements of B-density for elements of the form $[x, y)$. As all the orderings we consider have finite first invariant, proving the B-density of L amounts to verifying two conditions for every $b = [x, y)$ and every $m \in \mathbb{N}$:

- (1) $m < inv_1(b) \Rightarrow \exists a \leq b (inv(a) = \langle m, \infty, 0 \rangle)$, and
- (2) $inv_2(b) = \infty \Rightarrow \exists a \leq b (inv_1(a) = inv_1(b) = inv_1(b - a) \ \& \ inv_2(a) = \infty = inv_2(b - a))$.

Lemma 5.16. *If K and L are linear orderings with first elements, then $K + L$ is B-dense if and only if both K and L are B-dense.*

Proof. Assume K and L are B-dense. We verify the requirements for the B-density of $K + L$ for an arbitrary $[x, y)$. If x and y are in the same component of $K + L$ (including $y = 0_L$ which plays the role of ∞ in K), then the requirements follow from the B-density of the component and Lemma 5.13. So suppose that $x \in K$ and $y \in L - \{0_L\}$ so that $[x, y) = [x, 0_L) \cup [0_L, y)$. Now the result follows from our assumptions and Corollary 5.12.

For the other direction, assume that $K + L$ is B-dense. Consider any $b = [x, y)$ with x and y in the same component (we identify ∞_K and 1_L). If b satisfies the hypotheses of one of the conditions defining B-density in its own component, then it

does so in $K + L$ as well by 5.13. The witness a , assured by the B-density of $K + L$, is smaller than b and so contained in the same component. Again, its invariants in that component are the same as in $K + L$, and so it witnesses satisfaction of the requirement for the B-density of the component. \square

Theorem 5.17. *Every $L \in \mathcal{L}_n$ is B-dense for every n .*

Proof. We begin with P itself and consider an arbitrary $b = [x, y)$. The only ones such that $\text{inv}_1(b) = 1$ are those with $y = \infty$. For these $\text{inv}(b) = \langle 1, 1, 0 \rangle$; all the others have $\text{inv}_1(b) = 0$. So, if $\text{inv}_1(b) = 1$, we must verify Condition (1) with $k = 0$. Any element a corresponding to the suborder $\mathbf{1} + \omega \cdot \eta$ of any component of $P = \sum_{k \in \omega} (\mathbf{1} + \eta + \mathbf{1} + \omega \cdot \eta)$ in the final segment b has $\text{inv}(a) = \langle 0, \infty, 0 \rangle$ as required. If $\text{inv}_1(b) = 0$, then we only have to verify Condition (2). If $\text{inv}_2(b) = \infty$, i.e. b contains infinitely many pairs of points that are immediate successors, then it must contain some suborder of type $\omega \cdot \eta$. It can then be split into two such suborderings as required.

We should next consider P^2 and indeed all the P^n but their B-density follows from the general argument for Case 3 so we proceed by induction on n beginning with Case 3. We phrase the proof to cover the P^n as well. It also then covers Case 2 with $n = 1$.

Note that, in every case, by Lemma 5.16, we need only prove the B-density of orderings of the form $L = \sum_{k \in P} L_k$ for $L_k \in \mathcal{L}_n$. Note that by Lemma 5.13 and the inductively assumed B-density of the L_k , we only need to verify the conditions for B-density for $b = [x, y)$ with $y = \infty$ and with $x \in L_i$ and $y \in L_j$ for $i <_P j$. Also note that if j is the immediate successor of i in P , then $b = [x, 1_{L_j}) \cup [1_{L_j}, y)$. Now Corollary 5.12, together with the assumptions that L_i and L_j are B-dense, shows that b satisfies the requirements for B-density. Thus we may assume that there is a k such that $i <_P k <_P j$.

3. We want to show that if $L = \sum_{k \in P} L_k$ and, for every k , L_k is B-dense, $\text{inv}_1(K) = n + 1$ and $K^{(n+1)} = \mathbf{1}$ for every final segment K of L_k , then L is B-dense. We already know that $\text{inv}_1(b) = n + 2$ only if $b = [x, \infty)$ and that in this case $\text{inv}_2(b) = 1$. Thus, for these b , we need only verify Condition (1). By induction and Lemma 5.13, we only have to verify the case that $m = n + 1$. Here we know that $L^{[n+1]} = P$. Thus any element a whose image in $L^{[n+1]}$ corresponds to the suborder $\mathbf{1} + \omega \cdot \eta$ of any component of P in the image of b has $\text{inv}(a) = \langle n + 1, \infty, 0 \rangle$ as required. We are now left with the case that $x \in L_i$ and $y \in L_j$ with $i <_P j$. In this case $\text{inv}_1(b) = n + 1 = \text{inv}_1^{L_i}(x, \infty)$ so there is an $a = [w, z) \in L_i$ as required by Condition (1) for the B-density of L_i . This element satisfies the same condition for b in L by Lemma 5.13. Finally, we must verify Condition (2) for such b with $\text{inv}_2(b) = \infty$. Since $L^{[n+1]} = P$, the argument for P can be applied to the image of b in $L^{[n+1]}$ and the inverse image of the element so produced provides the desired witness.

2. ($n > 1$) Here $L^{[n]}$ is atomic and so $\text{inv}_1(L) = n$. Thus $\text{inv}_1(b) \leq n$. As noted above we may assume that b contains all of L_k for some k . Thus $\text{inv}_1(b) = n$ and there are w, z strictly between x and y such that $[w, z) \cong P^n$, so there are smaller elements satisfying Condition (1). The only way to have $\text{inv}_2(b) = \infty$ is for the image of $[x, y)$ in P (in the sum $\sum_{i \in P} \mathbf{m}_k = L^{[n]}$) to contain a subinterval of type η or $\omega \cdot \eta$. In either case, it is easy to split this image into two parts both of which contain infinitely many points. Each of these points of P corresponds to

a component of $L^{[n]}$ with at least one atom, and so its inverse image in L satisfies Condition (2).

1. In this case, there is a final segment of L which has the same properties as all of L did in Case 3, and so we have covered the situation in which $y = \infty$. Now $inv_1(b) = n + 1$ if and only if b contains a finite segment of one of the L_k with $inv_1(L_k) = n + 1$. Any witness for Condition (1) for this final segment in L_k works for b in L . As $L^{[n+1]} = P$ (with perhaps finitely many points removed) the analysis for P shows that Condition (2) is also satisfied for b if $inv_1(b) = n + 1$. Finally, if $inv_1(b) \leq n$, then $inv_1(b) = n$, since b contains some L_k with $inv_1(L_k) = n$. Condition (1) is satisfied by the density of the L_k contained in b . As for Condition (2), note that the only way that b does not contain some L_k with $inv_1(L_k) = n + 1$ is if it is contained in a finite sum of successive L_i with $inv_1(L_i) = n$ plus perhaps a proper initial segment of one L_j with $inv_1(L_j) = n + 1$. As for each such i other than j , $L_i^{[n]}$ is atomic, Condition (2) can come into play only because of $[0_{L_j}, y)$. As $L_j^{[n]}$ has P as an initial segment, $inv_2([0_{L_j}, y) = \infty$ implies that in $L_j^{[n]}$ the image of y is beyond the initial segment of P of type $\mathbf{1} + \eta$. Thus, it contains an initial segment of the first copy of $\omega \cdot \eta$ in P which can then easily be split as required for Condition (2).

4. If $n = 1$, L is a sum over P of finitely many copies of P^2 and the rest copies of P . Now B-density follows easily from our previous arguments.

If $n > 1$, a final segment of L is of the form considered in Case (2). Thus we need only consider an initial segment of L of the form $\sum_{i \in Q} L_k$ where $Q = (\mathbf{1} + \eta + \mathbf{1} + \omega \cdot \eta)\mathbf{m}$ for some $m \in \omega$ in which, for finitely many k , $inv_1(L_k) = n + 1$ and, for the rest, $inv_1(L_k) = n$. In fact, by Lemma 5.16, it suffices to take $m = 1$. Now $inv_1(b) = n + 1$ if and only if b contains a final segment of one of the copies of an L_k with $inv_1(L_k) = n + 1$. In this case, Condition (1) is immediate from the B-density of L_k . As $L^{[n+1]}$ is finite, Condition (2) is vacuous for such b . The only remaining case is that b contains only L_k with $inv_1(L_k) = n$ and at least one such L_k so that $inv_1(b) = n$. In particular, the image of b in $L^{[n]}$ is $\sum_{i \in Q} \mathbf{m}_k$ for some subinterval Q of P . In this case Condition (1) follows from the B-density of L_k (by induction) and Condition (2) can apply only if Q contains an interval of the form η or $\omega \cdot \eta$ and so is easily satisfied. \square

Remark 5.18. All the orderings of Proposition 5.7 (Morozov [1982]) can now be seen to be dense. We have directly proved it for all the powers P^n . The others all follow from Lemma 5.16 as all of the orderings with first invariant n are either sums of P^n or components of P^{n+1} or sums of each of these. (For example, $P^{n+1} = P^n(\mathbf{1} + \eta) + P^n(\mathbf{1} + \omega \cdot \eta)$, and so both $P^n(\mathbf{1} + \eta)$ and $P^n(\mathbf{1} + \omega \cdot \eta)$ are dense.)

We have now proven that the recursion theoretic complexity of the existence of isomorphism invariants for recursive sequences of dense Boolean algebras is also $0^{(\omega)}$.

Theorem 5.19. *There is a uniformly recursive sequence $\langle A_i \rangle$ of dense Boolean algebras such that $0^{(\omega)}$ is recursive in any witness $\langle n_i \rangle$ to the instance of $\psi(\mathcal{DB}, \cong)$ corresponding to $\langle A_i \rangle$.*

Proof. Theorem 5.17 and Remark 5.18 show that all the Boolean algebras appearing in the proof of Theorem 5.10 are dense. Goncharov [1997, 2.3] (as described in

Theorem 4.4) shows that for dense Boolean algebras, elementary equivalence is the same as isomorphism. \square

From a recursion theoretic point of view these results can be viewed as types of hardness results for index sets. In the terminology defined in Soare [1987, IV.3.1] we have shown, for example, that $(\Sigma_n, \Pi_n) \leq_m (\mathcal{B}_n, \mathcal{B}_{n+1})$ where $\mathcal{B}_n = \{B \in \mathcal{B} \mid \text{inv}_1(B) = n\}$. The first natural question is whether this is the best one can do. As mentioned above a simple quantifier count shows only that \mathcal{B}_n is in Σ_{4n} . One could therefore hope or even expect that it is Σ_{4n} complete. At one point, we thought that such a result would be useful for the reverse mathematical arguments we now wish to pursue. That does not seem to be the case, and so this seems to be an instance where the recursion theoretic questions are finer than those of reverse mathematics. Similar questions arise about the index sets for all the values of inv . We deal with these purely recursion theoretic issues along with determining the index set for the Boolean algebras with finite first invariant in Csima, Montalban and Shore [2005]. For now we turn to the issue of the complexity of the existence of these invariants in the sense of reverse mathematics.

6. REVERSALS

Our primary goal is to show that the strength, in the sense of reverse mathematics, of the existence of elementary equivalence invariants for Boolean algebras is that of ACA_0^+ . As we argued in §1, it is sufficient to prove that $RCA_0 \vdash \psi(\mathcal{B}, \equiv) \rightarrow ACA_0^+$. As usual, establishing ACA_0^+ amounts to proving the existence of $0^{(\omega)}$ (by a relativizable proof). Our recursion theoretic results seem to have provided us with such a proof already. In particular, Theorem 5.10 provides us with a uniformly recursive sequence $\langle A_i \rangle$ of Boolean algebras such that any witness $\langle n_i \rangle$ for the instance of $\psi(\mathcal{B}, \equiv)$ corresponding to $\langle A_i \rangle$ computes $0^{(\omega)}$. The issue is, of course, whether our proof of Theorem 5.10 takes place in, or can be revised to work in, RCA_0 .

The first problem is that even some of the simple algebraic arguments naturally take place in ACA_0 . This problem is easy to handle in the style of the results of the previous section. The only issue is that dealing with elementary equivalence in RCA_0 requires some care since one cannot assume that a given structure has a full satisfaction function (that would itself give ACA_0^+) nor even one for the set of Σ_1 sentences (as that in general would give ACA_0). Thus we must use structures for which we can prove the existence of a partial satisfaction covering the sentences we need.

Proposition 6.1. $RCA_0 \vdash \psi(\mathcal{B}, \equiv) \rightarrow ACA_0$. *Indeed, $RCA_0 \vdash \psi(\mathcal{DB}, \cong) \rightarrow ACA_0$.*

Proof. We begin with two nonisomorphic recursive dense Boolean algebras $C_0 = \text{Intalg}(\mathbf{1} + \eta)$ and $C_1 = \text{Intalg}(\mathbf{1} + \eta + \mathbf{1})$. For any natural choice of recursive presentations of C_0 and C_1 it is easy to recursively define inv on these algebras and prove in RCA_0 that they are dense. We can easily produce a uniformly recursive sequence $\langle C_i \rangle$ of Boolean algebras such that $\forall i (C_i \cong C_0 \text{ or } C_i \cong C_1)$ and $(\forall i > 1)(i \in 0' \Leftrightarrow C_i \cong C_0)$ where each of the isomorphisms is recursive. Let $\langle n_i \rangle$ be a witness for the instance of $\psi(\mathcal{B}, \equiv)$ or $\psi(\mathcal{DB}, \cong)$ corresponding to $\langle C_i \rangle$. If $\langle n_i \rangle$ is a witness for $\psi(\mathcal{DB}, \cong)$ it clearly computes $0'$. Suppose it witnesses $\psi(\mathcal{B}, \equiv)$. As isomorphism implies elementary equivalence (Proposition 4.8), $i \in 0' \Rightarrow n_i = n_0$

and $i \notin 0' \Rightarrow n_i = n_1$. Thus we only need to prove that $n_0 \neq n_1$, i.e. $C_0 \not\cong C_1$. Consider the sentence $\sigma = \forall x \exists y (x = 0 \vee 0 < y < x)$. It is true in C_0 and false in C_1 . Moreover, it is easy to define partial satisfaction functions for both structures on the atomic sentences (with constants for the elements of C_0, C_1) and the (propositional combinations of) substitution instances of subformulas of σ . (The only basic ones not covered by the atomic cases are of the form $\exists y (b = 0 \vee 0 < y < b)$. These are true of 0 (in both algebras) but false for every nonzero $b \in C_0$ and of every nonzero $b \in C_1$ except for the final segment of C_1 consisting of a single atom.) \square

In view of this proposition, we may work over ACA_0 from now on. We begin by discussing the properties of Boolean algebras developed in §§3 and 4. There is no problem with the definition of the basic notions dealing with atomicity or atomlessness and of the Tarski ideal and derivative. All make sense and can be proven to exist for every Boolean algebra B in ACA_0 . Our first problem arises when we try to iterate the Tarski derivative to define $B^{[n]}$ and the function inv on B . These are not provably total operations in ACA_0 . Of course, it only takes induction to define them and an induction argument to prove that they exist. Thus, in ACA , we can prove that $B^{[n]}$ exists for every n . Even in ACA , however, we cannot prove the existence of either the sequence $\langle B^{[n]} \rangle$ or the function inv on B unless $B^{[n]} = 0$ for some n . Note that we need much less than full induction to get these results. For example, let ACA_0^* be $\text{ACA}_0 + \forall X \forall n \exists (Y_0, \dots, Y_n) (Y_0 = X \ \& \ (\forall i < n) (Y_{i+1} = Y_i'))$. Clearly $\text{ACA} \vdash \text{ACA}_0^*$ and ACA_0^* suffices to prove that, for every Boolean algebra B and every natural number n , $B^{[n]}$ exists and if for some n , $B^{[n]} = 0$, then there is a function inv on B from which we can define the predicates of Definition 4.2 so that they satisfy the axioms listed there. We can also say that $\text{inv}(B) = \langle \infty, 0, 0 \rangle$ with the intended interpretation by saying that $\forall n \neg (\text{inv}_1(B) = n)$ (i.e. that the sequence of derivatives of length $n + 1$ exists and that B does not become 0 in any of them).

Note that the algebras we needed in the last section all had finite first invariant. Of course, for specifically constructed Boolean algebras, we may be able to prove even in ACA_0 that a function inv exists with the required properties.

One route to our main theorem might now be to prove an effective quantifier elimination result for Boolean algebra with these extra predicates (as Tarski apparently did). This would show that two Boolean algebras are elementary equivalent if and only if they have the same invariant. (One point being that the quantifier elimination result would show that they have total satisfaction functions.) Although something along these lines is possible, there is no direct classical style proof of quantifier elimination in the published literature. The closest is the proof of model completeness in Ershov [1964] which would also suffice to prove the existence of total satisfaction functions for each of our algebras in ACA_0 . His decision procedure for the one quantifier sentences would then give the decidability of the theories and show that they are elementary equivalent if and only if they have the same invariant. That would still leave us with the (inductive) task of establishing that our algebras have the desired invariants. We have chosen, instead, to follow a path of analysis that exploits the density of our algebras, and so establishes isomorphism results rather than elementary equivalence ones. Nonetheless, we will have to deal with many of the same issues of the apparent calls on induction needed to define the invariant functions for our algebras.

Turning then to the development of the theory of dense Boolean algebras we immediately hit the issue of the definition of density of B which, *prima facie*, assumes the existence of the function inv on B . Again, as we will only need ones with $inv_1 < \infty$, we could assume ACA_0^* and keep the original definition. As we will in the end want to work only in ACA_0 , it is better to follow a route like that taken for elementary equivalence.

Definition 6.2 (ACA_0). A Boolean algebra B is *dense* if, for every function inv and associated predicates $\mathcal{I}_n, \mathcal{A}_n, \mathcal{B}_n$ and \mathcal{C}_n on B as given in Definition 4.2 that satisfy the axioms listed there and the defining equations for inv of Definition 4.1, the conditions defining density (Definition 4.3) hold.

Note that, given a function h on B purporting to be a candidate for inv which gives $inv_1(B) = n$ for some n , the properties that need to be verified to show that it is really the function inv are all arithmetical, indeed, of fixed quantifier level (in h and B). (First define the extra predicates directly from inv and then just look at the axioms in Definition 4.2.) Thus it makes sense to hope that we can verify that h is the desired inv function in ACA_0 and that it witnesses the density of B even if an induction on inv_1 , or equivalently, on the levels of the predicates, is needed.

Next, it is easy to check that our basic definitions and lemmas about dense Boolean algebras, interval algebras and the connections between them make sense and can be carried out in ACA_0 . In particular, there are no problems with 5.1–5.5 and 5.8. (In Lemma 5.7 we understand the assertion that $inv_1(L_k) \geq 1$ to mean just that L_k is not a finite sum of atomic and dense suborderings.) So too there are no problems in proving 5.11–5.16 in ACA_0 whenever the invariant functions assumed in the statements exist. We also note again that Goncharov’s [1997, 2.3.2] proof that any two dense Boolean algebras with the same invariant are isomorphic works in ACA_0 as long as the function inv exists for both algebras.

Finally, we are left to consider the proofs of Theorem 5.9 where the Boolean algebras coding $0^{(\omega)}$ are constructed and analyzed and Theorem 5.17 where they are all proven to be dense. There is no problem with the definition of the orderings (and so algebras) $L_{k,l,m}$ and those in the \mathcal{L}_n hierarchy as they are given uniformly recursively. As presented in the last section, the proofs of both of these theorems are straightforward ones by induction. They can be routinely carried out in ACA . Thus we have proven our reversals over ACA ; actually, by Proposition 6.1, over RCA .

Theorem 6.3. $RCA \vdash \psi(\mathcal{B}, \equiv) \rightarrow ACA^+$ and $RCA \vdash \psi(\mathcal{DB}, \cong) \rightarrow ACA^+$.

One could also prove these results now in ACA_0^* by showing that all is well for each n . Here one can prove the existence of the functions inv for the orderings at each level \mathcal{L}_n directly from the definitions of the predicates up to level $n + 1$ which exist for an arbitrary Boolean algebra B since their definition only requires the existence of $B^{(4n+4)}$. However, we want to work over ACA_0 alone. The first step is to give the “right” proof in ACA_0^* that only needs the existence of $0^{(n)}$ to prove the existence of the function inv on the orderings in \mathcal{L}_n . We then plan on doing an induction argument that simultaneously proves the existence of $0^{(n)}$ and that the orderings in \mathcal{L}_n have the properties established in §5. (A subtle point here is that we cannot do an induction that establishes the existence of $0^{(n)}$ and then simply asserts that therefore there are inv functions for orderings in \mathcal{L}_n as, in the

abstract, this requires $0^{(4n+4)}$ whose existence we cannot prove in ACA_0 from the existence of $0^{(n)}$.)

We begin with a formal description of what it means to be the function inv_1 on a Boolean algebra. Of course, the full invariant function inv is arithmetically definable from inv_1 and so if the latter has been shown to exist in ACA_0 , then so does the former.

Definition 6.4 (ACA_0). If B is a Boolean algebra, we say that a function $h : B \rightarrow \mathbb{N}$ satisfies the conditions to be inv_1 on B if $\forall n (\mathcal{I}_{n+1} = \{x \in B \mid \text{the image of } x \text{ in } B/\mathcal{I}_n \text{ is a finite sum of atomic and atomless elements of } B/\mathcal{I}_n\})$ where $\mathcal{I}_k = \{x \in B \mid h(x) \leq k\}$.

Proposition 6.5 (ACA_0). Fix a natural number m and assume that $0^{(m)}$ exists, i.e. there is a sequence of sets $\langle Y_i \mid i \leq m \rangle$ such that $Y_0 = \emptyset$ and $(\forall i < n)(Y_{i+1} = Y'_i)$. Then there is a family of functions $h_{n,j}^\Gamma$ for $n < m$ and $j \in \mathbb{N}$ uniformly recursive in $0^{(m)}$ such that they satisfy the conditions to be inv_1 on $\text{Intalg}(L_{n,j}^\Gamma)$ for $n < m$, $j \in \mathbb{N}$ and $\Gamma = \Sigma$ or Π . Moreover, the properties established in Theorems 5.9 and 5.17 all hold.

Proof. We begin by defining the functions $h_{n,j}^\Gamma$ by induction on $n < m$ uniformly recursively in $0^{(m)}$. In every case, we define the functions directly on subintervals $[x, y)$ and extend to the whole algebra (i.e. to finite sums) by taking the max. When defining h on a sum over P of orderings L_k for which we already have defined functions h_k , we set $h([x, y)) = h_k([x, y))$ if x and y are in the same component L_k (with the usual understanding that we identify the first element of L_j with ∞_{L_k} if j is the immediate successor of k in P). Similarly, if $x \in L_i$, $y \in L_j$ and j is the immediate successor of i in P , then we let $h([x, y)) = \max\{h_{L_i}([x, \infty)), h_{L_j}([1, y))\}$. Thus we only need to specify the value of h on intervals of the form $[x, \infty)$ and $[x, y)$ with $x \in L_i$, $y \in L_j$ when there is some k such that $i <_P k <_P j$.

For $n = 1$, $L_{n,j}^\Gamma$ is either P or P^2 and $0'$ can tell which one effectively. The corresponding functions $h_{n,j}^\Gamma$ are then just set to be what we know are the corresponding functions inv_1 on P or P^2 . For P , $h([x, y))$ is 1 if $y = \infty$ and 0 otherwise. For $P^2 = \sum_{i \in P} P$, we let $h([x, y)) = 2$ if $y = \infty$ and otherwise 1, for x and y in different component components.

For $1 < n + 1 < m$, we divide into cases as in Theorem 5.9. Note that it suffices to define h on $\sum_{i \in P} L_i$ as we can then extend it to $P^{n+1} + P^{n+1} + \sum_{i \in P} L_i$ by our algebraic rules as long as we have a definition for P^{n+1} . This definition is given by the procedures in Case 3 for $\sum_{i \in P} L_i$ when each $L_i = P^n$. It also covers Case 2 for $n = 1$.

3. $h([x, y)) = n + 2$ if $y = \infty$ and otherwise $n + 1$ for x and y in different components.

1. As $j \in 0^{(n+1)}$, there are only finitely many k such that $f(n, j, k) \in 0^{(n)}$ and recursively in $0^{(n+1)}$ we can find (the canonical index of) the set F consisting of these k . Of course, we set $h([x, \infty)) = n + 2$ for every x . If $x \in L_i$ and $y \in L_j$ for $i <_P j$, we set $h([x, y)) = n + 1$ if there is a $k \notin F$ such that $i \leq_P k < j$; otherwise, $h([x, y)) = n$.

2. ($n > 1$) We set $h([x, \infty)) = n + 1$ for every x . For the other case that needs a definition, we set $h([x, y)) = n$.

4. As $j \in 0^{(n+1)}$, there are only finitely many k such that $f(n, j, k) \in 0^{(n)}$ and, recursively in $0^{(n+1)}$, we can find (the canonical index of) the set F consisting of

these k . We set $h([x, \infty)) = n + 1$ for every x . For the other case required, we set $h([x, y)) = n + 1$ if there is a $k \in F$ such that $i \leq_P k <_P j$; otherwise, $h([x, y)) = n$.

It is now routine to prove by induction on $n < m$ that these functions satisfy the conditions to be inv_1 on the corresponding algebras and (following the previous proofs) that they have the properties established in Theorems 5.9 and 5.17. \square

We can now prove the reversal part of our main theorem.

Theorem 6.6. $RCA_0 \vdash \psi(\mathcal{B}, \equiv) \rightarrow ACA_0^+$.

Proof. Let $\langle n_1 \rangle$ be the witness to $\psi(\mathcal{B}, \equiv)$ for $\langle A_i \rangle$. Define $\langle Y_i \rangle$ recursively in the jump of $\langle n_i \rangle$ by $j \in Y_i \Leftrightarrow \exists k, l, m, p, q (A_p = B_{k,l,m} \ \& \ A_q = L_{i,j}^\Sigma \ \& \ n_p = n_q)$. We claim that $Y_{i+1} = Y_i'$ for every i . If not, let m be the least counterexample. Thus, $0^{(m)}$ exists and, since $RCA_0 \vdash \psi(\mathcal{B}, \equiv) \rightarrow ACA_0$ (Proposition 6.1), so does $0^{(m+2)}$. Proposition 6.5 now guarantees that all the orderings up to level $m + 1$ have all the desired properties, and so the proofs of Theorems 5.10 and 5.19 work up to level $m + 1$ to show that $Y_{m+1} = Y_m'$, for the required contradiction. \square

Finally, we want to get the same result for $\psi(\mathcal{DB}, \cong)$. The problem is that we cannot prove in advance of applying $\psi(\mathcal{DB}, \cong)$ that all the orderings in the \mathcal{L}_n are B-dense. We need one more twist to strengthen our inductive hand. We also show that if any $L \in \mathcal{L}_{n+1}$ with a function satisfying the conditions to be inv_1 (or equivalently that the function inv with the required properties on L) exists, then $0^{(n)}$ exists.

Definition 6.7. We define the members of $\hat{\mathcal{L}}_n$ by induction. $\hat{\mathcal{L}}_1 = \mathcal{L}_1$. For $L_{n+1,j}^\Gamma \in \mathcal{L}_{n+1}$ we let $\hat{L}_{n+1,j}^\Gamma \in \hat{\mathcal{L}}_{n+1}$ be $\sum_{m \leq n} \sum_{i \in \omega \cdot \eta} L_{m,i}^\Sigma + \sum_{m \leq n} \sum_{i \in \omega \cdot \eta} L_{m,i}^\Pi + L_{n+1,j}^\Gamma$.

Proposition 6.8 (ACA_0). *If a function $h : \hat{L} \rightarrow \mathbb{N}$ satisfies the conditions to be inv_1 on $\hat{L} \in \hat{\mathcal{L}}_n$, then $0^{(n)}$ exists.*

Proof. Let h be the assumed function on \hat{L} . Each $L_{i,j}^\Sigma$ is an interval $[x_{i,j}, y_{i,j})$ in \hat{L} that can be found recursively. Define Y_i for $i \leq n$ by $j \in Y_i \Leftrightarrow h[x_{i,j}, y_{i,j}) = i + 1$. Again, it is at this point straightforward to prove by induction that the $L_{m,k}^\Gamma$ for $m \leq n$ all have the properties previously established, and so $Y_{i+1} = Y_i'$ for $i \leq n$. \square

Theorem 6.9. $RCA_0 \vdash \psi(\mathcal{DB}, \cong) \rightarrow ACA_0^+$.

Proof. The $\hat{L}_{n,j}$ are all dense, and so we can argue as in Theorem 6.10 that applying $\psi(\mathcal{DB}, \cong)$ to the same sequence as there, except that we substitute $\hat{L}_{n,j}$ for $L_{n,j}$, we obtain the same results. \square

We have thus completed the proofs of our main theorems of reverse mathematics.

Theorem 6.10. *The existence of number invariants for elementary equivalence of countable Boolean algebras or for isomorphism of dense countable Boolean algebras has the same complexity in the sense of reverse mathematics as ACA_0^+ .*

REFERENCES

[1987] Blass, A. R., Hirst, J. L. and Simpson, S. G., Logical analysis of some theorems of combinatorics and topological dynamics, in *Logic and Combinatorics*, S. G. Simpson, ed., Contemporary Mathematics, American Mathematical Society, Providence, 125-156. MR0891245 (88d:03113)

- [2005] Csima, B., Montalban, A. and Shore, R.A., Boolean algebras, Tarski invariants and index sets, *Notre Dame Journal of Formal Logic*, to appear.
- [1977] Chang, C. C. and Keisler, H. J., *Model Theory 2nd ed.*, North Holland, Amsterdam. MR0532927 (58:27177)
- [1964] Ershov, Yu. L., Decidability of the elementary theory of distributive lattices with relative complements and the theory of filters, *Algebra i Logika* **3**, 17-38. MR0180490 (31:4725)
- [1998] Ershov, Yu. L., Goncharov, S. S., Nerode, A. and Remmel, J. B., eds., *Handbook of Recursive Mathematics*, Studies in Logic and the Foundations of Mathematics, vols. 138-139, Elsevier, Amsterdam, vols. 1-2. MR1673617 (99k:03003); MR1673582 (99k:03004)
- [1983] Friedman, H., Simpson, S. G. and Smith, R., Countable algebra and set existence axioms, *Annals of Pure and Applied Logic* **25**, 141-181. MR0725732 (85i:03157)
- [1997] Goncharov, S. S., *Countable Boolean Algebras and Decidability*, Consultants Bureau, New York. MR1444819 (98h:03044b)
- [1988] Doner, J. and Hodges, W., Alfred Tarski and decidable theories, *The Journal of Symbolic Logic* **53**, 20-35. MR0929372 (89j:01048)
- [1972] Jensen, R. B., The fine structure of the constructible hierarchy, *Annals of Mathematical Logic* **4**, 229-308. MR0309729 (46:8834)
- [1989] Monk, J. D., ed., *Handbook of Boolean Algebras*, North Holland, Amsterdam, vols. 1-3. MR0991565 (90k:06002); MR0991595 (90k:06003); MR0991607 (90k:06004)
- [1982] Morozov, A. S., Strong constructivizability of countable saturated Boolean algebras, *Algebra i Logika* **21**, 193-203; English version in *Algebra and Logic* **21**, 130-137. MR0700992 (85b:03052)
- [1999] Simpson, S. G., *Subsystems of Second Order Arithmetic*, Springer, Berlin. MR1723993 (2001i:03126)
- [1987] Soare, R. I., *Recursively Enumerable Sets and Degrees*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin. MR0882921 (88m:03003)
- [1949] Tarski, A., Arithmetical classes and types of Boolean algebras, *Bulletin of the American Mathematical Society* **55**, 63.
- [1954] Vaught, R. L., *Topics in the Theory of Arithmetical Classes and Boolean Algebras*, Ph.D. Thesis, University of California, Berkeley.
- [2000] White, W. M., *Characterizations for Computable Structures*, Ph.D. Thesis, Cornell University.

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NEW YORK 14853
E-mail address: shore@math.cornell.edu