

MARTINGALES AND CHARACTER RATIOS

JASON FULMAN

ABSTRACT. Some general connections between martingales and character ratios of finite groups are developed. As an application we sharpen the convergence rate in a central limit theorem for the character ratio of a random representation of the symmetric group on transpositions. A generalization of these results is given for Jack measure on partitions. We also give a probabilistic proof of a result of Burnside and Brauer on the decomposition of tensor products.

1. INTRODUCTION

The Plancherel measure of a finite group G is a probability measure on the set of irreducible representations of G which chooses a representation ρ with probability $\frac{\dim(\rho)^2}{|G|}$, where $\dim(\rho)$ denotes the dimension of ρ . For instance if G is the symmetric group, the irreducible representations are parameterized by partitions λ of n , and the Plancherel measure chooses a partition λ with probability $\frac{n!}{\prod_{x \in \lambda} h(x)^2}$, where the product is over boxes in the partition and $h(x)$ is the hooklength of a box. The hooklength of a box x is defined as 1 + number of boxes in the same row as x and to the right of x + number of boxes in same column of x and below x . For example we have filled in each box in the partition of 7 with its hooklength,

$$\begin{array}{|c|} \hline 6 \\ \hline 3 \\ \hline 1 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 4 \\ \hline 1 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 2 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 1 \\ \hline \end{array} ,$$

and the Plancherel measure would choose this partition with probability $\frac{7!}{(6 \cdot 4 \cdot 3 \cdot 2)^2}$. Recently there has been interest in the statistical properties of partitions chosen from Plancherel measure. As it is beyond the scope of this paper to survey the topic, we refer the reader to the surveys [AID], [De] and the seminal papers [J], [O1], [BOO] for a glimpse of the remarkable recent work on Plancherel measure.

Let λ be a partition of n chosen from the Plancherel measure of the symmetric group S_n and let $\chi^\lambda(12)$ be the irreducible character parameterized by λ evaluated on the transposition (12). The quantity $\frac{\chi^\lambda(12)}{\dim(\lambda)}$ is called a character ratio and is crucial for analyzing the convergence rate of the random walk on the symmetric group generated by transpositions [DSh]. In fact Diaconis and Shahshahani prove that the eigenvalues for this random walk are the character ratios $\frac{\chi^\lambda(12)}{\dim(\lambda)}$ each

Received by the editors February 25, 2004 and, in revised form, September 21, 2004.

2000 *Mathematics Subject Classification*. Primary 05E10; Secondary 60C05.

Key words and phrases. Martingale, central limit theorem, character ratio, symmetric group, Jack polynomial.

©2006 American Mathematical Society
 Reverts to public domain 28 years from publication

occurring with multiplicity $\dim(\lambda)^2$. Character ratios on transpositions also play an essential role in work on the moduli space of curves [EO], [OP].

Given these motivations, it is natural to study the distribution of the character ratio $\frac{\chi^\lambda(12)}{\dim(\lambda)}$. Kerov [K1] outlined a proof of the following central limit theorem.

Theorem 1.1 ([K1]). *Let λ be a partition of n chosen from the Plancherel measure of the symmetric group S_n . Then the random variable $\frac{n-1}{\sqrt{2}} \frac{\chi^\lambda(12)}{\dim(\lambda)}$ is asymptotically normal with mean 0 and variance 1.*

The details of the proof appeared in the paper [IO], which gave a wonderful development of Kerov's work. Another proof, due to Hora [Ho], exploited the fact that the k th moment of a Plancherel distributed character ratio is the chance that the random walk generated by random transpositions is at the identity after k steps. Both of these proofs were essentially combinatorial in nature and used the method of moments (so information about all moments of the character ratio) to prove Theorem 1.1.

A more probabilistic approach to Kerov's central limit theorem appeared in [F1], which proved the following result. In its statement and throughout this paper, we use the notation that if A is an event, $P(A)$ denotes the probability of A .

Theorem 1.2 ([F1]). *For $n \geq 2$ and all real x_0 ,*

$$\left| P\left(\frac{n-1}{\sqrt{2}} \frac{\chi^\lambda(12)}{\dim(\lambda)} \leq x_0\right) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \leq 40.1n^{-1/4}.$$

The proof technique involved a random walk on the set of irreducible representations of the symmetric group and a technique known as Stein's method, which is fundamentally different from the method of moments. Note that unlike Kerov's original result, Theorem 1.2 included an error term (of order $n^{-1/4}$).

In this paper we give a completely different approach to central limit theorems for character ratios. To begin we describe the general setting. Let G be a finite group and C a conjugacy class on which all irreducible characters of G are real valued (this occurs for instance if $C = C^{-1}$). Let ρ be a random representation of G , chosen from the Plancherel measure of G , and let χ^ρ denote the character of ρ . From the orthogonality relations of irreducible characters it follows that if C is a nonidentity conjugacy class, then the random variable $\frac{|C|^{1/2} \chi^\rho(C)}{\dim(\rho)}$ has mean 0 and variance 1, and it is natural to investigate how close it is to a standard normal random variable with mean 0 and variance 1.

To see the connection with martingales, instead of investigating $\frac{|C|^{1/2} \chi^\rho(C)}{\dim(\rho)}$, consider $\frac{|C| \chi^\rho(C)}{\dim(\rho)}$. Given a subgroup chain $H_1 = id \subseteq H_2 \subseteq \cdots \subseteq H_n = G$, together with representations $\rho(j) \in Irr(H_j)$, and assuming that $C \cap H_j$ is at most one conjugacy class of H_j , one can write

$$\begin{aligned} & \frac{|C| \chi^\rho(C)}{\dim(\rho)} \\ &= \sum_{j=2}^n \left(\frac{|C \cap H_j| \chi^{\rho(j)}(C \cap H_j)}{\dim(\rho(j))} - \frac{|C \cap H_{j-1}| \chi^{\rho(j-1)}(C \cap H_{j-1})}{\dim(\rho(j-1))} \right), \end{aligned}$$

where we set $\chi^{\rho(j)}(C \cap H_j) = 0$ if $|C \cap H_j| = 0$. Starting from $\rho(1)$ equal to the trivial representation of H_1 , there is a way (see Section 2 for details) to grow a

series of random representations $\rho(2), \dots, \rho(n)$ such that each $\rho(i)$ is chosen from the Plancherel measure of H_i . Thus $\frac{|C|\chi^\rho(C)}{\dim(\rho)}$ is expressed as a sum of random variables, which one might hope to be small and nearly independent of each other, so that after renormalization a central limit theorem would hold.

Section 2 proves that under certain conditions the sequence of partial sums in the above decomposition of $\frac{|C|\chi^\rho(C)}{\dim(\rho)}$ is a martingale. It also shows that (due to the extra structure coming from representation theory) this martingale has special properties not shared by general martingales. There is a significant literature on central limit theorems for martingales, and in Section 6 it is applied to prove Theorem 1.3 (actually we just prove Theorem 1.5 below, which generalizes Theorem 1.3).

Theorem 1.3. *For any $s < \frac{1}{2}$, there is a constant A_s so that for all n and real x_0 ,*

$$\left| P \left(\frac{\sqrt{\binom{n}{2}} \chi^\lambda(12)}{\dim(\lambda)} \leq x_0 \right) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \leq A_s n^{-s}.$$


This sharpens the convergence rate in Theorem 1.2 and very nearly proves the conjecture in [F1] that in Theorem 1.3 one could take $s = \frac{1}{2}$. An essential ingredient in the proof of Theorem 1.3 is an understanding of the conditional moments of the summands in the decomposition of $\frac{|C|\chi^\rho(C)}{\dim(\rho)}$. For general conjugacy classes this is a difficult problem. A fortunate and remarkable fact is that work of Kerov [K3] in the combinatorics literature implies that asymptotically these conditional moments are simply the moments of the semicircle distribution. Section 3 explains this, and discusses some other phenomena which occur only for the class of transpositions.

It should be pointed out that both Kerov [K1] and Hora [Ho] use the method of moments to prove central limit theorems (with no error term) for character ratios evaluated on cycles of various lengths—even obtaining a multidimensional central limit theorem showing these to be asymptotically independent. Martingales do blend well with multidimensional central limit theorems, but it is unclear whether the methods of this paper can be pushed through. Hora [Ho] also shows that there are many conjugacy classes of the symmetric group where central limit theorems do not hold.

This paper also considers central limit theorems for Jack_α measure, a generalization of Plancherel measure of the symmetric group on partitions of size n . Here $\alpha > 0$, and this measure chooses a partition λ of size n with probability

$$\frac{\alpha^n n!}{\prod_{s \in \lambda} (\alpha a(s) + l(s) + 1)(\alpha a(s) + l(s) + \alpha)},$$

where the product is over all boxes in the partition. Here $a(s)$ denotes the number of boxes in the same row as s and to the right of s (the “arm” of s) and $l(s)$ denotes the number of boxes in the same column as s and below s (the “leg” of s). For example the partition of 5,



would have Jack_α measure

$$\frac{60\alpha^2}{(2\alpha + 2)(3\alpha + 1)(\alpha + 2)(2\alpha + 1)(\alpha + 1)}.$$

Note that when $\alpha = 1$, Jack measure reduces to Plancherel measure of the symmetric group. The papers [O2], [BO] emphasize that for α fixed the study of Jack_α measure is an important open problem, about which relatively little is known for general values of α (the three values $\alpha = 1/2, 1, 2$ are tractable). It is a discrete analog of ensembles from random matrix theory, and like Jack polynomials [GHJ], they should also be relevant to the moduli space of curves.

Recently the author proved a central limit related to Jack measure, for general values of α . We remark that when $\alpha = 2$ the statistic W_α is a spherical function for the Gelfand pair (S_{2n}, H_{2n}) , where S_{2n} is a symmetric group and H_{2n} is the hyperoctahedral group of size $2^n n!$. In Theorem 1.4, $n(\lambda)$ is the quantity $\sum_{i \geq 1} (i-1)\lambda_i$, where λ_i is the length of the i th row of λ . Also λ' is the transpose of λ .

Theorem 1.4 ([F2]). *Suppose that $\alpha \geq 1$. Let $W_\alpha(\lambda) = \frac{\alpha n(\lambda') - n(\lambda)}{\sqrt{\alpha \binom{n}{2}}}$. For $n \geq 2$ and all real x_0 ,*

$$\left| P_\alpha(W_\alpha \leq x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \leq A_\alpha n^{-1/4},$$

where A_α depends on α but not on n .

Section 4 develops connections between Jack symmetric functions and martingales, showing how statistics such as W_α can be written as a sum of martingale differences. Although the results are analogous to those of Section 2 (and generalize them for the case of the symmetric group), the proofs are completely different, as representation theory of finite groups can no longer be used. Section 5 then focuses on the summands of the specific statistic W_α . Again one finds that certain simplifications occur which do not hold in general for the martingales in Section 4. In Section 6 the results of Sections 4 and 5 are used to prove the following result.

Theorem 1.5. *Suppose that $\alpha \geq 1$. Let $W_\alpha(\lambda) = \frac{\alpha n(\lambda') - n(\lambda)}{\sqrt{\alpha \binom{n}{2}}}$. Then for any $s < \frac{1}{2}$, there is a constant $A_{s,\alpha}$ (depending only on s and α) such that for all $n \geq 2$ and real x_0 ,*

$$\left| P_\alpha(W_\alpha \leq x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \leq A_{s,\alpha} n^{-s}.$$

The assumption that $\alpha \geq 1$ in Theorems 1.4 and 1.5 is merely for convenience. Indeed, from the definition of Jack measure it is clear that the Jack_α probability of λ is equal to the $\text{Jack}_{\frac{1}{\alpha}}$ probability of the transpose of λ . From this one concludes that the Jack_α probability that $W_\alpha = w$ is equal to the $\text{Jack}_{\frac{1}{\alpha}}$ probability that $W_{\frac{1}{\alpha}} = -w$, so that a central limit theorem holds for α if and only if it holds for $\frac{1}{\alpha}$.

The original version of this paper stated the following conjecture:

Conjecture 1. *Theorem 1.5 also holds with $s = \frac{1}{2}$.*

Since this article was written, two proofs of Conjecture 1 have been given. The first proof used a refinement of Stein's method [ShSu]. A second, more recent, proof used Bolthausen's variation of Stein's method [F4]. A recent development which uses results from this paper is a law of iterated logarithm for character ratios [Su].

The construction of an exchangeable pair used in [F1] for a Stein's method approach to character ratios involved a random walk on the set of irreducible representations of a finite group and was shown in [F3] to have some implications for the

decomposition of tensor products. In Section 7 of this paper, it is shown how these random walks give a probabilistic proof of a theorem of Burnside and later Brauer (page 49 of [Is]) which states that if a faithful character η of a finite group assumes m distinct values, then every irreducible character of G occurs as a constituent of some η^j , where $0 \leq j < m$. Although the proof is less general than their result (it only works for real-valued characters), the conceptual framework is more appealing.

The organization of this paper is as follows. Section 2 develops connections between martingales and character ratios of finite groups. Section 3 then focuses on the case of transpositions, where certain simplifications occur. Section 4 develops general connections between Jack symmetric functions and martingales and is highly combinatorial in nature. Section 5 then focuses on combinatorics related to the statistic W_α , where again there are attractive simplifications. (The reader will observe that parts of Section 5 generalize parts of Section 3, but there are real differences, and since both sections are brief and Section 5 might be unreadable by group theorists who work on character ratios but not on Jack polynomials, they have been separated.) Section 6 uses central limit theory for martingales to prove Theorem 1.5 (and so also Theorem 1.3). Finally, Section 7 gives a probabilistic proof of a result of Burnside and Brauer on the decomposition of tensor products, thereby showing that the machinery underlying a Stein's method approach to Theorems 1.2 and 1.4 is of ongoing interest. This is also illustrated in the recent preprint [F5], which applies Stein's method to other algebraic structures.

2. CHARACTER RATIOS AND MARTINGALES

Let G be a finite group and C a conjugacy class of G on which all irreducible characters of G are real (this holds for instance if $C = C^{-1}$). Suppose there is a chain of subgroups $H_1 = id \subseteq H_2 \subseteq \cdots \subseteq H_n = G$. Given a representation $\rho(j)$ of H_j , one obtains a random representation $\rho(j+1)$ of H_{j+1} by choosing $\rho(j+1)$ with probability

$$\frac{|H_j| \dim(\rho(j+1)) \kappa(\rho(j), \rho(j+1))}{|H_{j+1}| \dim(\rho(j))}.$$

Here $\kappa(\rho(j), \rho(j+1))$ denotes the multiplicity of $\rho(j+1)$ in the induction of $\rho(j)$ from H_j to H_{j+1} , or by Frobenius reciprocity the multiplicity of $\rho(j)$ in the restriction of $\rho(j+1)$ from H_{j+1} to H_j . Thus starting from $\rho(1)$ (the trivial representation of a one-element group) one obtains a random sequence $(\rho(1), \dots, \rho(n))$ of representations. It is proved in [F3] that each $\rho(j)$ is distributed according to the Plancherel measure of H_j (this was known earlier for the symmetric group [K2]).

A sequence (Y_1, \dots, Y_n) of real-valued random variables is called a martingale with respect to the sequence of random variables (X_1, \dots, X_n) if two properties hold. First, for each $n \geq 1$ there is a function f_n such that $Y_n = f_n(X_1, \dots, X_n)$. Second, the martingale identity

$$E(Y_j | X_1, \dots, X_{j-1}) = Y_{j-1}$$

must hold for all $j \geq 1$. Throughout this section we also define Y_0 to be identically equal to 0.

Theorem 2.1. *Let G be a finite group and $C = C^{-1}$ a self-inverse conjugacy class of G . Let $\rho(1), \dots, \rho(n)$ be the sequence of random representations associated to a subgroup chain $H_1 = id \subseteq \cdots \subseteq H_n = G$. Suppose that $C_j = C \cap H_j$ is either empty*

or a single conjugacy class of H_j . Let $Y_j = \frac{|C_j|\chi^{\rho(j)}(C_j)}{\dim(\rho(j))}$, where this is understood to be 0 if $|C_j| = 0$. Then the sequence (Y_1, \dots, Y_n) is a martingale.

Proof. Let $Irr(H)$ denote the set of irreducible representations of a finite group H . Observe that the expected value of Y_{j+1} given the earlier ρ 's is

$$\begin{aligned} & \sum_{\tau \in Irr(H_{j+1})} \frac{|H_j|\dim(\tau)\kappa(\rho(j), \tau)}{|H_{j+1}|\dim(\rho(j))} \frac{|C_{j+1}|\chi^\tau(C_{j+1})}{\dim(\tau)} \\ &= \frac{|C_{j+1}||H_j|}{\dim(\rho(j))|H_{j+1}|} \sum_{\tau \in Irr(H_{j+1})} \kappa(\rho(j), \tau)\chi^\tau(C_{j+1}) \\ &= \frac{|C_{j+1}||H_j|}{\dim(\rho(j))|H_{j+1}|} Ind_{H_j}^{H_{j+1}}(\chi^{\rho(j)})(C_{j+1}). \end{aligned}$$

Here Ind stands for induction of characters. If C_{j+1} is empty, then $Y_{j+1} = Y_j = 0$. Otherwise since C_j is either empty or a single conjugacy class of H_j , it follows from a general formula for induced characters (page 30 of [Se]) that

$$Ind_{H_j}^{H_{j+1}}(\chi^{\rho(j)})(C_{j+1}) = \frac{|H_{j+1}||C_j|}{|H_j||C_{j+1}|} \chi^{\rho(j)}(C_j).$$

Substituting this in implies that the expected value of Y_{j+1} given the earlier ρ 's is equal to Y_j . □

Remarks. (1) The conditions of Theorem 2.1 may seem restrictive but they do apply in many cases of interest, most notably the symmetric groups and for many classes in finite classical groups.

(2) Note also that from page 52 of [Se] it follows that the random variables Y_1, \dots, Y_n are algebraic integers. Since the characters of the symmetric groups are integer valued, in this case Y_1, \dots, Y_n must be rational algebraic integers, and hence integers.

Proposition 2.2 computes certain conditional probabilities related to the martingale sequence (Y_1, \dots, Y_n) , which is not possible for general martingales. It will not be needed in the proof of central limit theorems in this paper, although it is applied in [F4]. We use the notation that $E^W(Z)$ denotes the expected value of Z given W .

Proposition 2.2. *Assume the hypotheses of Theorem 2.1. Then for all $1 \leq j \leq n$,*

$$E^{Y_n}(Y_j) = \frac{|C_j|}{|C_n|} Y_n.$$

Proof. In fact we show that

$$E^{\rho(n)}(Y_j) = \frac{|C_j|}{|C_n|} \frac{|C_n|\chi^{\rho(n)}(C_n)}{\dim(\rho(n))}$$

which is stronger.

For $j = n$ this is clear, so consider $j = n - 1$. For ease of notation we let λ denote $\rho(n)$. Then

$$E^{\rho(n)}(Y_{n-1}) = \sum_{\tau \in Irr(H_{n-1})} \frac{P(\rho(n-1) = \tau, \rho(n) = \lambda)}{P(\rho(n) = \lambda)} \frac{|C_{n-1}|\chi^\tau(C_{n-1})}{\dim(\tau)}.$$

Now since $\rho(n)$ has the distribution of Plancherel measure of H_n , $P(\rho(n) = \lambda) = \frac{\dim(\lambda)^2}{|H_n|}$. Similarly

$$\begin{aligned} P(\rho(n-1) = \tau, \rho(n) = \lambda) &= \frac{\dim(\tau)^2 \kappa(\tau, \lambda) \dim(\lambda) |H_{n-1}|}{|H_{n-1}| \dim(\tau) |H_n|} \\ &= \frac{\dim(\tau) \dim(\lambda) \kappa(\tau, \lambda)}{|H_n|}. \end{aligned}$$

Substituting this into the expression for $E^{\rho(n)}(Y_{n-1})$ shows that as desired, the latter is equal to

$$\frac{|C_{n-1}|}{\dim(\lambda)} \sum_{\tau \in \text{Irr}(H_{n-1})} \kappa(\tau, \lambda) \chi^\tau(C_{n-1}) = \frac{|C_{n-1}| \chi^\lambda(C_n)}{\dim(\lambda)}.$$

Next we indicate how to treat the case $j = n-2$, from which the general argument is completely clear. By considerations similar to those in the previous paragraph, $E^{\rho(n)}(Y_{n-2})$ is equal to

$$\begin{aligned} &\sum_{\tau \in \text{Irr}(H_{n-2})} \frac{P(\rho(n-2) = \tau, \rho(n) = \lambda)}{P(\rho(n) = \lambda)} \frac{|C_{n-2}| \chi^\tau(C_{n-2})}{\dim(\tau)} \\ &= \sum_{\tau \in \text{Irr}(H_{n-2})} \sum_{\eta \in \text{Irr}(H_{n-1})} \frac{P(\rho(n-2) = \tau, \rho(n-1) = \eta, \rho(n) = \lambda)}{P(\rho(n) = \lambda)} \\ &\quad \cdot \frac{|C_{n-2}| \chi^\tau(C_{n-2})}{\dim(\tau)} \\ &= \frac{|C_{n-2}|}{\dim(\lambda)} \sum_{\tau \in \text{Irr}(H_{n-2})} \sum_{\eta \in \text{Irr}(H_{n-1})} \kappa(\tau, \eta) \kappa(\eta, \lambda) \chi^\tau(C_{n-2}) \\ &= \frac{|C_{n-2}| \chi^\lambda(C_n)}{\dim(\lambda)}. \end{aligned}$$

Note that the last equality used the fact that restriction of characters is transitive (i.e. to compute the restriction of χ to elements in H_{n-2} , one can first restrict to H_{n-1} and then to H_{n-2}). □

Proposition 2.3 derives an exact expression (which will be needed) for $E(Y_j - Y_{j-1})^2$.

Proposition 2.3. *Assume the hypotheses of Theorem 2.1. Then for $j \geq 2$, $E(Y_j - Y_{j-1})^2 = |C_j| - |C_{j-1}|$.*

Proof. Clearly $E(Y_j - Y_{j-1})^2 = E(Y_j^2) + E(Y_{j-1}^2) - 2E(Y_j Y_{j-1})$. But

$$E(Y_j Y_{j-1}) = E(E^{Y_{j-1}}(Y_j Y_{j-1})) = E(Y_{j-1} E^{Y_{j-1}}(Y_j)) = E(Y_{j-1}^2).$$

Thus $E(Y_j - Y_{j-1})^2$ is equal to $E(Y_j^2) - E(Y_{j-1}^2)$. Now observe that

$$E(Y_j^2) = \sum_{\tau \in \text{Irr}(H_j)} \frac{\dim(\tau)^2 |C_j|^2 \chi^\tau(C_j)^2}{|H_j| \dim(\tau)^2}.$$

After canceling the factors of $\dim(\tau)^2$, it follows from the orthogonality relations of the characters of H_j that $E(Y_j^2) = |C_j|$ for all j . This implies the proposition. □

3. TRANSPOSITIONS

This section illustrates and sharpens the results in Section 2 for the case of character ratios on transpositions. Along the way we make some links with the combinatorics literature. Moreover, certain properties will be established for the class of transpositions which fail for other conjugacy classes; for instance it will be proved that not only the sequence (Y_1, \dots, Y_n) , but also the sequence

$$(Y_1^2, Y_2^2 - \binom{2}{2}, \dots, Y_n^2 - \binom{n}{2})$$

is a martingale.

Throughout this section $G = S_n$, $H_j = S_j$ for $1 \leq j \leq n$, and C is the conjugacy class of transpositions. Recall that the irreducible representations of the symmetric group are parameterized by partitions λ of n (see [Sa] for a friendly introduction to representation theory of the symmetric group). Frobenius [Fr] found the following explicit formula for the character ratio of the symmetric group on transpositions:

$$\frac{\chi^\lambda(12)}{\dim(\lambda)} = \frac{1}{\binom{n}{2}} \sum_i \left(\binom{\lambda_i}{2} - \binom{\lambda'_i}{2} \right),$$

where λ_i is the length of row i of λ and λ'_i is the length of column i of λ .

From the branching rules of the symmetric group, the multiplicity $\kappa(\mu, \lambda)$ is positive (and in fact is 1) if and only if λ is obtained from μ by adding a box to the diagram of μ . From Frobenius' formula it follows that if μ is such a partition, then

$$\frac{|C_n| \chi^\lambda(12)}{\dim(\lambda)} - \frac{|C_{n-1}| \chi^\mu(12)}{\dim(\mu)} = c(x),$$

where x is the box added to μ to obtain λ , and the content of a box $c(x)$ is defined as column number of box $-$ row number of box.

Given the above discussion, one sees that Theorem 2.1 implies that for any partition $\lambda = \rho(j)$ of size j , the expected value of the content of the box added to λ in moving from $\rho(j)$ to $\rho(j+1)$ is 0. Rather remarkably this combinatorial fact was known to Kerov [K2],[K3], who had also proved that for *any* λ of size j , the variance of the content of the box added to λ is equal to j . Note that this is stronger than Proposition 2.3, which only establishes this for a random Plancherel distributed λ . For conjugacy classes other than transpositions, it need not be the case that $E^{\rho(j)}(Y_{j+1} - Y_j)^2$ depends only on $\rho(j)$. Indeed the reader can check that this fails for the class of 3-cycles and $j = 3$.

For the purpose of proving Theorem 1.3, it will be necessary to have upper bounds on all moments of the variable $E(Y_{j+1} - Y_j)$. In very recent work Lassalle found formulas for all $E^{\rho(j)}(Y_{j+1} - Y_j)^r$, from which one can obtain $E(Y_{j+1} - Y_j)^r$ by averaging with respect to Plancherel measure. The formulas are quite complicated, but for instance for $r = 1, 2, 3, 4$ one has the following special case of Theorem 8.1 of [La].

Theorem 3.1 ([La]). *Let $d_k(\lambda)$ denote $\sum_{x \in \lambda} c(x)^k$. Let $s_r(\lambda)$ be the r th moment of the content of the box added to $\rho(j) = \lambda$ when transitioning to $\rho(j+1)$. Then*

$$\begin{aligned} s_1(\lambda) &= 0, \\ s_2(\lambda) &= |\lambda|, \\ s_3(\lambda) &= 2d_1(\lambda), \\ s_4(\lambda) &= 3d_2(\lambda) + \binom{|\lambda|+1}{2}. \end{aligned}$$

In Section 5 it is shown (in the Jack setting which is more general) how Lassalle’s formulas lead to the necessary upper bounds on $E(Y_{j+1} - Y_j)^r$. But in the case of Plancherel measure the exact asymptotics of $E(Y_{j+1} - Y_j)^r$ are known. Indeed, one simply obtains the moments of the semi-circle distribution. We only state the result for even moments, since the odd moments all vanish by symmetry of Plancherel measure under transposing the partition. Theorem 3.2 is due to Kerov [K3].

Theorem 3.2 ([K3]). *Let λ be chosen from the Plancherel measure on partitions of size n . Then*

$$\lim_{n \rightarrow \infty} \frac{E(s_{2r}(\lambda))}{n^r} = \frac{\binom{2r}{r}}{r + 1}.$$

The final results of this section are not needed for proving central limit theorems, but are of interest. Corollary 3.3 shows that for the class of transpositions, Proposition 2.2 has a curious combinatorial consequence. We remind the reader that a standard Young tableau of shape λ is a filling of the boxes of λ with the numbers $1, \dots, |\lambda|$ each occurring once such that the numbers increase going across rows (from left to right) and down columns.

Corollary 3.3. *Let T be a standard Young tableau of shape λ chosen uniformly at random. Then the expected value of the content of box j is equal to $\frac{j-1}{\binom{n}{2}} \sum_{x \in \lambda} c(x)$.*

Proof. From the discussion at the beginning of this section, $Y_j - Y_{j-1}$ is the content of box j . By Proposition 2.2, the expected value of $Y_j - Y_{j-1}$ given that $\rho(n) = \lambda$ is equal to

$$\frac{\binom{j}{2} - \binom{j-1}{2}}{\binom{n}{2}} \sum_{x \in \lambda} c(x) = \frac{j-1}{\binom{n}{2}} \sum_{x \in \lambda} c(x).$$

Now observe that given that $\rho(n) = \lambda$, all sequences $(\rho(1), \dots, \rho(n) = \lambda)$ occur with probability

$$\frac{1}{\dim(\lambda)^2/n!} \prod_{j=1}^{n-1} \frac{j! \dim(\rho(j+1))}{(j+1)! \dim(\rho(j))} = \frac{1}{\dim(\lambda)}.$$

Thus such sequences correspond to standard Young tableaux T under the uniform distribution. □

Finally, we prove that for the conjugacy class of transpositions the sequence $(Y_1^2, Y_2^2 - \binom{2}{2}, \dots, Y_n^2 - \binom{n}{2})$ is a martingale.

Proposition 3.4. *Let C be the conjugacy class of transpositions in S_n , and let (Y_1, \dots, Y_n) be the martingale of Theorem 2.1. Then $(Y_1^2, Y_2^2 - \binom{2}{2}, \dots, Y_n^2 - \binom{n}{2})$ is also a martingale.*

Proof. Observe that for all μ of size j ,

$$\begin{aligned} & -\binom{j+1}{2} + \sum_{|\tau|=j+1} P(\rho(j+1) = \tau | \rho(j) = \mu) \left(\sum_{x \in \tau} c(x) \right)^2 \\ &= -\binom{j+1}{2} + \sum_{|\tau|=j+1} P(\rho(j+1) = \tau | \rho(j) = \mu) \\ & \quad \cdot \left(\sum_{x \in \mu} c(x) \right)^2 + c(y)^2 + 2c(y) \sum_{x \in \mu} c(x), \end{aligned}$$

where y is the box of τ not in μ . Since the expected value of $c(y)$ given μ is 0 and the expected value of $c(y)^2$ given μ is j , it follows that the last expression is equal to $Y_j^2 - \binom{j}{2}$, as desired. \square

4. JACK POLYNOMIALS AND MARTINGALES

The purpose of this section is to extend to Jack measure the connections with martingales in Section 2. The arguments are (by necessity) combinatorial as opposed to the algebraic arguments used in Section 2, and we assume that the reader is familiar with symmetric functions as in [Ma].

As in the Introduction, given a box s in the diagram of λ , let $a(s)$ and $l(s)$ denote the arm and leg of s , respectively. One defines quantities

$$(1) \quad c_\lambda(\alpha) = \prod_{s \in \lambda} (\alpha a(s) + l(s) + 1),$$

$$(2) \quad c'_\lambda(\alpha) = \prod_{s \in \lambda} (\alpha a(s) + l(s) + \alpha).$$

Let $m_i(\lambda)$ be the number of parts (i.e. rows) of λ of size i and let $l(\lambda)$ denote the total number of parts of λ . The symbol z_λ denotes $\prod_{i \geq 1} i^{m_i(\lambda)} m_i(\lambda)!$, the size of the centralizer of a permutation of cycle type λ in the symmetric group.

Let $\theta_\mu^\lambda(\alpha)$ be the coefficient of the power sum symmetric function p_μ in $J_\lambda^{(\alpha)}$. These will be the analogs of $\frac{|C|X^\lambda(C)}{\dim(\lambda)}$ studied in Section 2. In fact when $\alpha = 1$ they specialize to $\frac{|C|X^\lambda(C)}{\dim(\lambda)}$, where C is a conjugacy class of the symmetric group of type μ . When $\alpha = 2$ the $\theta_\mu^\lambda(\alpha)$ are proportional to the spherical functions for the Gelfand pair (S_{2n}, H_{2n}) , where S_{2n} is a symmetric group and H_{2n} is the hyperoctahedral group of size $2^n n!$.

Next we consider the ring of symmetric functions, with the inner product defined by the orthogonality condition $\langle p_\nu, p_\mu \rangle_\alpha = \delta_{\nu, \mu} z_\mu \alpha^{l(\mu)}$. For a symmetric function f , its adjoint f^\perp is defined by the condition $\langle fg, h \rangle_\alpha = \langle g, f^\perp h \rangle_\alpha$ for all g, h in the ring of symmetric functions. It is straightforward to check using the basis of power sum symmetric functions that $p_1^\perp = \alpha \frac{\partial}{\partial p_1}$ (for the case $\alpha = 1$ see page 76 of [Ma]).

Let

$$\psi'_{\lambda/\tau} = \prod_{s \in C_{\lambda/\tau} - R_{\lambda/\tau}} \frac{(\alpha a_\lambda(s) + l_\lambda(s) + 1) (\alpha a_\tau(s) + l_\tau(s) + \alpha)}{(\alpha a_\lambda(s) + l_\lambda(s) + \alpha) (\alpha a_\tau(s) + l_\tau(s) + 1)},$$

where $C_{\lambda/\tau}$ is the union of columns of λ that intersect $\lambda - \tau$ and $R_{\lambda/\tau}$ is the union of rows of λ that intersect $\lambda - \tau$.

The following lemma will be useful.

Lemma 4.1. (1) ([Ma], page 382)

$$\sum_{|\rho|=n} \frac{\theta_\mu^\rho(\alpha)\theta_\eta^\rho(\alpha)}{c_\rho(\alpha)c'_\rho(\alpha)} = \delta_{\mu,\eta} \frac{1}{z_\mu \alpha^{l(\mu)}}.$$

(2) ([Ma], page 340)

$$p_1 J_\lambda^{(\alpha)} = \sum_{|\Lambda|=|\lambda|+1} \frac{c_\lambda(\alpha)}{c_\Lambda(\alpha)} \psi'_{\Lambda/\lambda}(\alpha) J_\Lambda^{(\alpha)}.$$

(3) [F2] $p_1^\perp J_\lambda^{(\alpha)} = \sum_{|\tau|=|\lambda|-1} \frac{c'_\lambda(\alpha)\psi'_{\lambda/\tau}(\alpha)}{c'_\tau(\alpha)} J_\tau^{(\alpha)}$.

Using Lemma 4.1 we establish some relations concerning the $\theta_\mu^\lambda(\alpha)$'s. We use the notation that $\mu + 1^b$ denotes the partition of $|\mu| + b$ given by adding b parts of size 1 to μ , and that $\mu - 1^b$ denotes the partition of $|\mu| - b$ given by removing b parts of size 1 from μ (if this is possible). If μ has fewer than b parts of size 1, we define $\theta_{\mu-1^b}^\lambda(\alpha)$ to be 0 for all λ . We let $m_1(\mu)$ denote the number of parts of μ of size 1.

Lemma 4.2. (1) Let $|\lambda| = n$ and $|\mu| = n + 1$. Then

$$\theta_{\mu-1}^\lambda(\alpha) = \sum_{|\Lambda|=n+1} \frac{c_\lambda(\alpha)}{c_\Lambda(\alpha)} \psi'_{\Lambda/\lambda}(\alpha) \theta_\mu^\Lambda(\alpha).$$

(2) Let $|\lambda| = |\mu| = n$. Then

$$\alpha m_1(\mu) \theta_\mu^\lambda(\alpha) = \sum_{|\tau|=n-1} \frac{c'_\lambda(\alpha)}{c'_\tau(\alpha)} \psi'_{\lambda/\tau}(\alpha) \theta_{\mu-1}^\tau(\alpha).$$

Proof. For the first assertion, consider the inner product $\langle p_1 J_\lambda^{(\alpha)}, p_\mu \rangle$. On one hand, by part (2) of Lemma 4.1 it is equal to

$$\begin{aligned} & \left\langle \sum_{|\Lambda|=n+1} \frac{c_\lambda(\alpha)}{c_\Lambda(\alpha)} \psi'_{\Lambda/\lambda}(\alpha) J_\Lambda^{(\alpha)}, p_\mu \right\rangle \\ &= \sum_{|\Lambda|=n+1} \frac{c_\lambda(\alpha)}{c_\Lambda(\alpha)} \psi'_{\Lambda/\lambda}(\alpha) \theta_\mu^\Lambda(\alpha) z_\mu \alpha^{l(\mu)}. \end{aligned}$$

On the other hand it is equal to

$$\langle J_\lambda^{(\alpha)}, p_1^\perp p_\mu \rangle = \langle J_\lambda^{(\alpha)}, \alpha \frac{\partial}{\partial p_1} p_\mu \rangle.$$

If $m_1(\mu) = 0$ this is 0, so the assertion is true. Otherwise it is

$$\langle J_\lambda^{(\alpha)}, \alpha(m_1(\mu)) p_{\mu-1} \rangle = \alpha(m_1(\mu)) z_{\mu-1} \alpha^{l(\mu)-1} \theta_{\mu-1}^\lambda(\alpha).$$

Comparing these two expressions and using the fact that $z_\mu = (m_1(\mu)) z_{\mu-1}$ proves the first assertion.

For the second assertion, note that both sides vanish if $m_1(\mu) = 0$. Otherwise, consider the inner product $\langle p_1^\perp J_\lambda^{(\alpha)}, p_{\mu-1} \rangle$. On one hand, by part (3) of Lemma 4.1

it is equal to

$$\begin{aligned} & \left\langle \sum_{|\tau|=n-1} \frac{c'_\lambda(\alpha)}{c'_\tau(\alpha)} \psi'_{\lambda/\tau}(\alpha) J_\tau^{(\alpha)}, p_{\mu-1} \right\rangle \\ &= \sum_{|\tau|=n-1} \frac{c'_\lambda(\alpha)}{c'_\tau(\alpha)} \psi'_{\lambda/\tau}(\alpha) \theta_{\mu-1}^\tau(\alpha) z_{\mu-1} \alpha^{l(\mu)-1}. \end{aligned}$$

On the other hand, it is equal to

$$\langle J_\lambda^{(\alpha)}, p_\mu \rangle = \theta_\mu^\lambda(\alpha) z_\mu \alpha^{l(\mu)}.$$

Comparing these two expressions and using the fact that $z_\mu = m_1(\mu) z_{\mu-1}$ proves the second assertion. \square

In order to have analogs of the results of Section 2, it is necessary to have an analog of the growth process for representations of the symmetric group. Fortunately, such a process has been developed by Kerov [K4]. His process is best understood in terms of harmonic functions on Bratelli diagrams, but rather than going into this (see [K4] or Sections 2.1 and 4 of [F2] for details) we simply describe in the next paragraph the process and state the properties we need.

It is convenient to define $dim_\alpha(\lambda) = \frac{n! \alpha^n}{c'_\lambda(\alpha)}$, which in the case $\alpha = 1$ reduces to the dimension of the irreducible representation of the symmetric group parameterized by λ . A result of Stanley [St] is that $dim_\alpha(\lambda) = \sum_{|\tau|=n-1} \psi'_{\lambda/\tau}(\alpha) dim_\alpha(\tau)$. In Kerov's growth process the chance of transitioning from a partition λ of size n to a partition Λ of size $n + 1$ is equal to

$$\frac{\psi'_{\Lambda/\lambda}(\alpha) dim_\alpha(\lambda) Jack_\alpha(\Lambda)}{dim_\alpha(\Lambda) Jack_\alpha(\lambda)} = \frac{c_\lambda(\alpha)}{c_\Lambda(\alpha)} \psi'_{\Lambda/\lambda}(\alpha).$$

If λ is chosen from $Jack_\alpha$ measure on partitions of size n , then after this transition Λ is chosen from $Jack_\alpha$ measure on partitions of size $n + 1$. Moreover, there is a way of transitioning from a partition λ of size n to a partition τ of size $n - 1$. The transition to τ occurs with probability

$$\frac{dim_\alpha(\tau) \psi'_{\lambda/\tau}(\alpha)}{dim_\alpha(\lambda)} = \frac{\psi'_{\lambda/\tau}(\alpha) c'_\lambda(\alpha)}{\alpha n c'_\tau(\alpha)}.$$

If λ is chosen from $Jack_\alpha$ measure on partitions of size n , then after this transition τ is chosen from $Jack_\alpha$ measure on partitions of size $n - 1$.

Now we can define analogs of the random variables Y_1, \dots, Y_n of Section 2. Let μ be a partition of n , and let $\lambda(1), \dots, \lambda(n)$ be a sequence of random partitions generated by Kerov's growth process starting from $\lambda(1)$ equal to the only partition of size 1. Now define $Y_j^{(\alpha)} = \theta_{\mu-1^{n-j}}^{\lambda(j)}(\alpha)$, where this is understood to be 0 if μ has fewer than $n - j$ parts of size 1, or if $j = 0$.

Theorem 4.3. *For any partition μ of n , the sequence $(Y_1^{(\alpha)}, \dots, Y_n^{(\alpha)})$ is a martingale.*

Proof. It is necessary to show that for all λ of size j ,

$$\sum_{|\Lambda|=j+1} \psi'_{\Lambda/\lambda}(\alpha) \frac{c_\lambda(\alpha)}{c_\Lambda(\alpha)} \theta_{\mu-1^{n-j-1}}^\Lambda(\alpha) = \theta_{\mu-1^{n-j}}^\lambda(\alpha).$$

But this is clear from part (1) of Lemma 4.2. \square

Proposition 4.4 is interesting because it shows that the martingale of Theorem 4.3 has special properties. However it will not be needed in what follows.

Proposition 4.4. *Let μ be a partition of n . Let $x_{\downarrow i}$ denote $(x)(x-1)\cdots(x-i+1)$ or 1 in the case that $i = 0$. Then for all $1 \leq j \leq n$,*

$$E^{Y_n^{(\alpha)}}(Y_j^{(\alpha)}) = Y_n^{(\alpha)} \left(\frac{m_1(\mu)_{\downarrow n-j}}{n_{\downarrow n-j}} \right).$$

Proof. In fact we show that

$$E^{\lambda(n)}(Y_j^{(\alpha)}) = \theta_{\mu}^{\lambda(n)}(\alpha) \left(\frac{m_1(\mu)_{\downarrow n-j}}{n_{\downarrow n-j}} \right),$$

which is stronger.

As the case $j = n$ is clear, we first treat the case $j = n - 1$. Observe that

$$E^{\lambda(n)}(Y_{n-1}^{(\alpha)}) = \sum_{|\tau|=n-1} \frac{P(\lambda(n-1) = \tau, \lambda(n) = \lambda)}{P(\lambda(n) = \lambda)} \theta_{\mu-1}^{\tau}(\alpha).$$

From the description of Kerov’s transition mechanism before the proof of Theorem 4.3, it is clear that $\frac{P(\lambda(n-1)=\tau, \lambda(n)=\lambda)}{P(\lambda(n)=\lambda)}$ is simply the probability of transitioning down from λ to τ , which is $\frac{\psi'_{\lambda/\tau}(\alpha)c'_{\lambda}(\alpha)}{\alpha n c'_{\tau}(\alpha)}$. Substituting this into the expression for $E^{\lambda(n)}(Y_{n-1}^{(\alpha)})$ shows that the latter is equal to

$$\sum_{|\tau|=n-1} \theta_{\mu-1}^{\tau}(\alpha) \frac{\psi'_{\lambda/\tau}(\alpha)c'_{\lambda}(\alpha)}{\alpha n c'_{\tau}(\alpha)}.$$

The case $j = n - 1$ now follows from part 2 of Lemma 4.2.

Next we indicate how to treat the case $j = n - 2$, from which the general argument is clear. By considerations similar to those in the previous paragraph, $E^{\lambda(n)}(Y_{n-2}^{(\alpha)})$ is equal to

$$\begin{aligned} & \sum_{|\tau|=n-2} \frac{P(\rho(n-2) = \tau, \rho(n) = \lambda)}{P(\rho(n) = \lambda)} \theta_{\mu-1^2}^{\tau}(\alpha) \\ &= \sum_{|\tau|=n-2} \sum_{|\eta|=n-1} \frac{P(\rho(n-2) = \tau, \rho(n-1) = \eta, \rho(n) = \lambda)}{P(\rho(n-1) = \eta, \rho(n) = \lambda)} \\ & \quad \cdot \frac{P(\rho(n-1) = \eta, \rho(n) = \lambda)}{P(\rho(n) = \lambda)} \theta_{\mu-1^2}^{\tau}(\alpha). \end{aligned}$$

Arguing as in the previous paragraph, one sees that

$$\begin{aligned} & \sum_{|\tau|=n-2} \frac{P(\rho(n-2) = \tau, \rho(n-1) = \eta, \rho(n) = \lambda)}{P(\rho(n-1) = \eta, \rho(n) = \lambda)} \theta_{\mu-1^2}^{\tau}(\alpha) \\ &= \sum_{|\tau|=n-2} \frac{\psi'_{\eta/\tau}(\alpha)c'_{\eta}(\alpha)}{\alpha(n-1)c'_{\tau}(\alpha)} \theta_{\mu-1^2}^{\tau}(\alpha) \\ &= \left(\frac{m_1(\mu) - 1}{n - 1} \right) \theta_{\mu-1}^{\eta}(\alpha). \end{aligned}$$

The result now follows by the fact in the previous paragraph that

$$\sum_{|\eta|=n-1} \frac{P(\rho(n-1) = \eta, \rho(n) = \lambda)}{P(\rho(n) = \lambda)} \theta_{\mu-1}^\eta(\alpha) = \frac{m_1(\mu)}{n} \theta_\mu^\lambda(\alpha).$$

□

Proposition 4.5. *Let μ be any partition of n . Then*

$$E(Y_j^{(\alpha)} - Y_{j-1}^{(\alpha)})^2 = \begin{cases} 0 & \text{if } m_1(\mu) < n - j, \\ \frac{\alpha^{n-l(\mu)}(j-1)!}{z_{\mu-1}^{n-j}} (n - m_1(\mu)) & \text{otherwise.} \end{cases}$$

Proof. If $m_1(\mu) < n - j$ the result is clear since then by definition $Y_j^{(\alpha)} = Y_{j-1}^{(\alpha)} = 0$. As in the proof of Proposition 2.3, the martingale property of $(Y_1^{(\alpha)}, \dots, Y_n^{(\alpha)})$ implies that

$$E(Y_j^{(\alpha)} - Y_{j-1}^{(\alpha)})^2 = E(Y_j^{(\alpha)})^2 - E(Y_{j-1}^{(\alpha)})^2.$$

Now one uses part (1) of Lemma 4.1 to compute that

$$E(Y_j^{(\alpha)})^2 = \sum_{|\tau|=j} \frac{\alpha^j j!}{c_\tau(\alpha) c'_\tau(\alpha)} (\theta_{\mu-1}^\tau(\alpha))^2 = \frac{\alpha^{n-l(\mu)} j!}{z_{\mu-1}^{n-j}}.$$

Similarly one has that $E(Y_{j-1}^{(\alpha)})^2 = 0$ if $m_1(\mu) = n - j$ and computes that otherwise it is equal to

$$\frac{\alpha^{n-l(\mu)}(j-1)!}{z_{\mu-1}^{n-j+1}} = \frac{m_1(\mu) - (n - j)}{j} \frac{\alpha^{n-l(\mu)} j!}{z_{\mu-1}^{n-j}}.$$

□

5. “TRANSPOSITIONS”

This section examines and sharpens the results of Section 4 in the case when $\mu = (2, 1^{n-2})$. We call this partition the class of “transpositions” even though the algebraic structure of conjugacy classes is not present. As in Section 3, there are certain phenomena which occur only for the partition $\mu = (2, 1^{n-2})$.

To begin note that there is an explicit formula ([Ma], page 384)

$$\theta_{(2,1^{n-2})}^\lambda(\alpha) = \sum_i \left(\alpha \binom{\lambda_i}{2} - \binom{\lambda'_i}{2} \right),$$

where λ_i is the length of row i of λ and λ'_i is the length of column i of λ . From this formula it follows that if λ is obtained by adding a box x to μ , then

$$\theta_{(2,1^{n-2})}^\lambda(\alpha) - \theta_{(2,1^{n-3})}^\mu(\alpha) = c_\alpha(x),$$

where $c_\alpha(x)$ is the α -content of x defined as $\alpha(\text{column number of } x - 1) - (\text{row number of } x - 1)$.

Consequently Theorem 4.3 implies that for any partition $\lambda(j)$ of j , the expected value of the α -content of the box added to λ in moving from $\lambda(j)$ to $\lambda(j+1)$ is 0. This combinatorial fact was first established by Kerov [K4], whose proof method was completely different. In the same paper Kerov also proved that for any λ of size j , the variance of the α -content of the box added to λ is equal to $\alpha|\lambda|$. This is stronger than Proposition 4.5, which implies this for a random Jack_α distributed λ , but unlike Kerov’s result has the merit of being true for arbitrary partitions μ .

Theorem 5.1 is a recent result of Lassalle.

Theorem 5.1 ([La], Theorem 8.1). *Let $s_{r,\alpha}(\lambda)$ be the r th moment of the α -content of the box added to $\lambda(j) = \lambda$ when transitioning to $\lambda(j + 1)$. Let*

$$d_\rho(\lambda) = \prod_{i \geq 1} \left(\sum_{x \in \lambda} c_\alpha(x)^i \right)^{m_i(\rho)}.$$

Then $s_{r,\alpha}(\lambda)$ is equal to

$$\alpha^r \sum_{i=0}^{\lfloor r/2 \rfloor} \sum_{h=0}^{r-2i} \sum_{k=0}^{\min(i,h)} \frac{1}{\alpha^{i+h}} \left(1 - \frac{1}{\alpha}\right)^{r-2i-h} \binom{|\lambda| + i - 1}{i - k} \sum_{\substack{|\rho|=h \\ l(\rho) \leq k}} u_{ihk}^\rho(r) \frac{d_\rho(\lambda)}{z_\rho},$$

where the coefficients $u_{ihk}^\rho(r)$ are certain positive integers discussed in [La].

For instance one has that

$$\begin{aligned} s_{1,\alpha}(\lambda) &= 0, \\ s_{2,\alpha}(\lambda) &= \alpha|\lambda|, \\ s_{3,\alpha}(\lambda) &= 2\alpha d_{1,\alpha}(\lambda) + \alpha(\alpha - 1)|\lambda|, \\ s_{4,\alpha}(\lambda) &= 3\alpha d_{2,\alpha}(\lambda) + 3\alpha(\alpha - 1)d_{1,\alpha}(\lambda) + \alpha^2 \binom{|\lambda|+1}{2} + \alpha(\alpha - 1)^2|\lambda|. \end{aligned}$$

In order to use Theorem 5.1 to analyze the Jack_α average of $s_{r,\alpha}(\lambda)$, it is necessary to upper bound the Jack_α average of $d_\rho(\lambda)$. This is done in Lemma 5.2.

Lemma 5.2. *Suppose that $\alpha \geq 1$. Then if ρ is a partition of size h with at most k parts, there is a constant $C_{k,h,\alpha}$ (depending only on k, h, α) such that if λ is chosen from Jack_α measure on partitions of size n , then*

$$E(d_\rho(\lambda)) \leq C_{k,h,\alpha} n^{k+\frac{h}{2}}.$$

Proof. From Lemma 6.6 of [F2] it follows that for α fixed, the Jack_α probability that $\lambda_1 \geq 2e\sqrt{\frac{n}{\alpha}}$ or $\lambda'_1 \geq 2e\sqrt{\alpha n}$ decays exponentially as a function of \sqrt{n} . Thus excluding this unlikely event, one sees that the α -content of any of the n boxes of λ is at most $2e\sqrt{\alpha n}$ in absolute value. The lemma follows easily, since $d_i(\lambda) \leq n(2e\sqrt{\alpha n})^i$. \square

Next we obtain an upper bound on the moments $E(Y_{j+1}^{(\alpha)} - Y_j^{(\alpha)})^{2r}$.

Proposition 5.3. *Suppose that $\alpha \geq 1$. Then there is a constant $C_{2r,\alpha}$ (depending only on $2r$ and α) such that $E(Y_{j+1}^{(\alpha)} - Y_j^{(\alpha)})^{2r} \leq C_{2r,\alpha} j^r$ for all $j \geq 1$.*

Proof. By Theorem 5.1 and Lemma 5.2, it follows that for fixed values of i, h, k , the Jack_α average of the corresponding term in Lassalle’s formula for $s_{2r,\alpha}(\lambda)$ where λ has size j is at most

$$C'_{2r,\alpha} j^{i-k} j^{k+\frac{h}{2}} \leq C'_{2r,\alpha} j^r.$$

Here $C'_{2r,\alpha}$ is a constant depending only on $2r, \alpha$. The result follows. \square

To close this section, we give a generalization of Proposition 3.4.

Proposition 5.4. *Let $(Y_1^{(\alpha)}, \dots, Y_n^{(\alpha)})$ be the martingale of Theorem 4.3 corresponding to the choice $\mu = (2, 1^{n-2})$. Then*

$$\left((Y_1^{(\alpha)})^2, (Y_2^{(\alpha)})^2 - \alpha \binom{2}{2}, \dots, (Y_n^{(\alpha)})^2 - \alpha \binom{n}{2} \right)$$

is a martingale.

Proof. The proof method is the same as in Proposition 3.4; one replaces all occurrences of $c(x)$ by $c_\alpha(x)$ and $\binom{j}{2}$ by $\alpha\binom{j}{2}$. One also needs the facts that the expected value of $c_\alpha(y)$ given μ is 0 and the expected value of $c_\alpha(y)^2$ given μ is $\alpha|\mu|$. \square

6. CENTRAL LIMIT THEORY FOR MARTINGALES

This section uses martingale theory to prove a central limit theorem for $\frac{\theta^\lambda_{(2,1^{n-2})}(\alpha)}{\sqrt{\alpha\binom{n}{2}}}$ under Jack_α measure. Namely we prove Theorem 1.5 (and hence also Theorem 1.3 which is a special case).

We apply the following recent result of Haeusler [Ha]. For general background on central limit theorems for martingales, see the references in the introduction of [Ha], most notably [Bol] and [HH].

Theorem 6.1 ([Ha]). *Let the real-valued random variables X_1, \dots, X_n be a square integrable martingale difference sequence with respect to the σ -fields $F_0 \subset F_1 \cdots \subset F_n$. In other words we suppose that $E(X_j^2) < \infty$ and $E(X_j|F_{j-1}) = 0$ for all j . For $\delta > 0$ let*

$$L_{n,2\delta} = \sum_{j=1}^n E(|X_j|^{2+2\delta})$$

and

$$N_{n,2\delta} = E \left(\left| \sum_{j=1}^n E(X_j^2|F_{j-1}) - 1 \right|^{1+\delta} \right).$$

Then there is a constant C_δ depending only on δ such that for all real x_0 ,

$$\left| P(X_1 + \dots + X_n \leq x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \leq C_\delta(L_{n,2\delta} + N_{n,2\delta})^{1/(3+2\delta)}.$$

To prove Theorem 1.5 we will apply Theorem 6.1 with $X_j = \frac{Y_j^{(\alpha)} - Y_{j-1}^{(\alpha)}}{\sqrt{\alpha\binom{n}{2}}}$ and $\mu = (2, 1^{n-2})$. In this case (the Jack analog of the character ratio on transpositions), two major simplifications will occur: the quantity $N_{n,2\delta}$ is in fact 0, and the quantity $L_{n,2\delta}$ can be bounded using Proposition 5.3. Bounding these quantities for other partitions seems to be a difficult problem, but as mentioned in the Introduction the class of transpositions seems to have unique importance.

Proof of Theorem 1.5. We apply Theorem 6.1 with $X_j = \frac{Y_j^{(\alpha)} - Y_{j-1}^{(\alpha)}}{\sqrt{\alpha\binom{n}{2}}}$ and $\mu = (2, 1^{n-2})$. Let δ be a positive integer. By work of Kerov [K4], $s_{2,\alpha}(\lambda) = \alpha|\lambda|$ is independent of λ , so $N_{n,2\delta} = 0$. By Proposition 5.3,

$$L_{n,2\delta} \leq \left(\frac{1}{\sqrt{\alpha\binom{n}{2}}} \right)^{2+2\delta} \sum_{j=1}^n C_{2\delta+2,\alpha} j^{\delta+1} \leq \frac{C'_{2\delta+2,\alpha}}{n^\delta}.$$

Here $C'_{2\delta+2,\alpha}$ is a constant depending only on δ and α . Now observe that for any $s < \frac{1}{2}$, one can find δ large enough so that $(\frac{1}{n^\delta})^{1/(3+2\delta)} < n^{-s}$. \square

Remark. The reader might be inclined to conjecture that any martingale sequence (for instance ours) with $N_{n,2\delta} = 0$ and with $\|X_j\|_2, \|X_j\|_3$ uniformly bounded would yield a central limit theorem with convergence rate of order $n^{-1/2}$. Grams [Gr] shows that such sequences give central limit theorems with convergence rate of order $n^{-1/4}$, but Bolthausen [Bol] gives examples showing that even with these hypotheses one cannot in general beat the $n^{-1/4}$ rate.

7. PROBABILISTIC PROOF OF BURNSIDE-BRAUER

In order to study the distribution of character ratios and their Jack analogs by Stein's method, the paper [F2] showed how any representation of a finite group G with real-valued character can be used to construct a natural Markov chain on the set of irreducible representations of G . The purpose of this section is to show how this Markov chain gives a probabilistic proof of the following result of Burnside and then Brauer (see [Is]), in the special case that χ is real valued.

Theorem 7.1 (Burnside, Brauer). *Let η be the character of a faithful representation of a finite group G . Let m be the number of distinct values assumed by η . Then every irreducible character of G occurs as a constituent of some η^j , where $0 \leq j < m$.*

In fact it will be shown that this result is closely related to a generalization of the following fact from algebraic graph theory.

Theorem 7.2 ([Big]). *A connected graph with diameter d has at least $d+1$ distinct eigenvalues.*

To explain this, a dictionary is needed to go between representation theory and graph theory. This was developed in [F2] and we recall it. Let η be a real-valued character of a finite group G , and let $\langle \phi, \theta \rangle$ be the usual inner product on class functions of G defined as $\frac{1}{|G|} \sum_{g \in G} \phi(g)\overline{\theta(g)}$. Then one can define a Markov chain L_η on the set of irreducible representations of G which transitions from λ to ρ with probability

$$L_\eta(\lambda, \rho) = \frac{\dim(\rho)}{\eta(1)\dim(\lambda)} \langle \chi^\rho, \eta\chi^\lambda \rangle.$$

Note that this transition probability is nonnegative since $\langle \chi^\rho, \eta\chi^\lambda \rangle$ is the multiplicity of ρ in the tensor product of λ and the representation with character η .

Recall that a Markov chain on a finite set X with transition probability $K(x, y)$ is said to be reversible with respect to a probability measure π on X if $\pi(x)K(x, y) = \pi(y)K(y, x)$ for all x, y . It is straightforward to see that this condition implies that π is a stationary distribution for K (i.e. that $\pi(y) = \sum_x \pi(x)K(x, y)$ for all y). Moreover, consider the space of real-valued functions $\ell^2(\pi)$ with the norm

$$\|f\|_2 = \left(\sum_x |f(x)|^2 \pi(x) \right)^{1/2}$$

and let K denote the Markov operator on $\ell^2(\pi)$ defined by

$$Kf(x) = \sum_y K(x, y)f(y).$$

Then the operator K is self-adjoint and has real eigenvalues $-1 \leq \beta_{\min} = \beta_{|X|-1} \leq \dots \leq \beta_0 = 1$. If $\{\psi_i\}$ is an orthonormal basis of eigenfunctions with eigenvalues β_i ,

elementary linear algebra implies that the chance that the chain started at x is at y after r steps is $\sum_{i=0}^{|X|-1} \beta_i^r \psi_i(x) \psi_i(y) \pi(y)$.

Theorem 7.3 ([F2]).

- (1) The transition probabilities of L_η sum to 1, and the Markov chain L_η is reversible with respect to the Plancherel measure π of G .
- (2) The eigenvalues of the chain L_η are indexed by conjugacy classes C of G . The eigenvalue parameterized by C is the character ratio $\frac{\eta(C)}{\eta(1)}$, and an orthonormal basis of eigenfunctions ψ_C in $L^2(\pi)$ is defined by $\psi_C(\rho) = \frac{|C|^{1/2} \chi^\rho(C)}{\dim(\rho)}$.

One can think of any reversible Markov chain on a finite set as inducing a graph structure on that set, with the unordered edge (x, y) having weight $\pi(x)K(x, y)$. In the case at hand, the vertices of the graph are the irreducible representations of G , and an edge (λ, ρ) is given weight $\pi(\lambda)L_\eta(\lambda, \rho)$, where π is the Plancherel measure of G .

Now we give a proof of Theorem 7.1, in the case that η is real valued.

Proof. Suppose that η is real valued, so that Theorem 7.3 is applicable. By Theorem 7.3, the number of distinct eigenvalues of L_η is equal to the number of distinct character values of η . Moreover since η is faithful, the eigenvalue 1 occurs with multiplicity one, and it follows that the weighted graph associated to L_η is connected (since otherwise each connected component could be used to construct an eigenvalue 1 eigenfunction supported on that component). Thus by a straightforward generalization of the proof of Theorem 7.2 to graphs with edge weights, it follows that the diameter of the graph associated to η is less than m , the number of distinct character values assumed by η .

To complete the proof it is sufficient to show that χ^ρ occurs as a constituent in η^j if and only if there is a path of length j from the trivial representation to ρ in the weighted graph corresponding to L_η . But Theorem 7.3 gives a complete diagonalization of the Markov chain L_η , so by the sentence preceding the statement of Theorem 7.3, it follows that the chance that L_η transitions from the trivial representation to ρ in j steps is equal to

$$\begin{aligned} & \sum_C \left(\frac{\eta(C)}{\eta(1)} \right)^j |C| \frac{\chi^\rho(C)}{\dim(\rho)} \frac{\dim(\rho)^2}{|G|} \\ &= \frac{\dim(\rho)}{\eta(1)^j} \frac{1}{|G|} \sum_{g \in G} \eta^j(g) \chi^\rho(g). \end{aligned}$$

The result now follows since

$$\frac{1}{|G|} \sum_{g \in G} \eta^j(g) \chi^\rho(g)$$

is the multiplicity of χ^ρ in η^j . □

ACKNOWLEDGEMENTS

The author was partially supported by National Security Agency grant MDA904-03-1-0049. The author thanks Michel Lassalle for correspondence.

REFERENCES

- [AID] Aldous, D. and Diaconis, P., Longest increasing subsequences: from patience sorting to the Baik-Deift-Johansson theorem, *Bull. AMS (N.S.)* **36** (1999), 413-432. MR1694204 (2000g:60013)
- [Big] Biggs, N., *Algebraic graph theory*, Second edition, Cambridge Univ. Press, Cambridge, 1993. MR1271140 (95h:05105)
- [Bol] Bolthausen, E., Exact convergence rates in some martingale central limit theorems, *Annals of Probab.* **10** (1982), 672-688. MR0659537 (84a:60056)
- [BOO] Borodin, A., Okounkov, A., and Olshanski, G., Asymptotics of Plancherel measures for symmetric groups, *J. Amer. Math. Soc.* **13** (2000), 481-515. MR1758751 (2001g:05103)
- [BO] Borodin, A. and Olshanski, G., Z-measures on partitions and their scaling limits, *European J. Combin.* **26** (2005), 795-834. MR2143199
- [De] Deift, P., Integrable systems and combinatorial theory, *Notices Amer. Math. Soc.* **47** (2000), 631-640. MR1764262 (2001g:05012)
- [DSh] Diaconis, P. and Shahshahani, M., Generating a random permutation with random transpositions, *Z. Wahr. Verw. Gebiete* **57** (1981), 159-179. MR0626813 (82h:60024)
- [EO] Eskin, A. and Okounkov, A., Asymptotics of branched coverings of a torus and volumes of moduli spaces of holomorphic differentials, *Invent. Math.* **145** (2001), 59-103. MR1839286 (2002g:32018)
- [Fr] Frobenius, F., Uber die charaktere der symmetrischen gruppe, *Sitz. Konig. Preuss. Akad. Wissen.* (1900), 516-534; *Gesammelte abhandlungen III*, Springer-Verlag, Heidelberg, 1968, 148-166.
- [F1] Fulman, J., Stein's method and Plancherel measure of the symmetric group, *Trans. Amer. Math. Soc.* **357** (2005), no. 2, 555-570. MR2095623 (2005e:05156)
- [F2] Fulman, J., Stein's method, Jack measure, and the Metropolis algorithm, *J. Combin. Theory Ser. A.* **108** (2004), no. 2, 275-296. MR2098845
- [F3] Fulman, J., Card shuffling and the decomposition of tensor products, *Pacific J. Math.* **217** (2004), no. 2, 247-262. MR2109933
- [F4] Fulman, J., An inductive proof of the Berry-Esseen theorem for character ratios, to appear in *Annals of Combin.*
- [F5] Fulman, J., Stein's method and random character ratios, preprint math.CO/0508291 at <http://xxx.lanl.gov>.
- [GHJ] Goulden, I., Harer, J., and Jackson, D., A geometric parametrization for the virtual Euler characteristic of the moduli spaces of real and complex algebraic curves, *Trans. Amer. Math. Soc.* **353** (2001), 4405-4427. MR1851176 (2002g:14035)
- [Gr] Grams, W., *Rates of convergence in the central limit theorem for dependent variables*, Florida State University Thesis, 1972.
- [Ha] Haeusler, E., On the rate of convergence in the central limit theorem for martingales with discrete and continuous time, *Annals of Prob.* **16** (1988), 275-299. MR0920271 (89a:60060)
- [HH] Hall, P. and Heyde, C., *Martingale limit theory and its applications*, Academic Press, New York, 1980. MR0624435 (83a:60001)
- [Ho] Hora, A., Central limit theorem for the adjacency operators on the infinite symmetric group, *Comm. Math. Phys.* **195** (1998), 405-416. MR1637801 (99i:46058)
- [Is] Isaacs, I.M., *Character theory of finite groups*, Dover Publications, Inc., New York, 1976.
- [IO] Ivanov, V. and Olshanski, G., Kerov's central limit theorem for the Plancherel measure on Young diagrams, in *Symmetric Functions 2001: Surveys of developments and perspectives*, Kluwer Academic Publishers, Dordrecht, 2002. MR2059361 (2005d:05148)
- [J] Johansson, K., Discrete orthogonal polynomial ensembles and the Plancherel measure, *Ann. of Math. (2)* **153** (2001), 259-296. MR1826414 (2002g:05188)
- [K1] Kerov, S.V., Gaussian limit for the Plancherel measure of the symmetric group, *Compt. Rend. Acad. Sci. Paris, Serie I* **316** (1993), 303-308. MR1204294 (93k:20106)
- [K2] Kerov, S.V., The boundary of Young lattice and random Young tableaux, in *Formal power series and algebraic combinatorics*, DIMACS Ser. Discrete Math. Theoret. Comput. Sci. **24**, Amer. Math. Soc., Providence, RI (1996), 133-158. MR1363510 (96i:05177)
- [K3] Kerov, S.V., Transition probabilities of continual Young diagrams and the Markov moment problem, *Funct. Anal. Appl.* **27** (1993), 104-117. MR1251166 (95g:82045)

- [K4] Kerov, S.V., Anisotropic Young diagrams and Jack symmetric functions, *Funct. Anal. Appl.* **34** (2000), 41-51. MR1756734 (2001f:05158)
- [La] Lassalle, M., Jack polynomials and some identities for partitions, *Trans. Amer. Math. Soc.* **356** (2004), 3455-3476. MR2055741
- [Ma] Macdonald, I., *Symmetric functions and Hall polynomials*, Second edition, Oxford University Press, New York, 1995. MR1354144 (96h:05207)
- [O1] Okounkov, A., Random matrices and random permutations, *Internat. Math. Res. Notices* **20** (2000), 1043-1095. MR1802530 (2002c:15045)
- [O2] Okounkov, A., The uses of random partitions, preprint math-ph/0309015 at <http://xxx.lanl.gov>.
- [OP] Okounkov, A. and Pandaripandhe, R., Gromov-Witten theory, Hurwitz numbers, and Matrix models, I, preprint math.AG/0101147 at <http://xxx.lanl.gov>.
- [Sa] Sagan, B., *The symmetric group. Representations, combinatorial algorithms, and symmetric functions*, Springer-Verlag, New York, 1991.
- [Se] Serre, J.-P., *Linear representations of finite groups*, Springer-Verlag, New York, 1977. MR0450380 (56:8675)
- [ShSu] Shao, Q. and Su, Z., The Berry-Esseen bound for character ratios, preprint (2004).
- [St] Stanley, R., Some combinatorial properties of Jack symmetric functions, *Advances in Math.* **77**, 76-115. MR1014073 (90g:05020)
- [Su] Su, Z., The law of the iterated logarithm for character ratios, *Statist. Probab. Lett.* **71** (2005), 337-346. MR2145501 (2006b:60054)

DEPARTMENT OF MATHEMATICS, 301 THACKERAY HALL, UNIVERSITY OF PITTSBURGH, PITTSBURGH, PENNSYLVANIA 15260

E-mail address: fulman@math.pitt.edu