

CROSSINGS AND NESTINGS OF MATCHINGS AND PARTITIONS

WILLIAM Y. C. CHEN, EVA Y. P. DENG, ROSENA R. X. DU, RICHARD P. STANLEY,
AND CATHERINE H. YAN

ABSTRACT. We present results on the enumeration of crossings and nestings for matchings and set partitions. Using a bijection between partitions and vacillating tableaux, we show that if we fix the sets of minimal block elements and maximal block elements, the crossing number and the nesting number of partitions have a symmetric joint distribution. It follows that the crossing numbers and the nesting numbers are distributed symmetrically over all partitions of $[n]$, as well as over all matchings on $[2n]$. As a corollary, the number of k -noncrossing partitions is equal to the number of k -nonnesting partitions. The same is also true for matchings. An application is given to the enumeration of matchings with no k -crossing (or with no k -nesting).

1. INTRODUCTION

A (complete) matching on $[2n] = \{1, 2, \dots, 2n\}$ is a partition of $[2n]$ of type $(2, 2, \dots, 2)$. It can be represented by listing its n blocks as $\{(i_1, j_1), (i_2, j_2), \dots, (i_n, j_n)\}$, where $i_r < j_r$ for $1 \leq r \leq n$. Two blocks (also called arcs) (i_r, j_r) and (i_s, j_s) form a *crossing* if $i_r < i_s < j_r < j_s$; they form a *nesting* if $i_r < i_s < j_s < j_r$. It is well known that the number of matchings on $[2n]$ with no crossings (or with no nestings) is given by the n -th Catalan number

$$C_n = \frac{1}{n+1} \binom{2n}{n}.$$

See [25, Exercise 6.19] for many combinatorial interpretations of Catalan numbers, where item (o) is for noncrossing matchings, and item (ww) can be viewed as nonnesting matchings, in which the blocks of the matching are the columns of the standard Young tableaux of shape (n, n) . Nonnesting matchings are also one of the items of [26].

Let $k \geq 2$ be an integer. A k -crossing of a matching M is a set of k arcs $(i_{r_1}, j_{r_1}), (i_{r_2}, j_{r_2}), \dots, (i_{r_k}, j_{r_k})$ of M such that $i_{r_1} < i_{r_2} < \dots < i_{r_k} < j_{r_1} < j_{r_2} < \dots < j_{r_k}$. A matching without any k -crossing is a k -noncrossing matching. Similarly, a k -nesting is a set of k arcs $(i_{r_1}, j_{r_1}), (i_{r_2}, j_{r_2}), \dots, (i_{r_k}, j_{r_k})$ of M such that $i_{r_1} < i_{r_2}$

Received by the editors January 14, 2005.

2000 *Mathematics Subject Classification*. Primary 05A18; Secondary 05E10, 05A15.

Key words and phrases. Crossing, nesting, partition, vacillating tableau.

The first author was supported by the 973 Project on Mathematical Mechanization, the National Science Foundation, the Ministry of Education and the Ministry of Science and Technology of China.

The fourth author was supported in part by NSF grant #DMS-9988459.

The fifth author was supported in part by NSF grant #DMS-0245526 and a Sloan Fellowship.

$< \cdots < i_{r_k} < j_{r_k} < \cdots < j_{r_2} < j_{r_1}$. A matching without any k -nesting is a k -nonnesting matching.

Enumeration on crossings/nestings of matchings has been studied for the cases $k = 2$ and $k = 3$. For $k = 2$, in addition to the above results on Catalan numbers, the distribution of the number of 2-crossings has been studied by Touchard [29], and later more explicitly by Riordan [21], who gave a generating function. M. de Sainte-Catherine [8] proved that 2-crossings and 2-nestings are identically distributed over all matchings of $[2n]$, i.e., the number of matchings with r 2-crossings is equal to the number of matchings with r 2-nestings.

The enumeration of 3-nesting matchings was first studied in [11] by Gouyou-Beauschamps, where he gave a bijection between involutions with no decreasing sequence of length 6 and pairs of noncrossing Dyck left factors by a recursive construction. His bijection is essentially a correspondence between 3-nesting matchings and pairs of noncrossing Dyck paths, where a matching can also be considered as a fixed-point-free involution. We observed that the number of 3-nesting matchings also equals the number of pairs of noncrossing Dyck paths, and a one-to-one correspondence between 3-nesting matchings and pairs of noncrossing Dyck paths can be built recursively.

In this paper, we extend the above results. Let $\text{cr}(M)$ be maximal i such that M has an i -crossing, and $\text{ne}(M)$ the maximal j such that M has a j -nesting. Denote by $f_n(i, j)$ the number of matchings M on $[2n]$ with $\text{cr}(M) = i$ and $\text{ne}(M) = j$. We shall prove that $f_n(i, j) = f_n(j, i)$. As a corollary, the number of matchings on $[2n]$ with $\text{cr}(M) = k$ equals the number of matchings M on $[2n]$ with $\text{ne}(M) = k$.

Our construction applies to a more general structure, viz., partitions of a set. Given a partition P of $[n]$, denoted by $P \in \Pi_n$, we represent P by a graph on the vertex set $[n]$ whose edge set consists of arcs connecting the elements of each block in numerical order. Such an edge set is called the *standard representation* of the partition P . For example, the standard representation of 1457-26-3 is $\{(1, 4), (4, 5), (5, 7), (2, 6)\}$. Here we always write an arc e as a pair (i, j) with $i < j$, and say that i is the *left-hand endpoint* of e and j is the *right-hand endpoint* of e .

Let $k \geq 2$ and $P \in \Pi_n$. Define a k -crossing of P as a k -subset $(i_1, j_1), (i_2, j_2), \dots, (i_k, j_k)$ of the arcs in the standard representation of P such that $i_1 < i_2 < \cdots < i_k < j_1 < j_2 < \cdots < j_k$. Let $\text{cr}(P)$ be the maximal k such that P has a k -crossing. Similarly, define a k -nesting of P as a k -subset $(i_1, j_1), (i_2, j_2), \dots, (i_k, j_k)$ of the set of arcs in the standard representation of P such that $i_1 < i_2 < \cdots < i_k < j_k < \cdots < j_2 < j_1$, and $\text{ne}(P)$ the maximal j such that P has a j -nesting. Note that when restricted to complete matchings, these definitions agree with the ones given before.

Let $g_n(i, j)$ be the number of partitions P of $[n]$ with $\text{cr}(P) = i$ and $\text{ne}(P) = j$. We shall prove that $g_n(i, j) = g_n(j, i)$, for all i, j and n . In fact, our result is much stronger. We present a generalization which implies the symmetric distribution of $\text{cr}(P)$ and $\text{ne}(P)$ over all partitions in Π_n , as well over all complete matchings on $[2n]$.

To state the main result, we need some notation. Given $P \in \Pi_n$, define

$$\begin{aligned} \min(P) &= \{\text{minimal block elements of } P\}, \\ \max(P) &= \{\text{maximal block elements of } P\}. \end{aligned}$$

For example, for $P = 135-26-4$, $\min(P) = \{1, 2, 4\}$ and $\max(P) = \{4, 5, 6\}$. The pair $(\min(P), \max(P))$ encodes some useful information about the partition P . For example, the number of blocks of P is $|\min(P)| = |\max(P)|$; number of singleton blocks is $|\min(P) \cap \max(P)|$; P is a (partial) matching if and only if $\min(P) \cup \max(P) = [n]$, and P is a complete matching if in addition, $\min(P) \cap \max(P) = \emptyset$.

Fix $S, T \subseteq [n]$ with $|S| = |T|$. Let $P_n(S, T)$ be the set $\{P \in \Pi_n : \min(P) = S, \max(P) = T\}$, and let $f_{n,S,T}(i, j)$ be the cardinality of the set $\{P \in P_n(S, T) : \text{cr}(P) = i, \text{ne}(P) = j\}$.

Theorem 1.1.

$$(1.1) \quad f_{n,S,T}(i, j) = f_{n,S,T}(j, i).$$

In other words,

$$(1.2) \quad \sum_{P \in P_n(S,T)} x^{\text{cr}(P)} y^{\text{ne}(P)} = \sum_{P \in P_n(S,T)} x^{\text{ne}(P)} y^{\text{cr}(P)}.$$

That is, the statistics $\text{cr}(P)$ and $\text{ne}(P)$ have a symmetric joint distribution over each set $P_n(S, T)$. □

Summing over all pairs (S, T) in (1.1), we get

$$(1.3) \quad g_n(i, j) = g_n(j, i).$$

We say that a partition P is k -noncrossing if $\text{cr}(P) < k$. It is k -nonnesting if $\text{ne}(P) < k$. Let $\text{NCN}_{k,l}(n)$ be the number of partitions of $[n]$ that are k -noncrossing and l -nonnesting. Summing over $1 \leq i < k$ and $1 \leq j < l$ in (1.3), we get the following corollary.

Corollary 1.2. $\text{NCN}_{k,l}(n) = \text{NCN}_{l,k}(n)$. □

Letting $l > n$, Corollary 1.2 becomes the following result.

Corollary 1.3. $\text{NC}_k(n) = \text{NN}_k(n)$, where $\text{NC}_k(n)$ is the number of k -noncrossing partitions of $[n]$, and $\text{NN}_k(n)$ is the number of k -nonnesting partitions of $[n]$.

Theorem 1.1 also applies to complete matchings. A partition P of $[2n]$ is a complete matching if and only if $|\min(P)| = |\max(P)| = n$ and $\min(P) \cap \max(P) = \emptyset$. (It follows that $\min(P) \cup \max(P) = [2n]$.) Restricting Theorem 1.1 to disjoint pairs (S, T) of $[2n]$ with $|S| = |T| = n$, we get the following result on the crossing and nesting number of complete matchings.

Corollary 1.4. Let M be a matching on $[2n]$.

1. The statistics $\text{cr}(M)$ and $\text{ne}(M)$ have a symmetric joint distribution over $P_{2n}(S, T)$, where $|S| = |T| = n$, and S, T are disjoint.
2. $f_n(i, j) = f_n(j, i)$, where $f_n(i, j)$ is the number of matchings on $[2n]$ with $\text{cr}(M) = i$ and $\text{ne}(M) = j$.
3. The number of matchings on $[2n]$ that are k -noncrossing and l -nonnesting is equal to the number of matchings on $[2n]$ that are l -noncrossing and k -nonnesting.
4. The number of k -noncrossing matchings on $[2n]$ is equal to the number of k -nonnesting matchings on $[2n]$.

The paper is arranged as follows. In Section 2 we introduce the concept of vacillating tableau of general shape, and give a bijective proof for the number of vacillating tableaux of shape λ and length $2n$. In Section 3 we apply the bijection of Section 2 to vacillating tableaux of empty shape, and characterize crossings and

nestings of a partition by the corresponding vacillating tableau. The involution on the set of vacillating tableaux defined by taking the conjugate to each shape leads to an involution on partitions which exchanges the statistics $cr(P)$ and $ne(P)$ while preserving $min(P)$ and $max(P)$, thus proving Theorem 1.1. Then we modify the bijection between partitions and vacillating tableaux by taking isolated points into consideration, and give an analogous result on the enhanced crossing number and nesting number. This is the content of Section 4. Finally in Section 5 we restrict our bijection to the set of complete matchings and oscillating tableaux, and study the enumeration of k -noncrossing matchings. In particular, we construct bijections from k -noncrossing matchings for $k = 2$ or 3 to Dyck paths and pairs of noncrossing Dyck paths, respectively, and present the generating function for the number of k -noncrossing matchings.

2. A BIJECTION BETWEEN SET PARTITIONS AND VACILLATING TABLEAUX

Let Y be *Young's lattice*, that is, the set of all partitions of all integers $n \in \mathbb{N}$ ordered component-wise, i.e., $(\mu_1, \mu_2, \dots) \leq (\lambda_1, \lambda_2, \dots)$ if $\mu_i \leq \lambda_i$ for all i . We write $\lambda \vdash k$ or $|\lambda| = k$ if $\sum \lambda_i = k$. A vacillating tableau is a walk on the Hasse diagram of Young's lattice subject to certain conditions. The main tool in our proof of Theorem 1.1 is a bijection between the set of set partitions and the set of vacillating tableaux of empty shape \emptyset .

Definition 2.1. A *vacillating tableau* V_λ^{2n} of shape λ and length $2n$ is a sequence $\lambda^0, \lambda^1, \dots, \lambda^{2n}$ of integer partitions such that (i) $\lambda^0 = \emptyset$, and $\lambda^{2n} = \lambda$, (ii) λ^{2i+1} is obtained from λ^{2i} by doing nothing (i.e., $\lambda^{2i+1} = \lambda^{2i}$) or deleting a square, and (iii) λ^{2i} is obtained from λ^{2i-1} by doing nothing or adding a square.

In other words, a vacillating tableau of shape λ is a walk on the Hasse diagram of Young's lattice from \emptyset to λ where each step consists of either (i) doing nothing twice, (ii) do nothing then adding a square, (iii) removing a square then doing nothing, or (iv) removing a square and then adding a square. Note that if the length is larger than 0, $\lambda^1 = \emptyset$. If the vacillating tableau is of empty shape, then $\lambda^{2n-1} = \emptyset$ as well.

Example 2.2. Abbreviate $\lambda = (\lambda_1, \lambda_2, \dots)$ by $\lambda_1 \lambda_2 \dots$. There are 5 vacillating tableaux of shape \emptyset and length 6. They are

$$(2.1) \quad \begin{array}{ccccccc} \lambda^0 & \lambda^1 & \lambda^2 & \lambda^3 & \lambda^4 & \lambda^5 & \lambda^6 \\ \emptyset & \emptyset & \emptyset & \emptyset & \emptyset & \emptyset & \emptyset \\ \emptyset & \emptyset & \emptyset & \emptyset & 1 & \emptyset & \emptyset \\ \emptyset & \emptyset & 1 & \emptyset & 1 & \emptyset & \emptyset \\ \emptyset & \emptyset & 1 & \emptyset & \emptyset & \emptyset & \emptyset \\ \emptyset & \emptyset & 1 & 1 & 1 & \emptyset & \emptyset \end{array}$$

Example 2.3. An example of a vacillating tableau of shape 11 and length 10 is given by

$$\emptyset, \emptyset, 1, 1, 1, 1, 2, 2, 21, 11, 11.$$

Theorem 2.4. (i) Let $g_\lambda(n)$ be the number of vacillating tableaux of shape $\lambda \vdash k$ and length $2n$. By a standard Young tableau (SYT) of shape λ , we mean an array T of shape λ whose entries are distinct positive integers that increase in every row

and column. The content of T is the set of positive integers that appear in it. (We do not require that $\text{content}(\lambda) = [k]$, where $\lambda \vdash k$.) We then have

$$g_\lambda(n) = B(n, k)f^\lambda,$$

where f^λ is the number of SYT's of shape λ and content $[k]$, and $B(n, k)$ is the number of partitions of $[n]$ with k blocks distinguished.

(ii) The exponential generating function of $B(n, k)$ is given by

$$(2.2) \quad \sum_{n \geq 0} B(n, k) \frac{x^n}{n!} = \frac{1}{k!} (e^x - 1)^k \exp(e^x - 1).$$

To prove Theorem 2.4, we construct a bijection between the set \mathcal{V}_λ^{2n} of vacillating tableaux of shape λ and length $2n$, and pairs (P, T) , where P is a partition of $[n]$, and T is an SYT of shape λ such that $\text{content}(T) \subseteq \max(P)$. In the next section we apply this bijection to vacillating tableaux of empty shape, and relate it to the enumeration of crossing and nesting numbers of a partition. In the following we shall assume familiarity with the RSK algorithm, and use row-insertion $P \leftarrow k$ as the basic operation of the RSK algorithm. For the notation, as well as some basic properties of the RSK algorithm, see e.g. [25, Chapter 7]. In general we shall apply the RSK algorithm to a sequence w of distinct integers, denoted by $w \xrightarrow{\text{RSK}} (A(w), B(w))$, where $A(w)$ is the (row)-insertion tableau and $B(w)$ the recording tableau. The shape of the SYT's $A(w)$ and $B(w)$ is also called the *shape* of the sequence w .

The bijection ψ from vacillating tableaux to pairs (P, T) . Given a vacillating tableau $V = (\emptyset = \lambda^0, \lambda^1, \dots, \lambda^{2n} = \lambda)$, we will recursively define a sequence $(P_0, T_0), (P_1, T_1), \dots, (P_{2n}, T_{2n})$, where P_i is a set of ordered pairs of integers in $[n]$, and T_i is an SYT of shape λ^i . Let P_0 be the empty set, and let T_0 be the empty SYT (on the empty alphabet).

- (1) If $\lambda^i = \lambda^{i-1}$, then $(P_i, T_i) = (P_{i-1}, T_{i-1})$.
- (2) If $\lambda^i \supset \lambda^{i-1}$, then $i = 2k$ for some integer $k \in [n]$. In this case let $P_i = P_{i-1}$ and T_i is obtained from T_{i-1} by adding the entry k in the square $\lambda^i \setminus \lambda^{i-1}$.
- (3) If $\lambda^i \subset \lambda^{i-1}$, then $i = 2k - 1$ for some integer $k \in [n]$. In this case let T_i be the unique SYT (on a suitable alphabet) of shape λ^i such that T_{i-1} is obtained from T_i by row-inserting some number j . Note that j must be less than k . Let P_i be obtained from P_{i-1} by adding the ordered pair (j, k) .

It is clear from the above construction that (i) $P_0 \subseteq P_1 \subseteq \dots \subseteq P_{2n}$, (ii) for each integer i , it appears at most once as the first component of an ordered pair in P_{2n} , and appears at most once as the second component of an ordered pair in P_{2n} . Let $\psi(V) = (P, T_{2n})$, where P is the partition on $[n]$ whose standard representation is P_{2n} .

Note that if an integer i appears in T_{2n} , then P_{2n} cannot contain any ordered pair (i, j) with $i < j$. It follows that i is the maximal element in the block containing it. Hence the content of T_{2n} is a subset of $\max(P)$.

Example 2.5. As an example of the map ψ , let the vacillating tableau be

$$\emptyset, \emptyset, 1, 1, 2, 2, 2, 2, 2, 2, 21, 21, 211, 21, 21, 11, 21.$$

Then the pairs (B_i, T_i) (where B_i is the pair added to P_{i-1} to obtain P_i) are given by

i	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
T_i	\emptyset	\emptyset	1	1	12	12	12	12	$\frac{12}{4}$	$\frac{12}{4}$	$\frac{12}{5}$	$\frac{14}{5}$	$\frac{14}{5}$	$\frac{1}{5}$	$\frac{17}{5}$
B_i													(2, 6)	(4, 7)	

Hence

$$T = \frac{1}{5} 7, \quad P = 1-26-3-47-5.$$

The map ψ is bijective since the above construction can be reversed. Given a pair (P, T) , where P is a partition of $[n]$ and T is an SYT whose content consists of maximal elements of some blocks of P , let $E(P)$ be the standard representation of P , and $T_{2n} = T$. We work our way backwards from T_{2n} , reconstructing the preceding tableaux and hence the sequence of shapes. If we have the SYT T_{2k} for some $k \leq n$, we can get the tableaux T_{2k-1}, T_{2k-2} by the following rules:

- (1) $T_{2k-1} = T_{2k}$ if the integer k does not appear in T_{2k} . Otherwise T_{2k-1} is obtained from T_{2k} by deleting the square containing k .
- (2) $T_{2k-2} = T_{2k-1}$ if $E(P)$ does not have an edge of the form (i, k) . Otherwise there is a unique $i < k$ such that $(i, k) \in E(P)$. In that case let T_{2k-2} be obtained from T_{2k-1} by row-inserting i , or equivalently, $T_{2k-2} = (T_{2k-1} \leftarrow i)$.

Proof of Theorem 2.4. Part (i) follows from the bijection ψ , where a block of P is distinguished if its maximal element belongs to $\text{content}(T)$. For part (ii), simply note that to get a structure counted by $B(n, k)$, we can partition $[n]$ into two subsets, S and T , and then partition S into k blocks and put a mark on each block, and partition T arbitrarily. The generating function of $B(n, k)$ then follows from the well-known generating functions for $S(n, k)$, the Stirling number of the second kind, and for the Bell number $B(n)$,

$$\sum_{n \geq k} S(n, k) \frac{x^n}{n!} = \frac{1}{k!} (e^x - 1)^k, \quad \sum_{n \geq 0} B(n) \frac{x^n}{n!} = \exp(e^x - 1). \quad \square$$

Remark 2.6. (1) Restricting to vacillating tableaux of empty shape, the map ψ provides a bijection between the set $\mathcal{V}_\emptyset^{2n}$ of vacillating tableaux of empty shape and length $2n$ and the set of partitions of $[n]$. In particular, $g_\emptyset(n)$, the cardinality of $\mathcal{V}_\emptyset^{2n}$, is equal to the n -th Bell number $B(n)$.

(2) Note that there is a symmetry between the four types of movements in the definition of vacillating tableaux. Thus any walk from \emptyset to \emptyset in $m + n$ steps can be viewed as a walk from \emptyset to some shape λ in n steps, then followed by the reverse of a walk from \emptyset to λ in m steps. It follows that

$$(2.3) \quad \sum_{\lambda} g_{\lambda}(n) g_{\lambda}(m) = g_{\emptyset}(m + n) = B(m + n).$$

For the case $m = n = k$, the identity (2.3) is proved by Halverson and Lowandowski [16], who gave a bijective proof using similar procedures as those in ψ .

(3) The *partition algebra* \mathfrak{P}_n is a certain semisimple algebra, say over \mathbb{C} , whose dimension is the Bell number $B(n)$ (the number of partitions of $[n]$). (The algebra \mathfrak{P}_n depends on a parameter x which is irrelevant here.) See [15, 16] for a survey

of this topic. Vacillating tableaux are related to irreducible representations of \mathfrak{P}_n in the same way that SYT of content $[n]$ are related to irreducible representations of the symmetric group \mathfrak{S}_n . In particular, the irreducible representations I_λ of \mathfrak{P}_n are indexed by partitions λ for which there exists a vacillating tableau of shape λ and length $2n$, and $\dim I_n$ is the number of such vacillating tableaux. This result is equivalent to [15, Thm. 2.24(b)], but that paper does not explicitly define the notion of vacillating tableau. Combinatorial identities arising from partition algebra and its subalgebras are discussed in [16], where the authors used the notion of vacillating tableau after the distribution of a preliminary version of this paper.

3. CROSSINGS AND NESTINGS OF PARTITIONS

In this section we restrict the map ψ to vacillating tableaux of empty shape, for which ψ provides a bijection between the set of vacillating tableaux of empty shape and length $2n$ and the set of partitions of $[n]$. To make the bijection clear, we restate the inverse map from the set of partitions to vacillating tableaux.

The map ϕ from partitions to vacillating tableaux. Given a partition $P \in \Pi_n$ with the standard representation, we construct the sequence of SYT's, hence the vacillating tableau $\phi(P)$ as follows: Start from the empty SYT by letting $T_{2n} = \emptyset$, read the number $j \in [n]$ one by one from n to 1, and define T_{2j-1}, T_{2j-2} for each j . There are four cases.

1. If j is the right-hand endpoint of an arc (i, j) , but not a left-hand endpoint, first do nothing, then insert i (by the RSK algorithm) into the tableau.
2. If j is the left-hand endpoint of an arc (j, k) , but not a right-hand endpoint, first remove j , then do nothing.
3. If j is an isolated point, do nothing twice.
4. If j is the right-hand endpoint of an arc (i, j) , and the left-hand endpoint of another arc (j, k) , then delete j first, and then insert i .

The vacillating tableau $\phi(P)$ is the sequence of shapes of the above SYT's.

Example 3.1. Let P be the partition 1457-26-3 of [7].




FIGURE 1. The standard representation of the partition 1457-26-3.

Starting from \emptyset on the right, go from 7 to 1. The seven steps are: (1) do nothing, then insert 5, (2) do nothing, then insert 2, (3) delete 5 and insert 4, (4) delete 4 and insert 1, (5) do nothing twice, (6) remove 2 then do nothing, (7) remove 1 then do nothing. Hence the corresponding SYT's, constructed from right to left, are

$$\emptyset \quad \emptyset \quad 1 \quad 1 \quad \frac{1}{2} \quad \frac{1}{2} \quad \frac{1}{2} \quad 2 \quad 24 \quad 2 \quad \frac{2}{5} \quad 5 \quad 5 \quad \emptyset \quad \emptyset$$

The vacillating tableau is

$$\emptyset, \emptyset, 1, 1, 11, 11, 11, 1, 2, 1, 11, 1, 1, \emptyset, \emptyset.$$

The relation between $\text{cr}(P), \text{ne}(P)$ and the vacillating tableau is given in the next theorem.

Theorem 3.2. *Let $P \in \Pi_n$ and $\phi(P) = (\emptyset = \lambda^0, \lambda^1, \dots, \lambda^{2n} = \emptyset)$. Then $\text{cr}(P)$ is the most number of rows in any λ^i , and $\text{ne}(P)$ is the most number of columns in any λ^i .*

Proof. We prove Theorem 3.2 in four steps. First, we interpret a k -crossing/ k -nesting of P in terms of entries of SYT's T_i in $\phi(P)$. Then, we associate to each SYT T_i a sequence σ_i whose terms are entries of T_i . We prove that T_i is the insertion tableau of σ_i under the RSK algorithm, and apply Schensted's theorem to conclude the proof.

Step 1. Let $T(P) = (T_0, T_1, \dots, T_{2n})$ be the sequence of SYT's associated to the vacillating tableau $\phi(P)$. By the construction of ψ and ϕ , a pair (i, j) is an arc in the standard representation of P if and only if i is an entry in the SYT's $T_{2i}, T_{2i+1}, \dots, T_{2j-2}$. We say that the integer i is *added* to $T(P)$ at step i and *leaves* at step j .

First we prove that the arcs $(i_1, j_1), \dots, (i_k, j_k)$ form a k -crossing of P if and only if there exists a tableau T_i in $T(P)$ such that the integers $i_1, i_2, \dots, i_k \in \text{content}(T_i)$, and i_1, i_2, \dots, i_k leave $T(P)$ in increasing order according to their numerical values. Given a k -crossing $((i_1, j_1), \dots, (i_k, j_k))$ of P , where $i_r < j_r$ for $1 \leq r \leq k$ and $i_1 < i_2 < \dots < i_k < j_1 < j_2 < \dots < j_k$, the integer i_r is added to $T(P)$ at step i_r and leaves at step j_r . Hence all i_r are in T_{2j_1-2} , and they leave $T(P)$ in increasing order. The converse is also true: if there are k integers $i_1 < i_2 < \dots < i_k$, all appearing in the same tableau at some step, and then leave in increasing order, say at steps $j_1 < j_2 < \dots < j_k$, then $i_k < j_1$ and the pairs $((i_1, j_1), \dots, (i_k, j_k)) \in P$ form a k -crossing. By a similar argument arcs $(i_1, j_1), \dots, (i_k, j_k)$ form a k -nesting of P if and only if there exists a tableaux T_i in $T(P)$ such that the integers $i_1, i_2, \dots, i_k \in \text{content}(T_i)$, and i_1, i_2, \dots, i_k leave $T(P)$ in decreasing order.

Step 2. For each $T_i \in T(P)$, we define a permutation σ_i of $\text{content}(T_i)$ (backward) recursively as follows. Let σ_{2n} be the empty sequence. (1) If $T_i = T_{i-1}$, then $\sigma_{i-1} = \sigma_i$. (2) If T_{i-1} is obtained from T_i by row-inserting some number j , then $\sigma_{i-1} = \sigma_i j$, the juxtaposition of σ_i and j . (3) If T_i is obtained from T_{i-1} by adding the entry $i/2$, (where i must be even), then σ_{i-1} is obtained from σ_i by deleting the number $i/2$. Note that in the last case, $i/2$ must be the largest entry in σ_i .

Clearly σ_i is a permutation of the entries in $\text{content}(T_i)$. If $\sigma_i = w_1 w_2 \dots w_r$, then the entries of $\text{content}(T_i)$ leave $T(P)$ in the order w_r, \dots, w_2, w_1 .

Step 3. Claim: If $\sigma_i \xrightarrow{\text{RSK}} (A_i, B_i)$, then $A_i = T_i$.

We prove the claim by backward induction. The case $i = 2n$ is trivial, as both A_{2n} and T_{2n} are the empty SYT. Assume the claim is true for some i , $1 \leq i \leq 2n$. We prove that the claim holds for $i - 1$.

If $T_{i-1} = T_i$, the claim holds by the inductive hypothesis. If T_{i-1} is obtained from T_i by inserting the number j , then the claim holds by the definition of the RSK algorithm. It is only left to consider the case that T_{i-1} is obtained from T_i by removing the entry $j = i/2$.

Let us write σ_i as $u_1 u_2 \dots u_s j v_1 \dots v_t$, and σ_{i-1} as $u_1 u_2 \dots u_s v_1 \dots v_t$, where $j > u_1, \dots, u_s, v_1, \dots, v_t$. We need to show that the insertion tableau of σ_{i-1} is the same as the insertion tableau of σ_i deleting the entry j , i.e., $A_{i-1} = A_i \setminus \{j\}$. Proof by induction on t . If $t = 0$, then it is true by the RSK algorithm that A_i is obtained from A_{i-1} by adding j at the end of the first row. Assume it is true for $t - 1$, i.e., $A(u_1 \dots u_s v_1 \dots v_{t-1}) = A(u_1 \dots u_s j v_1 \dots v_{t-1}) \setminus \{j\}$. Note that in $A(u_1 \dots u_s j v_1 \dots v_{t-1})$, if j is in position (x, y) , then there is no element in

positions $(x, y + 1)$ or $(x + 1, y)$. Now we insert the entry v_t by the RSK algorithm. Consider the insertion path $I = I(A(u_1 \cdots u_s j v_1 \cdots v_{t-1}) \leftarrow v_t)$. If j does not appear on this path, then we would have the exact same insertion path when inserting v_t into $A(u_1 \cdots u_s v_1 \cdots v_{t-1})$. This insertion path results in the same change to $A(u_1 \cdots u_s j v_1 \cdots v_{t-1})$ and $A(u_1 \cdots u_s v_1 \cdots v_{t-1})$, which does not touch the position (x, y) of j . So $A(u_1 \cdots u_s v_1 \cdots v_{t-1} v_t) = A(u_1 \cdots u_s j v_1 \cdots v_{t-1} v_t) \setminus \{j\}$. On the other hand, if j appears in the insertion path I , i.e., $(x, y) \in I$, then since j is the largest element, it must be bumped into the $(x + 1)$ -th row, and become the last entry in the $(x + 1)$ -th row without bumping any number further. Then the insertion path of v_t into $A(u_1 \cdots u_s v_1 \cdots v_{t-1})$ is I minus the last position $\{(x + 1, *)\}$, and again we have $A(u_1 \cdots u_s v_1 \cdots v_{t-1} v_t) = A(u_1 \cdots u_s j v_1 \cdots v_{t-1} v_t) \setminus \{j\}$. This finishes the proof of the claim.

Step 4. We shall need the following theorem of Schensted [22], [25, Thms. 7.23.13, 7.23.17], which gives the basic connection between the RSK algorithm and the increasing and decreasing subsequences.

Schensted’s Theorem. *Let σ be a sequence of integers whose terms are distinct. Assume $\sigma \xrightarrow{RSK} (A, B)$, where A and B are SYT’s of the shape λ . Then the length of the longest increasing subsequence of σ is λ_1 (the number of columns of λ), and the length of the longest decreasing subsequence is λ'_1 (the number of rows of λ).*

Now we are ready to prove Theorem 3.2. By Steps 1 and 2, a partition P has a k -crossing if and only if there exists i such that σ_i has a decreasing subsequence of length k . The Claim in Step 3 implies that the shape of the sequence σ_i is exactly the diagram of the i -th partition λ^i in the vacillating tableau $\phi(P)$. By Schensted’s Theorem, σ_i has a decreasing subsequence of length k if and only if the partition λ^i in $\phi(P)$ has at least k rows. This proves the statement for $\text{cr}(P)$ in Theorem 3.2. The statement for $\text{ne}(P)$ is proved similarly. \square

The symmetric joint distribution of statistics $\text{cr}(P)$ and $\text{ne}(P)$ over $P_n(S, T)$ follows immediately from Theorem 3.2.

Proof of Theorem 1. From Theorem 3.2, a partition $P \in \Pi_n$ has $\text{cr}(P) = k$ and $\text{ne}(P) = j$ if and only if for the partitions $\{\lambda^i\}_{i=0}^{2^n}$ of the vacillating tableau $\phi(P)$, the maximal number of rows of the diagram of any λ^i is k , and the maximal number of columns of the diagram of any λ^i is j . Let τ be the involution defined on the set $\mathcal{V}_\emptyset^{2^n}$ by taking the conjugate to each partition λ^i . For $i \in [n]$, $i \in \min(P)$ (resp. $\max(P)$) if and only if $\lambda^{2^{i-1}} = \lambda^{2^i-2}$ and $\lambda^{2^i} \setminus \lambda^{2^i-1} = \square$ (resp. $\lambda^{2^{i-2}} \setminus \lambda^{2^i-1} = \square$ and $\lambda^{2^i} = \lambda^{2^i-1}$). Since τ preserves $\min(P)$ and $\max(P)$, it induces an involution on $P_n(S, T)$ which exchanges the statistics $\text{cr}(P)$ and $\text{ne}(P)$. This proves Theorem 1.1. \square

Let $\lambda = (\lambda_1, \lambda_2, \dots)$ be the shape of a sequence w of distinct integers. Schensted’s Theorem provides a combinatorial interpretation of the terms λ_1 and λ'_1 : they are the length of the longest increasing and decreasing subsequences of w . In [14] C. Greene extended Schensted’s Theorem by giving an interpretation of the rest of the diagram of $\lambda = (\lambda_1, \lambda_2, \dots)$.

Assume w is a sequence of length n . For each $k \leq n$, let $d_k(w)$ denote the length of the longest subsequence of w which has no increasing subsequences of length

$k + 1$. It can be shown easily that any such sequence is obtained by taking the union of k decreasing subsequences. Similarly, define $a_k(w)$ to be the length of the longest subsequence consisting of k ascending subsequences.

Theorem 3.3 (Greene). *For each $k \leq n$,*

$$\begin{aligned} a_k(w) &= \lambda_1 + \lambda_2 + \cdots + \lambda_k, \\ d_k(w) &= \lambda'_1 + \lambda'_2 + \cdots + \lambda'_k, \end{aligned}$$

where $\lambda' = (\lambda'_1, \lambda'_2, \dots)$ is the conjugate of λ .

We may consider the analogue of Greene's Theorem for set partitions. Let $P \in \Pi_n$ with the standard representation $\{(i_1, j_1), (i_2, j_2), \dots, (i_t, j_t)\}$, where $i_r < j_r$ for $1 \leq r \leq t$. Let $e_r = (i_r, j_r)$. We define the *crossing graph* $\text{Cr}(P)$ of P as follows. The vertex set of $\text{Cr}(P)$ is $\{e_1, e_2, \dots, e_t\}$. Two arcs e_r and e_s are adjacent if and only if the edges e_r and e_s are crossing, that is, $i_r < i_s < j_r < j_s$. Clearly a k -crossing of P corresponds to a k -clique of $\text{Cr}(P)$. Let $\text{cr}_r(P)$ be the maximal number of vertices in a union of r cliques of $\text{Cr}(P)$. In other words, $\text{cr}_r(P)$ is the maximal number of arcs in a union of r crossings of P . Similarly, let $\text{Ne}(P)$ be the graph defined on the vertex set $\{e_1, \dots, e_t\}$, where two arcs e_r and e_s are adjacent if and only if $i_r < i_s < j_s < j_r$. Let $\text{ne}_r(P)$ be the maximal number of vertices in a union of r cliques of $\text{Ne}(P)$. In other words, $\text{ne}_r(P)$ is the maximal number of arcs in a union of r nestings of P .

Proposition 3.4. *Let $P = ((i_1, j_1), (i_2, j_2), \dots, (i_t, j_t))$ be the standard representation of a partition of $[n]$, where $i_r < j_r$ for all $1 \leq r \leq t$ and $j_1 < j_2 < \dots < j_t$. Let $\alpha(P)$ be the sequence $i_1 i_2 \cdots i_t$. Then there is a one-to-one correspondence between the set of nestings of P and the set of decreasing subsequences of $\alpha(P)$.*

Proof. Let $\phi(P)$ be the vacillating tableau corresponding to P , and $T(P)$ the sequence of SYT's constructed in the bijection. Then $\alpha(P)$ records the order in which the entries of T_i 's leave $T(P)$. Let $\sigma = i_t \cdots i_2 i_1$ be the reverse of $\alpha(P)$, and $\{\sigma_i : 1 \leq i \leq 2n\}$ the permutation of $\text{content}(T_i)$ defined in Step 2 of the proof of Theorem 3.2. Then the σ_i 's are subsequences of σ .

From Steps 1 and 2 of the proof of Theorem 3.2, nestings of P are represented by the increasing subsequences of σ_i , $1 \leq i \leq 2n$, and hence by the increasing subsequences of σ . Conversely, let $i_{r_1} < i_{r_2} < \dots < i_{r_t}$ be an increasing subsequence of σ . Being a subsequence of σ means that its terms leave $T(P)$ in reverse order, so i_{r_t} leaves first in step j_{r_t} . Thus all the entries i_{r_1}, \dots, i_{r_t} appear in the SYT's T_i with $2i_{r_t} \leq i \leq 2j_{r_t} - 2$. Therefore $i_{r_1} i_{r_2} \cdots i_{r_t}$ is also an increasing subsequence of σ_i , for $2i_{r_t} \leq i \leq 2j_{r_t} - 2$. □

Combining Proposition 3.4 and Greene's Theorem, we have the following corollary describing $\text{ne}_t(P)$.

Corollary 3.5. *Let P and $\alpha(P)$ be as in Proposition 3.4. Then*

$$\text{ne}_r(P) = \lambda'_1 + \lambda'_2 + \cdots + \lambda'_r,$$

where λ is the shape of $\alpha(P)$, and λ' is the conjugate of λ .

The situation for $\text{cr}_r(P)$ is more complicated. We do not have a result similar to Proposition 3.4. Any crossing of P uniquely corresponds to an increasing subsequence of $\alpha(P)$. But the converse is not true. An increasing subsequence

$i_{r_1}i_{r_2}\cdots i_{r_t}$ corresponds to a t -crossing of P only if we have the additional condition $i_{r_t} < j_{r_1}$. It would be interesting to get a result for $\text{cr}_r(P)$ analogous to Corollary 3.5.

To conclude this section we discuss the enumeration of noncrossing partitions. The following theorem is a direct corollary of the bijection between vacillating tableaux and partitions.

Theorem 3.6. *Let ϵ_i denote the i -th unit coordinate vector in \mathbb{R}^{k-1} . The number of k -noncrossing partitions of $[n]$ equals the number of closed lattice walks in the region*

$$V_k = \{(a_1, a_2, \dots, a_{k-1}) : a_1 \geq a_2 \geq \dots \geq a_{k-1} \geq 0, a_i \in \mathbb{Z}\}$$

from the origin to itself of length $2n$ with steps $\pm\epsilon_i$ or $(0, 0, \dots, 0)$, with the property that the walk goes backwards (i.e., with step $-\epsilon_i$) or stands still (i.e., with step $(0, 0, \dots, 0)$) after an even number of steps, and goes forwards (i.e., with step $+\epsilon_i$) or stands still after an odd number of steps.

Recall that a partition $P \in \Pi_n$ is k -noncrossing if $\text{cr}(P) < k$, and is k -nonnesting if $\text{ne}(P) < k$. A partition P has no k -crossings and no j -nestings if and only if for all the partitions λ^i of $\phi(P)$, the diagram fits into a $(k-1) \times (j-1)$ rectangle. Taking the conjugate of each partition, we get bijective proofs of Corollaries 1.2 and 1.3.

Theorem 1.1 asserts the symmetric distribution of $\text{cr}(P)$ and $\text{ne}(P)$ over $P_n(S, T)$, for all $S, T \subseteq [n]$ with $|S| = |T|$. Not every $P_n(S, T)$ is nonempty. A set $P_n(S, T)$ is nonempty if and only if for all $i \in [n]$, $|S \cap [i]| \geq |T \cap [i]|$. Another way to describe the nonempty $P_n(S, T)$ is to use lattice paths. Associate to each pair (S, T) a lattice path $L(S, T)$ with steps $(1, 1)$, $(1, -1)$ and $(1, 0)$: start from $(0, 0)$, read the integers i from 1 to n one by one, and move two steps for each i .

1. If $i \in S \cap T$, move $(1, 0)$ twice.
2. If $i \in S \setminus T$, move $(1, 0)$ then $(1, 1)$.
3. If $i \in T \setminus S$, move $(1, -1)$ then $(1, 0)$.
4. If $i \notin S \cup T$, move $(1, -1)$ then $(1, 1)$.

This defines a lattice path $L(S, T)$ from $(0, 0)$ to $(2n, 0)$. Conversely, the path uniquely determines (S, T) . Then $P_n(S, T)$ is nonempty if and only if the lattice path $L(S, T)$ is a Motzkin path, i.e., never goes below the x -axis.

There are existing notions of noncrossing partitions and nonnesting partitions, e.g., [25, Ex. 6.19]. A *noncrossing partition of $[n]$* is a partition of $[n]$ in which no two blocks “cross” each other, i.e., if $a < b < c < d$ and a, c belong to a block B and b, d to another block B' , then $B = B'$. A *nonnesting partition of $[n]$* is a partition of $[n]$ such that if a, e appear in a block B and b, d appear in a different block B' where $a < b < d < e$, then there is a $c \in B$ satisfying $b < c < d$.


It is easy to see that P is a noncrossing partition if and only if the standard representation of P has no 2-crossing, and P is a nonnesting partition if and only if the standard representation of P has no 2-nesting. Hence the vacillating tableau correspondence, in the case of 1-row/column tableaux, gives bijections between noncrossing partitions of $[n]$, nonnesting partitions of $[n]$ (both are counted by Catalan numbers) and sequences $0 = a_0, a_1, \dots, a_{2n} = 0$ of nonnegative integers such that $a_{2i+1} = a_{2i}$ or $a_{2i} - 1$, and $a_{2i} = a_{2i-1}$ or $a_{2i-1} + 1$. These sequences a_0, \dots, a_{2n} give a new combinatorial interpretation of Catalan numbers.

Replacing a term $a_{i+1} = a_i + 1$ with a step $(1, 1)$, a term $a_{i+1} = a_i - 1$ with a step $(1, -1)$, and a term $a_{i+1} = a_i$ with a step $(1, 0)$, we get a Motzkin path,

so we also have a bijection between noncrossing/nonnesting partitions and certain Motzkin paths. The Motzkin paths are exactly the ones defined as $L(S, T)$, where $S = \min(P)$ and $T = \max(P)$. Conversely, given a Motzkin path of the form $L(S, T)$, we can recover uniquely a noncrossing partition and a nonnesting partition. Write the path as $\{(i, a_i) : 0 \leq i \leq 2n\}$. Let $A = [n] \setminus T$ and $B = [n] \setminus S$. Clearly $|A| = |B|$. Assume $A = \{i_1, i_2, \dots, i_t\}_<$ and $B = \{j_1, j_2, \dots, j_t\}_<$, where elements are listed in increasing order. Then to get the standard representation of the noncrossing partition, pair each j_r with $\max\{i_s \in A : i_s < j_r, a_{2i_s} = a_{2j_r-2}\}$. To get the standard representation of the nonnesting partition, pair each j_r with i_r , for $1 \leq r \leq t$.

Remark 3.7. In our definition, a k -crossing is defined as a set of k mutually crossing arcs in the standard representation of the partition. There exist some other definitions. For example, in [17] M. Klazar defined the *3-noncrossing partition* as a partition P which does not have 3 mutually crossing blocks. It can be seen that P is 3-noncrossing in Klazar’s sense if and only if there do not exist 6 elements $a_1 < b_1 < c_1 < a_2 < b_2 < c_2$ in $[n]$ such that $a_1, a_2 \in A, b_1, b_2 \in B, c_1, c_2 \in C$, and A, B, C are three distinct blocks of P .

Klazar’s definition of 3-noncrossing partitions is different from ours. For example, let P be the partition 15-246-37 of [7], with standard representation as follows:



According to Klazar’s definition, P has a 3-crossing, since we have $1 < 2 < 3 < 5 < 6 < 7$ and $\{1, 5\}, \{2, 6\}$ and $\{3, 7\}$ belong to three different blocks, respectively. On the other hand, P has no 3-crossing on our sense.

For general k , these three notions of k -noncrossing partitions, i.e., (1) no k -crossing in the standard representation of P , (2) no k mutually crossing arcs in distinct blocks of P , and (3) no k mutually crossing blocks, are all different, with the first being the weakest, and the third the strongest.

4. A VARIANT: PARTITIONS AND HESITATING TABLEAUX

We may also consider the *enhanced* crossing/nesting of a partition by taking isolated points into consideration. For a partition P of $[n]$, let the *enhanced representation* of P be the union of the standard representation of P and the loops $\{(i, i) : i \text{ is an isolated point of } P\}$. An *enhanced k -crossing* of P is a set of k edges $(i_1, j_1), (i_2, j_2), \dots, (i_k, j_k)$ of the enhanced representation of P such that $i_1 < i_2 < \dots < i_k \leq j_1 < j_2 < \dots < j_k$. In particular, two arcs of the form (i, j) and (j, l) with $i < j < l$ are viewed as crossing. Similarly, an *enhanced k -nesting* of P is a set of k edges $(i_1, j_1), (i_2, j_2), \dots, (i_k, j_k)$ of the enhanced representation of P such that $i_1 < i_2 < \dots < i_k \leq j_k < \dots < j_2 < j_1$. In particular, an edge (i, k) and an isolated point j with $i < j < k$ form an enhanced 2-nesting.

Let $\overline{\text{cr}}(P)$ be the size of the largest enhanced crossing, and $\overline{\text{ne}}(P)$ the size of the largest enhanced nesting. Using a variant form of vacillating tableau, we again obtain a symmetric joint distribution of the statistics $\overline{\text{cr}}(P)$ and $\overline{\text{ne}}(P)$.

The variant tableau is a *hesitating tableau* of shape \emptyset and length $2n$, which is a path on the Hasse diagram of Young’s lattice from \emptyset to \emptyset , where each step consists of a pair of moves, where the pair is either (i) doing nothing then adding a square, (ii)

removing a square then doing nothing, or (iii) adding a square and then removing a square.

Example 4.1. There are 5 hesitating tableaux of shape \emptyset and length 6. They are

$$(4.1) \quad \begin{array}{cccccc} \emptyset & 1 & \emptyset & 1 & \emptyset & 1 & \emptyset \\ \emptyset & 1 & \emptyset & \emptyset & 1 & \emptyset & \emptyset \\ \emptyset & \emptyset & 1 & 11 & 1 & \emptyset & \emptyset \\ \emptyset & \emptyset & 1 & \emptyset & \emptyset & 1 & \emptyset \\ \emptyset & \emptyset & 1 & 2 & 1 & \emptyset & \emptyset \end{array}$$

To see the equivalence with vacillating tableaux, let U be the operator that takes a shape to the sum of all shapes that cover it in Young’s lattice (i.e., by adding a square), and similarly D takes a shape to the sum of all shapes that it covers in Young’s lattice (i.e., by deleting a square). Then, as is well known, $DU - UD = I$ (the identity operator). See, e.g., [23], [25, Exer. 7.24]. It follows that

$$(4.2) \quad (U + I)(D + I) = DU + ID + UI.$$

Iterating the left-hand side generates vacillating tableaux, and iterating the right-hand side gives the hesitating tableaux defined above.

A bijective map between partitions of $[n]$ and hesitating tableaux of empty shape has been given by Korn [18], based on growth diagrams. Here by modifying the map ϕ defined in Section 3 we get a more direct bijection $\bar{\phi}$ between partitions and hesitating tableaux of empty shape, which leads to the symmetric joint distribution of $\overline{\text{cr}}(P)$ and $\overline{\text{ncr}}(P)$. The construction and proofs are very similar to the ones given in Sections 2 and 3, and hence are omitted here. We will only state the definition of the map $\bar{\phi}$ from partitions to hesitating tableaux, to be compared with the map ϕ in Section 3.

The bijection $\bar{\phi}$ from partitions to hesitating tableaux. Given a partition $P \in \Pi_n$ with the enhanced representation, we construct the sequence of SYT’s, and hence the hesitating tableau $\bar{\phi}(P)$, as follows: start from the empty SYT by letting $T_{2n} = \emptyset$, read the numbers $j \in [n]$ one by one from n to 1, and define two SYT’s T_{2j-1}, T_{2j-2} for each j . When j is a left-hand endpoint only, or a right-hand endpoint only, the construction is identical to that of the map ϕ . Otherwise,

1. If j is an isolated point, first insert j , then delete j .
2. If j is the right-hand endpoint of an arc (i, j) , and the left-hand endpoint of another arc (j, k) , then insert i first, and then delete j .

Example 4.2. For the partition 1457-26-3 of [7] in Figure 2, the corresponding SYT’s are

$$\emptyset \quad \emptyset \quad 1 \quad 1 \quad \frac{1}{2} \quad \frac{13}{2} \quad \frac{1}{2} \quad \frac{14}{2} \quad 24 \quad \frac{24}{5} \quad \frac{2}{5} \quad 5 \quad 5 \quad \emptyset \quad \emptyset$$

The hesitating tableau $\bar{\phi}(P)$ is

$$\emptyset, \emptyset, 1, 1, 11, 21, 11, 21, 2, 21, 11, 1, 1, \emptyset, \emptyset.$$

The conjugation of shapes does not preserve $\min(P)$ or $\max(P)$. Instead, it preserves $\min(P) \setminus \max(P)$ and $\max(P) \setminus \min(P)$. Let S, T be disjoint subsets of $[n]$ with the same cardinality, $\bar{P}_n(S, T) = \{P \in \Pi_n : \min(P) \setminus \max(P) = S, \max(P) \setminus \min(P) = T\}$, and $f_{n,S,T}(i, j) = \#\{P \in \bar{P}_n(S, T) : \overline{\text{cr}}(P) = i, \overline{\text{ncr}}(P) = j\}$.




FIGURE 2. The enhanced representation of the partition 1457-26-3.

Theorem 4.3. *We have*

$$\bar{f}_{n,S,T}(i,j) = \bar{f}_{n,S,T}(j,i).$$

As a consequence, Corollaries 1.2 and 1.3 remain valid if we define k -noncrossing (or k -nonnesting) by $\bar{c}r(P) < k$ (or $\bar{n}e(P) < k$).

Remark 4.4. As done for vacillating tableaux, one can extend the definition of hesitating tableaux by considering moves from \emptyset to λ , and denote by $f_\lambda(n)$ the number of such hesitating tableaux of length $2n$. Identity (4.2) implies that $f_\lambda(n) = g_\lambda(n)$, the number of vacillating tableaux from \emptyset to λ . It follows that $f_\emptyset(n) = B(n)$ and

$$\sum_{\lambda} f_\lambda(n)f_\lambda(m) = B(m+n),$$

where $B(n)$ is the n -th Bell number. For further discussion of the number $f_\lambda(n)$, see [26, Problem 33] (version of 17 August 2004).

5. ENUMERATION OF k -NONCROSSING MATCHINGS

Restricting Theorem 1.1 to disjoint subsets (S, T) of $[n]$, where $n = 2m$ and $|S| = |T| = m$, we get the symmetric joint distribution of the crossing number and nesting number for matchings, as stated in Corollary 1.4 in Section 1.

In a complete matching, an integer is either a left endpoint or a right endpoint in the standard representation. In applying the map ϕ to complete matchings on $[2m]$, if we remove all steps which do nothing, we obtain a sequence $\emptyset = \lambda^0, \lambda^1, \dots, \lambda^{2m} = \emptyset$ of partitions such that for all $1 \leq i \leq 2m$, the diagram of λ^i is obtained from that of λ^{i-1} by either adding one square or removing one square. Such a sequence is called an *oscillating tableau* (or *up-down tableau*) of empty shape and length $2m$. Thus we get a bijection between complete matchings on $[2m]$ and oscillating tableaux of empty shape and length $2m$. This bijection was originally constructed by the fourth author, and then extended by Sundaram [27] to arbitrary shapes to give a combinatorial proof of the Cauchy identity for the symplectic group $\text{Sp}(2m)$. The explicit description of the bijection has appeared in [27] and was included in [25, Exercise 7.24]. Oscillating tableaux first appeared (though not with that name) in [5].


Recall that the ordinary RSK algorithm gives a bijection between the symmetric group \mathfrak{S}_m and pairs (P, Q) of SYTs of the same shape $\lambda \vdash m$. This result and Schensted's Theorem can be viewed as a special case of what we do. Explicitly, identify an SYT T of shape λ and content $[m]$ with a sequence $\emptyset = \lambda^0, \lambda^1, \dots, \lambda^m = \lambda$ of integer partitions, where λ^i is the shape of the SYT T^i obtained from T by deleting all entries $\{j : j > i\}$. Let w be a permutation of $[m]$, and form the matching M_w on $[2m]$ with arcs between $w(i)$ and $2m - i + 1$. We get an oscillating tableau O_w that increases to the shape $\lambda \vdash m$ and then decreases to the empty shape. Assume $w \xrightarrow{\text{RSK}} (A(w), B(w))$, where $A(w)$ is the (row)-insertion tableau

and $B(w)$ the recording tableau. Then $A(w)$ is given by the first m steps of O_w , and $B(w)$ the reverse of the last m steps. The size of the largest crossing (resp. nesting) of M_w is exactly the length of the longest decreasing (resp. increasing) subsequence of w .

Example 5.1. Let $w = 231$. Then

$$A(w) = \begin{matrix} 1 & 3 \\ 2 & \end{matrix}, \quad B(w) = \begin{matrix} 1 & 2 \\ 3 & \end{matrix}.$$

The matching M_w and the corresponding oscillating tableau are given below:



Remark 5.2. The Brauer algebra \mathfrak{B}_m is a certain semisimple algebra, say over \mathbb{C} , whose dimension is the number $1 \cdot 3 \cdots (2m - 1)$ of matchings on $[2m]$. (The algebra \mathfrak{B}_m depends on a parameter x which is irrelevant here.) Oscillating tableaux of length $2m$ are related to irreducible representations of \mathfrak{B}_m in the same way that SYT of content $[m]$ are related to irreducible representations of the symmetric group \mathfrak{S}_m and that vacillating tableaux of length $2m$ are related to irreducible representations of the partition algebra \mathfrak{P}_m . In particular, the irreducible representations J_λ of \mathfrak{B}_m are indexed by partitions λ for which there exists an oscillating tableau of shape λ and length $2m$, and $\dim J_\lambda$ is the number of such oscillating tableaux. See, e.g., [4, Appendix B6] for further information.

Next we use the bijection between complete matchings and oscillating tableaux to study the enumeration of k -noncrossing matchings. All the results in the following hold for k -nonnesting matchings as well.

For complete matchings, Theorem 3.6 becomes the following.

Corollary 5.3. *The number of k -noncrossing matchings of $[2m]$ is equal to the number of closed lattice walks of length $2m$ in the set*

$$V_k = \{(a_1, a_2, \dots, a_{k-1}) : a_1 \geq a_2 \geq \dots \geq a_{k-1} \geq 0, a_i \in \mathbb{Z}\}$$

from the origin to itself with unit steps in any coordinate direction or its negative.

Restricted to the cases $k = 2, 3$, Corollary 5.3 leads to some nice combinatorial correspondences. Recall that a Dyck path of length $2m$ is a lattice path in the plane from the origin $(0, 0)$ to $(2m, 0)$ with steps $(1, 1)$ and $(1, -1)$ that never passes below the x -axis. A pair (P, Q) of Dyck paths is *noncrossing* if they have the same origin and the same destination, and P never goes below Q .

Corollary 5.4. (1) *The set of 2-noncrossing matchings is in one-to-one correspondence with the set of Dyck paths.*

(2) *The set of 3-noncrossing matchings is in one-to-one correspondence with the set of pairs of noncrossing Dyck paths.*

Proof. By Corollary 5.3, 2-noncrossing matchings are in one-to-one correspondence with closed lattice paths $\{\vec{v}_i = (x_i)\}_{i=0}^{2m}$ with $x_0 = x_{2m} = 0$, $x_i \geq 0$ and $x_{i+1} - x_i = \pm 1$. Given such a 1-dimensional lattice path, define a lattice path in the plane by letting $P = \{(i, x_i) \mid i = 0, 1, \dots, 2m\}$. Then P is a Dyck path, and this gives the desired correspondence.

For $k = 3$, 3-noncrossing matchings are in one-to-one correspondence with 2-dimensional lattice paths $\{\vec{v}_i = (x_i, y_i)\}_{i=0}^{2m}$ with $(x_0, y_0) = (x_{2m}, y_{2m}) = (0, 0)$, $x_i \geq y_i \geq 0$ and $(x_{i+1}, y_{i+1}) - (x_i, y_i) = (\pm 1, 0)$ or $(0, \pm 1)$. Given such a lattice path, define two lattice paths in the plane by setting $P = \{(i, x_i + y_i) \mid i = 0, 1, \dots, 2m\}$ and $Q = \{(i, x_i - y_i) \mid i = 0, 1, \dots, 2m\}$. Then (P, Q) is a pair of noncrossing Dyck paths. It is easy to see that this is a bijection. \square

Example 5.5. We illustrate the bijections between 3-noncrossing matchings, oscillating tableaux, and pairs of noncrossing Dyck paths. The oscillating tableau is

$$\emptyset, 1, 2, 21, 31, 21, 11, 21, 2, 1, \emptyset.$$

The sequence of SYT's as defined in the bijection is

$$\emptyset, 1, 12, \begin{matrix} 1 & 2 \\ 3 \end{matrix}, \begin{matrix} 1 & 2 & 4 \\ 3 \end{matrix}, \begin{matrix} 1 & 2 \\ 3 \end{matrix}, \begin{matrix} 1 \\ 3 \end{matrix}, \begin{matrix} 1 & 7 \\ 3 \end{matrix}, 37, 3, \emptyset.$$

The corresponding matching and the pair of noncrossing Dyck paths are given in Figure 3.




FIGURE 3. The matching and the pair of noncrossing Dyck paths.

Let $f_k(m)$ be the number of k -noncrossing matchings of $[2m]$. By Corollary 5.3 it is also the number of lattice paths of length $2m$ in the region V_k from the origin to itself with step set $\{\pm\epsilon_1, \pm\epsilon_2, \dots, \pm\epsilon_{k-1}\}$. Set

$$F_k(x) = \sum_m f_k(m) \frac{x^{2m}}{(2m)!}.$$

It turns out that a determining expression for $F_k(x)$ has been given by Grabiner and Magyar [12]. It is simply the case $\lambda = \eta = (m, m - 1, \dots, 1)$ of equation (38) in [12], giving

$$(5.1) \quad F_k(x) = \det [I_{i-j}(2x) - I_{i+j}(2x)]_{i,j=1}^{k-1},$$

where

$$I_m(2x) = \sum_{j \geq 0} \frac{x^{m+2j}}{j!(m+j)!},$$

the hyperbolic Bessel function of the first kind of order m [30]. One can easily check that when $k = 2$, the generating function of 2-noncrossing matchings equals

$$F_2(x) = I_0(2x) - I_2(2x) = \sum_{j \geq 0} C_j \frac{x^{2j}}{(2j)!},$$

where C_j is the j -th Catalan number. When $k = 3$, we have

$$\begin{aligned} f_3(m) &= \frac{3!(2m+2)!}{m!(m+1)!(m+2)!(m+3)!} \\ &= C_m C_{m+2} - C_{m+1}^2. \end{aligned}$$

This result agrees with the formula on the number of pairs of noncrossing Dyck paths due to Gouyou-Beauchamps in [11].

Remark 5.6. The determinant formula (5.1) has been studied by Baik and Rains in [2, Eqs. (2.25)]. One simply puts $i - 1$ for j and $j - 1$ for k in (2.25) of [2] to get our formula. The same formula was also obtained by Goulden [10] as the generating function for fixed-point-free permutations with no decreasing subsequence of length greater than $2k$. See Theorem 1.1 and 2.3 of [10] and specialize h_i to be $x^i/i!$, so g_l becomes the hyperbolic Bessel function. The asymptotic distribution of $\text{cr}(M)$ follows from another result of Baik and Rains. In Theorem 3.1 of [3] they obtained the limit distribution for the length of the longest decreasing subsequence of fixed-point-free involutions w . But representing w as a matching M , the condition that w has no decreasing subsequence of length $2k + 1$ is equivalent to the condition that M has no $k + 1$ -nesting, and we already know that $\text{cr}(M)$ and $\text{ne}(M)$ have the same distribution. Combining the above results, one has

$$\lim_{m \rightarrow \infty} \Pr \left(\frac{\text{cr}(M) - \sqrt{2m}}{(2m)^{1/6}} \leq \frac{x}{2} \right) = F_1(x),$$

where

$$F_1(x) = \sqrt{F(x)} \exp \left(\frac{1}{2} \int_x^\infty u(s) ds \right),$$

where $F(x)$ is the Tracy-Widom distribution and $u(x)$ the Painlevé II function.

Similarly one can try to enumerate complete matchings of $[2m]$ with no $(k + 1)$ -crossing and no $(j + 1)$ -nesting. By the oscillating tableau bijection this is just the number of walks of length $2m$ from $\hat{0}$ to $\hat{0}$ in the Hasse diagram of the poset $L(k, j)$, where $\hat{0}$ denotes the unique bottom element (the empty partition) of $L(k, j)$, the lattice of integer partitions whose shape fits in a $k \times j$ rectangle, ordered by inclusion. Let $g_{k,j}(m)$ be this number, and let $G_{k,j}(x) = \sum_m g_{k,j}(m)x^{2m}$ be the generating function.

For $j = 1$, the number $g_{k,1}(m)$ counts lattice paths from $(0, 0)$ to $(2m, 0)$ with steps $(1, 1)$ or $(1, -1)$ that stay between the lines $y = 0$ and $y = k$. The evaluation of $g_{k,1}(m)$ was first considered by Takács in [28] by a probabilistic argument. Explicit formula and generating function for this case are well known. For example, in [19] one obtains the explicit formula by applying the reflection principle repeatedly, viz.,

$$g_{k,1}(m) = \sum_i \left[\binom{2m}{m-i(k+2)} - \binom{2m}{m+i(k+2)+k+1} \right].$$

The generating function $G_{k,1}(x)$ is a special case of the one for the duration of the game in the classical ruin problem, that is, restricted random walks with absorbing barriers at 0 and a , and initial position z . See, for example, Equation (4.11) of Chapter 14 of [9]: let

$$U_z(x) = \sum_{m=0}^\infty u_{z,m} x^m,$$

where $u_{z,n}$ is the probability that the process ends with the n -th step at the barrier 0. Then

$$U_z(x) = \left(\frac{q}{p}\right)^z \frac{\lambda_1^{a-z}(x) - \lambda_2^{a-z}(x)}{\lambda_1^a(x) - \lambda_2^a(x)},$$

where

$$\lambda_1(x) = \frac{1 + \sqrt{1 - 4pqx^2}}{2px}, \quad \lambda_2(x) = \frac{1 - \sqrt{1 - 4pqx^2}}{2px}.$$

The generating function $G_{k,1}(x)$ is just $U_1(2x)/x$ with $a = k + 2$, $z = 1$, and $p = q = 1/2$.

In general, by the transfer matrix method [24, §4.7]

$$(5.2) \quad G_{k,j}(x) = \frac{\det(I - xA_{k,j}(0))}{\det(I - xA_{k,j})}$$

is a rational function, where $A_{k,j}$ is the adjacency matrix of the Hasse diagram of $L(k, j)$, and $A_{k,j}(0)$ is obtained from $A_{k,j}$ by deleting the row and the column corresponding to $\hat{0}$. $A_{k,j}(0)$ is also the adjacency matrix of the Hasse diagram of $L(k, j)$ with its bottom element (the empty partition) removed. Note that $\det(I - xA_{k,j})$ is a polynomial in x^2 since $L(k, j)$ is bipartite [6, Thm. 3.11]. Let $\det(I - xA_{k,j}) = p_{k,j}(x^2)$. The following is a table of $p_{k,j}(x)$ for the values of $1 \leq k \leq j \leq 4$. (We only need to list those with $k \leq j$ since $p_{k,j}(x) = p_{j,k}(x)$.)

(k, j)	$p_{k,j}(x) = \det(I - \sqrt{x}A_{k,j})$
(1, 1)	$1 - x$
(1, 2)	$1 - 2x$
(1, 3)	$1 - 3x + x^2$
(1, 4)	$(1 - x)(1 - 3x)$
(2, 2)	$(1 - x)(1 - 5x)$
(2, 3)	$(1 - x)(1 - 3x)(1 - 8x + 4x^2)$
(2, 4)	$(1 - 14x + 49x^2 - 49x^3)(1 - 6x + 5x^2 - x^3)$
(3, 3)	$(1 - x)(1 - 19x + 83x^2 - x^3)(1 - 5x + 6x^2 - x^3)^2$
(3, 4)	$(1 - 2x)^2(1 - 8x + 8x^2)(1 - 4x + 2x^2)^2(1 - 16x + 60x^2 - 32x^3 + 4x^4)$ $\cdot (1 - 24x + 136x^2 - 160x^3 + 16x^4)$
(4, 4)	$(1 - x)^2(1 - 18x + 81x^2 - 81x^3)^2(1 - 27x + 99x^2 - 9x^3)(1 - 9x + 18x^2 - 9x^3)^2$ $\cdot (1 - 27x + 195x^2 - 361x^3)(1 - 6x + 9x^2 - x^3)^2(1 - 9x + 6x^2 - x^3)^2$

The polynomial $p_{k,j}(x)$ seems to have a lot of factors. We are grateful to Christian Krattenthaler for explaining (5.4) below, from which we can explain the factorization of $p_{k,j}(x)$. By an observation [13, §5] of Grabiner, $g_{k,j}(n)$ is equal to the number of walks with n steps $\pm e_i$ from $(j, j - 1, \dots, 2, 1)$ to itself in the chamber $j + k + 1 > x_1 > x_2 > \dots > x_j > 0$ of the affine Weyl group \tilde{C}_n . Write $m = j + k + 1$. By [13, (23)] there follows

$$\sum_n g_{k,j}(n) \frac{x^{2n}}{(2n)!} = \det \left[\frac{1}{m} \sum_{r=0}^{2m-1} \sin(\pi r a / m) \sin(\pi r b / m) \cdot \exp(2x \cos(\pi r / m)) \right]_{a,b=1}^j.$$

When this determinant is expanded, we obtain a linear combination of terms of the form

$$(5.3) \quad \begin{aligned} &\exp(2x(\cos(\pi r_1 / m) + \dots + \cos(\pi r_j / m))) \\ &= \sum_{n \geq 0} 2^n (\cos(\pi r_1 / m) + \dots + \cos(\pi r_j / m))^n \frac{x^n}{n!}, \end{aligned}$$

where $0 \leq r_i \leq 2m - 1$ for $1 \leq i \leq j$. In fact, the case $\eta = \lambda$ of Grabiner's formula [13, (23)] shows that the number of walks of length n in the Weyl chamber from *any* integral point to itself is again a linear combination of terms

$$2^n (\cos(\pi r_1/m) + \cdots + \cos(\pi r_j/m))^n.$$

It follows that every eigenvalue of $A_{k,j}$ has the form

$$(5.4) \quad \theta = 2(\cos(\pi r_1/m) + \cdots + \cos(\pi r_j/m)).$$

(In particular, the Galois group over \mathbb{Q} of every irreducible factor of $p_{k,j}(x)$ is abelian.) Note that *a priori* not every such θ may be an eigenvalue, since it may appear with coefficient 0 after the linear combinations are taken. The algebraic integer $z = 2(\cos(\pi r_1/m) + \cdots + \cos(\pi r_j/m))$ lies in the field $\mathbb{Q}(\cos(\pi/m))$, an extension of \mathbb{Q} of degree $\phi(2m)/2$, where ϕ is the Euler phi-function. To see this, let z be a primitive $2m$ -th root of unity. Then z is a root of $x + 1/x = 2 \cos(\pi/m)$. Hence the field $L = \mathbb{Q}(z)$ is quadratic or linear over $K = \mathbb{Q}(\cos(\pi/m))$. Since K is real and L is not for $m > 1$, we cannot have $K = L$. Hence $[L : K] = 2$. Since $[L : \mathbb{Q}] = \phi(2m)$, we have $[K : \mathbb{Q}] = \phi(2m)/2$. It follows that the minimal polynomial over \mathbb{Q} of z has degree dividing $\phi(2m)/2$. Thus every irreducible factor of $\det(I - Ax)$ has degree dividing $\phi(2m)/2$, explaining why $p_{k,j}(x)$ has many factors. A more careful analysis should yield more precise information about the factors of $p_{k,j}(x)$, but we will not attempt such an analysis here.

An interesting special case of determining $p_{k,j}(x)$ is determining its degree, since the number of eigenvalues of $A_{k,j}$ equal to 0 is given by $\binom{j+k}{j} - 2 \cdot \deg p_{k,j}(x)$. Equivalently, since $A_{k,j}$ is a symmetric matrix, $2 \cdot \deg p_{k,j}(x) = \text{rank}(A_{k,j})$. We have observed the following:

- (1) For $k + j \leq 12$ and $1 \leq k \leq j$, $A_{k,j}$ is invertible exactly for $(k, j) = (1, 1), (1, 3), (1, 5), (1, 7), (1, 9), (1, 11), (3, 3), (3, 7), (3, 9), (5, 5)$ and $(5, 7)$.
- (2) $A_{1,j}$ is invertible if and only if j is odd. This is true because $L(1, j)$ is a path of length j , whose determinant satisfies the recurrence $\det(A_{1,j}) = -\det(A_{1,j-2})$. The statement follows from the initial conditions $\det(A_{1,1}) = -1$ and $\det(A_{1,2}) = 0$.
- (3) If $A_{k,j}$ is invertible, then kj is odd. To see this, let $X_0(X_1)$ be the set of integer partitions of even (odd) n whose shape fits in a $k \times j$ rectangle. Since $L(k, j)$ is bipartite graph with vertex partition (X_0, X_1) , a necessary condition for $A_{k,j}$ to be invertible is $|X_0| = |X_1|$. That is, the generating function $\sum_n p(k, j, n)q^n = \binom{\mathbf{k}+\mathbf{j}}{\mathbf{k}}$ must have a root at $q = -1$, where $p(k, j, n)$ is the number of integer partitions on n whose shape fits into a $k \times j$ rectangle. But the multiplicity of $1 + q$ in the Gaussian polynomial $\binom{\mathbf{k}+\mathbf{j}}{\mathbf{k}}$ is $\lfloor \frac{k+j}{2} \rfloor - \lfloor \frac{k}{2} \rfloor - \lfloor \frac{j}{2} \rfloor$, which is 0 unless both j and k are odd.

Item (3) is also proved independently by Jason Burns, who found a counterexample for the converse: for $k = 3$ and $j = 11$, $A_{3,11}$ is not invertible, in fact its corank is 6. The invertibility of $A_{k,j}$ for kj being odd is currently under investigation.

ACKNOWLEDGMENTS

The authors would like to thank Professor Donald Knuth for carefully reading the manuscript and providing many helpful comments.

REFERENCES

1. J. Baik, P. Deift and K. Johansson, *On the distribution of the length of the longest increasing subsequence of random permutations*. J. Amer. Math. Soc., **12** (1999), No. 4, 1119–1178. MR1682248 (2000e:05006)
2. J. Baik and E. Rains, *Algebraic aspects of increasing subsequences*, Duke Math. J., **109** (2001), No. 1, 1–65. MR1844203 (2002i:05119)
3. J. Baik and E. Rains, *The asymptotics of monotone subsequences of involutions*, Duke Math. J., **109** (2001), 205–282. MR1845180 (2003e:60016)
4. H. Barcelo and A. Ram, *Combinatorial representation theory*, in New Perspectives in Algebraic Combinatorics (Berkeley, CA, 1996–97), MSRI Publ. **38**, Cambridge University Press, Cambridge, 1999, pp. 23–90. MR1731814 (2000j:05125)
5. A. Berele, *A Schensted-type correspondence for the symplectic group*, J. Combinatorial Theory (A), **43** (1986), 320–328. MR0867655 (88b:20027)
6. D. M. Cvetković, M. Doob, and H. Sachs, *Spectra of Graphs*. 3rd ed., Johann Ambrosius Barth, Heidelberg, 1995. MR1324340 (96b:05108)
7. H. Davenport and A. Schinzel, *A combinatorial problem connected with differential equations*, Amer. J. Math., **87** (1965), 684–694. MR0190010 (32:7426)
8. M. de Sainte-Catherine, *Couplages et Pfaffiens en combinatoire, physique et informatique*, Ph.D. Thesis, University of Bordeaux I, 1983.
9. W. Feller, *An Introduction to Probability Theory and Its Applications*, 3rd edition, John Wiley and Sons, Inc., New York, 1968. MR0228020 (37:3604)
10. I. P. Goulden, *A linear operator for symmetric functions and tableaux in a strip with given trace*, Discrete Math., **99** (1992), 69–77. MR1158782 (93f:05100)
11. D. Gouyou-Beauchamps, *Standard Young tableaux of height 4 and 5*, Europ. J. Combin., **10** (1989), 69–82. MR0977181 (90b:05013)
12. D. Grabiner and P. Magyar, *Random walks in Weyl chambers and the decomposition of tensor powers*, J. Alg. Combinatorics, **2** (1993), 239–260. MR1235279 (94m:05202)
13. D. Grabiner, *Random walk in an alcove of an affine Weyl group, and non-colliding random walks on an interval*, J. Combin. Theory Ser. A, **97** (2002), 285–306. MR1883868 (2002k:60032)
14. C. Greene, *An extension of Schensted's theorem*, Adv. Math., **14** (1974), 254–265. MR0354395 (50:6874)
15. T. Halverson and A. Ram, *Partition algebras*, European J. of Combinatorics, **26** (2005), No. 6, 869–921. MR2143201 (2006g:05228)
16. T. Halverson and T. Lewandowski, *RSK insertion for set partitions and diagram algebras*, Electron. J. Combin. **11** (2004/06), no. 2, Research Paper 24, 24 pp. MR2195430
17. M. Klazar, *Bell numbers, their relatives, and algebraic differential equations*, J. Combin. Theory Ser. A, **102** (2003), 63–87. MR1970977 (2004d:11014)
18. M. Korn, Personal communication.
19. S. G. Mohanty, *Lattice Path Counting and Applications*, Academic Press, New York, 1979. MR0554084 (81f:60020)
20. R. C. Mullin and R. C. Stanton, *A map-theoretic approach to Davenport-Schinzel sequences*, Pacific J. Math., **40** (1972), 167–172. MR0302601 (46:1745)
21. J. Riordan, *The distribution of crossings chords joining pairs of $2n$ points on a circle*, Math. Computation, **29** (1975), 215–222. MR0366686 (51:2933)
22. C. E. Schensted, *Longest increasing and decreasing subsequences*, Canad. J. Math., **13** (1961), 179–191. MR0121305 (22:12047)
23. R. Stanley, *Differential posets*, J. Amer. Math. Soc., **1** (1988), 919–961. MR0941434 (89h:06005)
24. R. Stanley, *Enumerative Combinatorics*, vol. 1, Wadsworth and Brooks/Cole, Pacific Grove, CA, 1986; second printing, Cambridge University Press, Cambridge, 1996. MR1442260 (98a:05001)
25. R. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, Cambridge, 1999. MR1676282 (2000k:05026)
26. R. Stanley, *Supplementary Exercises for Chapter 7 of Enumerative Combinatorics*, available at <http://www-math.mit.edu/~rstan/ec>.

27. S. Sundaram, *The Cauchy identity for $Sp(2n)$* , J. Combin. Theory Ser. A, **53** (1990), 209–238. MR1041446 (91i:05122)
28. L. Takács, *Ballot problems*, Z. Wahrsch. Verw. Gebiete, **1** (1962), 154–158. MR0145601 (26:3131)
29. J. Touchard, *Sur un problème de configuration et sur les fractions continues*, Canad. J. Math., **14** (1952), 2–25. MR0046325 (13:716c)
30. E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, Cambridge University Press, Cambridge, 1927. MR1424469 (97k:01072)

CENTER FOR COMBINATORICS, LPMC, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

E-mail address: chen@nankai.edu.cn

CENTER FOR COMBINATORICS, LPMC, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

Current address: Department of Applied Mathematics, Dalian University of Technology, Dalian, Liaoning 116024, People's Republic of China

E-mail address: dengyp@eyou.com

CENTER FOR COMBINATORICS, LPMC, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

Current address: Department of Mathematics, East China Normal University, Shanghai 200062, People's Republic of China

E-mail address: rxdu@math.ecnu.edu.cn

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MASSACHUSETTS 02139

E-mail address: rstan@math.mit.edu

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843-3368

E-mail address: cyan@math.tamu.edu