

Quadrature Formulas for Semi-Infinite Integrals*

By Ravindra Kumar and M. K. Jain

Abstract. Polynomials orthogonal on $[0, \infty]$ with regard to the weight function $w(x) = x^\alpha(1+x)^{-\beta}$ are obtained, recurrence relations are found and the differential equation, which is satisfied by them, is given. Formulas for weights and abscissas in the corresponding quadrature formula are given.

1. Introduction. Harper [1] developed a quadrature formula on $[-\infty, \infty]$ with the algebraic weight function $w(x) = (1+x^2)^{-k-1}$. Even though the degree of the polynomial approximation to $f(x)$ is limited, he illustrated the superiority of his formula over that due to Gauss-Hermite, when $f(x)$ is a particular algebraic function.

In this paper we have worked out a similar formula on $[0, \infty]$ with the algebraic weight function $w(x) = x^\alpha(1+x)^{-\beta}$. While the Gauss-Laguerre formula is expected to work well in situations where the integrand function behaves asymptotically like $e^{-\gamma x}$, our formula would be more suitable where the integrand function has both the following features:

- (a) asymptotic behaviour like $x^{-\beta+n}$,
- (b) algebraic singularity like $(1+x)^{-\beta}$ at $x = -1$.

This fact has been illustrated with the help of an example in Section 3.

2. Derivation of Formulas. The existence of the formula

$$(1) \quad \int_0^\infty x^\alpha(1+x)^{-\beta} f(x) dx = \sum_{i=1}^n H_i f(x_i) + E_{n,\alpha,\beta}$$

requires that

$$(2) \quad \int_0^\infty x^\alpha(1+x)^{-\beta} dx \geq 0$$

and that the growth of the function $f(x)$, as x approaches infinity, be restricted by the condition

$$(3) \quad |f(x)| \leq M \cdot |x|^p$$

where M is a finite constant and p is a real number such that $p + \alpha < \beta$.

In (1) the abscissas x_i are the zeros of the n th degree polynomial $\phi_{n,\alpha,\beta}(x)$ which satisfies the orthogonality condition

Received February 22, 1973.

AMS (MOS) subject classifications (1970). Primary 65D30.

Key words and phrases. Orthogonal polynomials, weight function, recurrence relation, quadrature formula, weight coefficients, error.

* Part of this work was done at the University of Lancaster, England.

Copyright © 1974, American Mathematical Society

$$(4) \quad \int_0^\infty x^\alpha (1+x)^{-\beta} \phi_{m,\alpha,\beta}(x) \phi_{n,\alpha,\beta}(x) dx = 0, \quad m \neq n.$$

Following Hildebrand [2] it is found that such a system of orthogonal polynomials is given by the formula

$$\phi_{n,\alpha,\beta}(x) = c_n x^{-\alpha} (1+x)^\beta \frac{d^n}{dx^n} [x^{n+\alpha} (1+x)^{n-\beta}].$$

If these polynomials are standardised such that the coefficient of x^n is unity, we require

$$c_n = (-1)^n \Gamma(\beta - 2n - \alpha) / \Gamma(\beta - n - \alpha).$$

Thus, the Rodrigues formula for $\phi_{n,\alpha,\beta}(x)$ is

$$(5) \quad \phi_{n,\alpha,\beta}(x) = (-1)^n \frac{\Gamma(\beta - 2n - \alpha)}{\Gamma(\beta - n - \alpha)} x^{-\alpha} (1+x)^\beta \frac{d^n}{dx^n} [x^{n+\alpha} (1+x)^{n-\beta}]$$

$$[\alpha > 0, \beta > \alpha + 1, n < \tfrac{1}{2}(\beta - \alpha)].$$

Here α, β are any real numbers such that $\alpha > 0, \beta > \alpha + 1$ and n is the greatest integer less than $\frac{1}{2}(\beta - \alpha)$. These restrictions are imposed by condition (2) and by the requirement for the existence of orthogonal polynomials.

The following recurrence relations are found to be satisfied by (5):

$$(6) \quad \phi_{n,\alpha,\beta}(x) = \frac{1}{2} \left[(2x+1) - \frac{\beta^2 - \alpha^2}{(\beta - 2n - \alpha)(\beta - 2n - \alpha + 2)} \right] \phi_{n-1,\alpha,\beta}(x)$$

$$- \frac{(n-1)(\beta - n + 1)(n + \alpha - 1)(\beta - n - \alpha + 1)}{(\beta - 2n - \alpha + 3)(\beta - 2n - \alpha + 2)^2(\beta - 2n - \alpha + 1)} \phi_{n-2,\alpha,\beta}(x),$$

$$n = 2, 3, 4, \dots,$$

$$(7) \quad \phi_{0,\alpha,\beta}(x) = 1, \quad \phi_{1,\alpha,\beta}(x) = x - (\alpha + 1)/(\beta - \alpha - 2),$$

$$x(1+x)\phi'_{n,\alpha,\beta}(x) + (\beta - \alpha - n - 1) \left[\frac{n + \alpha + 1}{\beta - 2n - \alpha - 2} - x \right] \phi_{n,\alpha,\beta}(x)$$

$$+ (\beta - 2n - \alpha - 1) \phi_{n+1,\alpha,\beta}(x) = 0,$$

$$(8) \quad x(1+x)\phi'_{n,\alpha,\beta}(x) = n \left[\frac{\beta - n}{\beta - 2n - \alpha} + x \right] \phi_{n,\alpha,\beta}(x)$$

$$+ \frac{(\beta - n)(n + \alpha)n(\beta - n - \alpha)}{(\beta - 2n - \alpha)^2(\beta - 2n - \alpha + 1)} \phi_{n-1,\alpha,\beta}(x),$$

$$(9) \quad \phi_{n,\alpha,\beta-1}(x) = \frac{(\beta - n - 2)(\beta - n - \alpha - 1)\phi_{n,\alpha,\beta}(x) + (\beta - 2n - \alpha - 1)(\beta - 2n - \alpha - 2)\phi_{n+1,\alpha,\beta}(x)}{(1+x)(\beta - 2n - \alpha - 1)(\beta - 2n - \alpha - 3)},$$

$$(10) \quad \phi_{n,\alpha+1,\beta}(x) = \frac{1}{x} \left[\frac{(n + \alpha + 1)(\beta - n - \alpha + 1)}{(\beta - 2n - \alpha - 1)(\beta - 2n - \alpha - 2)} \phi_{n,\alpha,\beta}(x) + \phi_{n+1,\alpha,\beta}(x) \right].$$

The differential equation satisfied by the polynomial $\phi_{n,\alpha,\beta}(x)$ is

$$(11) \quad x(1+x)y'' + [1 + \alpha - (\beta - \alpha - 2)x]y' + n(\beta - \alpha - n - 1)y = 0.$$

The polynomials (5) are related to the Jacobi polynomials $P_n^{(a,b)}(x)$ and to the hypergeometric functions $F(a, b; c; x)$ as follows:

$$(12) \quad \phi_{n,\alpha,\beta}(x) = n! \frac{\Gamma(\beta - 2n - \alpha)}{\Gamma(\beta - n - \alpha)} P_n^{(-\beta, \alpha)}[-(2x + 1)],$$

$$(13) \quad \phi_{n,\alpha,\beta}(x) = \frac{\Gamma(\beta)\Gamma(\beta - 2n - \alpha)}{\Gamma(\beta - n)\Gamma(\beta - n - \alpha)} x^n \cdot F\left(-n, -n - \alpha; -\beta + 1; \frac{x+1}{x}\right).$$

The weight coefficients and the error term in (1) are computed to be given by

$$(14) \quad H_i = n! \Gamma(n + \alpha + 1)(\Gamma(\beta - 2n - \alpha))^2 [x_i(1 + x_i)]^{-1} [\phi'_{n,\alpha,\beta}(x_i)]^{-2},$$

$$(15) \quad \begin{aligned} E_{n,\alpha,\beta} &= \frac{f^{(2n)}(\xi)}{(2n)!} \int_0^\infty x^\alpha (1+x)^{-\beta} [\phi_{n,\alpha,\beta}(x)]^2 dx \\ &= \frac{n! \Gamma(n + \alpha + 1)(\Gamma(\beta - 2n - \alpha))^2}{\Gamma(\beta - n - \alpha)\Gamma(\beta - n)(\beta - 2n - \alpha - 1)(2n)!} f^{(2n)}(\xi) \\ &\quad [n < \tfrac{1}{2}(\beta - \alpha) - \tfrac{1}{2}, 0 < \xi < \infty]. \end{aligned}$$

3. Illustration. For $\alpha = 1$, and $\beta = 13$, the following set of polynomials is generated:

$$\begin{aligned} \phi_0 &= 1, \\ \phi_1 &= x - \tfrac{1}{8}, \\ \phi_2 &= x^2 - \tfrac{3}{4}x + \tfrac{1}{12}, \\ \phi_3 &= x^3 - 2x^2 + \tfrac{6}{7}x + \tfrac{1}{14}, \\ \phi_4 &= x^4 - 5x^3 + 6x^2 - 2x + \tfrac{1}{7}, \\ \phi_5 &= x^5 - 15x^4 + 50x^3 - 50x^2 + 15x - 1. \end{aligned}$$

For the quadrature formula

$$(16) \quad \int_0^\infty \frac{x}{(1+x)^{13}} f(x) dx = \sum_{i=1}^n H_i f(x_i),$$

the values of x_i and H_i are given in Table 1.

As an illustration, the integral

$$(17) \quad I(f) = \int_0^\infty \frac{x(1+x)^{1/2}}{(1+x)^{13}} dx = 0.0082 \ 8157$$

has been approximated using formula (16) and also using a straightforward application of the Gauss-Laguerre formula. This example is 'hand-tailored' for the use of formula (16) and, since it does not behave asymptotically like e^{-x} , the Gauss-Laguerre formula is quite inappropriate. The numerical results listed in Table 2 show that for this example, formula (16) is much more accurate than the Gauss-Laguerre formula.

TABLE 1
Abscissas and Weights for Quadrature (17)

x	x_i	H_i
1	0.2000 0000	(-2) 0.7575 7575
2	0.6143 5678 0.1356 4322	(-2) 0.1018 4615 (-2) 0.6557 2960
3	0.1100 2615 0.4512 2182 (1) 0.1438 7520	(-2) 0.5645 9170 (-2) 0.1906 3812 (-4) 0.2345 9366
4	(-1) 0.9786 2600 0.3868 4283 (1) 0.1107 2489 (1) 0.3408 0457	(-3) 0.1039 5084 (-4) 0.4906 0878 (-5) 0.1594 2659 (-8) 0.1309 1096
5	(-1) 0.9276 2478 0.3616 1054 1.0000 0000 (1) 0.2765 4061 (2) 0.1078 0221	(-2) 0.3982 9769 (-2) 0.1066 6655 (-4) 0.2976 1905 (-6) 0.8624 1404 (-9) 0.2354 0418

TABLE 2
Error in the Evaluation of (17)

n	Gauss-Laguerre	Formula (16)	Upper bound (15)
1		-1725×10^{-8}	-2525×10^{-8}
2	54 6101×10^{-8}	-34×10^{-8}	-129×10^{-8}
3	24 7277×10^{-8}	-2×10^{-8}	-58×10^{-8}
4	1 1412×10^{-8}		
5	-13 6969×10^{-8}		
6	-21 4710×10^{-8}		
7	-24 4831×10^{-8}		
8	-24 5982×10^{-8}		
16	-7 8818×10^{-8}		
24	-1 6744×10^{-8}		
32	- 3506×10^{-8}		

The last column of Table 2 contains the upper bounds on errors calculated through the estimate (15) after replacing $f^{(2n)}(\xi)$ by $\max_{0 \leq x \leq \infty} |f^{(2n)}(x)|$.

4. Acknowledgements. We are grateful to Professor L. Fox for his suggestions and to Professor C. W. Clenshaw for his help and comments. We are indebted to

the referee for his suggested improvement of the text in Sections 1 and 3, especially for a better layout of Table 2.

Department of Mathematics
Indian Institute of Technology
Hauz Khas, New Delhi—29, India

1. W. M. HARPER, "Quadrature formulas for infinite integrals," *Math. Comp.*, v. 16, 1962, pp. 170–175. MR 26 #3185.
2. F. B. HILDEBRAND, *Introduction to Numerical Analysis*, McGraw-Hill, New York, 1956, pp. 258–367. MR 17, 788.
3. A. H. STROUD & DON SECREST, *Gaussian Quadrature Formulas*, Prentice-Hall, Englewood Cliffs, N.J., 1966, p. 254. MR 34 #2185.
4. G. SZEGÖ, *Orthogonal Polynomials*, 2nd rev. ed., Amer. Math. Soc. Colloq. Publ., vol. 23, Amer. Math. Soc., Providence, R.I., 1959, pp. 58–72. MR 21 #5029.