

Chebyshev Approximation of $(1 + 2x)\exp(x^2)\operatorname{erfc} x$ in $0 \leq x < \infty$

By M. M. Shepherd and J. G. Laframboise

Abstract. We have obtained a single Chebyshev expansion of the function $f(x) = (1 + 2x)\exp(x^2)\operatorname{erfc} x$ in $0 \leq x < \infty$, accurate to 22 decimal digits. The presence of the factors $(1 + 2x)\exp(x^2)$ causes $f(x)$ to be of order unity throughout this range, ensuring that the use of $f(x)$ for approximating $\operatorname{erfc} x$ will give uniform relative accuracy for all values of x .

I. Introduction. The functions $\operatorname{erfc} x = (2/\sqrt{\pi})\int_x^\infty \exp(-t^2) dt$ and $\exp(x^2)\operatorname{erfc} x$ occur frequently in kinetic theory of gases and related subjects. Calculation of these functions using the identity $\operatorname{erfc} x = 1 - \operatorname{erf} x$, together with available approximations [1] for $\operatorname{erf} x$, usually results in large relative errors for large x because $\operatorname{erf} x \rightarrow 1$ as $x \rightarrow \infty$. To overcome this difficulty, Clenshaw [2], Luke [3], [4], and Schonfelder [5] have presented Chebyshev approximations in which the range $0 \leq x < \infty$ is split into two ranges $0 \leq x \leq c$ and $c \leq x < \infty$, with $\operatorname{erf} x$ being Chebyshev-approximated in $0 \leq x \leq c$, and $x \exp(x^2)\operatorname{erfc} x$ being Chebyshev-approximated in $c \leq x < \infty$. Clenshaw [2] uses $c = 4$, 33 terms for $x \leq 4$ and 18 terms for $x > 4$, and obtains an accuracy of twenty decimal places (20D). Corresponding figures for Luke [3], [4] and Schonfelder [5] are $c = 3$, 25 and 22 terms, and 20D; and $c = 2$, 27 and 43 terms, and 30D. These authors use various transformations $t(x)$ to map $c \leq x < \infty$ into $-1 \leq t < 1$. Use of the identity $\operatorname{erfc}(-x) = 2 - \operatorname{erfc} x$ eliminates the need to approximate $\operatorname{erfc} x$ for negative x .

Schonfelder [5] has also presented a single 43-term Chebyshev expansion of $\exp(x^2)\operatorname{erfc} x$ for the entire interval $0 \leq x < \infty$, using a relation of the form $t = (x - k)/(x + k)$ to map this interval into $-1 \leq t < 1$. Oldham [6] has presented a simple approximation of $\sqrt{\pi}x \exp(x^2)\operatorname{erfc} x$, having a maximum relative error of one part in 7000 and suitable for hand calculation.

Whenever a function to be Chebyshev-approximated has a zero within its interval of definition or at either end of it, such an approximation is likely to give large relative errors near such a zero because the usual procedures for calculating Chebyshev coefficients minimize maximum *absolute* error. Accordingly, it is advantageous to multiply $\operatorname{erfc} x$ by factors which yield a product of order unity for all x in $(0, \infty)$ and then to Chebyshev-approximate this product function, because one will then obtain good uniformity of relative as well as absolute error. Our chosen function, $f(x) = (1 + 2x)\exp(x^2)\operatorname{erfc} x$, satisfies this criterion. It has limiting values

Received April 14, 1980.

AMS (MOS) subject classifications (1970). Primary 33-04, 33A20, 41-04, 41A10, 41A20, 41A50, 41A60, 42-04, 42A56, 65A05, 65B05, 65D20.

© 1981 American Mathematical Society
0025-5718/81/0000-0021/\$02.25

of 1 and $2/\sqrt{\pi} \approx 1.13$ at $x = 0$ and $x \rightarrow \infty$, respectively. In comparison, Schonfelder's function $\exp(x^2)\operatorname{erfc} x$ approaches 0 as $x \rightarrow \infty$. Furthermore, our choice contains no irrational coefficients, in contrast with the more obvious choice $(1 + \sqrt{\pi}x)\exp(x^2)\operatorname{erfc} x$, which $\rightarrow 1$ at $x = 0$ and $x \rightarrow \infty$. A graph of $f(x)$ appears in Figure 1.




FIGURE 1

Graph of the function $F(t)$ defined by the relation $f(x) = (1 + 2x)\exp(x^2)\operatorname{erfc} x$ together with the mapping $t = (x - 3.75)/(x + 3.75)$.

We have used the same transformation as that of Schonfelder [5], i.e. $t = (x - k)/(x + k)$ with $k = 3.75$, to map $0 < x < \infty$ into $-1 < t < 1$. Tests of various k values for our $f(x)$ yielded results similar to his, namely that this value gives near-optimum convergence of the resulting Chebyshev series over the precision range of greatest interest, i.e. 8D to 18D. Our calculations were done in IBM quadruple precision, which yields a machine precision of 34D.

II. Calculation of Chebyshev Coefficients. We have used the usual [7] form of an m th-order Chebyshev expansion. Thus, the Chebyshev polynomials $T_j(t)$ are given by

$$(1) \quad T_j(t) = \cos(j \arccos t); \quad j = 0, 1, 2, \dots$$

The above-mentioned $f(x)$ and transformation from x to t define a function $F(t)$ which is expanded as follows:

$$(2) \quad F(t) = \sum_{j=0}^m c_j T_j(t),$$

where

$$(3) \quad c_j = \frac{\sum_{k=0}^m F(t_k) T_j(t_k)}{\|T_j\|^2},$$

$$(4) \quad t_k = \cos\left(\frac{2k + 1}{m + 1} \frac{\pi}{2}\right); \quad k = 0, 1, 2, \dots, m,$$

$$(5) \quad \|T_0\|^2 = m + 1; \quad \|T_i\|^2 = \frac{1}{2}(m + 1) \quad \text{for } i > 0.$$

In order to calculate the required values of $f(x)$, we note that the Taylor expansion

$$(6) \quad \operatorname{erfc} x = 1 - \frac{2}{\sqrt{\pi}} \left(t - \frac{t^3}{3} + \frac{t^5}{5 \cdot 2!} - \frac{t^7}{7 \cdot 3!} + \frac{t^9}{9 \cdot 4!} - \dots \right)$$

can be rearranged [8] into the form

$$(7) \quad \exp(x^2)\operatorname{erfc} x = \exp(x^2) - \frac{2x}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{2^n x^{2n}}{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n + 1)},$$

the use of which is less sensitive to roundoff errors.

The asymptotic expansion

$$(8) \quad \exp(x^2)\operatorname{erfc} x = \frac{1}{x\sqrt{\pi}} \left(1 - \frac{1}{2x^2} + \frac{1 \cdot 3}{(2x^2)^2} - \frac{1 \cdot 3 \cdot 5}{(2x^2)^3} + \dots \right)$$

is of limited use when x is large. The continued-fraction expansion

$$(9) \quad \sqrt{\pi} x \exp(x^2)\operatorname{erfc} x = \cfrac{1}{1} + \cfrac{\cfrac{1}{2x^2}}{\cfrac{1}{1}} + \cfrac{\cfrac{2}{2x^2}}{\cfrac{1}{1}} + \cfrac{\cfrac{3}{2x^2}}{\cfrac{1}{1}} + \dots$$

(Perron [9]) yields better precision. Perron [10] gives the following algorithm for use of (9).

We define

$$(10) \quad \begin{aligned} A_{-1} &= 1, & A_0 &= b_0, \\ B_{-1} &= 0, & B_0 &= 1; \\ a_i &= i / (2x^2), & b_i &= 1 \quad \text{for } i = 0, 1, 2, 3, \dots; \\ b_0 + \cfrac{a_1}{b_1} + \cfrac{a_2}{b_2} + \dots + \cfrac{a_n}{b_n} &= \cfrac{A_n}{B_n}. \end{aligned}$$

Then A_n and B_n are given recursively by the relations

$$(11) \quad \left. \begin{aligned} A_n &= b_n A_{n-1} + a_n A_{n-2} \\ B_n &= b_n B_{n-1} + a_n B_{n-2} \end{aligned} \right\}, \quad n = 1, 2, 3, \dots$$

At smaller values of x , the convergence of (9) becomes slower. To overcome this, we have used double Aitken δ^2 extrapolation (Burden et al. [11, pp. 56–57]) as follows. If y_j is the approximation obtained by taking j terms of (9), then the sequences of numbers

$$(12) \quad \begin{aligned} y'_j &= y_j - (y_j - y_{j-1})^2 / (y_j - 2y_{j-1} + y_{j-2}), \\ y''_j &= y'_j - (y'_j - y'_{j-1})^2 / (y'_j - 2y'_{j-1} + y'_{j-2}), \end{aligned}$$

converge progressively faster to (9). Use of (12) with (9)–(11) improved the fit of the Chebyshev approximation by about five orders of magnitude.

We have used (7) for $x \leq 2.83$ and (9)–(12) for $x > 2.83$. This yielded values of $f(x)$ accurate to at least 23D for all x . The resulting Chebyshev coefficients,

generated by (3)–(5), are shown in Table 1. Use of these in (2) gives an approximation which reproduces $f(x)$ to at least 22D for all x . Schonfelder [5] generates Chebyshev coefficients using a different method [12], [13] in which an expression equivalent to (2) is substituted into a linear differential equation satisfied by the given function. Together with the boundary conditions satisfied by the same function, this procedure generates an infinite set of simultaneous linear equations for the c_j , a truncated version of which is then solved.

TABLE 1

$$Y = (1+2X) * \text{EXP}(X*X) * \text{ERFC}(X) \quad X = (0, \text{INF})$$

$$T = (X-K)/(X+K) \quad K = 3.75$$

ORD	C(N)
0	0.11775789345674017540800+01
1	-0.4590054580646477331-02
2	-0.84249133366517915584-01
3	0.59209939998191890498-01
4	-0.26658668435305752277-01
5	0.9074997670705265094-02
6	-0.2413163540417608191-02
7	0.490775836525808632-03
8	-0.69169733025012064-04
9	0.4139027986073010-05
10	0.774038306619849-06
11	-0.218864010492344-06
12	0.10764999465671-07
13	0.4521959811218-08
14	-0.775440020883-09
15	-0.63180883409-10
16	0.28687950109-10
17	0.194558685-12
18	-0.965469675-12
19	0.32525481-13
20	0.33478119-13
21	-0.1864563-14
22	-0.1250795-14
23	0.74182-16
24	0.50681-16
25	-0.2237-17
26	-0.2187-17
27	0.27-19
28	0.97-19
29	0.3-20
30	-0.4-20

III. Acknowledgments. We wish to thank Y. L. Luke, G. Dahlquist, and S.-Å. Gustavson for valuable discussions and comments. This work was supported by the Natural Sciences and Engineering Research Council of Canada under grants A-4638 and A-4749.

Department of Computer Science
York University
Toronto, Ontario, Canada M3J 1P3

Department of Physics
York University
Toronto, Ontario, Canada M3J 1P3

1. C. HASTINGS, JR., *Approximations for Digital Computers*, Princeton Univ. Press, Princeton, N. J., 1955, p. 169.

2. C. W. CLENSHAW, "Chebyshev series for mathematical functions," in *National Physical Laboratory Mathematical Tables*, Vol. 5, Her Majesty's Stationery Office, London, 1962.

3. Y. L. LUKE, *The Special Functions and their Approximations* (2 Volumes; see especially Vol. 2, pp. 323–324), Academic Press, New York, 1969.

4. Y. L. LUKE, *Mathematical Functions and their Approximations*, Academic Press, New York, 1975, pp. 123–124.
5. J. L. SCHONFELDER, “Chebyshev expansions for the error and related functions,” *Math. Comp.*, v. 32, 1978, pp. 1232–1240.
6. K. B. OLDHAM, “Approximations for the $x \exp x^2 \operatorname{erfc} x$ function,” *Math. Comp.*, v. 22, 1968, p. 454.
7. G. DAHLQUIST & Å. BJÖRCK, *Numerical Methods*, Prentice-Hall, Englewood Cliffs, N. J., 1974, pp. 104–108.
8. S.-Å. GUSTAVSON, Private communication, 1979.
9. O. PERRON, *Die Lehre von den Kettenbrüchen*, Band II, Teubner, Berlin, 1929, p. 103.
10. O. PERRON, *Die Lehre von den Kettenbrüchen*, Band I, Teubner, Berlin, 1929, p. 4.
11. R. L. BURDEN, J. D. FAIRES & A. C. REYNOLDS, *Numerical Analysis*, Prindle, Weber & SCHMIDT, BOSTON, 1978, pp. 56–57.
12. L. FOX & I. B. PARKER, *Chebyshev Polynomials in Numerical Analysis*, Oxford Univ. Press, London and New York, 1968.
13. J. L. SCHONFELDER, “Generation of high precision Chebyshev expansions.” (In preparation.)