

## A Monte Carlo Factoring Algorithm With Linear Storage

By C. P. Schnorr\* and H. W. Lenstra, Jr.

**Abstract.** We present an algorithm which will factor an integer  $n$  quite efficiently if the class number  $h(-n)$  is free of large prime divisors. The running time  $T(n)$  (number of compositions in the class group) satisfies  $\text{prob}[T(m) \leq n^{1/2r}] \geq (r-2)^{-(r-2)}$  for random  $m \in [n/2, n]$  and  $r \geq 2$ . So far it is unpredictable which numbers will be factored fast. Running the algorithm on all discriminants  $-ns$  with  $s \leq r'$  and  $r = \sqrt{\ln n / \ln \ln n}$ , every composite integer  $n$  will be factored in  $o(\exp \sqrt{\ln n \ln \ln n})$  bit operations. The method requires an amount of storage space which is proportional to the length of the input  $n$ . In our analysis we assume a lower bound on the frequency of class numbers  $h(-m)$ ,  $m \leq n$ , which are free of large prime divisors.

**1. Introduction.** The problem of factoring an integer  $n$  into its prime power divisors is computationally equivalent to determining all ambiguous, reduced positive forms  $ax^2 + bxy + cy^2$  (notation  $(a, b, c)$ ),  $a, b, c \in \mathbb{Z}$ , with discriminant  $b^2 - 4ac = -n$  ( $b^2 - 4ac = +n$ , resp.). In fact these ambiguous forms correspond to the relatively prime factorizations of  $n$ , i.e. to the pairs  $(n_1, n_2)$  with  $n = n_1 n_2$ ,  $\gcd(n_1, n_2) = 1$ .

According to Gauss [5] the equivalence classes of forms with fixed discriminant  $\Delta$  form a group under composition, the *class group*  $G(\Delta)$ . The order  $h(\Delta)$  of this group is the *class number*. Multiplication in  $G(\Delta)$  can be done efficiently working with representatives of classes. The *ambiguous* classes are the classes  $H$  with  $H^2 = 1$ .

In case of negative discriminant  $\Delta < 0$  there is a unique reduced form in each class, and this form can be efficiently calculated from any other class representative. Therefore, factoring  $n$  is computationally equivalent to determining representatives of all ambiguous classes in  $G(-n)$ . The reduced forms of these classes correspond to the relatively prime factorizations  $n_1 n_2 = n$  of  $n$ .

In case of positive discriminants  $\Delta > 0$ , under a different concept of reduction, there are  $O(\sqrt{\Delta} \ln \Delta)$  reduced forms in each class. They form a cycle under the reduction operation. Composition of forms yields a group-like structure on the principal cycle. A reduced form  $(a, b, c)$  is *ambiguous* if its square under composition yields the unit form. Again, the ambiguous, reduced forms with discriminant  $n > 0$  correspond to the relatively prime factorizations  $n_1 n_2 = n$  of  $n$ .

Several factoring algorithms have been developed on this basis. Here we are only concerned with negative discriminants. For positive discriminants, see the algorithms of Shanks as described in Monier [11] and Wagstaff-Wunderlich [22]. For negative

---

Received March 21, 1983.

1980 *Mathematics Subject Classification*. Primary 10A30, 10C07, 68C25.

\*Research supported by Bundesminister für Forschung und Technologie under grant 08 30108.

©1984 American Mathematical Society  
0025-5718/84 \$1.00 + \$.25 per page

discriminants Shanks [17], after guessing generators for  $G(-n)$ , computes the class number  $h(-n)$  by exploiting the group structure. Then Shanks computes  $H(k) := H^{h(-n)/2^k}$  for the smallest  $k$  such that  $H(k) \neq 1$  with  $H \in G(-n)$  chosen arbitrarily. Clearly  $H(k)$  is ambiguous. Under reasonable assumptions it takes  $O(n^{1/4})$  steps to factor  $n$  in this way. This method can be speeded up to an  $O(n^{1/5})$ -algorithm by approximating  $h(-n)$  via the class formula (let  $(\frac{m}{p})$  denote the Kronecker symbol):

$$\lim_m \frac{\sqrt{n}}{\pi} \prod_{p \leq m} \frac{p}{p - (\frac{-n}{p})} = h(-n),$$

which by the generalized Riemann hypothesis has an error term  $O(\ln(mn)n^{1/2}m^{-1/2})$ . For this algorithm the amount of storage will be proportional to the running time.

In Schnorr [19] a method was proposed to generate ambiguous forms which is similar to the Morrison-Brillhart factoring algorithm. We collect equations

$$H_i^2 = \prod_p H_p^{a_{p,i}}, \quad a_{p,i} \in \mathbb{Z},$$

with  $H_i \in G(-n)$  chosen at random and  $H_p = [(p, b_p, c_p)]$  for small primes  $p$ . By combining these equations one obtains

$$H = \prod_i H_i^{h_i}, \quad F = \prod_p H_p^{(\sum_i h_i a_{p,i})/2}$$

such that  $H^2 = F^2$ ,  $H \neq F$ . Then  $HF^{-1}$  is ambiguous. Under reasonable assumptions  $n$  will be factored with  $o(\exp \sqrt{\frac{3}{2}} \ln n \ln \ln n)$  steps and  $o(\exp \sqrt{\frac{2}{3}} \ln n \ln \ln \ln n)$  storage.

The new algorithm, given the first  $t$  primes  $p_1 = 2, p_2 = 3, \dots, p_t = n^{1/2^r}$ , needs only to store a fixed number of forms which takes  $O(\log n)$  bits. Let  $e_i = \max\{v: p_i^v \leq p_i^2\}$ . Then Stage 1 of the new algorithm computes

$$H = H_0^{\prod_{i=2}^t p_i^{e_i}}$$

for an arbitrarily chosen  $H_0 \in G(-n)$ . Then compute  $H^{2^k}$  for the smallest  $k \leq \log_2 \sqrt{n}$  such that  $H^{2^k} = 1$ . Clearly  $H^{2^{k-1}}$  is ambiguous.  $n$  will be factored by Stage 1 if  $h(-n)$  divides  $2^k \prod_{i=2}^t p_i^{e_i}$  for some  $k$ . If Stage 1 fails then Stage 2 does a random walk through the group generated by  $H$ .

Stage 2 will factor  $n$  if  $\text{ord}(H^{2^k}) < p_i^2$  for some  $k$ , i.e. if  $h(-n)$  divides  $2^k \prod_{i=2}^t p_i^{e_i} q$  for some  $q \leq p_i^2$ . With  $p_t = n^{1/2^r}$  Stage 1 of the algorithm takes  $O(p_t)$  compositions and for random composite  $m \in [0, n]$  with probability  $\geq r^{-r}$  detects a proper divisor of  $m$ . Stage 2 also takes  $O(p_t) = O(n^{1/2^r})$  compositions and with probability  $\geq (r-2)^{-(r-2)}$ ,  $r \geq 2$ , detects a proper divisor of  $m$ . Running Stage 1 on the integers  $ns$  for  $s \leq r'$ ,  $r = \sqrt{\ln n / \ln \ln n}$ , every composite integer  $n$  will be factored within  $o(\exp \sqrt{\ln n \ln \ln n})$  bit operations. The latter bound already takes into account the cost of the arithmetic. The cost for a composition in  $G(-n)$  is proportional to the cost of the extended Euclidean algorithm, which given integers  $u, v \leq \sqrt{n}$  computes  $r, s \in \mathbb{N}$  with  $ru + sv = \gcd(u, v)$ . Using standard algorithms for multiplication and division this takes  $O(\ln n)^2$  bit operations, i.e. binary Boolean operations; see Knuth [7, 4.5.2, exercise 30 and algorithm X].

The particular features of the new factoring algorithm are:

(1) it can easily be operated with  $O(\log n)$  bit storage,

(2) it is Monte Carlo in the sense that every 1000th integer will be factored about 1000 times faster than average time,

(3) the integers which will be factored very fast are randomly distributed; there is no way to predict whether a given  $m$  will be factored fast,

(4) the algorithm is of the parallel type, e.g. 1000 processors will factor 1000 times faster.

Properties (2), (3) seem to endanger the RSA-cryptoscheme, see Rivest et al. [15]. In particular no methods are known that generate class numbers with large prime divisors.

Stages 1 and 2 of the algorithm are presented in Sections 2 and 3. The main algorithm which factors arbitrary integers is given in Section 4. Some computational experience with the factoring algorithm is reported in Section 5. In Appendix I we collect basic theorems and algorithms on quadratic forms. Appendix II contains various tables which demonstrate the performance of the algorithm. We exemplify the distribution of class numbers and integers which are free of large prime divisors, the frequency of class numbers divisible by small primes, and the performance of various pseudo-random functions used in Stage 2 of the algorithm.

**2. Stage 1 of the Algorithm.** Let  $n$  be the integer to be factored.  $-n$  is the discriminant of some quadratic form if and only if  $-n \equiv 1 \pmod{4}$  or  $-n \equiv 0 \pmod{4}$ . The purpose of Stages 1, 2 is to find a nontrivial divisor of  $n$ , provided  $-n$  is a discriminant and  $h(-n)$  is a product of small primes. In order to factor general integers  $n$ , the main algorithm in Section 4 applies Stages 1, 2 to multiples  $ns$  with  $-ns \equiv 0, 1 \pmod{4}$ . If  $-n$  is a discriminant, we can easily construct forms  $(a, b, c)$  with discriminant  $-n$ : choose a small odd prime  $p$  with  $(\frac{-n}{p}) = 1$  and solve  $b^2 \equiv -n \pmod{4p}$  which yields  $b^2 = -n + 4pc$  for some  $c \in \mathbb{Z}$ . Hence  $(p, b, c)$  has discriminant  $-n$ .

Throughout Sections 2, 3 we restrict ourselves to the case  $-n \equiv 1 \pmod{4}$ ; consult Theorem III, Appendix I for the case  $-n \equiv 0 \pmod{4}$ . Then the unit  $1 \in G(-n)$  is represented by the form  $(1, 1, (1+n)/4)$ . This ambiguous class yields the improper factorization  $1 \cdot n = n$ . The other ambiguous classes correspond in a 1-1 way to the relatively prime factorizations of  $n$  with nontrivial divisors.

*Stage 1.* Let  $n \in \mathbb{N}$ ,  $-n \equiv 1 \pmod{4}$ , be given.

1. For some  $t \in \mathbb{N}$  compute the  $t$  first primes  $p_1 = 2, p_2 = 3, \dots, p_t$ ,
2. choose  $H_0 \in G(-n)$  arbitrarily,
3.  $H := H_0^{\prod_{i=1}^t p_i^{e_i}}$  with  $e_i := \max\{v: p_i^v \leq p_i^2\}$ ; if  $H = 1$ , then stop (in this case  $\text{ord}(H_0)$  is odd and another  $H_0$  must be chosen),
4.  $\bar{H} := H, e_* := \lfloor \log_2 \sqrt{n} \rfloor$ ,
5. for  $v = 1, 2, \dots, e_*$  do [ $S := H, H := H^2$ , if  $H = 1$  go to 7],
6. go to Stage 2,
7. (at this point  $S$  is ambiguous and yields some divisor  $d$  of  $n$ ).

Stage 1 by itself is the core of the new factoring algorithm. The improvements resulting from Stage 2 are important for practical applications, but they scarcely influence the asymptotic time bound of the main algorithm.

*Fact 1.* Suppose  $h(-n) \mid \prod_{i=1}^t p_i^{e_i}$  and  $\text{ord}(H_0)$  is even, then Stage 1 generates an ambiguous class  $S \neq 1$ .

In case  $-n \equiv 1 \pmod{4}$  every ambiguous class  $S \neq 1$  yields a proper divisor of  $n$ . In particular, when  $n$  has  $d$  odd prime divisors, then  $2^{d-1} \mid h(-n)$ , and there are exactly  $2^{d-1}$  ambiguous classes corresponding to the  $2^{d-1}$  pairs  $\{n_1, n_2\}$  with  $n_1 n_2 = n$ ,  $n_1 < n_2$ ,  $\gcd(n_1, n_2) = 1$ . Moreover, when  $n$  is composite and  $H_0 \in G(-n)$  is chosen at random, then  $\text{prob}[\text{ord}(H_0) \text{ even}] \geq 1/2$ . Hence Stage 1 has a chance  $\geq 1/2$  to find a proper divisor of  $n$ , provided  $h(-n) \mid \prod_{i=1}^t p_i^{e_i}$ . A few repetitions of Stage 1 almost surely generate a proper divisor of  $n$ , provided  $h(-n) \mid \prod_{i=1}^t p_i^{e_i}$  and  $n$  is composite:

*Fact 2.* Suppose  $h(-n) \mid \prod_{i=1}^t p_i^{e_i}$  and  $n$  is composite. If Stage 1 is passed with  $H_0$  chosen independently  $k$  times, then with probability  $\geq 1 - 2^{-k}$  a proper divisor of  $n$  has been found.

Next consider the chance that for random  $m \leq n$ :

$$h(-m) \mid \prod_{i=1}^t p_i^{e_i}, \quad e_i = \max\{v: p_i^v \leq p_i^2\}.$$

Siegel [21] proved:

$$\forall \varepsilon: \exists n_\varepsilon: \forall m \geq n_\varepsilon: h(-m) \in [m^{1/2-\varepsilon}, m^{1/2+\varepsilon}].$$

We will base the analysis of Stage 1 on the following hypothesis

For all  $n$  and  $t$ :

$$\begin{aligned} (2.1) \quad & \# \left\{ m \leq n: h(-m) \mid \prod_{i=1}^t p_i^{e_i} \right\} / (0.5n) \\ & \geq \# \left\{ \bar{m} \leq \sqrt{n}: \bar{m} \mid \prod_{i=1}^t p_i^{e_i} \right\} / \sqrt{n}. \end{aligned}$$

H. G. Franke has tested this hypothesis experimentally, see Tables 1–3, Appendix II for  $n \approx 4.7 \cdot 10^8$ . In general the first term in (2.1) is considerably larger than the second. Note that the frequency of class numbers  $h(-n)$  of fundamental discriminants (and a fortiori of general discriminants) which are divisible by  $p$  is larger than  $1/p$  and is close to  $1/(p-1)$  provided  $p$  is small with respect to  $n$ . These experimental data and some recent calculations of Cohen and H. W. Lenstra, Jr. indicate that class groups  $G(-n)$  of fundamental discriminants are distributed like random Abelian groups of order  $O(\sqrt{n} \log n)$ . Cohen and Lenstra have calculated  $\text{prob}[m \text{ divides } |G|]$  for  $m = 2, 3, \dots$  and for random Abelian groups, where the probability weight of  $G$  is proportional to  $1/|\text{Aut}(G)|$ . The values of Cohen and Lenstra completely match with our experimental data. A corresponding observation with respect to prime discriminants has been made by Leopoldt as cited in Zimmer [23].

Recently Canfield, Erdős, and Pomerance improved the theoretical lower bound on the second term in (2.1). We refer in particular to the proof in Pomerance [14]:

**THEOREM 3.** Let  $\Psi(n, v) := \#\{x \leq n: x \text{ free of primes } > v\}$ . For every  $\varepsilon > 0$  there exists  $c_\varepsilon$  such that for all  $n \geq 10$  and all  $r$  with  $n^{1/r} \geq (\ln n)^{1+\varepsilon}$ :  $\Psi(n, n^{1/r})/n \geq (c_\varepsilon r \ln r)^{-r}$ .

In practice, however,  $\Psi(n, n^{1/r})/n$  is larger than the bound stated in Theorem 3. From experimental data, see Table 2, Appendix II, we conclude

$$(2.2) \quad \text{for all } n \text{ and } r \leq \sqrt{\ln n / \ln \ln n}: \Psi(n, n^{1/r})/n \geq r^{-r}.$$

**COROLLARY 4.** Assume (2.2) and (2.1). Then for all  $n \geq 10^{20}$  and all  $p_t = n^{1/2r}$  with  $r \leq \sqrt{\ln n / \ln \ln n}$ :

$$\# \left\{ m \leq n: h(-m) \mid \prod_{i=1}^t p_i^{e_i} \right\} / (0.5n) \geq 0.83 r^{-r}.$$

*Proof.*

$$\begin{aligned} & \# \left\{ m \leq n: h(-m) \mid \prod_{i=1}^t p_i^{e_i} \right\} / (0.5n) \\ (2.1) \quad & \geq \Psi(\sqrt{n}, n^{1/2r}) / \sqrt{n} - \sum_{i=1}^t p_i^{e_i-1} \\ (2.2) \quad & \geq r^{-r} - n^{-1/r_t} \\ & \geq r^{-r} - 1.1 n^{-1/r} n^{1/2r} / \ln n^{1/2r} \quad (\text{since } t = \pi(n^{1/2r}) \leq 1.1 n^{1/2r} / \ln n^{1/2r}) \\ & \geq r^{-r} - 2.2 n^{-1/2r} r / \ln n \\ & \geq r^{-r} (1 - 2.2r / \ln n) \\ & \quad (\text{in fact } r \leq \sqrt{\ln n / \ln \ln n} \text{ implies } r^{-r} \geq n^{-1/2r}) \\ & \geq 0.83 r^{-r} \quad \text{for } n \geq 10^{20}, r \leq \sqrt{\ln n / \ln \ln n}. \quad \square \end{aligned}$$

*Runtime of Stage 1.* If  $H^{p_i^{e_i}}$  is computed by the binary method (see Knuth [7, §4.63]), this takes

$$2 \log_2 p_i^{e_i} \leq 2 \log_2 p_i^2 \leq 4 \log_2 p_i$$

compositions in  $G(-n)$ . Since there are about

$$t \leq p_t / \ln p_t$$

primes  $\leq p_t$ , this yields a worst case bound of

$$\frac{4}{\ln 2} p_t \approx 5.8 p_t \quad \text{compositions in total.}$$

On the average, the binary method is somewhat more efficient. It takes about  $1.5 \log p_i^{e_i}$  compositions to compute  $H^{p_i^{e_i}}$  and therefore Stage 1 will only take about  $4.4 p_t$  compositions in total. All together we have proved the following

**THEOREM 5.** Assume (2.1), (2.2), and that for every discriminant  $m \leq n$  (a single)  $H_0 \in G(-n)$  in Stage 1 is chosen at random. Then for all  $n \geq 10^{20}$  and all  $p_t = n^{1/2r}$  with  $r \leq \sqrt{\ln n / \ln \ln n}$ : Stage 1 factors at least a  $0.83 n r^{-r}$  fraction of the discriminants  $\leq n$  and takes about  $4.4 p_t$  compositions in  $G(-n)$ .

*Remark.* The discriminants which will be factored are “randomly” distributed in  $[0, n]$ .

For *practical applications* we advise to choose somewhat smaller exponents  $e'_i$  instead of the  $e_i$ :

$$e'_i := \max \{ v: p_i^v \leq p_t \} \quad \text{with } p_t = n^{1/2r}.$$

We used the larger  $e_i$  for proving Corollary 4 by a crude argument. Assuming  $\Psi(n, n^{1/r})/n = O(r^{-r})$  one obtains Corollary 4 for the  $e'_i$ :

$$\begin{aligned} & \# \left\{ m \leq n : h(-m) \mid \prod_{i=1}^t p_i^{e'_i} \right\} / (0.5n) \\ & \quad (\text{assuming that (2.1) holds for the } e_i) \\ & \geq r^{-r} - \sum_{i=1}^t p_i^{-e'_i-1} \frac{\Psi(n^{1/2-1/2r}, n^{1/2r})}{n^{1/2-1/2r}} \\ & \geq^* r^{-r} - O\left(n^{-1/2r} \frac{n^{1/2r}}{\ln(n^{1/2r})} (r-1)^{-(r-1)}\right) \\ & \geq r^{-r} - O\left(\frac{2r}{\ln n} (r-1)^{-(r-1)}\right) \\ & \geq r^{-r} - O\left((r-1)^{-(r-1)} / (r \ln \ln n)\right) = r^{-r} (1 - O(1/\ln \ln n)) \\ & \quad (\text{since } r \leq \sqrt{\ln n / \ln \ln n}). \end{aligned}$$

The choice of the  $e'_i$  are justified by our data in Appendix II. Tables 1, 2 show that there are only a few discriminants  $h(-m)$ ,  $m \approx 4.7 \cdot 10^8$ , such that  $h(-m) = \prod_{i=1}^t p_i^{\bar{e}_i}$  with  $\bar{e}_i > e'_i$  for some  $i \geq 2$ .

Table 4, Appendix II, by Odlyzko considers large integers. This table shows that for  $r \leq \sqrt{\ln n / \ln \ln n}$

$$\# \{ m \leq n : m \text{ even, } m \mid \text{lcm}(2, 3, \dots, p_t) \} / n > r^{-r}.$$

Note that  $m \mid \text{lcm}(2, 3, \dots, p_t) \Leftrightarrow m = \prod_i p_i^{\bar{e}_i}$  with  $\bar{e}_i \leq e'_i$  for all  $i$ . Hence in practice Corollary 4 even holds when the  $e'_i$  are taken for the  $e_i$ , and the constant 0.83 can be replaced by some constant  $> 1$ . If we use the  $e'_i$  then Stage 1 will only take about  $2.2 p_t$  compositions.

**3. Using a Pollard-Brent Recursion in Stage 2.** If  $h(-n) + 2^{e_*} \prod_{i=2}^t p_i^{e'_i}$  with  $e_i = \lfloor \log_2 n / \log_2 p_i \rfloor$ ,  $e_* = \lfloor \log_2 \sqrt{n} \rfloor$ , then Stage 1 fails to factor  $n$  and computes

$$\bar{H} := H_0^{\prod_{i=2}^t p_i^{e'_i}}, \quad H := \bar{H}^{2^{e_*}}.$$

Stage 2 uses  $H, \bar{H}$  and will most likely find a proper divisor of  $n$  within  $O(p_t)$  steps, provided that  $\text{ord}(H) \leq p_t^2$  and  $\text{ord}(H_0)$  is even.

Stage 2 generates a random walk through the cyclic group  $\langle H \rangle$  with generator  $H$ . With some function  $f: \langle H \rangle \rightarrow \langle H \rangle$  let

$$H_1 := H, \quad H_{i+1} := f(H_i).$$

The function  $f$  must be chosen such that

(3.1)  $f$  is easy to compute,

(3.2)  $f$  is sufficiently random,

(3.3) every relation  $H_j = H_k$  with  $j \neq k$  yields an ambiguous class  $S$ , depending on  $H, f, j, k$ .

It is known (see Knuth [7, Exercise 3.1.12]) that some  $j < k \leq \sqrt{\pi/2} p_t$  with  $H_j = H_k$  can be expected if  $f$  is sufficiently random and  $\text{ord}(H) \leq p_t^2$ .

We have two methods to design  $f$  and to associate the ambiguous class  $S$  to  $H, f, j, k$ . Both methods will produce ambiguous classes  $S$  with  $S \neq 1$  whenever  $\text{ord}(H)$  is even. Experience must decide which of the methods is more efficient.

*Method 1.* For some  $q \in \mathbb{N}$  choose random integers  $a_i \in [p_t^2, 2p_t^2]$  for  $i = 1, \dots, q$ . Precompute  $F_i := H^{a_i}$ ,  $i = 1, \dots, q$ . For some random function  $g: \langle H \rangle \rightarrow \{1, \dots, q\}$  and recursively compute:

$$H_1 = H, \quad H_{i+1} = H_i F_{g(H_i)}.$$

Use the procedure *search* below in order to find some  $j < k$  with  $H_j = H_k$ . Then

$$H_k H_j^{-1} = H^T = 1 \quad \text{with } T = \sum_{i=j}^{k-1} g(H_i).$$

Most likely we will have  $k \leq 2p_t$  which implies  $T \leq 4p_t^3$ .

Now suppose that  $\text{ord}(\bar{H}) \equiv 2^e \pmod{2^{e+1}}$ . We can easily compute  $\bar{e}$  with  $T \equiv 2^{\bar{e}} \pmod{2^{\bar{e}+1}}$ . Then  $\bar{H}^{T2^{-\bar{e}}}$  has order  $2^e$  and yields an ambiguous class

$$S := \bar{H}^{T2^{e-\bar{e}-1}} \quad \text{with } S \neq 1 \text{ provided } e \geq 1.$$

*Comment.* A theoretical analysis of this method has been done by Sattler and Schnorr [16]. For small values of  $q$ , e.g.  $q = 2, 3, 4$ , the commutativity of the recursion steps increases the number of recursion steps as compared with a pure random recursion  $f: \langle H \rangle \rightarrow \langle H \rangle$ . By experience this slow-down is negligible as soon as  $q \geq 16$ . We have tested this recursion scheme in class groups (see Table 5, Appendix II) and in cyclic groups  $\langle H \rangle = \mathbb{Z}/n\mathbb{Z}$ , in particular with  $n$  prime. Method 1 even works well for nonrandom  $a_i$  like  $a_i = c^i$  with  $c$  fixed. The advantage of Method 1 over Method 2 is that it explicitly yields a multiple  $T$  of the order of  $H$ . Also, Method 1 only takes a single group operation (i.e. composition in the case of class groups) per recursion step.

*Method 2.* Choose a random function  $g: \langle H \rangle \rightarrow \{1, \dots, q\}$ , choose random values  $a_1, \dots, a_q \in [p_t, 2p_t^2]$  and precompute  $F_i := H^{a_i}$ ,  $i = 1, \dots, q$ .

*Recursion on  $H$ .* (We compute  $H_i = H^{c_i}$  and  $d_i$  with  $d_i \equiv c_i \pmod{2^{32}}$ .)

$$H_1 := H, \quad d_1 := 1;$$

for  $i = 1, 2, \dots$  till *search* finds some  $j < k$  with  $H_j = H_k$  do

$$(H_{i+1}, d_{i+1}) := \begin{cases} (H_i^3, 3d_i \pmod{2^{32}}) & \text{if } g(H_i) \leq q/2, \\ (H_i F_{g(H_i)}, d_i + a_{g(H_i)} \pmod{2^{32}}) & \text{otherwise.} \end{cases}$$

Use the procedure *search* below in order to find some  $j < k$  with  $H_j = H_k$ . Since  $H_i = H^{c_i}$  and  $H = \bar{H}^{2^{e*}}$ , it follows that  $\bar{H}^{2^{e*(c_j - c_k)}} = 1$ . We compute  $t$  such that  $d_j - d_k \equiv 2^t \pmod{2^{t+1}}$ . Almost surely  $t$  will be less than 32, and this implies  $c_j - c_k = 2^t m$  for some odd  $m$ . It remains to compute  $\bar{H}^m$ , since  $\bar{H}^{m/2^v}$  is ambiguous for some  $v \leq t$ . We do not compute  $m$  explicitly, but we retrace the above recursion on  $\bar{H}$ . In the following assume  $t \geq 1$ . If  $t = 0$ , then  $\bar{H}^m$  can easily be computed from the  $g(H_i)$ .

*Recursion on  $\bar{H}$ .* (We compute  $\bar{H}_i = \bar{H}^{\lfloor c_i/2^t \rfloor}$  and  $r_i = c_i \pmod{2^t}$  for  $t \geq 1$ .)

$$\bar{H}_1 := 1 \text{ (the unit class),} \quad r_1 := 1;$$

for  $i = 1, 2, \dots, k$  do

$$(\bar{H}_{i+1}, r_{i+1}) := \begin{cases} (\bar{H}_i^3 \bar{H}^s, 3r_i - s2^t) \text{ with } s = \lfloor 3r_i/2^t \rfloor & \text{if } g(H_i) \leq q/2, \\ (\bar{H}_i \bar{H}^s, r_i + a_{g(H_i)} - s2^t) & \\ \text{with } s = \lfloor (r_i + a_{g(H_i)})/2^t \rfloor & \text{if } g(H_i) > q/2. \end{cases}$$

It can easily be verified that  $\bar{H}_i = \bar{H}^{\lfloor c_i/2^t \rfloor}$ . Hence  $\bar{H}_j \bar{H}_k^{-1} = \bar{H}^{(c_j - c_k)/2^t} = \bar{H}^m$  with  $m$  odd. This yields

*Fact 6.* Let  $\text{ord}(\bar{H}) \equiv 2^e \pmod{2^{e+1}}$ ,  $e < 32$ , and  $H_j H_k^{-1} = 1$ , then  $\text{ord}(\bar{H}_j \bar{H}_k^{-1}) = 2^e$ .

Therefore  $S = (\bar{H}_j \bar{H}_k^{-1})^{2^{e-1}}$  is an ambiguous class with  $S \neq 1$  whenever  $e \neq 0$ .

*Comment.* We have tested Method 2 in class groups and in cyclic groups  $\langle H \rangle = \mathbb{Z}/n\mathbb{Z}$ , in particular with  $n$  prime. For random functions  $g: \langle H \rangle \rightarrow \{1, \dots, q\}$  we obtained average values of about  $\sqrt{\pi/2} \sqrt{n}$  for the smallest index  $k$  such that there exists some  $j < k$  with  $H_j = H_k$ , see Table 6, Appendix II. On the average, Method 2 takes 1.5 group operations (i.e. compositions in the case of the class group) per recursion step. A recursion step takes 2 compositions if  $H_{i+1} = H_i^3$  and 1 composition if  $H_{i+1} = H_i F_{g(H_i)}$ . By reducing the frequency of the  $H_{i+1} = H_i^3$ -steps the average number of compositions per recursion step can still be reduced. Method 2 also works well with nonrandom  $a_i$  like  $a_i = c^i$ ,  $i = 1, \dots, q$ , with  $c$  fixed. Because of the noncommutativity of the recursion steps, Method 2 works with a smaller number  $q$  of multipliers  $F_i = H^{a_i}$  than Method 1. We successfully applied Method 2 with  $q = 4$ .

The following pseudo-random function  $g: \langle H \rangle \rightarrow \{1, \dots, q\}$  works well for both methods (let  $(a, b, c)$  be the reduced form in  $H$ ):

$$g(H) = \lfloor (b^2 \bmod p) q/p \rfloor + 1$$

with  $p$  a prime,  $q < p < \sqrt{\Delta}$ ; see Tables 5, 6, Appendix II, for  $p = 2^{13} - 1$ .

*The search for  $H_k = H_j$  with  $j < k$ .* Let  $H_1 = H$ ,  $H_{i+1} = f(H_i)$ . We follow an idea of Brent [1] and do not store all the  $H_i$  but only a fixed number of them. When computing  $H_i$ , the stored classes

$$H_{\sigma(\nu)}, \quad \nu = 1, 2, \dots, 7,$$

for sufficiently large  $i$ , will be such that

$$\sigma(\nu) \approx \sigma(1) 1.1^\nu, \quad \nu = 1, \dots, 7,$$

$$\text{with } 1.1^7 \sigma(1) < i < 1.1^8 \sigma(1) \approx 2.14 \sigma(1).$$

The recursion for  $H_i$  is continued until some  $H_k = H_{\sigma(\nu)}$  has been found. The corresponding program looks like

*Search.*  $H_1 := H$ ,  $\sigma(\nu) := 1$  for  $\nu = 1, \dots, 7$ ; for  $i = 2, \dots$  do

$$\left[ \begin{array}{l} \text{compute } H_i \text{ from } H_{i-1} \\ \text{if } \exists \nu: H_{\sigma(\nu)} = H_i \text{ then } [j := \sigma(\nu), k := i \text{ stop}] \\ \text{if } 1.1^8 \sigma(1) < i + 1 \text{ then} \\ \quad \left[ \begin{array}{l} \text{store } H_i \text{ instead of } H_{\sigma(1)} \\ \sigma(\nu) := \begin{cases} \sigma(\nu + 1) & \text{for } \nu \neq 7 \\ i & \text{for } \nu = 7 \end{cases} \end{array} \right] \end{array} \right]$$



Let  $\lambda$  be the *period* and  $\mu$  the *length of the nonperiodic segment* of the sequence  $H_i$ , e.g.

$$H_\mu = H_{\mu+\lambda}, \quad H_i \neq H_l \quad \text{for } i < l < \mu + \lambda.$$

*Fact 7.* The procedure *search* finds some  $j < k$  with  $H_k = H_j$  within  $\leq 1.1m + \lambda$  recursion steps,  $m := \max(\lambda, \mu)$ .

*Proof.* Since  $\sigma$  increases by the factor 1.1,  $\sigma$  will take some value  $\sigma(\nu)$  with  $m \leq \sigma(\nu) \leq 1.1m$ . Hence the for-loop stops, at the latest, with

$$k = \sigma(\nu) + \lambda \leq 1.1m + \lambda, \quad j = \sigma(\nu),$$

and finds the equality  $H_k = H_j$ .  $\square$

Under the assumption that each of the  $\text{ord}(H)^{\text{ord}(H)}$  functions  $f: \langle H \rangle \rightarrow \langle H \rangle$  has probability  $\text{ord}(H)^{-\text{ord}(H)}$ , the stochastic behavior of  $\mu, \lambda$  have been well analyzed (see Knuth [7, Exercise 3.1.12]).

The expected values of  $\mu$  and  $\lambda$  are

$$1 + E(\mu) = E(\lambda) \approx \sqrt{\frac{\pi \text{ord}(H)}{8}} + 1/3,$$

$$E(\mu + \lambda) \approx 1.25\sqrt{\text{ord}(H)} - 1/3,$$

$$\text{Prob}\left[\mu + \lambda \leq \sqrt{\frac{\pi}{2} \text{ord}(H)}\right] \approx e^{-\pi/4} \approx 0.46.$$

We conclude from Fact 7 that the number of recursion steps in *search* will be about

$$1.1 (E(\mu) + E(\lambda)) \approx 1.32\sqrt{\text{ord}(H)},$$

provided that  $f: \langle H \rangle \rightarrow \langle H \rangle$  is sufficiently random.

If in Stage 2 we compute the  $H_i$  for  $i \leq 1.32 p_t$ , then most likely some relation  $H_j = H_k$ ,  $j < k$ , will be found, provided  $\text{ord}(H) \leq p_t^2$ . It remains to analyze the chance that  $\text{ord}(H) \leq p_t^2$ . For each prime  $p$ ,  $p_t < p \leq p_t^2$ , we assume that the frequency of class numbers  $h(-m)$ ,  $m \leq n$ , which are divisible by  $p$  is  $\geq p^{-1}$ , and we assume that  $h(-m)/p$  factors like random integers of size  $\sqrt{n}/p$ . By retracing the proof of Corollary 4, we conclude from the assumptions (2.1), (2.2):

$$\begin{aligned} & \text{For all } r, n, t \text{ with } n \geq n_0, p_t = n^{1/2r}, r \leq \sqrt{\ln n / \ln \ln n}: \\ & \text{and for all primes } p = n^{1/2s} < p_t^2 \\ (3.4) \quad & \# \left\{ m \leq n: h(-m) \mid p \prod_{i=1}^t p_i^{e_i} \right\} / (0.5 n) \\ & \geq 0.83 p^{-1} (r - r/s)^{-(r-r/s)}. \end{aligned}$$

Summing over all  $p$ ,  $n^{r^{-1}/(1+\epsilon)} < p < n^{1/r}$ , this yields

$$\begin{aligned} & \# \left\{ m \leq n: h(-m) \mid p \prod_{i=1}^t p_i^{e_i} \text{ with } p < p_t^2 \right\} / (0.5 n) \\ & \geq 0.83 \sum_{n^{r^{-1}/(1+\epsilon)} < p < n^{1/r}} p^{-1} (r - 2/(1+\epsilon))^{-(r-2/(1+\epsilon))} \\ & \quad \text{(using Theorem 4.27 in Hardy and Wright [6] there follows)} \\ & \geq 0.83 \ln(1+\epsilon) (r - 2/(1+\epsilon))^{-(r-2/(1+\epsilon))}. \end{aligned}$$

**Conclusion.** Assume (2.1), (2.2) and that for every discriminant  $m \leq n$ ,  $H_0$  in  $G(-m)$  is chosen at random. Then  $\forall \epsilon > 0: \exists c_\epsilon > 0: \forall n \geq n_0$  and all  $p_t = n^{1/2r}$ ,  $r \leq \sqrt{\ln n / \ln \ln n}$ : Stages 1 and 2 with  $O(p_t)$  compositions, factor at least  $c_\epsilon(r-2+\epsilon)^{-(r-2+\epsilon)}n$  discriminants  $\leq n$ .

If one assumes that very large class numbers  $h(-n)$  factor like even integers of size  $\sqrt{n}$ , then we can compare the efficiency of Stages 1 and 2 by Odlyzko's Table 4, Appendix II. The table indicates that for class numbers  $h(-m) \approx 10^l$ ,  $l = 15, 20, 25, 30$ , the success frequency of Stages 1 and 2 is at least  $r^{-r}$  and is at most  $er^2$  times the success frequency of Stage 1. Note that  $(r-2)^{-(r-2)}/r^{-r}$  approaches  $er^2$  for large  $r$ .

**Remark.** There is a well-known deterministic method for doing Stage 2 within  $\sqrt{2}p_t$  compositions and with  $O(p_t)$  storage. The method is explained in Shanks [17, p. 419] in terms of "baby" and "giant" steps. In our situation we can even speed this method by a constant factor if we exploit the fact that  $\text{ord}(H)$  will most likely have no prime divisor  $\leq p_t$ .

**4. The Main Algorithm.** The new algorithm can be used for factoring any composite integer  $n$ . We apply Stage 1 to multiples  $ns$  of  $n$  such that  $-ns$  is a discriminant. Here we exploit the observation that class numbers  $h(-ns)$  of fundamental discriminants  $-ns$  are uncorrelated for distinct values of  $s$ . The nonfundamental discriminants  $-ns$  should be discarded as far as possible. The discriminant  $\Delta$  is *fundamental* if

$$\neg \exists w \in \mathbf{N}, w \neq 1: \Delta/w^2 \text{ is a discriminant.}$$

In fact, the class number formula (see Dirichlet [8]),

$$h(-m) = \frac{\sqrt{m}}{\pi} \prod_{p \text{ prime}} \left( 1 - \frac{1}{p} \left( \frac{-m}{p} \right) \right)^{-1} \quad \text{for } m < -4,$$

implies for  $\gcd(w, m) = 1$  and  $w$  square free:

$$h(-mw^2)/h(-m) = w \prod_{p|w} \left( 1 - \frac{1}{p} \left( \frac{-m}{p} \right) \right) = \prod_{p|w} \left( p - \left( \frac{-m}{p} \right) \right).$$

Hence for small  $w$ ,  $h(-m)$  and  $h(-mw^2)$  have the same large prime divisors, provided  $\gcd(m, w) = 1$ .

**Main Algorithm.** Let  $n$  be the number to be factored and  $p_1 = 2, p_2 = 3, \dots, p_t$  the first  $t$  primes,  $p_t = n^{1/2r}$  (the appropriate choice of  $t, r$  will be determined by the subsequent analysis).

1.  $s := 0$
2. take the next  $s$  with  $\gcd(n, s) = 1$ ,  $-ns \equiv 0, 1 \pmod{4}$  and  $\neg[\exists w \in \mathbf{N}: w^2 | s, w \neq 1, -ns/w^2 \equiv 0, 1 \pmod{4}]$
3. run Stage 1 on  $ns$ , which takes  $O(p_t)$  compositions. If Stage 1 yields an ambiguous class  $S$  then go to 4, otherwise return to 2 and take the next  $s$
4. if  $S$  yields a factorization of  $n$  then stop, otherwise go to 5
5. return to 3 and repeat Stage 1 on  $ns$  with independently chosen classes  $H_0 \in G(-ns)$  until some factorization of  $n$  has been found. In order to prevent that merely useless ambiguous classes are generated, continue to build up the 2-Sylow group  $S_2(-ns)$  of  $G(-ns)$ . Use Stage 1 to generate classes in

$S_2(-ns)$ . Apply the recursion step of the method below whenever a new  $A \in S_2(-ns)$  has been found.

The 2-Sylow group  $S_2(-ns)$  is a direct product of cyclic groups of order  $2^{m_i}$ ,  $m_i > 0$ :  $S_2(-ns) \cong \bigoplus_{i=1}^{\lambda} \mathbf{Z}/2^{m_i}\mathbf{Z}$ . Let  $\lambda$  be the number of cyclic components, then  $S_2(-ns)$  has  $2^\lambda$  ambiguous classes. Let  $d$  be the number of odd prime factors of  $ns$ . Then by Theorem III, Appendix I,  $\lambda$  is  $d-1$ ,  $d$  or  $d+1$  depending on the maximal power of 2 which divides  $ns$ . The ambiguous classes that do not yield a factorization of  $n$  form a subgroup  $S'_2(-ns)$  of  $S_2(-ns)$ . Let  $d_n, d_s$  be the numbers of distinct odd primes of  $n$  and  $s$ . Since  $\gcd(n, s) = 1$ , we have  $d = d_n + d_s$ . It follows immediately from Theorem III, Appendix I, that the number  $\lambda'$  of cyclic components of  $S'_2(-ns)$  is  $\leq \lambda - d_n \leq d_s + 1$ .

*Constructing  $S_2(-ns)$  till a factorization of  $n$  is found.* Given a procedure that generates elements of  $S_2(-ns)$  (this will be done by Stage 1) we recursively construct subsets  $\{A_1, \dots, A_{\bar{\lambda}}\} \subset S_2(-ns)$ ,  $\bar{\lambda} \leq \lambda'$  such that

$$(4.1) \quad |\langle A_1, \dots, A_{\bar{\lambda}} \rangle| = \prod_{i=1}^{\bar{\lambda}} \text{ord}(A_i).$$

Let  $\text{ord}(A_i) = 2^{\bar{m}_i}$ ,  $B_i = A_i^{2^{\bar{m}_i-1}}$ ,  $\langle A_1, \dots, A_{\bar{\lambda}} \rangle = \bar{S}_2(-ns)$ . Then  $\bar{S}_2(-ns) \cong \bigoplus_{i=1}^{\bar{\lambda}} \mathbf{Z}/2^{\bar{m}_i}\mathbf{Z}$  and  $B_1, \dots, B_{\bar{\lambda}}$  generate the subgroups of ambiguous classes of  $\bar{S}_2(-ns)$ . After each recursion step either a factorization of  $n$  has been found or the new group  $\bar{S}_2(-ns)$  will be the subgroup of  $S_2(-ns)$  which is generated by the previous  $\bar{S}_2(-ns)$  and the element  $A \in S_2(-ns)$  obtained in step 1.

(4.2) *Algorithm for  $S_2(-ns)$ .*

0.  $\bar{\lambda} := 0, A_1 := 1$  (= the unit class)
1. generate another  $A \in S_2(-ns)$ ,  $A \neq 1$
2. compute  $A, A^2, \dots, A^{2^{m-1}} \neq 1, A^{2^m} = 1$  and put  $B := A^{2^{m-1}}$
3. if  $B$  yields a factorization of  $n$  then stop
4. test whether  $B \in \langle B_1, \dots, B_{\bar{\lambda}} \rangle$ ; if  $B \notin \langle B_1, \dots, B_{\bar{\lambda}} \rangle$  then go to 5 else compute  $J \subset \{1, \dots, \bar{\lambda}\}$  with  $B = \prod_{j \in J} B_j$  and go to 6
5.  $A_{\bar{\lambda}+1} := A, B_{\bar{\lambda}+1} := B, \bar{\lambda} := \bar{\lambda} + 1$  return to 1
6. If  $\exists j \in J: m_j < m$  then select  $j \in J$  with  $m_j$  minimal and interchange  $A$  with  $A_j$ ,  $B$  with  $B_j$ , and  $m$  with  $m_j$
7. (we have  $A^{2^{m-1}} = \prod_{j \in J} A_j^{2^{m_j-1}}$ ,  $m \leq m_j$  for  $j \in J$ ). Put  $A := A \cdot \prod_{j \in J} A_j^{2^{m_j-m}}$ ; if  $A = 1$  go to 1 else go to 2 (the new  $m$  to be computed in 2. will be smaller than the present  $m$  since  $A^{2^{m-1}} = 1$  holds for the new  $A$ )

*Run Time Analysis of the Main Algorithm.* We separately bound

1. the number  $T(n)$  of bit operations to be done till some  $s$  has been reached with

$$h(-ns) \mid \prod_{i=1}^t p_i^{e_i}, \quad e_i = \max\{v: p_i^v \leq p_i^2\}.$$

2. The number  $\bar{T}(n)$  of bit operations for building up the 2-Sylow group  $S_2(-ns)$  of  $G(-ns)$  till a factorization of  $n$  is found.

1.  $T(n)$ . We will assume that Corollary 4 extends to multiples of  $n$ :

$$(4.3) \quad \begin{aligned} &\exists c, n_0 > 0: \forall m: \forall n \geq n_0: \forall p_i = (nm)^{1/2r} \text{ with } r \leq \sqrt{\ln n / \ln \ln n}: \\ &\# \left\{ ns: s \leq m \wedge h(-ns) \mid \prod_{i=1}^t p_i^{e_i} \right\} / (0.5m) \geq cr^{-r}. \end{aligned}$$

Our experimental data in fact confirm the lower bound  $r^{-r}$ . The assumption (4.3) implies

$$\begin{aligned} \forall n \geq n_0: \forall p_i &= (n3r^r/c)^{1/2r}, r \leq \sqrt{\ln n / \ln \ln n} \\ \exists s &\leq 3r^r/c: h(-ns) \mid \prod_{i=1}^t p_i^{e_i}. \end{aligned}$$

Since Stage 1 takes  $O(p_i)$  compositions, we have

$$T(n) = O(p_i r^r (\ln n)^2) = O(n^{1/2r} r^{r+1/2} (\ln n)^2).$$

Here  $O(\ln n)^2$  takes into account the costs for the arithmetic. We choose  $r = \sqrt{\ln n / \ln \ln n}$ ,  $p_i \approx (n3r^r/c)^{1/2r} = O(n^{1/2r} \sqrt{r})$ . Then all together (4.3) implies

$$T(n) = o(\exp \sqrt{\ln n \ln \ln n}).$$

2.  $\bar{T}(n)$ . In order to factor  $n$  we need only to find at most  $d_s + 2$  cyclic components of  $S_2(-ns)$ . If the passes through Stage 1 generate independent elements of  $S_2(-ns)$  then  $k$  passes of Stage 1 with probability  $\geq 1 - 2^{-k}$  detect a new cyclic component of  $S_2(-ns)$ . Hence almost surely we need at most  $O(d_s)$  passes through Stage 1, and each pass takes  $O(p_i)$  compositions. The number of steps for updating the information on  $S_2(-ns)$  can be bounded as  $O(s)$ : the most costly operation in Algorithm (4.2) is to check whether  $B \in \langle B_1, \dots, B_{\bar{\lambda}} \rangle$  (step 4). Since  $\bar{\lambda} \leq \lambda' \leq d_s + 1$  this can be done in a crude way by comparing  $B$  with each of the  $2^{\bar{\lambda}} \leq 2^{d_s+1} = O(s)$  elements of  $\langle B_1, \dots, B_{\bar{\lambda}} \rangle$ . This takes  $O(s)$  steps and is sufficient for our purposes. We obtain

$$\bar{T}(n) = O(d_s(p_i + s)(\ln n)^2) = O(\log s(n^{1/2r} \sqrt{r} + s)(\ln n)^2)$$

with  $s \leq r^r$ ,  $r \leq \sqrt{\ln n / \ln \ln n}$ . Here again  $O(\ln n)^2$  bounds the cost for the arithmetic. It follows immediately that  $\bar{T}(n) = o(T(n))$ .

*Conclusion.* If (4.3) holds, then the Main Algorithm, using only Stage 1, takes  $o(\exp \sqrt{\ln n \ln \ln n})$  bit operations to factor arbitrary, composite integers  $n$ .

If we also apply Stage 2, then  $s$  will be bounded as  $O((r-2)^{(r-2)})$ , and this will save a time factor of about  $r^2 = \ln n / \ln \ln n$ .

**5. Some Computational Experience.** The new factoring method has been programmed in Fortran on a DEC-1091 at Frankfurt University. The core of the algorithm is a subroutine for composition of quadratic forms written in machine language and based on the improved composition method proposed by Seysen [20]. The arithmetic operations and the gcd-calculations have been programmed for two-word integers, i.e. for integers  $\leq 2^{70}$ . This means that the program can factor integers  $\leq 2^{130} \approx 10^{39}$  using multipliers  $\leq 2^{10}$ . Stage 1 uses the exponents  $e'_i := \max\{v: p_i^v \leq p_i\}$ . Hence the number of compositions per multiplier for Stage 1 is about  $2.2 p_i$ . Both methods for Stage 2 have been tested and they are comparable in efficiency.

TABLE 0

	$n \approx 10^{30}$	$n \approx 10^{34}$	$n \approx 10^{38}$
$p_t$	4093	8191	16 381
$t = \#$ of primes	563	1027	1899
# compositions per multiplier in Stage 1	$10^4$	$2 \cdot 10^4$	$4 \cdot 10^4$
in Stage 2	$3 \cdot 10^3$	$3.7 \cdot 10^3$	$3 \cdot 10^4$
average number of multipliers	14.4	18	31
median of the numbers of multipliers	8	9	23
# of integers factored	50	20	20
seconds per composition	$1.4 \cdot 10^{-3}$	$1.5 \cdot 10^{-3}$	$1.57 \cdot 10^{-3}$
average time for factoring	4.4 min	10.7 min	57 min
median of the factoring time	2.4 min	5.3 min	42 min

The integers  $n$  which have been factored for Table 0 are products of two distinct primes  $p_1, p_2$  of nearly the same size. It turns out that the median of the factoring time is considerably smaller than the average factoring time. This is due to a small fraction of integers  $n$  which take extremely many multipliers. On the other hand there is a considerable fraction of integers which only take very few multipliers. For instance the seventh Fermat number  $F_7 = 2^{2^7} + 1 \approx 3.4 \cdot 10^{38}$  only took 7 multipliers and was factored in about 7 minutes. Here we used  $p_t = 16381$  but we run Stage 2 for only 7500 compositions, hence each multiplier took about 1 minute. The multiplier 15 has been successful.

We observed that the factoring method is somewhat faster for integers with more than 2 prime divisors. By our observation class numbers of discriminants with many prime divisors tend to have fewer large prime divisors compared with class numbers of discriminants which are prime or products of two primes. For instance, for  $n \approx 10^{30}$ ,  $n$  a product of 5 primes  $p_i \approx 10^6$ , the algorithm on the average only took 8.7 multipliers. The median of the number of multipliers has been 5, compared with 14.4 and 8 in Table 0. We have factored a sample of 200 of these integers  $n$ .

**Appendix I on Quadratic Forms.** We report classical theorems and algorithms on quadratic forms, see Gauss [5], Mathews [10]. A quadratic form  $ax^2 + bxy + cy^2$  with  $a, b, c \in \mathbb{Z}$  is denoted as  $(a, b, c)$ . Its discriminant is  $\Delta = b^2 - 4ac$ .  $(a, b, c)$  is *positive* if  $a > 0$ , *primitive* if  $\gcd(a, b, c) = 1$ . Two forms  $(a, b, c)$ ,  $(\bar{a}, \bar{b}, \bar{c})$  are equivalent if there exists a linear transformation with integer coefficients and determinant 1 transforming the one form into the other, i.e.

$$T' \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix} T = \begin{pmatrix} \bar{a} & \bar{b}/2 \\ \bar{b}/2 & \bar{c} \end{pmatrix}$$

for some integer matrix  $T$  with  $\det T = 1$ . Let  $[(a, b, c)]$  be the class represented by  $(a, b, c)$ . For negative discriminants we always restrict to positive forms.

Two classes  $[(a, b, c)]$ ,  $[(\bar{a}, \bar{b}, \bar{c})]$  yield a new class  $[(A, B, C)]$  by *composition* as follows (for explanation see Lenstra [9]):

$$\begin{aligned} d &:= \gcd(a, \bar{a}, (b + \bar{b})/2) \\ \text{Let } \alpha, \beta, \gamma &\in \mathbb{Z} \text{ be such that } \alpha a + \beta \bar{a} + \frac{\gamma}{2}(b + \bar{b}) = d. \\ (5.1) \quad A &= a\bar{a}/d^2 \\ B &= \bar{b} + \frac{2\bar{a}}{d} \left[ \left( \beta \frac{b - \bar{b}}{2} - \gamma \bar{c} \right) \bmod \frac{a}{d} \right] \\ C &= (-\Delta + B^2)/(4A). \end{aligned}$$

$[(A, B, C)]$  does not depend on the particular choice for  $\alpha, \beta, \gamma, B$ , and  $C$ .  $(A, B, C)$  will be primitive, if  $(a, b, c)$  and  $(\bar{a}, \bar{b}, \bar{c})$  are primitive.

**THEOREM I.** *The equivalence classes of primitive quadratic forms with discriminant  $\Delta$  form an Abelian group  $G(\Delta)$  under composition. Its order  $h(\Delta)$  is the **class number**.*

The unit class 1 in  $G(\Delta)$  is represented by the form

$$\begin{aligned} (1, 0, -\Delta/4) & \quad \text{if } \Delta \equiv 0 \pmod{4}, \\ (1, 1, (1 - \Delta)/4) & \quad \text{if } \Delta \equiv 1 \pmod{4}. \end{aligned}$$

The inverse of  $[(a, b, c)]$  is  $[(a, -b, c)]$ .

A class  $H \in G(\Delta)$  is *ambiguous* if  $H^2 = 1$ . The following assertions are equivalent:

- (1)  $H$  is ambiguous,
- (2) every form  $(a, b, c)$  in  $H$  is equivalent to  $(a, -b, c)$ ,
- (3) 
$$T \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix} T' = \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix}$$
 for some integer matrix  $T$  with  $\det T = -1$ ,
- (4) there is a form  $(a, b, c)$  in  $H$  with  $a|b$ .

For negative discriminants  $\Delta$ , classes in  $G(\Delta)$  correspond to reduced forms.  $(a, b, c)$  is *reduced* if (1)  $|b| \leq a \leq c$  and (2)  $b \geq 0$  if  $|b| = a$  or  $a = c$  (i.e. if  $[(a, b, c)]$  is ambiguous).

**THEOREM II (GAUSS, [5, Art. 172]).** *In every equivalence class with negative discriminant there is exactly one reduced form.*

Gauss also gave a gcd-like reduction algorithm which transforms a given form  $(a, b, c)$  into an equivalent reduced form:

(5.2) *reduction process for  $(a, b, c)$*

1. find  $\nu \in \mathbb{Z}$ :  $-|a| < b + 2\nu a \leq |a|$
2.  $b := b + 2\nu a, c := (b^2 - \Delta)/(4a)$
3. if  $(a, b, c)$  is not reduced then replace  $(a, b, c)$  by  $(c, -b, a)$  and go to 1.

The reduced forms of ambiguous classes with  $\Delta < 0$  are of either of the following types:

- (i)  $b = 0$ ,      (ii)  $a = b$ ,      (iii)  $a = c$ .

These forms are called *ambiguous*. Every ambiguous form corresponds to a factorization of the discriminant as

$$(i) \quad \Delta = -4ac, \quad (ii) \quad \Delta = b(b - 4c), \quad (iii) \quad \Delta = (b - 2a)(b + 2a).$$

In order to describe this correspondence more precisely, let

$$F(n) = \{(n_1, n_2) \in \mathbf{N}^2: n = n_1 n_2, \gcd(n_1, n_2) = 1\}$$

be the set of relatively prime, ordered factor pairs of  $n \in \mathbf{N}$ . We have  $\#F(n) = 2^d$ , where  $d$  is the number of distinct prime divisors of  $n$ .

Let  $\Delta \equiv 2^e \pmod{2^{e+1}}$ ,  $\Delta < 0$ , then the set  $A(\Delta)$  of ambiguous forms with discriminant  $\Delta$  is either in 2-1 or in 1-1 or in 1-2-correspondence with  $F(-\Delta/2^e)$ .

**THEOREM III.** *Let the discriminant  $\Delta < 0$  have  $d$  odd prime divisors. Then the set  $A(\Delta)$  of ambiguous (reduced, positive, primitive) forms with discriminant  $\Delta = -2^e n$ ,  $n$  odd, is obtained from the  $(n_1, n_2) \in F(n)$  with  $n_1 \leq n_2$  as follows:*

case $e = 0$ (i.e. $\Delta \equiv 1 \pmod{4}$ ):	$\#A(\Delta) = \frac{1}{2} \#F(n) = 2^{d-1}$
(ii) $(n_1, n_1, (n_1 + n_2)/4)$	if $3n_1 < n_2$
(iii) $((n_1 + n_2)/4, (n_2 - n_1)/2, (n_1 + n_2)/4)$	if $3n_1 \geq n_2$
case $e = 2, n \equiv 1 \pmod{4}, n \neq 1$ :	$\#A(\Delta) = \#F(n) = 2^d$
(i) $(n_1, 0, n_2)$	
(ii) $(2n_1, 2n_1, (n_1 + n_2)/2)$	if $3n_1 \leq n_2$
(iii) $((n_1 + n_2)/2, n_2 - n_1, (n_1 + n_2)/2)$	if $3n_1 > n_2$
case $e = 2, n \equiv -1 \pmod{4}$ :	$\#A(\Delta) = \frac{1}{2} \#F(n) = 2^{d-1}$
(i) $(n_1, 0, n_2)$	
case $e = 3, 4$ :	$\#A(\Delta) = \#F(n) = 2^d$
(i) $(\min(n_1 2^{e-2}, n_2), 0, \max(n_1 2^{e-2}, n_2))$ $(\min(n_2 2^{e-2}, n_1), 0, \max(n_2 2^{e-2}, n_1))$	
case $e \geq 5$ :	$\#A(\Delta) = 2 \#F(n) = 2^{d+1}$
(i) $(\min(n_1 2^{e-2}, n_2), 0, \max(n_1 2^{e-2}, n_2))$ $(\min(n_2 2^{e-2}, n_1), 0, \max(n_2 2^{e-2}, n_1))$	
(ii) $(4, 4, 1 + 2^{e-4} n_1 n_2)$	
(iii) $(2^{e-4} n_2 + n_1, 2^{e-3} n_2 - 2n_1, 2^{e-4} n_2 + n_1)$	if $3n_1 \geq n_2 2^{e-4}$
$(2^{e-4} n_1 + n_2, 2^{e-3} n_1 - 2n_2, 2^{e-4} n_1 + n_2)$	if $3n_1 < n_2 2^{e-4}$

We have listed pairwise inequivalent forms corresponding to distinct positive ambiguous classes. They have been arranged according to their types (i), (ii), (iii) as introduced above.

Theorem III can easily be obtained from Gauss [5, Art. 257–259]. Observe that our classes with discriminant  $\Delta \equiv 0 \pmod{4}$  ( $\Delta \equiv 1 \pmod{4}$ , resp.) correspond to primitive Gauss classes with determinant  $D = \Delta/4$  (improper primitive Gauss classes with determinant  $D = \Delta$ , resp.). The number of ambiguous classes has also been listed in Cassels [2, p. 342].

*The Efficiency of Composition.* An efficient composition algorithm is the main requirement for a satisfactory implementation of our factoring algorithm. All

calculations in  $G(\Delta)$  are done with reduced forms. Composition consists of two parts:

1. evaluation of (5.1):  $(a, b, c), (\bar{a}, \bar{b}, \bar{c}) \rightarrow (A, B, C)$  (this amounts to an extended gcd-calculation on integers of size  $O(\sqrt{|\Delta|})$ ),
2. reduction of  $(A, B, C)$ .

If the reduction is done as in (5.2) this corresponds to an extended gcd-calculation on integers of size  $O(|\Delta|)$ . However, M. Seysen [20] found a faster reduction algorithm for this particular situation. Reducing  $(A, B, C)$  by this algorithm corresponds to only half an extended gcd-calculation on integers of size  $O(\sqrt{|\Delta|})$ .

**Appendix II: Statistical Tables.** Table 1 shows the distribution of class numbers  $h(-m)$  without large prime divisors for discriminants  $-m$  in the interval  $I = [-472\ 650\ 003, -472\ 600\ 000]$ . There are 25 002 discriminants, the minimal, maximal, and average class numbers are 1518, 47 452, and 9 469.77. We put

$$\bar{e}_i(m) := \max\{v: p_i^v | h(-m)\}.$$

For every prime  $p_i = 2, 3, \dots, 89$  we record the percentage of those discriminants  $-m \in I$  satisfying the following conditions:

- |          |   |
|----------|---|
| column 1 | $h(-m)$ is free of primes $> p_t$ , i.e. $h(-m) = \prod_{i=1}^t p_i^{\bar{e}_i(m)}$ ,                 |
| column 2 | for all $i \geq 2$ : $\bar{e}_i(m) \leq e_i := \max\{v: p_i^v \leq p_t^2\}$ ,                         |
| column 3 | for all $i \geq 2$ : $\bar{e}_i(m) \leq e'_i := \max\{v: p_i^v \leq p_t\}$ ,                          |
| column 4 | $h(-m)   \prod_{i=1}^t p_i^{\bar{e}_i(m)} \cdot q$ for some $q \leq p_t^2$ , $q$ prime,               |
| column 5 | $\bar{h}(-m)   \prod_{i=2}^t p_i^{e_i} q \cdot 2^{\bar{e}_1(m)}$ for some $q \leq p_t^2$ , $q$ prime, |
| column 6 | $h(-m)   \prod_{i=2}^t p_i^{e'_i} q \cdot 2^{\bar{e}_1(m)}$ for some $q \leq p_t^2$ , $q$ prime.      |

Moreover we note in

- |          |                           |
|----------|---------------------------|
| column 7 | $r = \ln n / (2 \ln p_t)$ |
| column 8 | $10^2 \cdot r^{-r}$ .     |

Observe that the entries in columns 1–3 of Table 1 are always greater than  $10^2 r^{-r}$ , which confirms Corollary 4. For  $r \leq \sqrt{\ln n / \ln \ln n}$  (i.e.  $r \geq 2.58$ ,  $p_t \geq 53$ ) the entries in column 3 are only slightly smaller than those in columns 1, 2. This suggests that Stage 1 should be done with the smaller exponents  $e'_i$  instead of the  $e_i$ .

Table 2 has the same meaning as Table 1 but is restricted to fundamental discriminants in the same interval  $I$ . Minimal, maximal, and average class numbers are 1518, 47 425, and 10 033.9. There are 15195 fundamental discriminants in  $I$ .



TABLE 1

$p_i$	col 1	col 2	col 3	col 4	col 5	col 6	col 7	col 8
2	0.18	0.18	0.18	0.44	0.44	0.44	14.408	.000
3	2.08	0.82	0.44	5.39	2.82	1.46	9.090	.000
5	5.70	2.70	1.00	18.32	12.55	6.46	6.205	.001
7	10.21	7.55	2.00	32.52	28.95	14.45	5.132	.023
11	14.43	12.60	6.12	49.58	47.47	35.57	4.165	.263
13	18.87	17.64	8.92	58.26	56.95	43.27	3.894	.503
17	22.47	21.75	11.51	68.90	68.16	53.20	3.525	1.179
19	26.04	25.55	14.12	73.57	73.08	57.57	3.392	1.588
23	29.28	28.79	16.77	79.60	79.11	63.43	3.185	2.497
29	32.10	31.93	24.73	85.48	85.30	77.38	2.966	3.978
31	34.70	34.52	27.13	87.27	87.09	79.09	2.908	4.484
37	37.00	36.87	29.32	90.69	90.56	82.46	2.766	5.999
41	39.24	39.10	31.46	92.49	92.36	84.25	2.689	6.992
43	41.30	41.16	33.43	93.30	93.16	85.03	2.655	7.480
47	43.18	43.12	35.25	94.43	94.38	86.16	2.594	8.438
53	44.92	44.88	38.93	95.71	95.67	89.58	2.515	9.825
59	46.48	46.45	40.46	96.66	96.63	90.52	2.449	11.147
61	48.02	48.00	41.98	97.02	96.99	90.88	2.429	11.574
67	49.44	49.41	43.38	97.64	97.61	91.50	2.375	12.813
71	50.65	50.63	44.59	98.02	98.00	91.88	2.343	13.606
73	51.58	51.80	45.74	98.13	98.11	91.98	2.328	13.993
79	53.06	53.04	46.97	98.56	98.54	92.41	2.286	15.117
83	54.08	54.07	49.59	98.77	98.76	94.27	2.260	15.837
89	55.12	55.11	50.63	99.00	98.99	94.50	2.225	16.875

TABLE 2

$p_i$	col 1	col 2	col 3	col 4	col 5	col 6	col 7	col 8
2	0.17	0.17	0.17	0.31	0.31	0.31	14.408	.000
3	1.26	0.51	0.31	3.17	1.69	1.00	9.090	.000
5	3.69	1.65	0.67	12.93	8.90	4.62	6.205	.001
7	6.96	4.98	1.36	25.44	22.53	11.79	5.132	.023
11	10.22	8.68	3.96	41.54	39.70	29.83	4.165	.263
13	14.02	13.04	6.19	50.71	49.63	37.90	3.894	.503
17	17.09	16.46	8.37	61.77	61.11	48.16	3.525	1.179
19	20.14	19.69	10.57	67.04	66.57	53.10	3.392	1.588
23	23.09	22.64	12.89	74.04	73.58	59.89	3.185	2.497
29	25.73	25.55	19.23	81.03	80.85	73.64	2.966	3.978
31	28.21	28.03	21.47	83.22	83.04	75.73	2.908	4.484
37	30.42	30.29	23.57	87.39	87.27	79.86	2.766	5.999
41	32.66	32.53	25.69	89.78	89.65	82.22	2.689	6.992
43	34.68	34.56	27.61	90.82	90.69	83.24	2.655	7.480
47	36.59	36.54	29.44	92.28	92.23	84.69	2.594	8.438
53	38.37	38.33	33.02	94.01	93.97	88.48	2.515	9.825
59	39.94	39.91	34.56	95.26	95.23	89.73	2.449	11.147
61	41.46	41.43	36.06	95.75	95.72	90.22	2.429	11.574
67	42.84	42.81	37.42	96.63	96.60	91.10	2.375	12.813
71	44.02	44.01	38.59	97.16	97.14	91.62	2.343	13.606
73	45.08	45.07	39.63	97.29	97.28	91.75	2.328	13.993
79	46.45	46.44	40.97	97.90	97.89	92.36	2.286	15.117
83	47.60	47.59	43.35	98.17	98.16	93.89	2.260	15.837
89	48.69	48.69	44.44	98.50	98.49	94.22	2.225	16.875

TABLE 3

$\nu$	—	2	3	4	5	6	7	8	9	10
all discr.		93.64	51.25	82.53	24.70	48.18	16.92	63.44	20.30	23.18
fund. discr.		91.83	42.49	78.51	23.94	38.97	16.41	56.90	15.41	21.97
$\nu$	11	12	13	14	15	16	17	18	19	20
all d.	9.83	42.42	8.33	15.98	12.58	43.54	6.07	19.09	5.36	20.50
fund.	9.35	33.15	8.42	15.16	9.95	36.62	5.96	14.11	5.22	18.85
$\nu$	21	22	23	24	25	26	27	28	29	30
all d.	8.77	9.27	4.45	32.64	5.03	7.77	7.15	14.13	3.62	11.85
fund.	7.13	8.65	4.48	23.88	4.99	7.67	5.27	13.02	3.67	9.12
$\nu$	31	32	33	34	35	36	37	38	39	40
all d.	3.13	27.09	5.10	5.68	4.15	16.90	4.66	5.01	4.26	15.73
fund.	3.17	21.38	3.91	5.44	4.00	12.04	2.71	4.76	3.61	13.55
$\nu$	41	42	43	44	45	46	47	48	49	50
all d.	2.53	8.30	2.27	8.21	5.10	4.19	2.04	22.33	2.42	4.70
fund.	2.64	6.58	2.34	7.40	3.56	4.15	2.13	15.42	2.29	4.57
$\nu$	51	52	53	54	55	56	57	58	59	60
all d.	3.12	6.89	1.86	6.74	2.35	10.74	2.75	3.40	1.66	10.43
fund.	2.54	6.58	1.95	4.81	2.22	9.44	2.26	3.39	1.70	7.71
$\nu$	61	62	63	64	65	66	67	68	69	70
all d.	1.65	2.95	3.46	15.67	2.03	4.84	1.49	4.96	2.16	3.88
fund.	1.67	2.95	2.59	11.66	2.03	3.62	1.49	4.57	1.75	3.65
$\nu$	71	72	73	74	75	76	77	78	79	80
all d.	1.27	13.10	1.18	2.53	2.53	4.39	1.62	4.02	1.28	10.87
fund.	1.27	8.73	1.08	2.53	2.09	4.00	1.53	3.32	1.42	8.65
$\nu$	81	82	83	84	85	86	87	88	89	90
all d.	2.45	2.38	1.03	7.33	1.36	2.11	1.82	6.38	1.06	4.81
fund.	1.84	2.44	1.18	5.67	1.30	2.12	1.57	5.36	1.12	3.25
$\nu$	91	92	93	94	95	96	97	98	99	100
all d.	1.38	3.78	1.64	1.89	1.26	13.85	1.00	2.30	1.99	4.19
fund.	1.35	3.63	1.40	1.92	1.17	8.91	1.05	2.12	1.45	3.98

Table 3 shows the percentages of discriminants (fundamental discriminants, resp.)  $-m \in I$  such that  $\nu$  divides  $h(-m)$  for  $\nu = 2, \dots, 100$ . These percentages are always greater than  $100/\nu$ , which confirms hypothesis (2.1). For small primes  $p$  these frequencies are close to  $100/(p-1)$ .

Table 4 is due to A. Odlyzko. The entry  $a_{l,k,\nu}$  in the line starting with  $l, k$  and column headed with  $\nu$  ( $\nu = 2k, 8, \dots, 0$ ) is the number of integers  $m$  from among the first 100,000 even integers  $> 10^l$  which have the property that

$$m/\gcd(m, \text{lcm}(1, \dots, 2^k)) \leq 2^{2k-\nu}.$$

The last two columns record  $r = \ln 10^l / \ln 2^k$  and  $10^5 r^{-r}$ .

TABLE 4

$l$	$k$	$2k$	8	6	4	2	0	$r$	$10^5 r^{-r}$
15	6	0	0	0	0	0	0	8.305	.002
	7	0	0	1	2	2	3	7.118	.086
	8	1	2	6	15	36	49	6.229	1.127
	9	6	17	68	129	210	336	5.537	7.674
	10	27	162	316	530	859	1 282	4.983	33.460
	11	110	585	1 043	1 661	2 447	3 445	4.530	106.655
	12	326	1 806	2 805	4 059	5 565	7 388	4.152	270.747
	13	691	4 075	5 854	7 956	10 434	13 307	3.833	579.832
	14	1 425	7 716	10 462	13 538	16 981	20 804	3.559	1090.481
	15	2 416	12 853	16 521	20 531	24 871	29 519	3.322	1853.443
20	16	3 852	19 174	23 694	28 542	33 620	39 114	3.114	2907.505
	17	5 691	26 485	31 787	37 273	43 148	49 122	2.931	4276.365
	18	7 979	34 478	40 349	46 620	52 592	58 262	2.768	5968.217
	9	0	0	0	0	0	0	7.382	.039
	10	0	0	1	3	6	17	6.644	.344
	11	0	9	22	48	85	120	6.040	1.917
	12	6	70	118	190	271	384	5.537	7.674
	13	25	218	345	510	725	967	5.111	23.945
	14	71	625	907	1 258	1 672	2 182	4.746	61.742
	15	163	1 410	1 937	2 579	3 331	4 258	4.429	137.169
25	16	320	2 787	3 689	4 719	5 930	7 353	4.152	270.747
	17	604	4 952	6 257	7 800	9 590	11 639	3.908	485.833
	18	1 019	7 744	9 640	11 764	14 156	16 640	3.691	806.566
	11	0	0	0	0	0	0	7.550	.024
	12	0	2	3	3	3	3	6.921	.153
	13	0	7	9	16	21	33	6.388	.711
	14	2	35	48	63	93	134	5.932	2.590
	15	8	98	128	182	262	357	5.537	7.674
	16	21	240	339	474	635	809	5.191	19.395
	17	43	567	767	1 014	1 273	1 610	4.885	43.121
30	18	105	1 176	1 532	1 911	2 367	2 895	4.614	86.333
	12	0	0	0	0	0	0	8.305	.002
	13	0	0	0	0	0	1	7.666	.017
	14	0	0	0	0	1	1	7.118	.086
	15	0	1	1	3	5	11	6.644	.344
	16	0	9	15	22	35	56	6.229	1.127
	17	0	43	66	94	130	166	5.862	3.144
	18	6	122	170	234	297	375	5.537	7.674
35	15	0	0	0	0	0	0	7.751	.013
	16	0	2	3	3	3	3	7.267	.055
	17	1	8	8	9	9	13	6.839	.195
	18	3	21	22	22	27	34	6.459	.585

Table 4 also confirms our assumption (2.2). Note that  $a_{l,k,1}$  is the number of integers  $m$  among the first 100,000 even integers  $> 10^l$  such that

$$m = \prod_i p_i^{\tilde{e}_i(m)} \quad \text{with } p_i^{\tilde{e}_i(m)} \leq 2^k$$

(which implies  $m \mid \prod_{i=1}^t p_i^{e'_i}$  for the first prime  $p_t > 2^k$ ). The table shows

$$a_{l,k,1} > 10^5 r^{-r} \quad \text{for } r = \ln 10^l / \ln 2^k \leq \sqrt{\ln 10^l / \ln \ln 10^l}.$$

This suggests an even stronger assumption than (2.2):

$$\# \left\{ m \leq n : m \mid \prod_{i=1}^l p_i^{e'_i} \right\} / n \geq r^{-r}$$

for all  $n, r \leq \sqrt{\ln n / \ln \ln n}$  and  $p_i \leq n^{1/r}$ . Here  $e'_i := \max\{\nu : p_i^\nu \leq p_i\}$ .

Table 4 can be used to balance Stages 1 and 2. If we factor a discriminant  $n \approx 10^{2l}$ , then  $h(-n)$  will be about  $10^l$ . We choose  $p_i \approx 2^k$ . Then hypothesis (2.1) suggests that there is some  $s \leq 10^5 / a_{l,k,\nu}$  with

$$h(-ns) \mid \prod_{i=1}^l p_i^{e'_i} q \quad \text{and} \quad q \leq 2^{2k-\nu}.$$

Hence Stages 1 and 2 will run on at most  $10^5 / a_{l,k,\nu}$  multiples  $-ns$ . Stage 1 with the exponents  $e'_i$  takes about  $2.2p_i$  compositions. If we run Stage 2 with Method 2 for  $2^{k-\nu/2}$  recursion steps, then Stages 1 and 2 will most likely factor this particular  $ns$  (see Table 6, Appendix II, for the performance of Method 2 in Stage 2). In this way Stage 2 takes about  $1.5 \cdot 2^{k-\nu/2}$  compositions. Therefore the total number of compositions of the Main Algorithm will be bounded by

$$b_{l,k,\nu} := \frac{10^5}{a_{l,k,\nu}} 2^k (2.2 + 1.5 \cdot 2^{-\nu/2}).$$

*Examples.*  $n \approx 10^{30}$ . Choose  $k = 12, \nu = 0, a_{15,12,0} = 7 \cdot 388$ . We have  $\lfloor b_{15,12,0} \rfloor = 205132$ , and  $n$  will be factored in about  $2 \cdot 10^5 \approx n^{0.18}$  compositions.

$n \approx 10^{40}$ . Choose  $k = 14, \nu = 0, a_{20,14,0} = 2 \cdot 182$ . We have  $\lfloor b_{20,14,0} \rfloor = 2778221$ , and  $n$  will be factored in about  $2.8 \cdot 10^6 \approx n^{0.16}$  compositions.

$n \approx 10^{50}$ . Choose  $k = 17, \nu = 0, a_{25,17,0} = 1 \cdot 610$ . We have  $\lfloor b_{25,17,0} \rfloor = 30122136$ , and  $n$  will be factored in about  $2.9 \cdot 10^7 \approx n^{0.15}$  compositions.

$n \approx 10^{60}$ . Choose  $k = 18, \nu = 0, a_{30,18,0} = 375$ . We have  $\lfloor b_{30,18,0} \rfloor = 258648746$ , and  $n$  will be factored in about  $2.6 \cdot 10^8 \approx n^{0.14}$  compositions.

The examples show that the number of compositions while factoring  $n$  is smaller than  $\exp \sqrt{\ln n \ln \ln n}$ . For instance, for  $n \approx 10^{60}$  we have  $2.5 \cdot 10^8 \approx 0.00116 \cdot \exp \sqrt{\ln n \ln \ln n}$ . The examples indicate that our algorithm will be faster on integers  $n \geq 10^{40}$  than the Morrison-Brillhart algorithm. Wunderlich [22] reports that the Morrison-Brillhart algorithm for  $n \approx 10^{40}$  takes about  $322 n^{0.152} \approx 3.8 \cdot 10^8 \approx n^{0.21}$  divisions of  $Q_i, Q_i = O(\sqrt{n})$ , by small primes  $p$ . Meanwhile the above estimations have been verified by a program running on the DEC-1091 in Frankfurt; see Section 5.

Table 5 demonstrates the performance of Method 1 of Stage 2. We choose the pseudo-random function  $g: G(\Delta) \rightarrow \{1, \dots, 16\}$

$$g(H) = \lfloor [b^2 \bmod (2^{13} - 1)] 16 / (2^{13} - 1) \rfloor + 1,$$

where  $(a, b, c)$  is the reduced form corresponding to  $H$ . We consider the method with 6 distinct samples of exponents  $a_1, \dots, a_{16}$ : three samples with  $a_i$  chosen at random and three samples with regular  $a_i, a_i = c^{i+31} \bmod 2^{70} \ i = 1, \dots, 16$  with  $c = 2, 3, 5$ . We have the recursion

$$H_{i+1} := H_i H_0^{a_i(H_i)}$$

TABLE 5

$10^m$	$10^8$	$10^{10}$	$10^{12}$	$10^{14}$
sample 1	29	84	252	698
	16	47	99	281
	71	176	535	1401
	49	109	332	783
sample 2	28	87	237	605
	19	56	130	241
	53	171	531	1314
	38	109	413	930
sample 3	28	88	219	606
	20	49	117	266
	57	184	469	1501
	45	130	402	982
$a_i = 2^{i+31} \bmod 2^{70}$	19	61	176	504
	9	29	98	269
	44	121	319	1010
	31	85	163	692
$a_i = 3^{i+31} \bmod 2^{70}$	31	65	211	585
	20	28	102	380
	57	168	477	1485
	40	111	385	945
$a_i = 5^{i+31} \bmod 2^{70}$	28	79	217	819
	20	56	152	388
	59	174	480	1733
	44	113	414	1013

with  $H_0 \in G(\Delta)$  chosen at random. We apply this recursion to the 50 largest discriminants  $\Delta < -10^m$  with  $\Delta \equiv 1 \pmod{4}$  for  $m = 8, 10, 12, 14$ .

For every sample the  $10^m$ -column records four values

1. the average period length,
2. the median of the period lengths,
3. the average number of recursion steps till *search* finds some  $k$  with  $\exists j < k$ :  
 $H_j = H_k$ ,
4. the median of the number of recursion steps.

The particular favorable performance of  $a_i = 2^{i+31} \bmod 2^{70}$  and of  $a_i = 3^{i+31} \bmod 70$  can be explained by the fact that the order of most class groups is even and is a multiple of 3 for about half of the class groups. Despite this favorable performance for random class groups the choice of  $a_i = 2^{i+31} \bmod 2^{70}$  is unfavorable for the factoring algorithm since in the particular situation of Stage 2 the class numbers are free of small prime divisors.

Table 6 shows the performance of Method 2 in Stage 2 for the pseudo-random function  $g: G(\Delta) \rightarrow \{1, \dots, 4\}$

$$g(H) = \lfloor [b^2 \bmod (2^{13} - 1)] 4 / (2^{13} - 1) \rfloor + 1,$$

where  $(a, b, c)$  is the reduced form corresponding to  $H$ . We used  $a_i = d^i, i = 3, 4$ , with constants  $d = 2, 3, \dots, 8$ . The recursion scheme is

$$H_{i+1} = \begin{cases} H_i^3 & \text{if } g(H_i) \leq 2, \\ H_i H_0^{d_{g(H_i)}} & \text{otherwise,} \end{cases}$$

$H_0 \in G(\Delta)$  is chosen at random.

For every  $d = 2, 3, \dots, 8$  and  $m = 8, \dots, 14$  we applied this recursion to the 50 largest discriminants  $\Delta \approx -10^m$  with  $\Delta \equiv 1 \pmod{4}$ . For every  $d$  and  $m$  the table records four values:

1. the average period length,
2. the median of the period lengths,
3. the average number of recursion steps till *search* finds some  $k$  with  $\exists j < k$ :  
 $H_j = H_k$ ,
4. the median of the number of recursion steps.

TABLE 6

base $d$	$10^8$	$10^{10}$	$10^{12}$	$10^{14}$
2	21	63	178	684
	12	42	112	289
	51	158	395	1513
	43	92	322	1099
3	28	116	231	783
	17	53	86	555
	63	219	503	1519
	50	157	260	1118
4	24	76	185	780
	15	41	83	277
	48	165	360	1260
	31	123	285	688
5	34	103	273	746
	22	58	143	429
	74	224	570	1613
	50	161	357	1013
6	23	43	98	435
	12	24	79	196
	55	114	299	927
	35	72	188	684
7	28	90	276	713
	24	74	154	376
	63	224	516	1533
	51	159	386	925
8	26	76	209	763
	13	47	129	445
	53	144	391	1449
	41	94	228	918

**Acknowledgement.** Several members of the Frankfurt working group participated in implementing this algorithm. H. G. Franke produced Tables 1–3 and implemented large integer arithmetic and an efficient composition algorithm in machine language. J. Sattler produced Tables 5, 6 which compare Methods 1, 2 of Stage 2.

Thanks are due to the computer centres of Frankfurt and Amsterdam university for providing computing time on the DEC-1091 in Frankfurt and the CDC-Cyber in Amsterdam. We also like to thank A. Odlyzko for the permission to include in this paper Table 4 which is part of a larger statistic made at Bell Laboratories.

Fachbereich Mathematik  
Universität Frankfurt  
6 Frankfurt am Main, West Germany

Mathematisch Instituut  
Universiteit Amsterdam  
1018 WB Amsterdam, The Netherlands

1. R. P. BRENT, "An improved Monte Carlo factorization algorithm," *BIT*, v. 20, 1980, pp. 176–184.
2. J. W. S. CASSELS, *Rational Quadratic Forms*, Academic Press, London, New York, 1978.
3. H. COHEN & H. W. LENSTRA, JR., *Divisibility by Small Primes of Class Numbers*, Personal communication, 1982.
4. J. D. DIXON, "Asymptotically fast factorization of integers," *Math. Comp.*, v. 36, 1981, pp. 255–260.
5. C. F. GAUSS, *Disquisitiones Arithmeticae*, Leipzig, 1801; English transl. by A. A. Clarke, Yale University Press, New Haven and London, 1966.
6. G. H. HARDY & E. M. WRIGHT, *An Introduction to the Theory of Numbers*, 5th ed., Oxford Univ. Press, Oxford, 1979.
7. D. E. KNUTH, *The Art of Computer Programming*, Vol. 2, *Seminumerical Algorithms*, 2nd ed., Addison-Wesley, Reading, Mass., 1981.
8. P. G. LEJEUNE DIRICHLET & R. DEDEKIND, *Vorlesungen über Zahlentheorie*, Braunschweig, 1893; reprint, New York, 1968.
9. H. W. LENSTRA, JR., *On the Calculation of Regulators and Class Numbers of Quadratic Fields*, Journées Arithmétiques 1980 (J. V. Armitage, Ed.), Cambridge Univ. Press, Oxford, 1982, pp. 123–150.
10. G. B. MATHEWS, *Theory of numbers*, 1892; Reprint, Chelsea, New York, 1962.
11. L. MONIER, *Algorithmes de Factorisation d'Entiers*, Thèse d'informatique, Université Paris Sud, 1980.
12. M. A. MORRISON & J. BRILLHART, "A method of factorization and the factorization of  $F_7$ ," *Math. Comp.*, v. 29, 1975, pp. 183–205.
13. J. M. POLLARD, "A Monte Carlo method for factorization," *BIT*, v. 15, 1975, pp. 331–334.
14. C. POMERANCE, "Analysis and comparison of some integer factoring algorithms," *Computational Methods in Number Theory* (R. Tijdemen and H. Lenstra, Eds.), Mathematical Centrum, Amsterdam, 1981.
15. R. L. RIVEST, A. SHAMIR & L. ADLEMAN, "A method for obtaining digital signatures and public key cryptosystems," *Comm. ACM*, v. 21, 1978, pp. 120–126.
16. J. SATTLER & C. P. SCHNORR, "Ein Effizienzvergleich der Faktorisierungsverfahren von Morrison-Brillhart und Schroepfel," *Computing*, v. 30, 1983, pp. 91–110.
17. D. SHANKS, *Class Number, A Theory of Factorization, and Genera*, Proc. Sympos. Pure Math., vol. 20, Amer. Math. Soc., Providence, R. I., 1971, pp. 415–440.
18. J. SATTLER & C. P. SCHNORR, *Generating Random Walks in Groups*, Preprint, Universität Frankfurt, 1983; submitted for publication.
19. C. P. SCHNORR, "Refined analysis and improvements on some factoring algorithms," *J. Algorithms*, v. 3, 1982, pp. 101–127.
20. C. P. SCHNORR & M. SEYSEN, *An Improved Composition Algorithm*, Preprint, Universität Frankfurt, 1982; submitted for publication.
21. C. L. SIEGEL, "Über die Klassenzahl quadratischer Zahlkörper," *Acta Arith.*, v. 1, 1936, pp. 83–86.
22. S. S. WAGSTAFF & M. C. WUNDERLICH, *A Comparison of Two Factorization Methods*, Unpublished manuscript.
23. H. G. ZIMMER, *Computational Problems, Methods, and Results in Algebraic Number Theory*, Lecture Notes in Math., Vol. 262, Springer, Berlin and New York, 1972.