

ON THE REMAINDER OF GAUSSIAN QUADRATURE FORMULAS FOR BERNSTEIN-SZEGŐ WEIGHT FUNCTIONS

F. PEHERSTORFER

ABSTRACT. We give an explicit expression for the kernel of the error functional for Gaussian quadrature formulas with respect to weight functions of Bernstein-Szegő type, i.e., weight functions of the form $(1-x)^\alpha(1+x)^\beta/\rho(x)$, $x \in (-1, 1)$, where $\alpha, \beta \in \{-\frac{1}{2}, \frac{1}{2}\}$ and ρ is a polynomial of arbitrary degree which is positive on $[-1, 1]$. With the help of this result the norm of the error functional can easily be calculated explicitly for a wide subclass of these weight functions.

1. INTRODUCTION AND NOTATION

We consider Gaussian quadrature formulas with respect to a nonnegative weight function w on the interval $[-1, 1]$,

$$(1.1) \quad \int_{-1}^1 f(x)w(x)dx = \sum_{j=1}^n \lambda_j f(x_j) + R_n(f, w),$$

where $x_j = x_{j,n}$ are the zeros of the n th-degree monic orthogonal polynomial $P_n(\cdot, w)$ and $\lambda_j = \lambda_{j,n}$ are the corresponding Christoffel numbers. If f is analytic in a domain D which contains in its interior the interval $[-1, 1]$ and a contour Γ surrounding $[-1, 1]$, the remainder term can be represented as a contour integral (see, e.g., [3])

$$(1.2) \quad R_n(f, w) = \frac{1}{2\pi i} \int_{\Gamma} K_n(z, w) f(z) dz,$$

where the kernel $K_n(\cdot, w)$ is given by

$$(1.3) \quad K_n(z, w) = R_n\left(\frac{1}{z-\cdot}, w\right)$$

or, alternatively, by

$$(1.4) \quad K_n(z, w) = \frac{Q_n(z, w)}{P_n(z, w)},$$

where $Q_n(\cdot, w)$ is the n th function of the second kind, i.e.,

$$(1.5) \quad Q_n(z, w) = \int_{-1}^1 \frac{P_n(x, w)}{z-x} w(x) dx \quad \text{for } z \in \mathbb{C} \setminus [-1, 1].$$

Received by the editor October 30, 1990 and, in revised form, November 5, 1991 and February 10, 1992.

1991 *Mathematics Subject Classification.* Primary 65D32; Secondary 33C45.

©1993 American Mathematical Society
0025-5718/93 \$1.00 + \$.25 per page

Let us note that by (1.3), $K_n(\cdot, w)$ has the following series expansion:

$$(1.6) \quad K_n(z, w) = \sum_{k=2n}^{\infty} \frac{R_n(x^k, w)}{z^{k+1}} \quad \text{for } |z| > 1.$$

From (1.2) the following well-known estimate of the remainder, based on contour integration, follows:

$$(1.7) \quad |R_n(f, w)| \leq \frac{l(\Gamma)}{2\pi} \max_{z \in \Gamma} |K_n(z, w)| \max_{z \in \Gamma} |f(z)|,$$

where $l(\Gamma)$ denotes the length of Γ .

Another useful method to estimate the remainder for a function analytic in $C_r = \{z \in \mathbb{C}: |z| < r\}$, $r > 1$, has been suggested by Hämmerlin [4], namely: For a function $f(z) = \sum_{k=0}^{\infty} a_k(f)z^k$ analytic in C_r define

$$|f|_r := \sup\{|a_k(f)|r^k : k \in \mathbb{N}_0 \text{ and } R_n(x^k, w) \neq 0\}.$$

Then, $|\cdot|_r$ in the space

$$\mathfrak{X}_r := \{f: f \text{ analytic in } C_r \text{ and } |f|_r < \infty\}$$

is a seminorm. The error functional $R_n(f, w)$ is continuous in $(\mathfrak{X}_r, |\cdot|_r)$, and we have

$$|R_n(f, w)| \leq \|R_n\| |f|_r,$$

where $\|R_n\|$ can be estimated by

$$(1.8) \quad \|R_n\| \leq \sum_{k=2n}^{\infty} \frac{|R_n(x^k, w)|}{r^k}.$$

Equality holds (put $f(z) = 1/(r-z)$, resp. $1/(r+z)$) if for all $k \geq 2n$ the condition

$$(1.9) \quad R_n(x^k, w) \geq 0, \quad \text{resp. } (-1)^k R_n(x^k, w) \geq 0,$$

is fulfilled. Since by [3, Theorem 2.1], proved in [2], and the proof of Theorem 3.1 in [3], the condition

$$(1.10) \quad w(x)/w(-x) \text{ nondecreasing on } (-1, 1),$$

resp.

$$(1.11) \quad w(x)/w(-x) \text{ nonincreasing on } (-1, 1),$$

implies that condition (1.9) holds for all $k \in \mathbb{N}_0$, it follows by (1.6) (see [3, Theorem 3.1]) that

$$\max_{|z|=r} |K_n(z, w)| = \begin{cases} K_n(r, w) & \text{if } w \text{ satisfies (1.10),} \\ |K_n(-r, w)| & \text{if } w \text{ satisfies (1.11),} \end{cases}$$

and

$$\|R_n\| = \begin{cases} rK_n(r, w) & \text{if } w \text{ satisfies (1.10),} \\ -rK_n(-r, w) & \text{if } w \text{ satisfies (1.11).} \end{cases}$$

Thus, we see that for the estimation of the remainder it is very desirable to have an explicit expression for the kernel $K_n(z, w)$.

Very recently, Notaris [8] computed $\|R_n\|$ explicitly for weight functions of the form

$$(1.12) \quad w(x) = (1-x)^\alpha(1+x)^\beta/\rho_2(x) \quad \text{for } x \in (-1, 1),$$

where $\alpha, \beta \in \{-\frac{1}{2}, \frac{1}{2}\}$ and ρ_2 is a polynomial of degree at most two which is positive on $[-1, 1]$ and satisfies condition (1.10) or (1.11). For the special case when ρ_2 is a polynomial of degree one or a particular even polynomial of degree two, this has been done before by Akrivis [1] (see also Kumar [5, 6]). Let us also mention that a detailed study of the kernel function for the four Chebyshev weight functions, i.e., $\rho_2(x) \equiv 1$ in (1.12), can be found in Gautschi and Varga [3]. In this note we derive an explicit expression for the kernel $K_n(z, w)(\|R_n\|)$ for all Bernstein-Szegő weight functions w (which satisfy condition (1.10) or (1.11)), where a weight function is called a Bernstein-Szegő weight function if it is of the form

$$(1.13) \quad \pi w_{\alpha, \beta}(x, \rho_m) = (1-x)^\alpha(1+x)^\beta/\rho_m(x) \quad \text{for } x \in (-1, 1),$$

with $\alpha, \beta \in \{-\frac{1}{2}, \frac{1}{2}\}$ and ρ_m a polynomial of degree m , m arbitrary, which is positive on $[-1, 1]$.

2. MAIN RESULT

First let us recall the well-known fact that the so-called Joukowski transformation

$$(2.1) \quad y = \frac{1}{2}(z + z^{-1})$$

maps $\{z \in \mathbf{C}: |z| < 1\} \setminus \{0\}$ ($\{z \in \mathbf{C}: |z| > 1\}$) one-to-one onto $\mathbf{C} \setminus [-1, 1]$ and that the inverse transformation is given by

$$(2.2) \quad z = y_{(+)} \sqrt{y^2 - 1},$$

where that branch of $\sqrt{}$ is chosen for which $\sqrt{y^2 - 1} > 0$ for $y \in (1, \infty)$. Note that the transformation (2.1) maps the circumference $|z| = 1$ onto the interval $[-1, 1]$.

The following version of the Fejér-Riesz Theorem on the representation of positive trigonometric polynomials (compare Theorems 1.2.1 and 1.2.2 in [10]) will be needed.

Lemma. *Let ρ_m be a real positive polynomial on $[-1, 1]$ of exact degree m . Then there exists a unique real polynomial*

$$(2.3) \quad g_m(z) = \prod_{\nu=1}^m (z - z_\nu) \quad \text{with } 0 < |z_\nu| < 1 \text{ for } \nu = 1, \dots, m$$

such that

$$(2.4) \quad \rho_m(\cos \varphi) = c |g_m(e^{i\varphi})|^2 \quad \text{for } \varphi \in [0, 2\pi],$$

where $c \in \mathbf{R}^+$.

Proof. Let $\rho_m(x) = \tilde{c} \prod_{\nu=1}^m (\alpha_\nu - x)$, where $\tilde{c} \in \mathbf{R}$ and the α_ν 's are either in $\mathbf{R} \setminus [-1, 1]$ or appear in pairs of complex conjugate numbers. Hence, if we set

$$(2.5) \quad z_\nu = \alpha_\nu - \sqrt{\alpha_\nu^2 - 1} \quad \text{for } \nu = 1, \dots, m,$$

then

$$0 < |z_\nu| < 1 \quad \text{and} \quad \alpha_\nu = \frac{1}{2}(z_\nu + z_\nu^{-1}) \quad \text{for } \nu = 1, \dots, m.$$

Thus,

$$\rho_m(\cos \varphi) = \tilde{c} \prod_{\nu=1}^m \left(\frac{1 + z_\nu^2}{2z_\nu} - \cos \varphi \right) = c \prod_{\nu=1}^m |e^{i\varphi} - z_\nu|^2,$$

since

$$(e^{i\varphi} - z_\nu)(e^{-i\varphi} - z_\nu) = 2z_\nu \left(\frac{1 + z_\nu^2}{2z_\nu} - \cos \varphi \right)$$

and the z_ν 's are real or appear in pairs of complex conjugate numbers.

Now the uniqueness remains to be shown. Suppose that

$$\rho_m(\cos \varphi) = d \prod_{\nu=1}^m |e^{i\varphi} - v_\nu|^2,$$

where $d \in \mathbf{R}^+$, $v_\nu \in \{z \in \mathbf{C}: |z| < 1\} \setminus \{0\}$ for $\nu = 1, \dots, m$, and the v_ν 's are real or complex conjugate. Then it follows as above that $(v_\nu + v_\nu^{-1})/2$, $\nu = 1, \dots, m$, is a zero of $\rho_m(\cos \varphi)$ and thus, since $0 < |v_\nu| < 1$ and since the Joukowski transformation is one-to-one, the uniqueness follows. \square

Let us note that other representations of ρ_m of the form (2.4), but with g_m having $m-l$, resp. l , $l \in \{1, \dots, m\}$, zeros inside, resp. outside, of the unit disk, can be obtained by replacing (2.5) by

$$(2.6) \quad z_{\nu_j} = \alpha_{\nu_j} + \sqrt{\alpha_{\nu_j}^2 - 1} \quad \text{for } j = 1, \dots, l,$$

where $\{\nu_1, \dots, \nu_l\}$ is an arbitrary subset of $\{1, \dots, m\}$, and

$$z_\nu = \alpha_\nu - \sqrt{\alpha_\nu^2 - 1} \quad \text{for } \nu \in \{1, \dots, m\} \setminus \{\nu_1, \dots, \nu_l\}.$$

Now let us set

$$\pi w(x, \rho_m) = 1/(\sqrt{1-x^2} \rho_m(x)) \quad \text{for } x \in (-1, 1)$$

and let g_m be the unique polynomial from the above lemma. Then it follows by well-known results of Bernstein and Szegö (see, e.g., [10, p. 31] and set $h_m(z) = \sqrt{c\pi} z^m g_m(\frac{1}{z})$ there) that, with $z = e^{i\varphi}$ and $x = \cos \varphi$,

$$\begin{aligned} (2.7) \quad 2^{n-1} P_n(x, w(\cdot, \rho_m)) &= \sum_{j=0}^m a_j T_{n-j}(x) \\ &= \operatorname{Re}\{z^{n-m} g_m(z)\} \quad \text{for } 2n > m, \\ 2^{n-1} P_{n-1}(x, (1-x^2)w(\cdot, \rho_m)) &= \sum_{j=0}^m a_j U_{n-1-j}(x) \\ &= \operatorname{Im}\{z^{n-m} g_m(z)\}/\sin \varphi \quad \text{for } 2n > m, \\ 2^n P_n(x, (1+x)w(\cdot, \rho_m)) &= \sum_{j=0}^m a_j \frac{T_{n+1-j}(x) + T_{n-j}(x)}{x+1} \\ &= \operatorname{Re}\{z^{n-m+1/2} g_m(z)\}/\cos(\varphi/2) \quad \text{for } 2n+1 > m, \\ 2^n P_n(x, (1-x)w(\cdot, \rho_m)) &= \sum_{j=0}^m a_j \frac{T_{n+1-j}(x) - T_{n-j}(x)}{x-1} \\ &= \operatorname{Im}\{z^{n-m+1/2} g_m(z)\}/\sin(\varphi/2) \quad \text{for } 2n+1 > m, \end{aligned}$$

where $g_m(z) = \sum_{j=0}^m a_j z^{m-j}$ and T_j , resp. U_j , denotes the Chebyshev polynomial of degree j of the first, resp. second, kind on $[-1, 1]$.

We mention in passing that if g_m in (2.7) is replaced by a polynomial \tilde{g}_m which also satisfies (2.4) but does not have all zeros in the open unit disk, then the polynomials on the right-hand side in (2.7) are not orthogonal with respect to $w_{\alpha, \beta}(\cdot, \rho_m)$, $\alpha, \beta \in \{-\frac{1}{2}, \frac{1}{2}\}$. In fact (see [9, Corollary 5], corresponding results hold also for $\alpha = -\beta = \pm\frac{1}{2}$), they are orthogonal with respect to a functional Ψ of the form

$$\Psi(p) = \int_{-1}^{+1} p(x) w_{\alpha, \beta}(x, \rho_m) dx + L(p) \quad \text{for } p \in \mathbf{P},$$

where L is a functional given by

$$L(p) = \sum_{j=1}^{l^*} \sum_{\kappa=1}^{l_j} \mu_{\kappa, j} p^{(\kappa-1)}(\alpha_{\nu_j}),$$

and the α_{ν_j} 's are those zeros of ρ_m which correspond to the zeros of \tilde{g}_m lying outside of the unit disk by (2.6), l_j is the multiplicity of the zero α_{ν_j} , and the $\mu_{\kappa, j}$'s are certain real numbers.

We now give the announced explicit expression for the kernel function $K_n(z, w_{\alpha, \beta})$, $|\alpha| = |\beta| = \frac{1}{2}$.

Theorem. Let ρ_m be given by (2.4). Then we have for $y \in \mathbf{C} \setminus [-1, 1]$, on writing $y = \frac{1}{2}(z + z^{-1})$ with $|z| < 1$, i.e., $z = y - \sqrt{y^2 - 1}$, that

$$\begin{aligned} cK_n(y, w(\cdot, \rho_m)) &= \frac{4z^{2n+1}}{(1-z^2)g_m^*(z)\{z^{2n-m}g_m(z) + g_m^*(z)\}} \\ &\quad \text{for } 2n > m, \\ cK_n(y, (1 \begin{smallmatrix} + \\ - \end{smallmatrix} x)w(\cdot, \rho_m)) &= \frac{2z^{2n+1}(z \begin{smallmatrix} + \\ - \end{smallmatrix} 1)}{(1 \begin{smallmatrix} - \\ + \end{smallmatrix} z)g_m^*(z)\{z^{2n+1-m}g_m(z) \begin{smallmatrix} + \\ - \end{smallmatrix} g_m^*(z)\}} \\ &\quad \text{for } 2n+1 > m, \\ cK_n(y, (1-x^2)w(\cdot, \rho_m)) &= \frac{z^{2n+1}(z^2-1)}{g_m^*(z)\{z^{2n+2-m}g_m(z) - g_m^*(z)\}} \\ &\quad \text{for } 2n+2 > m, \end{aligned}$$

where $g_m^*(z) = z^m g_m(\frac{1}{z})$.

Proof. Let R and S be monic polynomials of degree at most two such that

$$R(y)S(y) = y^2 - 1,$$

and let us put for abbreviation

$$P_n(x) := P_n(x, R w(\cdot, \rho_m)) \quad \text{and} \quad \tilde{P}_n(x) := P_n(x, S w(\cdot, \rho_m)).$$

Using the simple fact that for $k \in \mathbf{Z}$ and $\varphi \in [0, 2\pi]$

$$(2.8) \quad [\operatorname{Re}\{e^{ik\varphi} g_m(e^{i\varphi})\}]^2 + [\operatorname{Im}\{e^{ik\varphi} g_m(e^{i\varphi})\}]^2 = |g_m(e^{i\varphi})|^2,$$

we get, using (2.7) and (2.4), that with $l = n + \partial R - 1$, where ∂R denotes the exact degree of R ,

$$(2.9) \quad RP_n^2 - S\tilde{P}_l^2 = k_n \rho_m,$$

where

$$k_n = \begin{cases} 2^{-2n+2}/c & \text{for } R(x) \equiv 1, \\ -2^{-2n}/c & \text{for } R(x) \equiv x^2 - 1, \\ \binom{+}{-} 2^{-2n+1}/c & \text{for } R(x) \equiv x + 1 \ (x - 1). \end{cases}$$

Furthermore, it follows from Theorem 3(a) of our paper [9] that for $2n \geq m + 1 - \partial R$

$$(2.10) \quad RP_n^2 - S(YP_n + \rho_m P_{n-1}^{(1)})^2 = d_n \rho_m,$$

where $P_{n-1}^{(1)}$ denotes the associated polynomial of P_n , i.e.,

$$P_{n-1}^{(1)}(y) = \int_{-1}^1 \frac{P_n(y) - P_n(x)}{y - x} R(x) w(x, \rho_m) dx,$$

and $Y \in \mathbf{P}_\mu$, $\mu = \max\{m - 1, \partial R - 1\}$, is uniquely determined by the conditions that at each zero α_ν of $\rho_m(x) = \tilde{c} \prod_{\nu=1}^{m^*} (\alpha_\nu - x)^{m_\nu}$, where $\tilde{c} \in \mathbf{R}$ and m_ν is the multiplicity of the zero α_ν ,

$$(2.11) \quad Y^{(j)}(\alpha_\nu) = (R/\sqrt{y^2 - 1})^{(j)}(\alpha_\nu) \quad \text{for } j = 0, \dots, m_\nu - 1,$$

and that for $y \rightarrow \infty$

$$(2.12) \quad \frac{(R/\sqrt{y^2 - 1} - Y)(y)}{\rho_m(y)} = O(y^{-1});$$

furthermore,

$$(2.13) \quad d_n = 2 \int_{-1}^{+1} P_n^2(x) R(x) w(x, \rho_m) dx.$$

(We note that in the definition of $1/h$ in [9, p. 461] $(-1)^{l-k}/\sqrt{-H}$ is to be replaced by $(-1)^{l-k}/\pi\sqrt{-H}$.) It now follows from [10, (2.6.5)] that the leading coefficient of the orthonormal polynomial of degree n with respect to $Rw(\cdot, \rho_m)$ is equal to $\sqrt{2/k_n}$ for $2n \geq m + 1 - \partial R$, which implies that $d_n = k_n$ and thus, in view of (2.9) and (2.10),

$$(2.14) \quad \pm \tilde{P}_l = YP_n + \rho_m P_{n-1}^{(1)} \quad \text{for } 2n \geq m + 1 - \partial R.$$

For a function f defined on $\mathbf{C} \setminus [-1, 1]$ and for $x \in (-1, 1)$ we write, provided the limits involved exist,

$$f^{(\pm)}(x) := \lim_{\substack{z \rightarrow x \\ z \in \mathbf{C}^{(\pm)}}} f(z),$$

where $\mathbf{C}^{(\pm)} := \{z \in \mathbf{C} : \operatorname{Im} z \gtrless 0\}$. Observing that by (2.11) and (2.12)

$$(2.15) \quad \Phi(y) := \frac{(R/\sqrt{y^2 - 1} - Y)(y)}{\rho_m(y)}$$

is analytic on $\mathbf{C} \setminus [-1, 1]$ and vanishes at infinity, and that the boundary values $\Phi^\pm(x)$, $x \in (-1, 1)$, from the upper (lower) half plane satisfy the relation

$$\Phi^+(x) - \Phi^-(x) = \frac{2}{i} \frac{R(x)}{\rho_m(x) \sqrt{1 - x^2}} \quad \text{for } x \in (-1, 1),$$

where we have used the fact that

$$(\sqrt{y^2 - 1})^+(x) = i\sqrt{1 - x^2} = -(\sqrt{y^2 - 1})^-(x),$$

we get by the Sochozki-Plemelj formula (see, e.g., [7])

$$(2.16) \quad \begin{aligned} \Phi(y) &= \frac{1}{\pi} \int_{-1}^{+1} \frac{1}{y-x} \frac{R(x)}{\rho_m(x)\sqrt{1-x^2}} dx \\ &= Q_0(y, R w(\cdot, \rho_m)) \quad \text{for } y \in \mathbb{C} \setminus [-1, 1]. \end{aligned}$$

Recalling the well-known fact (see, e.g., [10, §3.5]) that for sufficiently large $|y|$

$$\frac{P_{n-1}^{(1)}(y)}{P_n(y)} = Q_0(y, R w(\cdot, \rho_m)) + O(y^{-(2n+1)}),$$

we get, using (2.16) and (2.15), that

$$(Y P_n + \rho_m P_{n-1}^{(1)})(y) = P_n(y) R(y) / \sqrt{y^2 - 1} + O(y^{-(n+1)+m})$$

and thus, since for $2n \geq m + 1 - \partial R$

$$\lim_{y \rightarrow \infty} y^{-(n+\partial R-1)} \{P_n(y) R(y) / \sqrt{y^2 - 1} + O(y^{-(n+1)+m})\} = 1,$$

the polynomial $Y P_n + \rho_m P_{n-1}^{(1)}$, which by (2.14) is of exact degree $n + \partial R - 1$, has leading coefficient one. Hence, the plus sign holds in (2.14). Thus, the n th function of the second kind is of the form

$$(2.17) \quad \begin{aligned} Q_n(y, R w(\cdot, \rho_m)) &= -P_{n-1}^{(1)}(y) + P_n(y) Q_0(y) \\ &= \frac{\sqrt{R(y)} P_n(y) - \sqrt{S(y)} \tilde{P}_l(y)}{\sqrt{S(y)} \rho_m(y)} = \frac{k_n}{(\sqrt{y^2 - 1} P_n + S \tilde{P}_l)(y)}, \end{aligned}$$

where the second equality follows with the help of (2.16), (2.15), and (2.14), and the third equality with the help of (2.9). Now the following equalities hold on the circumference $|z| = 1$:

$$(2.18) \quad \begin{aligned} 2^n P_n \left(\frac{1}{2}(z + z^{-1}), w(\cdot, \rho_m) \right) &= z^{-n} (z^{2n-m} g_m(z) + g_m^*(z)), \\ 2^{n-1} P_{n-1} \left(\frac{1}{2}(z + z^{-1}), (1 - x^2) w(\cdot, \rho_m) \right) &= \frac{z^{-n} (z^{2n-m} g_m(z) - g_m^*(z))}{z - z^{-1}}, \\ 2^n P_n \left(\frac{1}{2}(z + z^{-1}), (1 \begin{smallmatrix} + \\ - \end{smallmatrix} x) w(\cdot, \rho_m) \right) &= \frac{z^{-n} (z^{2n+1-m} g_m(z) \begin{smallmatrix} + \\ - \end{smallmatrix} g_m^*(z))}{z \begin{smallmatrix} + \\ - \end{smallmatrix} 1}. \end{aligned}$$

Since all functions appearing in (2.18) are analytic in the domain $\mathbb{C} \setminus \{0\}$, it follows that in (2.18) equality holds also on $\mathbb{C} \setminus \{0\}$. Hence, we get from (2.17)

and (2.18) that for $y \in \mathbb{C} \setminus [-1, 1]$, on writing $y = \frac{1}{2}(z + z^{-1})$ with $|z| < 1$,

$$(2.19) \quad \begin{aligned} Q_n(y, w(\cdot, \rho_m)) &= \frac{2^{-n+2}}{c} \frac{z^{n+1}}{(1 - z^2)g_m^*(z)}, \\ Q_n(y, (1 - x^2)w(\cdot, \rho_m)) &= \frac{2^{-n}}{c} \frac{z^{n+1}}{g_m^*(z)}, \\ Q_n(y, (1_{(-)}^+ x)w(\cdot, \rho_m)) &= \frac{2^{-n+1}}{c} \frac{z^{n+1}}{(1_{(-)}^+ z)g_m^*(z)}, \end{aligned}$$

where we have used the fact that $\sqrt{y^2 - 1} = (z^{-1} - z)/2$. Relation (2.19) in conjunction with (2.18) and (1.4) gives the assertion. \square

In the remark below we state sufficient conditions on the weight function $w_{\alpha, \beta}(x, \rho_m)$, defined in (1.13), such that (1.10), resp. (1.11), is fulfilled. Since the product $w_1(x)w_2(x)$ of two weight functions w_1, w_2 satisfies condition (1.10), resp. (1.11) if w_1 and w_2 satisfy (1.10), resp. (1.11), we consider the behavior of $w(x, \rho_m)/w(-x, \rho_m)$ for $m \in \{1, 2\}$ only.

Remark. The ratio $w(x, \rho)/w(-x, \rho)$ is nondecreasing (nonincreasing) on $(-1, 1)$ if

$$\rho(x) = \begin{cases} {}_{(-)}^+(\alpha - x), & \alpha \in (1, \infty) \text{ } ((-\infty, -1)), \\ (\alpha - x)(x - \beta), & \alpha \in (1, \infty), \beta \in (-\infty, -1), \text{ and } -\beta \stackrel{\geq}{(\leq)} \alpha, \\ (x - \alpha)^2 + \beta^2, & \alpha \in \mathbf{R}^{(+)}_{(-)}, \beta \in \mathbf{R}, \text{ and } \alpha^2 + \beta^2 \geq 1, \end{cases}$$

where the expressions in parentheses refer to the case of nonincreasing ratio.

Setting in the preceding theorem

$$g_1(z) = z + \tilde{a}, \quad \tilde{a} \in (-1, 1), \quad \text{i.e., } |g_1(e^{i\varphi})|^2 = 1 + \tilde{a}^2 + 2\tilde{a}x,$$

resp. for $b > 0$

$$g_2(z) = z^2 + (1 + 2b)^{-1}, \quad \text{i.e., } |g_2(e^{i\varphi})|^2 = 4(b^2 + (1 + 2b)x^2)/(2b + 1)^2,$$

where $x = \cos \varphi$, we obtain the results of Kumar [5, 6] concerning the functions of the second kind, and the results of Akivis [1] on the norm of the error functional $R_n(\cdot, w_{\alpha, \beta}(\cdot, |g_j(e^{i\varphi})|^2))$, $j = 1, 2$. If we put

$$g_2(z) = z^2 + \frac{2\delta}{\beta}z + \left(1 - \frac{2\alpha}{\beta}\right)$$

with $0 < \alpha < \beta$, $\beta \neq 2\alpha$, and $|\delta| < \beta - \alpha$, which gives

$$\frac{\beta^2}{4}|g_2(e^{i\varphi})|^2 = \beta(\beta - 2\alpha)x^2 + 2\delta(\beta - \alpha)x + \alpha^2 + \delta^2,$$

we obtain the results of Notaris [8] on the norm of the error functional, using his conditions (2.3₁)–(2.4₂) on the parameters $\alpha, \beta, \gamma, \delta$ under which the function $w(x, |g_2(e^{i\varphi})|^2)/w(-x, |g_2(e^{i\varphi})|^2)$ is strictly increasing, resp. strictly decreasing.

ACKNOWLEDGMENT

I would like to thank the referee for a careful reading of the manuscript and for his valuable comments.

BIBLIOGRAPHY

1. G. Akrivis, *The error norm of certain Gaussian quadrature formulae*, Math. Comp. **45** (1985), 513–519.
2. W. Gautschi, *On Padé approximants associated with Hamburger series*, Calcolo **20** (1983), 111–127.
3. W. Gautschi and R. S. Varga, *Error bounds for Gaussian quadrature of analytic functions*, SIAM J. Numer. Anal. **20** (1983), 1170–1186.
4. G. Hämmerlin, *Fehlerabschätzungen bei numerischer Integration nach Gauss*, Methoden und Verfahren der Mathematischen Physik, vol. 6 (B. Brosowski and E. Martensen, eds.), Bibliographisches Institut, Mannheim, Wien, Zürich, 1972, pp. 153–163.
5. R. Kumar, *A class of quadrature formulas*, Math. Comp. **28** (1974), 769–778.
6. —, *Certain Gaussian quadratures*, J. Inst. Math. Appl. **14** (1974), 175–182.
7. N. I. Muschelischwili, *Singuläre Integralgleichungen*, Akademie Verlag, Berlin, 1965.
8. S. E. Notaris, *The error norm of Gaussian quadrature formulae for weight functions of Bernstein-Szegő type*, Numer. Math. **57** (1990), 271–283.
9. F. Peherstorfer, *On Bernstein-Szegő orthogonal polynomials on several intervals*, SIAM J. Math. Anal. **21** (1990), 461–482.
10. G. Szegő, *Orthogonal polynomials*, 3rd ed., Amer. Math. Soc. Colloq. Publ., vol. 23, Amer. Math. Soc., Providence, RI, 1967.

INSTITUT FÜR MATHEMATIK, JOHANNES KEPLER UNIVERSITÄT LINZ, A-4040 LINZ, AUSTRIA