

## SPHERICAL $t_\epsilon$ -DESIGNS FOR APPROXIMATIONS ON THE SPHERE

YANG ZHOU AND XIAOJUN CHEN

ABSTRACT. A spherical  $t$ -design is a set of points on the unit sphere that are nodes of a quadrature rule with positive equal weights that is exact for all spherical polynomials of degree  $\leq t$ . The existence of a spherical  $t$ -design with  $(t+1)^2$  points in a set of interval enclosures on the unit sphere  $\mathbb{S}^2 \subset \mathbb{R}^3$  for any  $0 \leq t \leq 100$  is proved by Chen, Frommer, and Lang (2011). However, how to choose a set of points from the set of interval enclosures to obtain a spherical  $t$ -design with  $(t+1)^2$  points is not given in loc. cit. It is known that  $(t+1)^2$  is the dimension of the space of spherical polynomials of degree at most  $t$  in 3 variables on  $\mathbb{S}^2$ . In this paper we investigate a new concept of point sets on the sphere named spherical  $t_\epsilon$ -design ( $0 \leq \epsilon < 1$ ), which are nodes of a positive, but not necessarily equal, weight quadrature rule exact for polynomials of degree  $\leq t$ . The parameter  $\epsilon$  is used to control the variation of the weights, while the sum of the weights is equal to the area of the sphere. A spherical  $t_\epsilon$ -design is a spherical  $t$ -design when  $\epsilon = 0$ , and a spherical  $t$ -design is a spherical  $t_\epsilon$ -design for any  $0 < \epsilon < 1$ . We show that any point set chosen from the set of interval enclosures (loc. cit.) is a spherical  $t_\epsilon$ -design. We then study the worst-case error in a Sobolev space for quadrature rules using spherical  $t_\epsilon$ -designs, and investigate a model of polynomial approximation with  $l_1$ -regularization using spherical  $t_\epsilon$ -designs. Numerical results illustrate the good performance of spherical  $t_\epsilon$ -designs for numerical integration and function approximation on the sphere.

### 1. INTRODUCTION

Let  $\mathbb{S}^d := \{\mathbf{x} = (x_1, \dots, x_{d+1})^T \in \mathbb{R}^{d+1} \mid \|\mathbf{x}\|^2 = 1\}$  be the unit sphere in  $\mathbb{R}^{d+1}$ ,  $d \geq 2$ , where  $\|\cdot\|$  denotes the Euclidean norm, provided with the surface area measure  $\omega_d$ . The surface area of  $\mathbb{S}^d$  is denoted as  $|\mathbb{S}^d|$ . A spherical  $t$ -design [20] for a given positive integer  $t$  is a set of  $N$  points  $X_N = \{\mathbf{x}_1, \dots, \mathbf{x}_N\} \subset \mathbb{S}^d$  such that

$$(1.1) \quad \frac{1}{N} \sum_{j=1}^N p(\mathbf{x}_j) = \frac{1}{|\mathbb{S}^d|} \int_{\mathbb{S}^d} p(\mathbf{x}) d\omega_d(\mathbf{x})$$

holds for all spherical polynomials  $p$  with degree  $\leq t$ . For a numerical integration rule on the sphere, we say that the rule is “polynomially exact of degree  $\leq t$ ”, if

---

Received by the editor December 28, 2015, and, in revised form, November 9, 2016, April 26, 2017, and June 14, 2017.

2010 *Mathematics Subject Classification*. Primary 65D30, 41A10, 65G30.

*Key words and phrases*. Spherical  $t$ -designs, polynomial approximation, interval analysis, numerical integration,  $l_1$ -regularization.

The first author’s work was supported in part by Department of Applied Mathematics, The Hong Kong Polytechnic University and Hong Kong Research Council Grant PolyU5002/13p and in part by NSFC grant No. 11626147.

The second author’s work was supported in part by Hong Kong Research Council Grant PolyU153001/14p.

the rule integrates all spherical polynomials with degree  $\leq t$  exactly. A spherical  $t$ -design can be seen as the node set of a positive equal weight quadrature rule with polynomial exactness of degree  $\leq t$ , which is also proved to perform well for numerical integration of spherical functions belonging to Sobolev spaces. In [15] it is shown that a sequence of spherical  $t$ -designs with order  $t^d$  points has so-called “infinite strength”; i.e., the convergence rate of the worst-case error for integrating functions from a Sobolev space over  $\mathbb{S}^d$  with smoothness parameter  $s$  is optimal for every  $s > d/2$ .

Delsarte, Goethals, and Seidel [20] provided a lower bound for the number of points of a spherical  $t$ -design on  $\mathbb{S}^d$  of order  $t^d$ . For arbitrary degree  $t$ , Seymour and Zaslavsky [32] proved that there is always a spherical  $t$ -design with  $N$  points. Consequently, a natural problem is to find the minimal number  $N(d, t)$  of points, such that (1.1) holds for fixed  $t$  and  $d$ . In 1993, Korevaar and Meyers [26] proved that  $N(d, t) \leq C_d t^{(d^2+d)/2}$  and conjectured that  $N(d, t) \leq C_d t^d$  for a sufficiently large positive constant  $C_d$  depending only on  $d$ . This conjecture was then proved by Bondarenko, Radchenko and Viazaovska [10] in 2011. For  $d = 2$ , there is an even stronger conjecture by Hardin and Sloane [23] saying that  $N(2, t) \leq \frac{1}{2}t^2 + o(t^2)$  as  $t \rightarrow \infty$ . Numerical evidence supporting the conjecture was given in [23, 33]. Spherical  $t$ -designs have been extensively studied from various viewpoints, among which the application to polynomial approximation and the number of points needed to construct a spherical  $t$ -design have received great attention; see [2, 3, 5–8, 17, 18, 23, 26, 33]. Moreover, numerical methods have been developed for finding spherical  $t$ -designs. In some methods, the problem of finding a spherical  $t$ -design is reformulated as nonlinear equations or optimization problems; see [2, 21, 22, 33]. Numerical results in these papers suggest that these methods can find approximate spherical  $t$ -designs with high precision.

There are examples of explicit constructions of spherical  $t$ -designs for some specific  $t$ ; see, e.g., Bannai and Bannai [6]. The numerical methods of [17] yield a set of “interval” enclosures in the form of a collection of small spherical caps that is guaranteed to include a spherical  $t$ -design. However, the methods in [17] do not provide the exact locations of spherical  $t$ -designs in those interval enclosures. In general, for any given  $t$  the exact location of a spherical  $t$ -design is unknown. In most cases, the best we know is that there is a set of points  $\{\hat{\mathbf{x}}_1, \dots, \hat{\mathbf{x}}_N\}$  on the sphere  $\mathbb{S}^2$  such that a set of narrow intervals defined by

$$\mathbb{X}_N := \{[\mathbf{x}]_i = \mathcal{C}(\hat{\mathbf{x}}_i, \gamma_i), i = 1, \dots, N, \hat{\mathbf{x}}_i \in \mathbb{S}^2, \gamma_i > 0\} \subset \mathbb{S}^2$$

can be computed to contain a spherical  $t$ -design for  $N = (t + 1)^2$  and  $t \leq 100$  in [17], where

$$\mathcal{C}(\hat{\mathbf{x}}_i, \gamma_i) := \{\mathbf{x} \in \mathbb{S}^2 \mid \cos^{-1}(\mathbf{x} \cdot \hat{\mathbf{x}}_i) \leq \gamma_i\}.$$

Among all the spherical polynomials with degree  $\leq t$ , if the zero polynomial is the only one that vanishes at each point in the set  $X_N \subset \mathbb{S}^2$ , then the point set  $X_N \subset \mathbb{S}^2$  is said to be fundamental with order  $t$ . In 2011, Chen, Frommer and Lang [17] provided a computational-assisted proof for the existence of fundamental spherical  $t$ -designs on  $\mathbb{S}^2$  with  $N = (t + 1)^2$  for all values of  $t \leq 100$ . An interval arithmetic based algorithm is proposed in [17] to compute a sequence of sets of polar coordinates type interval enclosures containing fundamental spherical  $t$ -designs. By choosing the center point of each interval enclosure, an approximate spherical  $t$ -design can be obtained and numerical results show that the *Weyl sums* of these

point sets are very close to 0. In this paper we relax the equal-weight requirements of spherical  $t$ -designs to that the weights are chosen in an interval with respect to a number  $0 \leq \epsilon < 1$ , with whose average is  $\frac{|\mathbb{S}^d|}{N}$ .

**Definition 1.1** (Spherical  $t_\epsilon$ -design). A spherical  $t_\epsilon$ -design with  $0 \leq \epsilon < 1$  on  $\mathbb{S}^d$  is a set of points  $X_N^\epsilon := \{\mathbf{x}_1^\epsilon, \dots, \mathbf{x}_N^\epsilon\} \subset \mathbb{S}^d$ , such that the quadrature rule

$$(1.2) \quad \sum_{i=1}^N w_i p(\mathbf{x}_i^\epsilon) = \int_{\mathbb{S}^d} p(\mathbf{x}) d\omega_d(\mathbf{x})$$

is exact for all spherical polynomials  $p$  of degree at most  $t$ , with the weight vector  $\mathbf{w} = (w_1, \dots, w_N)^T$  satisfying

$$(1.3) \quad \frac{|\mathbb{S}^d|}{N}(1 - \epsilon) \leq w_i \leq \frac{|\mathbb{S}^d|}{N}(1 - \epsilon)^{-1}, \quad i = 1, \dots, N.$$

Spherical  $t_\epsilon$ -designs serve as a bridge between true spherical  $t$ -designs and positive weight quadrature rules.

*Remark 1.2.* A spherical  $t$ -design is a spherical  $t_0$ -design with  $\epsilon = 0$ . By letting  $p(\mathbf{x}) \equiv 1$  in (1.2) we obtain  $\sum_{i=1}^N w_i = |\mathbb{S}^d|$  and thus  $0 < w_i < |\mathbb{S}^d|$  for  $i = 1, \dots, N$ .

Since the existence of spherical  $t$ -designs has been proved for arbitrary  $t$ , and a spherical  $t$ -design is also a spherical  $t_\epsilon$ -design for arbitrary  $0 \leq \epsilon < 1$ , then we have the existence of spherical  $t_\epsilon$ -designs. Due to the relaxation of the equal-weight requirement, an important advantage is that we can have a positive weight quadrature rule exact for polynomials of degree  $\leq t$ . Therefore, a criteria for positive quadrature rules provided by Reimer in [29] called “*quadrature regularity*” should also be satisfied by spherical  $t_\epsilon$ -designs. Moreover, our numerical experiments show that with the increase of the value of  $\epsilon$  we can get polynomially exact rules by using fewer points than spherical  $t$ -designs.

The rest of this paper is organized as follows. In Section 2, we discuss the relationship between spherical  $t_\epsilon$ -designs and spherical  $t$ -designs when they are fundamental systems with the same number of points. Based on these results we study sets of interval enclosures containing fundamental spherical  $t$ -designs in Section 3. We prove that all point sets arbitrarily chosen in those sets of interval enclosures computed in [17] are spherical  $t_\epsilon$ -designs. In Section 4, we analyze the worst-case errors of spherical  $t_\epsilon$ -designs for numerical integration on the unit sphere  $\mathbb{S}^2$ . Numerical results show that the worst-case errors can be improved by the relaxation of the equal-weight requirement. In Section 5, we investigate an  $l_2 - l_1$  regularized weighted least squares model for polynomial approximation on the two-sphere using spherical  $t_\epsilon$ -designs and present numerical results to demonstrate the efficiency of the  $l_2 - l_1$  model.

In this paper we concentrate on the case  $d = 2$ . Throughout the paper we assume that all the points in a point set on the unit sphere are distinct. The computation is implemented in Matlab 2012b and done on a Lenovo Thinkcenter PC equipped with Intel Core i7-3770 3.4G Hz CPU, 8 GB RAM running Windows 7.

## 2. SPHERICAL $t_\epsilon$ -DESIGNS: NEIGHBORHOOD OF SPHERICAL $t$ -DESIGNS

In this section we will study the relationship between spherical  $t_\epsilon$ -designs and spherical  $t$ -designs when they are both fundamental systems and have the same number of points. A spherical  $t$ -design defines an equal-weight quadrature rule

which is polynomially exact of degree  $\leq t$  while a spherical  $t_\epsilon$ -design defines a positive-weight quadrature rule which is polynomially exact of degree  $\leq t$ . Based on these two properties, in this section we study a neighborhood of spherical  $t$ -designs. Let

$$\begin{aligned} \mathbb{P}_t &:= \mathbb{P}_t(\mathbb{S}^2) = \{\text{spherical polynomials of degree } \leq t \text{ on } \mathbb{S}^2\} \\ &= \text{span}\{Y_{\ell,k} : \ell = 0, \dots, t, k = 1, \dots, 2\ell + 1\} \end{aligned}$$

denote the space of all spherical polynomials on  $\mathbb{S}^2$  of degree  $\leq t$ . Here,  $Y_{\ell,k}$  is a fixed  $\mathbb{L}_2$ -orthonormal real spherical harmonic of degree  $\ell$  and order  $k$ , which means

$$(2.1) \quad \int_{\mathbb{S}^2} Y_{\ell,k}(\mathbf{x})Y_{\ell',k'}(\mathbf{x})d\omega(\mathbf{x}) = \delta_{\ell,\ell'}\delta_{k,k'}, \quad \ell, \ell' = 0, \dots, t; k, k' = 1, \dots, 2\ell + 1,$$

where  $d\omega(\mathbf{x}) = d\omega_2(\mathbf{x})$ , and  $\delta_{\ell,\ell'}$  is the Kronecker delta. It is well known that

$$(2.2) \quad d_t := \dim(\mathbb{P}_t) = \sum_{\ell=0}^t (2\ell + 1) = (t + 1)^2.$$

By the addition theorem [27]

$$(2.3) \quad \sum_{k=1}^{2\ell+1} Y_{\ell,k}(\mathbf{x})Y_{\ell,k}(\mathbf{y}) = \frac{2\ell + 1}{4\pi} P_\ell(\mathbf{x} \cdot \mathbf{y}),$$

which implies  $\sum_{k=1}^{2\ell+1} Y_{\ell,k}^2(\mathbf{x}) = \frac{2\ell+1}{4\pi}$ , where  $P_\ell$ ,  $\ell \geq 0$ , denotes the *Legendre Polynomial* normalized to  $P_\ell(0) = 1$  and  $\mathbf{x} \cdot \mathbf{y}$  denotes the Euclidean inner product, we obtain

$$(2.4) \quad \|Y_{\ell,k}\|_{C(\mathbb{S}^2)} \leq \max_{\mathbf{x} \in \mathbb{S}^2} \left( \sum_{k=1}^{2\ell+1} Y_{\ell,k}^2(\mathbf{x}) \right)^{\frac{1}{2}} = \sqrt{\frac{2\ell + 1}{4\pi}} \quad \text{for } k = 1, \dots, 2\ell + 1, \ell \geq 0.$$

Real spherical harmonics can be expressed in spherical coordinates as follows (see [1, 4]):

$$(2.5) \quad Y_{\ell,k}(\mathbf{x}) = \begin{cases} \sqrt{2}N_{\ell,k}P_\ell^{+1-k}(\cos \theta) \cos k\varphi, & k = 1, \dots, \ell, \\ N_{\ell,k}P_\ell^0(\cos \theta), & k = \ell + 1, \\ \sqrt{2}N_{\ell,k}P_\ell^{k-\ell-1}(\cos \theta) \sin k\varphi, & k = \ell + 2, \dots, 2\ell + 1, \end{cases}$$

with

$$\mathbf{x} = \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix},$$

where  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi < 2\pi$ ,  $P_\ell^k$  is the associated Legendre polynomial of degree  $\ell$  and order  $k$ , and  $N_{\ell,k}$  are the normalization coefficients

$$N_{\ell,k} = \sqrt{\frac{2\ell + 1}{4\pi} \frac{(\ell - |k - \ell - 1|)!}{(\ell + |k - \ell - 1|)!}}, \quad k = 1, \dots, 2\ell + 1.$$

When taking  $k = \ell + 1$  and  $\theta = 0$  we obtain  $Y_{\ell,\ell+1}(\mathbf{x}) \equiv \sqrt{\frac{2\ell+1}{4\pi}}$  with  $\mathbf{x} = (0, 0, 1)^T$ . Therefore, (2.4) is a sharp upper bound of  $\|Y_{\ell,k}\|_{C(\mathbb{S}^2)}$ . With the fact that all

spherical harmonics of degree  $\leq t$  form a basis of  $\mathbb{P}_t$  we have that a finite point set  $X_N = \{\mathbf{x}_1, \dots, \mathbf{x}_N\}$  is a spherical  $t$ -design if and only if the *Weyl sums* satisfy

$$(2.6) \quad \sum_{i=1}^N Y_{\ell,k}(\mathbf{x}_i) = 0, \quad k = 1, \dots, 2\ell + 1, \ell = 1, \dots, t;$$

see [20, 33] for details. Let  $X_N = \{\mathbf{x}_1, \dots, \mathbf{x}_N\} \subset \mathbb{S}^2$  and  $X'_N = \{\mathbf{x}'_1, \dots, \mathbf{x}'_N\} \subset \mathbb{S}^2$  be two point sets on the sphere. To describe the relationship among points and point sets, we introduce the following definitions of distances.

1. The *geodesic distance* between two points  $\mathbf{x}, \mathbf{y} \in \mathbb{S}^2$  is given by

$$\text{dist}(\mathbf{x}, \mathbf{y}) := \cos^{-1}(\mathbf{x} \cdot \mathbf{y}).$$

2. The *separation distance* of a point set  $X_N$  is given by

$$\rho(X_N) := \min_{i \neq j} \cos^{-1}(\mathbf{x}_i \cdot \mathbf{x}_j),$$

which represents the minimal geodesic distance between two different points in  $X_N$ .

3. The *least distance* from a point  $\mathbf{x} \in \mathbb{S}^2$  to a point set  $X_N$  is given by

$$\text{dist}(\mathbf{x}, X_N) := \min_{\mathbf{x}_i \in X_N} \text{dist}(\mathbf{x}, \mathbf{x}_i) = \min_{\mathbf{x}_i \in X_N} \cos^{-1}(\mathbf{x} \cdot \mathbf{x}_i),$$

which can be seen as the geodesic distance between  $\mathbf{x}$  and the nearest point in  $X_N$ .

4. The *Hausdorff distance* between two point sets  $X_N$  and  $X'_N$  is given by

$$(2.7) \quad \begin{aligned} \sigma(X_N, X'_N) &:= \max\left\{ \max_{\mathbf{x}'_i \in X'_N} \text{dist}(\mathbf{x}'_i, X_N), \max_{\mathbf{x}_i \in X_N} \text{dist}(\mathbf{x}_i, X'_N) \right\} \\ &= \max\left\{ \max_{\mathbf{x}'_i \in X'_N} \min_{\mathbf{x}_j \in X_N} \cos^{-1}(\mathbf{x}'_i \cdot \mathbf{x}_j), \max_{\mathbf{x}_j \in X_N} \min_{\mathbf{x}'_i \in X'_N} \cos^{-1}(\mathbf{x}'_i \cdot \mathbf{x}_j) \right\}. \end{aligned}$$

Note that  $\sigma(X_N, X'_N) = \sigma(X'_N, X_N)$  and  $\sigma(X_N, X'_N) = 0$  if and only if  $X_N = X'_N$ .

A *spherical cap*  $\mathcal{C}(\mathbf{x}, r)$  with center  $\mathbf{x} \in \mathbb{S}^2$  and radius  $r \in \mathbb{R}_+$  is defined as

$$(2.8) \quad \mathcal{C}(\mathbf{x}, r) := \{\mathbf{y} \in \mathbb{S}^2 \mid \cos^{-1}(\mathbf{x} \cdot \mathbf{y}) \leq r\}.$$

*Remark 2.1.* Let  $X_N$  and  $X'_N$  be two  $N$ -point sets on  $\mathbb{S}^2$  with  $\sigma(X_N, X'_N) < \frac{1}{2}\rho(X_N)$ , then there exists for each  $\mathbf{x} \in X_N$  a unique  $\mathbf{x}' \in X'_N \cap \mathcal{C}(\mathbf{x}, \frac{1}{2}\rho(X_N))$ . In this situation,  $\mathbf{x}'_i$  shall always denote the unique point in  $X'_N \cap \mathcal{C}(\mathbf{x}_i, \frac{1}{2}\rho(X_N))$  associated with  $\mathbf{x}_i \in X_N$ .

For a point set  $X_N = \{\mathbf{x}_1, \dots, \mathbf{x}_N\} \subset \mathbb{S}^2$ , we define the matrix  $\mathbf{Y}(X_N) \in \mathbb{R}^{N \times d_t}$  with elements

$$(2.9) \quad \mathbf{Y}_{i, \ell^2+k}(X_N) = Y_{\ell,k}(\mathbf{x}_i), \quad i = 1, \dots, N, \quad k = 1, \dots, 2\ell + 1, \quad \ell = 0, \dots, t.$$

Note that  $|\mathbb{S}^2| = 4\pi$ . In [16], the following characterization of a spherical  $t_\epsilon$ -design, equivalent with Definition 1.1, is given: a point set  $X_N^\epsilon := \{\mathbf{x}_1^\epsilon, \dots, \mathbf{x}_N^\epsilon\} \subset \mathbb{S}^2$  is a spherical  $t_\epsilon$ -design if and only if

$$(2.10) \quad \mathbf{Y}(X_N^\epsilon)^T \mathbf{w} - \sqrt{4\pi} \mathbf{e}_1 = \mathbf{0} \quad \text{and} \quad \frac{4\pi(1-\epsilon)}{N} \mathbf{e} \leq \mathbf{w} \leq \frac{4\pi(1-\epsilon)^{-1}}{N} \mathbf{e},$$

where  $\mathbf{e}_1 = (1, 0, \dots, 0)^T \in \mathbb{R}^{(L+1)^2}$  and  $\mathbf{e} = (1, \dots, 1)^T \in \mathbb{R}^N$ .

For two matrices constructed by two near enough point sets on  $\mathbb{S}^2$ , we have the following property.

**Proposition 2.2.** For any two point sets  $X_N, X'_N \subset \mathbb{S}^2$  satisfying  $\sigma(X_N, X'_N) < \frac{1}{2}\rho(X_N)$ , there always holds

$$\begin{aligned} \|(\mathbf{Y}(X_N) - \mathbf{Y}(X'_N))^T\|_\infty &= \|\mathbf{Y}(X_N) - \mathbf{Y}(X'_N)\|_1 \\ (2.11) \qquad \qquad \qquad &\leq N(t+1)\sqrt{\frac{2t+1}{4\pi}}\sigma(X_N, X'_N), \end{aligned}$$

where  $\mathbf{Y}(X_N), \mathbf{Y}(X'_N) \in \mathbb{R}^{N \times d_t}$  are matrices defined by (2.9) which depend on  $t$ .

*Proof.* For a point  $\mathbf{x}_i \in X_N$ , by Remark 2.1 we let  $\mathbf{x}'_i$  be the unique point located in  $\mathcal{C}(\mathbf{x}_i, \frac{1}{2}\rho(X_N)) \cap X'_N$ . Let  $Q_{\ell,k}$  be the restriction of  $Y_{\ell,k}$  on the great circle through these two points. Then  $Q_{\ell,k}$  is a trigonometric polynomial on the sphere and by *Bernstein's inequality* [11] we obtain

$$\begin{aligned} |Y_{\ell,k}(\mathbf{x}_i) - Y_{\ell,k}(\mathbf{x}'_i)| &= |Q_{\ell,k}(\mathbf{x}_i) - Q_{\ell,k}(\mathbf{x}'_i)| \\ &\leq \cos^{-1}(\mathbf{x}_i \cdot \mathbf{x}'_i) \sup |Q'_{\ell,k}| \\ &\leq \cos^{-1}(\mathbf{x}_i \cdot \mathbf{x}'_i)(t+1) \sup |Q_{\ell,k}| \\ &\leq \cos^{-1}(\mathbf{x}_i \cdot \mathbf{x}'_i)(t+1)\|Y_{\ell,k}\|_{C(\mathbb{S}^2)} \\ (2.12) \qquad \qquad \qquad &\leq \sigma(X_N, X'_N)(t+1)\sqrt{\frac{2\ell+1}{4\pi}}, \end{aligned}$$

where the last inequality is obtained by (2.4). Together with (2.11) and (2.12) we have

$$\begin{aligned} \|\mathbf{Y}(X_N) - \mathbf{Y}(X'_N)\|_1 &= \max_{0 \leq \ell \leq t, 1 \leq k \leq 2\ell+1} \sum_{j=1}^N |Y_{\ell,k}(\mathbf{x}_j) - Y_{\ell,k}(\mathbf{x}'_j)| \\ &\leq N(t+1)\sqrt{\frac{2t+1}{4\pi}}\sigma(X_N, X'_N). \quad \square \end{aligned}$$

Let  $X_N^0 = \{\mathbf{x}_1^0, \dots, \mathbf{x}_N^0\} \subset \mathbb{S}^2$  be a fundamental spherical  $t$ -design. Given a number  $\sigma^* \geq 0$ , denote the neighborhood of  $X_N^0$  with radius  $\sigma^*$  by

$$(2.13) \qquad \mathbb{C}(X_N^0, \sigma^*) := \{X_N \subset \mathbb{S}^2 : \sigma(X_N, X_N^0) \leq \sigma^*\}.$$

The following lemma indicates that any point set contained in a small enough neighborhood of a fundamental spherical  $t$ -design is a fundamental spherical  $t_\epsilon$ -design.

**Lemma 2.3.** Let  $X_N^0$  be a fundamental spherical  $t$ -design with  $N = (t+1)^2$ . Set  $\tau = \sqrt{\frac{2t+1}{4\pi}}(t+1)^3$ . Suppose

$$(2.14) \qquad \sigma^* < \frac{1}{2} \min \left( \frac{1}{\tau \|\mathbf{Y}(X_N^0)^{-1}\|_1}, \rho(X_N^0) \right).$$

Then any point set  $X_N \in \mathbb{C}(X_N^0, \sigma^*)$  is a fundamental spherical  $t_\epsilon$ -design with

$$(2.15) \qquad \frac{\tau\sigma^* \|\mathbf{Y}(X_N^0)^{-1}\|_1}{1 - \tau\sigma^* \|\mathbf{Y}(X_N^0)^{-1}\|_1} \leq \epsilon < 1.$$

*Proof.* It suffices to show that for any point set  $X_N \in \mathbb{C}(X_N^0, \sigma^*)$ , we have

$$(2.16) \qquad \mathbf{Y}(X_N)^T \mathbf{w} = \sqrt{4\pi} \mathbf{e}_1 \text{ and } \|\mathbf{w} - \frac{4\pi}{N} \mathbf{e}\|_\infty < \frac{4\pi}{N},$$

where  $\mathbf{Y}(X_N)$  is nonsingular.

From Proposition 2.2 for any point set  $X_N$  with  $\sigma(X_N, X_N^0) < \sigma^*$  we have

$$\|\mathbf{Y}(X_N^0)^{-1}\|_1 \|\mathbf{Y}(X_N^0) - \mathbf{Y}(X_N)\|_1 < \|\mathbf{Y}(X_N^0)^{-1}\|_1 \tau \sigma^* < 1.$$

Hence,  $\mathbf{Y}(X_N)$  is nonsingular; i.e.,  $X_N$  is a fundamental system with order  $t$ .

By the fact that  $X_N^0$  is a fundamental spherical  $t$ -design and (2.6) we have

$$(2.17) \quad \frac{4\pi}{N} \mathbf{Y}(X_N^0)^T \mathbf{e} = \sqrt{4\pi} \mathbf{e}_1.$$

The well-known perturbation theorem for linear systems in [36, Theorem 2.3.9, pp. 135] and

$$\|\mathbf{I} - \mathbf{Y}(X_N^0)^{-1} \mathbf{Y}(X_N)^T\|_\infty \leq \|\mathbf{Y}(X_N^0)^{-1}\|_1 \|\mathbf{Y}(X_N^0) - \mathbf{Y}(X_N)\|_1 < 1,$$

yields

$$(2.18) \quad \frac{\|\mathbf{w} - \frac{4\pi}{N} \mathbf{e}\|_\infty}{\|\frac{4\pi}{N} \mathbf{e}\|_\infty} \leq \frac{\|(\mathbf{Y}(X_N^0)^T)^{-1}\|_\infty \|(\mathbf{Y}(X_N) - \mathbf{Y}(X_N^0))^T\|_\infty}{1 - \|(\mathbf{Y}(X_N^0)^T)^{-1}\|_\infty \|(\mathbf{Y}(X_N) - \mathbf{Y}(X_N^0))^T\|_\infty}.$$

Therefore, we obtain

$$(2.19) \quad \begin{aligned} \|\mathbf{w} - \frac{4\pi}{N} \mathbf{e}\|_\infty &= \frac{4\pi}{N} \frac{\|\mathbf{w} - \frac{4\pi}{N} \mathbf{e}\|_\infty}{\|\frac{4\pi}{N} \mathbf{e}\|_\infty} \\ &\leq \frac{4\pi}{N} \frac{\|\mathbf{Y}(X_N^0)^{-1}\|_1 \|\mathbf{Y}(X_N^0) - \mathbf{Y}(X_N)\|_1}{1 - \|\mathbf{Y}(X_N^0) - \mathbf{Y}(X_N)\|_1 \|\mathbf{Y}(X_N^0)^{-1}\|_1} \end{aligned}$$

$$(2.20) \quad \leq \frac{4\pi}{N} \frac{\tau \sigma^* \|\mathbf{Y}(X_N^0)^{-1}\|_1}{1 - \tau \sigma^* \|\mathbf{Y}(X_N^0)^{-1}\|_1} < \frac{4\pi}{N}.$$

Hence, the vector  $\mathbf{w}$  is positive which implies that  $X_N$  is spherical  $t_\epsilon$ -design with  $\epsilon$  satisfying (2.15). □

Lemma 2.3 enables us to prove an upper bound on the radius of a neighborhood of a fundamental spherical  $t$ -design (see (2.13) for the definition of a neighborhood) so that every point set in this neighborhood is a spherical  $t_\epsilon$ -design.

**Corollary 2.4.** *Let  $X_N^0$  be a fundamental spherical  $t$ -design with order  $t$  and  $N = (t + 1)^2$ . For any  $0 \leq \epsilon < 1$ , if*

$$(2.21) \quad \sigma(X_N, X_N^0) < \min \left( \frac{1}{2} \rho(X_N^0), \frac{\epsilon}{\tau(1 + \epsilon) \|\mathbf{Y}(X_N^0)^{-1}\|_1} \right),$$

*then  $X_N$  is a fundamental spherical  $t_\epsilon$ -design.*

*Proof.* By (2.21) it can be derived that there holds

$$\sigma(X_N, X_N^0) < \frac{1}{2} \min \left( \frac{1}{\tau \|\mathbf{Y}(X_N^0)^{-1}\|_1}, \rho(X_N^0) \right).$$

Therefore, by Lemma 2.3 we have  $\mathbf{Y}(X_N)$  is nonsingular and

$$\|\mathbf{w} - \frac{4\pi}{N} \mathbf{e}\|_\infty \leq \frac{4\pi}{N} \frac{\tau \sigma(X_N, X_N^0) \|\mathbf{Y}(X_N^0)^{-1}\|_1}{1 - \tau \sigma(X_N, X_N^0) \|\mathbf{Y}(X_N^0)^{-1}\|_1} < \frac{4\pi}{N} \epsilon.$$

Hence, from (2.21), we derive

$$(1 - \epsilon) \frac{4\pi}{N} \mathbf{e} < \mathbf{w} < (1 + \epsilon) \frac{4\pi}{N} \mathbf{e} \leq \frac{4\pi(1 - \epsilon)^{-1}}{N} \mathbf{e}.$$

This completes the proof. □

3. INTERVAL ANALYSIS OF SPHERICAL  $t_\epsilon$ -DESIGNS

In this section we will study sets of interval enclosures containing fundamental spherical  $t_\epsilon$ -designs. In the last section we describe a neighborhood of a fundamental spherical  $t$ -design in which any point set is a fundamental spherical  $t_\epsilon$ -design. The paper [17] obtains interval enclosures that are guaranteed to contain a spherical  $t$ -design for up to 100, but the exact location of spherical  $t$ -designs cannot be obtained. In the following we will show that any point set in sets of the interval enclosures given in [17] is a fundamental spherical  $t_\epsilon$ -design. Let

$$(3.1) \quad \mathbb{X}_N := \{[\mathbf{x}]_i = \mathcal{C}(\hat{\mathbf{x}}_i, \gamma_i) \subset \mathbb{S}^2, i = 1, \dots, N\}$$

be a set of spherical caps, with  $\hat{\mathbf{x}}_i$  as the center point and  $\gamma_i$  as the geodesic radius. Additionally, let  $\hat{X}_N = \{\hat{\mathbf{x}}_1, \dots, \hat{\mathbf{x}}_N\} \subset \mathbb{S}^2$  be the set of center points. Define the radius of  $\mathbb{X}_N$  by

$$\text{rad}(\mathbb{X}_N) := \max_{1 \leq i \leq N} \gamma_i,$$

and the separation distance of  $\mathbb{X}_N$  by

$$\rho(\mathbb{X}_N) := \min_{\substack{i \neq j \\ \mathbf{x}_i \in [\mathbf{x}]_i, \mathbf{x}_j \in [\mathbf{x}]_j}} \text{dist}(\mathbf{x}_i, \mathbf{x}_j).$$

We say that  $\mathbb{X}_N$  is an interval enclosure of a point set  $X_N = \{\mathbf{x}_1, \dots, \mathbf{x}_N\}$ , denoted as  $X_N \in \mathbb{X}_N$ , if for any  $\mathbf{x} \in X_N$  there exists a unique  $[\mathbf{x}] \in \mathbb{X}_N$  such that  $\mathbf{x} \in [\mathbf{x}]$ . In this situation,  $[\mathbf{x}]_i$  shall always denote the interval containing the point  $\mathbf{x}_i$ .

**Assumption 3.1.** Let  $\mathbb{X}_N$  defined by (3.1) be a set of spherical caps. Assume that:

- (1) there exists a spherical  $t$ -design  $X_N^0 \in \mathbb{X}_N$ ;
- (2)  $\mathbf{Y}(\hat{X}_N)$  is nonsingular.

The assumption that  $\mathbf{Y}(\hat{X}_N)$  is nonsingular implies that  $\mathbf{Y}(\hat{X}_N)$  is a square matrix with  $N = (t+1)^2$ . In the following theorem we show that under Assumption 3.1 if  $\text{rad}(\mathbb{X}_N)$  is smaller than a certain number, then  $\mathbf{Y}(X_N)$  is nonsingular and (2.21) holds for any  $X_N \in \mathbb{X}_N$ .

**Theorem 3.2.** Set  $\tau := \sqrt{\frac{2t+1}{4\pi}}(t+1)^3$ . Under Assumption 3.1, any point set  $X_N \in \mathbb{X}_N$  is a fundamental spherical  $t_\epsilon$ -design with

$$(3.2) \quad \frac{2\tau \text{rad}(\mathbb{X}_N) \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1}{1 - 4\tau \text{rad}(\mathbb{X}_N) \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1} \leq \epsilon < 1,$$

if

$$(3.3) \quad \text{rad}(\mathbb{X}_N) < \min \left( \frac{1}{4} \rho(\mathbb{X}_N), \frac{\epsilon}{2(1+2\epsilon)\tau \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1} \right).$$

*Proof.* Any  $X_N \in \mathbb{X}_N$  satisfies

$$(3.4) \quad \rho(X_N) \geq \rho(\mathbb{X}_N).$$

Hence, for any two point sets  $X_N, X'_N \in \mathbb{X}_N$  we have

$$(3.5) \quad \sigma(X_N, X'_N) \leq \max_{1 \leq i \leq N} \max_{\mathbf{x}_i, \mathbf{y}_i \in [\mathbf{x}]_i} \text{dist}(\mathbf{x}_i, \mathbf{y}_i) = 2 \text{rad}(\mathbb{X}_N).$$



First we show that  $\mathbf{Y}(X_N)$  is nonsingular for any  $X_N \in \mathbb{X}_N$ . By Proposition 2.2 we have

$$\begin{aligned} \|\mathbf{I} - \mathbf{Y}(\hat{X}_N)^{-1}\mathbf{Y}(X_N)\|_1 &\leq \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1\|\mathbf{Y}(\hat{X}_N) - \mathbf{Y}(X_N)\|_1 \\ &\leq \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1\tau\sigma(\hat{X}_N, X_N) \\ &\leq 2\tau\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1\text{rad}(\mathbb{X}_N) < 1. \end{aligned}$$

Hence, all the point sets  $X_N \in \mathbb{X}_N$  including  $X_N^0$  are fundamental systems with order  $t$ . Moreover, from (3.3), (3.4), and (3.5), we have

$$\sigma(X_N, X_N^0) < \frac{1}{2}\rho(X_N^0).$$

By (3.3) we can also have

$$2(1 + \epsilon)\tau\text{rad}(\mathbb{X}_N)\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1 < \epsilon - 2\epsilon\tau\text{rad}(\mathbb{X}_N)\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1.$$

By Corollary 2.7 in [35, pp. 119] it can be concluded that for arbitrary  $X_N \in \mathbb{X}_N$  we have

$$(3.6) \quad \|\mathbf{Y}(X_N)^{-1}\|_1 \leq \frac{\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1}{1 - \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1\|\mathbf{Y}(\hat{X}_N) - \mathbf{Y}(X_N)\|_1}.$$

From (3.6) and Proposition 2.2 we have

$$\begin{aligned} \sigma(X_N, X_N^0) &\leq 2\text{rad}(\mathbb{X}_N) \\ &< \frac{\epsilon(1 - 2\tau\text{rad}(\mathbb{X}_N)\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1)}{(1 + \epsilon)\tau\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1} \\ &\leq \frac{\epsilon(1 - \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1\|\mathbf{Y}(\hat{X}_N) - \mathbf{Y}(X_N^0)\|_1)}{(1 + \epsilon)\tau\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1} \\ (3.7) \quad &\leq \frac{\epsilon}{(1 + \epsilon)\tau\|\mathbf{Y}(X_N^0)^{-1}\|_1}. \end{aligned}$$

By Corollary 2.4 we have that any point set  $X_N \in \mathbb{X}_N$  is a fundamental spherical  $t_\epsilon$ -design. Additionally, from (2.19) and (3.5) there holds

$$\begin{aligned} \|\mathbf{w} - \frac{4\pi}{N}\mathbf{e}\|_\infty &\leq \frac{4\pi}{N} \frac{\|\mathbf{Y}(X_N^0)^{-1}\|_1\|\mathbf{Y}(X_N^0) - \mathbf{Y}(X_N)\|_1}{1 - \|\mathbf{Y}(X_N^0)^{-1}\|_1\|\mathbf{Y}(X_N^0) - \mathbf{Y}(X_N)\|_1} \\ &\leq \frac{4\pi}{N} \frac{2\tau\text{rad}(\mathbb{X}_N)\|\mathbf{Y}(X_N^0)^{-1}\|_1}{1 - 2\tau\text{rad}(\mathbb{X}_N)\|\mathbf{Y}(X_N^0)^{-1}\|_1}. \end{aligned}$$

Together with (3.6) we have

$$\begin{aligned} &\frac{2\tau\text{rad}(\mathbb{X}_N)\|\mathbf{Y}(X_N^0)^{-1}\|_1}{1 - 2\tau\text{rad}(\mathbb{X}_N)\|\mathbf{Y}(X_N^0)^{-1}\|_1} \\ &\leq \frac{2\tau\text{rad}(\mathbb{X}_N)\frac{\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1}{1 - \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1\|\mathbf{Y}(\hat{X}_N) - \mathbf{Y}(X_N)\|_1}}{1 - 2\tau\text{rad}(\mathbb{X}_N)\frac{\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1}{1 - \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1\|\mathbf{Y}(\hat{X}_N) - \mathbf{Y}(X_N)\|_1}} \\ &\leq \frac{2\tau\text{rad}(\mathbb{X}_N)\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1}{1 - 4\tau\text{rad}(\mathbb{X}_N)\|\mathbf{Y}(\hat{X}_N)^{-1}\|_1}. \end{aligned}$$

Then we have that any point set  $X_N \in \mathbb{X}_N$  is a spherical  $t_\epsilon$ -design with

$$\frac{2\tau \operatorname{rad}(\mathbb{X}_N) \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1}{1 - 4\tau \operatorname{rad}(\mathbb{X}_N) \|\mathbf{Y}(\hat{X}_N)^{-1}\|_1} \leq \epsilon < 1,$$

subject to the assumptions on  $\operatorname{rad}(\mathbb{X}_N)$ . □

Theorem 3.2 proves that an arbitrarily chosen point set in a set of interval enclosures of a fundamental spherical  $t$ -design is a spherical  $t_\epsilon$ -design if  $\operatorname{rad}(\mathbb{X}_N)$  is smaller than a certain number. The interval enclosures discussed in the theorem are spherical-caps defined as  $[\mathbf{x}]_i = \mathcal{C}(\hat{\mathbf{x}}_i, \gamma_i) = \{\mathbf{x} \in \mathbb{S}^2 \mid \cos^{-1}(\mathbf{x} \cdot \hat{\mathbf{x}}_i) \leq \gamma_i\}$ . However, in practice, to reduce the spherical constraint of points and the dimension of variables, the spherical coordinate form of the points are preferable to compute the interval enclosures; see [17]. For a point  $\mathbf{x}_i \in X_N \subset \mathbb{S}^2$ , denote  $\theta_i, \varphi_i$  as its spherical coordinate. Then in [17] a sequence of intervals  $[\theta]_i = [\underline{\theta}_i, \bar{\theta}_i], [\varphi]_i = [\underline{\varphi}_i, \bar{\varphi}_i]$  are computed such that  $\mathbb{Z}_N = \{[\mathbf{z}]_1, \dots, [\mathbf{z}]_N\}$  is a set of interval enclosures of a *well-conditioned spherical  $t$ -design* [2], in which each element in  $\mathbb{Z}_N$  is defined by

$$(3.8) \quad [\mathbf{z}]_i := \begin{pmatrix} \sin([\theta]_i) \cos([\varphi]_i) \\ \sin([\theta]_i) \sin([\varphi]_i) \\ \cos([\theta]_i) \end{pmatrix}, \quad i = 1, \dots, N.$$

In this sense, different from the interval enclosures defined by the spherical caps, each interval enclosure computed in [17] is a rectangle  $[\theta]_i \times [\varphi]_i$ . Therefore, there remains a gap between real computation of interval enclosures of spherical  $t$ -designs and our analysis above. Naturally, a strategy to overcome this gap is that for each spherical rectangle in [17] we construct a spherical cap which is as small as possible to cover it. For the spherical rectangle  $[\theta]_i \times [\varphi]_i = [\underline{\theta}_i, \bar{\theta}_i] \times [\underline{\varphi}_i, \bar{\varphi}_i]$ , its four vertices can be written as

$$\begin{aligned} \mathbf{x}_{i,1} &:= \begin{pmatrix} \sin(\underline{\theta}_i) \cos(\underline{\varphi}_i) \\ \sin(\underline{\theta}_i) \sin(\underline{\varphi}_i) \\ \cos(\underline{\theta}_i) \end{pmatrix}, & \mathbf{x}_{i,2} &:= \begin{pmatrix} \sin(\underline{\theta}_i) \cos(\bar{\varphi}_i) \\ \sin(\underline{\theta}_i) \sin(\bar{\varphi}_i) \\ \cos(\underline{\theta}_i) \end{pmatrix}, \\ \mathbf{x}_{i,3} &:= \begin{pmatrix} \sin(\bar{\theta}_i) \cos(\bar{\varphi}_i) \\ \sin(\bar{\theta}_i) \sin(\bar{\varphi}_i) \\ \cos(\bar{\theta}_i) \end{pmatrix}, & \mathbf{x}_{i,4} &:= \begin{pmatrix} \sin(\bar{\theta}_i) \cos(\underline{\varphi}_i) \\ \sin(\bar{\theta}_i) \sin(\underline{\varphi}_i) \\ \cos(\bar{\theta}_i) \end{pmatrix}. \end{aligned}$$

It can be shown that there exists a point  $\hat{\mathbf{x}}_i$  defined by  $[\theta, \varphi] \in [\theta]_i \times [\varphi]_i$  satisfying

$$(3.9) \quad \operatorname{dist}(\hat{\mathbf{x}}_i, \mathbf{x}_{i,j}) = \operatorname{dist}(\hat{\mathbf{x}}_i, \mathbf{x}_{i,k}) \quad \text{for } j, k = 1, 2, 3, 4,$$

and  $[\mathbf{z}]_i \subseteq \mathcal{C}(\hat{\mathbf{x}}_i, \gamma_i)$  with  $\gamma_i = \operatorname{dist}(\hat{\mathbf{x}}_i, \mathbf{x}_{i,1})$ . However, computing such a point  $\hat{\mathbf{x}}_i$  is time-consuming and imports large round-off errors when the radii of interval enclosures are small. Instead of computing  $\hat{\mathbf{x}}_i$ , we investigate another strategy to compute the spherical caps to cover spherical rectangles which is coarser but more

practical. For a spherical coordinate interval  $[\mathbf{z}]_i$ , we use the center point of the interval  $[\theta_i, \bar{\theta}_i] \times [\varphi_i, \bar{\varphi}_i]$  to define a point as

$$(3.10) \quad \tilde{\mathbf{x}}_i := \begin{pmatrix} \sin(\frac{1}{2}(\bar{\theta}_i + \theta_i)) \cos(\frac{1}{2}(\bar{\varphi}_i + \varphi_i)) \\ \sin(\frac{1}{2}(\bar{\theta}_i + \theta_i)) \sin(\frac{1}{2}(\bar{\varphi}_i + \varphi_i)) \\ \cos(\frac{1}{2}(\bar{\theta}_i + \theta_i)) \end{pmatrix}.$$

Note that the spherical coordinates of  $\tilde{\mathbf{x}}_i$  give the center point of the interval  $[\theta]_i \times [\varphi]_i$  but itself is not necessary to be the center point of  $[\mathbf{z}]_i$  in the form of the spherical coordinate. Still, we have that

$$(3.11) \quad \text{dist}(\tilde{\mathbf{x}}_i, \mathbf{x}_{i,1}) = \text{dist}(\tilde{\mathbf{x}}_i, \mathbf{x}_{i,2}), \quad \text{dist}(\tilde{\mathbf{x}}_i, \mathbf{x}_{i,3}) = \text{dist}(\tilde{\mathbf{x}}_i, \mathbf{x}_{i,4}).$$

It is obvious to obtain that the distance between  $\tilde{\mathbf{x}}_i$  and any point in  $[\mathbf{z}]_i$  does not exceed the maximum of the four distances in (3.11). Therefore, if we let

$$(3.12) \quad \gamma_i = \max\{\text{dist}(\tilde{\mathbf{x}}_i, \mathbf{x}_{i,1}), \text{dist}(\tilde{\mathbf{x}}_i, \mathbf{x}_{i,3})\},$$

then we have

$$(3.13) \quad [\mathbf{z}]_i \subseteq \mathcal{C}(\tilde{\mathbf{x}}_i, \gamma_i).$$

Consequently, we call the set of spherical caps,

$$\tilde{\mathbb{X}}_N := \{\mathcal{C}(\tilde{\mathbf{x}}_i, \gamma_i), i = 1, \dots, N\},$$

a *cap-cover* of  $\mathbb{Z}_N$  with  $\tilde{\mathbf{x}}_i, \gamma_i$  defined in (3.10) and (3.12). Similarly to the set of spherical caps  $\mathbb{X}_N$ , we define the radius and separation distance of  $\mathbb{Z}_N$  by

$$(3.14) \quad \text{rad}(\mathbb{Z}_N) := \max_{1 \leq i \leq N} \{ \max\{\text{dist}(\tilde{\mathbf{x}}_i, \mathbf{x}_{i,1}), \text{dist}(\tilde{\mathbf{x}}_i, \mathbf{x}_{i,3})\} \} = \text{rad}(\tilde{\mathbb{X}}_N),$$

and

$$(3.15) \quad \rho(\mathbb{Z}_N) := \min_{\substack{i \neq j \\ 1 \leq i, j \leq N}} \{ \text{dist}(\tilde{\mathbf{x}}_i, \tilde{\mathbf{x}}_j) - \gamma_i - \gamma_j \} = \rho(\tilde{\mathbb{X}}_N).$$

Then we have the following corollary for the lower bound of  $\epsilon$  for  $\mathbb{Z}_N$ .

**Corollary 3.3.** *Set  $\tau := \sqrt{\frac{2t+1}{4\pi}}(t+1)^3$ . Let  $\mathbb{Z}_N$  be a set of spherical rectangle and its cap-cover  $\tilde{\mathbb{X}}_N$  satisfy Assumption 3.1. Then any point set  $X_N \in \mathbb{Z}_N$  is a fundamental spherical  $t_\epsilon$ -design with*

$$(3.16) \quad \frac{2\tau \text{rad}(\mathbb{Z}_N) \|\mathbf{Y}(\tilde{X}_N)^{-1}\|_1}{1 - 4\tau \text{rad}(\mathbb{Z}_N) \|\mathbf{Y}(\tilde{X}_N)^{-1}\|_1} \leq \epsilon < 1$$

if

$$(3.17) \quad \text{rad}(\mathbb{Z}_N) < \min \left( \frac{1}{4} \rho(\mathbb{Z}_N), \frac{\epsilon}{2(1+2\epsilon)\tau \|\mathbf{Y}(\tilde{X}_N)^{-1}\|_1} \right).$$

The proof of this corollary is similar to the proof of Theorem 3.2.

Based on Corollary 3.3, the lower bounds of  $\epsilon$ , e.g., the left-hand side of inequality (3.16), denoted by  $\epsilon$ , for sets of interval enclosures provided in [17] can be computed. The data containing the sets of interval enclosures for the parameterization of the spherical  $t$ -designs and relative programs can be downloaded from the website <http://www-ai.math.uni-wuppertal.de/SciComp/SphericalTDesigns>. The computational results are shown in Figure 3.1 and Table 3.1.




FIGURE 3.1.  $\epsilon$  computed by (3.19) for  $t = 2, \dots, 100$

TABLE 3.1. Information for sets of interval enclosures  $\mathbb{Z}_N$  for some selected  $t$

$t$	$\text{rad}(\mathbb{Z}_N)$	$\rho(\mathbb{Z}_N)$	$\epsilon$	$\tilde{\epsilon}$ for $\tilde{X}_N$
10	1.843454e-12	3.396362e-01	6.748890e-08	6.694645e-14
20	1.515848e-11	1.805783e-01	5.480524e-06	1.783018e-13
30	5.588085e-11	1.249714e-01	7.888659e-05	2.480238e-13
40	1.044163e-10	9.203055e-02	4.043164e-04	5.339063e-13
50	2.199182e-10	7.638945e-02	1.862348e-03	5.057066e-13
60	4.006638e-10	6.302748e-02	6.502352e-03	6.747935e-13
70	6.143914e-10	5.421869e-02	1.820130e-02	8.820722e-13
80	1.220430e-09	4.771142e-02	6.050880e-02	1.151368e-12
90	2.089473e-09	4.264961e-02	2.066649e-01	1.228462e-12
100	2.273791e-09	3.846343e-02	4.420562e-01	1.880540e-12

In Figure 3.1, we report the values of  $\epsilon$  defined by the left-hand side of inequality (3.16), for sets of interval enclosures computed in [17] for  $t = 2, \dots, 100$  (for  $t = 1$  we have known that the regular tetrahedron is a spherical  $t$ -design so that  $\epsilon = 0$  and we would not consider this case here). In this figure we also plot the function

$$(3.18) \quad y = 10^{-14.4}(t + 1)^{6.9}$$

to approximately estimate the curve traced out by results for  $\epsilon$  as a function of discrete  $t$ 's. From the figure we can conclude that the lower bound of  $\epsilon$  grows with the increase of  $t$  in an order about 6.9. Additionally, by (3.16) it is known that the norm of  $\mathbf{Y}(\tilde{X}_N)^{-1}$  is also very important in the process of estimating  $\epsilon$ . Fortunately, since the sets of interval enclosures computed in [17] seek to include well-conditioned spherical  $t$ -designs [2], the growth of lower bounds of  $\epsilon$  keeps stable for all the  $t$  considered here.

We also report some information about the sets of interval enclosures and their theoretical lower bounds of  $\epsilon$  for some selected  $t$  in Table 3.1. Not only  $\epsilon$  but also the radii and separation distances of  $\mathbb{X}_N$  are shown in the table. We can see that the radius of each interval enclosure is far smaller than their separation distance, which means that the assumptions in the above lemmas and theorems are satisfied.

The lower bound  $\epsilon$  is computed using the left-hand side of inequality (3.16), namely,

$$(3.19) \quad \epsilon = \frac{2\tau \operatorname{rad}(\mathbb{Z}_N) \|\mathbf{Y}(\tilde{X}_N)^{-1}\|_1}{1 - 4\tau \operatorname{rad}(\mathbb{Z}_N) \|\mathbf{Y}(\tilde{X}_N)^{-1}\|_1}.$$

This is a theoretical lower bound for any arbitrarily chosen point set  $X_N \in \mathbb{Z}_N$ .

The value of  $\tilde{\epsilon}$  for  $\tilde{X}_N$  in the last column of Table 3.1 is the minimum value of  $\epsilon$  satisfying (2.10) with  $X_N^\epsilon = \tilde{X}_N$ . In particular, we first choose the center point  $\tilde{X}_N$  of  $\mathbb{Z}_N$ . Next, we solve

$$\mathbf{Y}^T(\tilde{X}_N)\mathbf{w} = \sqrt{4\pi}\mathbf{e}_1$$

and let the unique solution be  $\tilde{\mathbf{w}}$ . If  $\frac{N}{4\pi}\tilde{\mathbf{w}}_i < 1$ , we set  $\tilde{\epsilon}_i = 1 - \frac{N}{4\pi}\tilde{\mathbf{w}}_i$ , otherwise,  $\tilde{\epsilon}_i = 1 - 1/(\frac{N}{4\pi}\tilde{\mathbf{w}}_i)$ . Let  $\tilde{\epsilon} = \max_i\{\tilde{\epsilon}_i\}$ . Since this value is only for a special point set  $\tilde{X}_N \in \mathbb{Z}_N$ , generally it is smaller than  $\epsilon$ .

As is shown in Table 3.1, the values of  $\tilde{\epsilon}$  are all very small positive numbers, and grow with increasing  $t$ . This means that  $\tilde{X}_N$  which is selected properly from the interval enclosures can be a spherical  $t_\epsilon$ -design with  $\tilde{\mathbf{w}} \approx \frac{4\pi}{N}e$ .

*Remark 3.4.* To obtain an approximate spherical  $t$ -design as accurate as possible, in [17] the radius of the set of interval enclosures  $\operatorname{rad}(\mathbb{Z}_N)$  is computed to a small scale around  $10^{-10}$ . With the introduction of the concept spherical  $t_\epsilon$ -designs, it has been shown that any point set selected in  $\mathbb{Z}_N$  is a fundamental spherical  $t_\epsilon$ -design if  $\operatorname{rad}(\mathbb{Z}_N) < 1/(6\tau \operatorname{rad}(\mathbb{Z}_N) \|\mathbf{Y}(\tilde{X}_N)^{-1}\|_1)$ , which implies  $\epsilon < 1$  in (3.19).

#### 4. WORST-CASE ERRORS OF QUADRATURE RULES USING SPHERICAL $t_\epsilon$ -DESIGNS

In this section we will investigate the worst-case errors for quadrature rules using spherical  $t_\epsilon$ -designs in Sobolev spaces which are finite-dimensional rotationally invariant subspaces of  $C(\mathbb{S}^2)$ . The bizonal reproducing kernel will be used in the analysis, which has been widely applied to analyze approximations on the sphere [14, 15, 24, 25, 37].

In [12, 13, 15], a method to compute the worst-case errors for positive equal-weight quadrature rules in Sobolev spaces has been developed. In this section, we intend to extend their method to nonequal but still positive weight quadrature rules and show the performance of spherical  $t_\epsilon$ -designs in numerical integration.

We follow the notations given in [15] with the following exception: the surface measure, denoted here by  $\omega$ , is not normalized. Denote the space of square integrable functions on  $\mathbb{S}^2$  by  $\mathbb{L}_2(\mathbb{S}^2)$ . Then it is a Hilbert space with the inner product

$$(4.1) \quad \langle f, g \rangle_{\mathbb{L}_2(\mathbb{S}^2)} = \int_{\mathbb{S}^2} f(\mathbf{x})g(\mathbf{x})d\omega(\mathbf{x}), \quad f, g \in \mathbb{L}_2(\mathbb{S}^2),$$

and the induced norm as

$$(4.2) \quad \|f\|_{\mathbb{L}_2(\mathbb{S}^2)} = \left( \int_{\mathbb{S}^2} |f(\mathbf{x})|^2 d\omega(\mathbf{x}) \right)^{\frac{1}{2}}, \quad f \in \mathbb{L}_2(\mathbb{S}^2).$$

The Sobolev space  $\mathbb{H}^s(\mathbb{S}^2)$  can be defined for  $s \geq 0$  as the set of all functions  $f \in \mathbb{L}_2(\mathbb{S}^2)$  whose Laplace-Fourier coefficients,

$$(4.3) \quad \hat{f}_{\ell,k} = \langle f, Y_{\ell,k} \rangle_{\mathbb{L}_2(\mathbb{S}^2)} = \int_{\mathbb{S}^2} f(\mathbf{x}) Y_{\ell,k}(\mathbf{x}) d\omega(\mathbf{x}),$$

satisfy

$$(4.4) \quad \sum_{\ell=0}^{\infty} \sum_{k=1}^{2\ell+1} (1 + \lambda_{\ell})^s \left| \hat{f}_{\ell,k} \right|^2 < \infty,$$

where  $\lambda_{\ell} = \ell(\ell + 1)$ . Obviously, by letting  $s = 0$  we can obtain  $\mathbb{H}^0(\mathbb{S}^2) = \mathbb{L}_2(\mathbb{S}^2)$ . Then the norm of  $\mathbb{H}^s(\mathbb{S}^2)$  can be defined as

$$(4.5) \quad \|f\|_{\mathbb{H}^s} = \left[ \sum_{\ell=0}^{\infty} \sum_{k=1}^{2\ell+1} \frac{1}{\alpha_{\ell}^{(s)}} \hat{f}_{\ell,k}^2 \right]^{\frac{1}{2}},$$

where the positive parameters  $\alpha_{\ell}^{(s)}$  satisfy

$$(4.6) \quad \alpha_{\ell}^{(s)} \asymp (1 + \lambda_{\ell})^{-s} \asymp (\ell + 1)^{-2s}.$$

Here we say that  $a_n \asymp b_n$  if there exist positive constants  $c_1$  and  $c_2$  independent of  $n$  such that  $c_1 a_n \leq b_n \leq c_2 a_n$  for all  $n$ . Correspondingly, the inner product of  $\mathbb{H}^s(\mathbb{S}^2)$  can be defined as

$$(4.7) \quad \langle f, g \rangle_{\mathbb{H}^s} = \sum_{\ell=0}^{\infty} \sum_{k=1}^{2\ell+1} \frac{1}{\alpha_{\ell}^{(s)}} \hat{f}_{\ell,k} \hat{g}_{\ell,k}.$$

For a point set  $X_N$  and a weight vector  $\mathbf{w}$ , we define the numerical quadrature rule and the integral of a function  $f$  on  $\mathbb{S}^2$  as

$$(4.8) \quad Q[X_N, \mathbf{w}](f) := \sum_{j=1}^N \frac{w_j}{4\pi} f(\mathbf{x}_j), \quad I(f) := \frac{1}{4\pi} \int_{\mathbb{S}^2} f(\mathbf{x}) d\omega(\mathbf{x}),$$

The worst-case error of the quadrature rule  $Q[X_N, \mathbf{w}]$  on  $\mathbb{H}^s(\mathbb{S}^2)$  can be defined as [15, 24]

$$(4.9) \quad E_s(Q[X_N, \mathbf{w}]) := \sup \{ |Q[X_N, \mathbf{w}](f) - I(f)| : f \in \mathbb{H}^s(\mathbb{S}^2), \|f\|_{\mathbb{H}^s} \leq 1 \}.$$

The Riesz representation theorem and the additional theorem assure the existence of a reproducing kernel of the form

$$(4.10) \quad \begin{aligned} K_s(\mathbf{x}, \mathbf{y}) &= \sum_{\ell=0}^{\infty} (2\ell + 1) \alpha_{\ell}^{(s)} P_{\ell}(\mathbf{x} \cdot \mathbf{y}) \\ &= \sum_{\ell=0}^{\infty} \sum_{k=1}^{2\ell+1} \alpha_{\ell}^{(s)} Y_{\ell,k}(\mathbf{x}) Y_{\ell,k}(\mathbf{y}). \end{aligned}$$

Together with the property of reproducing kernel  $K_s(\cdot, \cdot)$  defined in (4.10) and the addition theorem, it is shown in [15, 24] that

$$\begin{aligned} (E_s(Q[X_N, \mathbf{w}]))^2 &= \left[ \sup_{\substack{f \in \mathbb{H}^s(\mathbb{S}^2) \\ \|f\|_{\mathbb{H}^s} \leq 1}} |Q[X_N, \mathbf{w}](f) - I(f)| \right]^2 \\ &= \left\| \sum_{i=1}^N \frac{w_i}{4\pi} K_s(\cdot, \mathbf{x}_i) - \int_{\mathbb{S}^2} K_s(\cdot, \mathbf{x}) d\omega(\mathbf{x}) \right\|_{\mathbb{H}^s}^2. \end{aligned}$$

Then with the following equality [15],

$$\int_{\mathbb{S}^2} K_s(\mathbf{x}, \cdot) d\omega(\mathbf{x}) = \alpha_0^{(s)},$$

the worst-case error could be reformulated as

$$\begin{aligned} (E_s(Q[X_N, \mathbf{w}]))^2 &= \left[ \sum_{\ell=1}^{\infty} \sum_{k=1}^{2\ell+1} \alpha_\ell^{(s)} \left( \sum_{i=1}^N \frac{w_i}{4\pi} Y_{\ell,k}(\mathbf{x}_i) \right)^2 \right] \\ (4.11) \quad &= \sum_{i=1}^N \sum_{j=1}^N \frac{w_i w_j}{16\pi^2} \sum_{\ell=1}^{\infty} \sum_{k=1}^{2\ell+1} \alpha_\ell^{(s)} Y_{\ell,k}(\mathbf{x}_i) Y_{\ell,k}(\mathbf{x}_j). \end{aligned}$$

Reproducing kernels for  $\mathbb{H}^s(\mathbb{S}^2)$  for  $s > 1$  can be constructed utilizing powers of distances, provided the power  $2s - 2$  is not an even integer. Indeed, it is known (cf., e.g., [9, 14]) that the signed power of the distance, with sign  $(-1)^{J+1}$  with  $J := J(s) := \lfloor s - 1 \rfloor$ , has the following Laplace-Fourier expansion:

$$(4.12) \quad (-1)^{J+1} |\mathbf{x} - \mathbf{y}|^{2s-2} = (-1)^{J+1} V_{2-2s}(\mathbb{S}^2) + \sum_{\ell=1}^{\infty} a_\ell^{(s)} (2\ell + 1) P_\ell(\mathbf{x} \cdot \mathbf{y}),$$

where

$$(4.13) \quad V_{2-2s}(\mathbb{S}^2) := \int_{\mathbb{S}^2} \int_{\mathbb{S}^2} |\mathbf{x} - \mathbf{y}|^{2s-2} d\omega(\mathbf{x}) d\omega(\mathbf{y}) = 2^{2s-1} \frac{\Gamma(3/2)\Gamma(s)}{\sqrt{\pi}\Gamma(1+s)},$$

$$(4.14) \quad a_\ell^{(s)} := V_{2-2s}(\mathbb{S}^2) \frac{(-1)^{J+1} (1-s)_\ell}{(1+s)_\ell}, \quad \ell \geq 1,$$

and

$$\frac{(1-s)_\ell}{(1+s)_\ell} := \frac{\Gamma(1+s)\Gamma(\ell+1-s)}{\Gamma(1-s)\Gamma(\ell+1+s)} \sim \frac{\Gamma(1-s)}{\Gamma(1+s)} \ell^{-2s} \sim \ell^{-2s}, \quad \text{as } \ell \rightarrow \infty.$$

Thus, we have

$$(4.15) \quad (-1)^{J+2} (V_{2-2s}(\mathbb{S}^2) - |\mathbf{x} - \mathbf{y}|^{2s-2}) = \sum_{\ell=1}^{\infty} a_\ell^{(s)} (2\ell + 1) P_\ell(\mathbf{x} \cdot \mathbf{y})$$

$$(4.16) \quad = \sum_{\ell=1}^{\infty} a_\ell^{(s)} \sum_{k=1}^{2\ell+1} Y_{\ell,k}(\mathbf{x}) Y_{\ell,k}(\mathbf{y}).$$

Note that for  $a_\ell^{(s)}$  we have

$$(4.17) \quad a_\ell^{(s)} \sim 2^{2s-1} \frac{\Gamma(\frac{3}{2})\Gamma(s)}{\sqrt{\pi}(-1)^{J+1}\Gamma(1+s)} \ell^{-2s} \quad \text{as } \ell \rightarrow \infty,$$

and when  $1 < s < 2$ , which means  $J = J(s) = 0$ , we have  $a_\ell^{(s)} > 0$  for all  $\ell = 1, \dots$ . Therefore, we regard the left-hand side of (4.15) as the reproducing kernel of  $\mathbb{H}^s(\mathbb{S}^2)$ , which is

$$K_s(\mathbf{x}, \mathbf{y}) = V_{2-2s}(\mathbb{S}^2) - |\mathbf{x} - \mathbf{y}|^{2s-2},$$

and then we obtain

$$(4.18) \quad (E_s(Q[X_N, \mathbf{w}]))^2 = \sum_{i=1}^N \sum_{j=1}^N \frac{w_i w_j}{16\pi^2} (V_{2-2s}(\mathbb{S}^2) - |\mathbf{x}_i - \mathbf{x}_j|^{2s-2}).$$

For the case  $s > 2$  and  $s$  is not an integer, we know that  $a_\ell^{(s)} > 0$  does not hold for all  $\ell = 1, \dots$ . In this situation, we let

$$K_s(\mathbf{x}, \mathbf{y}) = (1 - (-1)^{J+1})V_{2-2s}(\mathbb{S}^2) + \mathcal{Q}_J(\mathbf{x} \cdot \mathbf{y}) + (-1)^{J+1}|\mathbf{x} - \mathbf{y}|^{2s-2},$$

with

$$\mathcal{Q}_J(\mathbf{x} \cdot \mathbf{y}) := \sum_{\ell=1}^J ((-1)^{J+1-\ell} - 1) a_\ell^{(s)} (2\ell + 1) P_\ell(\mathbf{x} \cdot \mathbf{y}), \quad \mathbf{x}, \mathbf{y} \in \mathbb{S}^2,$$

which changes the signs of the negative coefficients  $a_\ell^{(s)}$  in (4.12). Hence the worst-case error on  $\mathbb{H}^s(\mathbb{S}^2)$  with  $s > 2$  can be represented as

$$(4.19) \quad (E_s(Q[X_N, \mathbf{w}]))^2 = \sum_{i=1}^N \sum_{j=1}^N \frac{w_i w_j}{16\pi^2} (\mathcal{Q}_J(\mathbf{x}_i \cdot \mathbf{x}_j) + (-1)^{J+1}|\mathbf{x}_i - \mathbf{x}_j|^{2s-2} - (-1)^{J+1}V_{2-2s}(\mathbb{S}^2)).$$

For quadrature rules using spherical  $t_\epsilon$ -designs as node sets, we have the following theorem.

**Theorem 4.1.** *Given  $s > 1$  and  $s$  is not an integer, assume that  $Q[X_N^\epsilon, \mathbf{w}]$  is polynomially exact of degree  $\leq t$ , where  $X_N^\epsilon$  is a spherical  $t_\epsilon$ -design with a weight vector  $\mathbf{w}$ . Then*

$$(4.20) \quad E_s(Q[X_N, \mathbf{w}]) = \left[ (-1)^{J+1} \sum_{i=1}^N \sum_{j=1}^N \frac{w_i w_j}{16\pi^2} (|\mathbf{x}_i - \mathbf{x}_j|^{2s-2} - V_{2-2s}(\mathbb{S}^2)) \right]^{\frac{1}{2}}.$$

*Proof.* By (2.3) it can be concluded that  $\sum_{i=1}^N \sum_{j=1}^N \frac{w_i w_j}{16\pi^2} \mathcal{Q}_J(\mathbf{x}_i \cdot \mathbf{x}_j)$  will vanish in (4.19). Then (4.20) can be directly deduced by (4.18) and (4.19).  $\square$

In what follows we will compute the worst-case errors of quadrature rules using spherical  $t_\epsilon$ -designs by using (4.20). In this experiment we choose  $\epsilon = 0.1$  for spherical  $t_\epsilon$ -designs and use (2.10) to find a spherical  $t_\epsilon$ -design, which is a system of nonlinear equations. The system can be solved by minimizing its least squares form using a smoothing trust-region filter method proposed in [16]. Note that the number of points needed for constructing spherical  $t_\epsilon$ -designs may decrease as  $\epsilon$  gets larger. Thus in the computation of spherical  $t_\epsilon$ -designs we always attempt to find the one with a possible minimal number of points, denoted as  $N(t, \epsilon)$ . The detailed process for finding spherical  $t_\epsilon$ -designs can be found in [16]. In the numerical test of computation of spherical  $t_\epsilon$ -designs it is found that a possible minimal number of points satisfies  $\lceil (t+1)^2/3 \rceil + 1 \leq N(t, \epsilon) \leq \lceil (t+2)^2/2 \rceil + 1$ . In the numerical test in the current and next sections, the spherical  $t_\epsilon$ -designs are chosen with  $N = N(t, \epsilon)$ .






FIGURE 4.1. Worst-case errors for spherical  $t_{0,1}$ -designs and spherical  $t$ -designs

The worst-case errors of quadrature rules using spherical  $t_{0,1}$ -designs in  $\mathbb{H}^s(\mathbb{S}^2)$  for  $s = 1.5$  are illustrated in Figure 4.1(a). For comparison, the worst-case errors for quadrature rules using approximate spherical  $t$ -designs computed in [33] will also be implemented. For all spherical  $t_{0,1}$ -designs, the worst-case error is calculated using (4.20) and the distance kernel, and for spherical  $t$ -designs the worst-case errors are calculated by formulas (42) and (46) in [15]. From the figure we can see that in this case, the computed worst-case errors of approximate spherical  $t$ -designs and spherical  $t_{0,1}$ -designs essentially lie on the same curve, which remains as a conjecture that the worst-case errors of both spherical  $t$ -designs and spherical  $t_\epsilon$ -designs decay in the same speed with respect to the number of points in the case  $1 < s < 2$  on  $\mathbb{S}^2$ . Figure 4.1(b) plots the worst-case errors for both spherical  $t$ -designs and spherical  $t_{0,1}$ -designs with  $s = 5.5$ . The optimal rate of convergence for the worst-case error for integrating functions from a Sobolev space of smoothness index  $s$  is known. The worst-case error of quasi-Monte Carlo methods defined by a sequence of spherical  $t$ -designs with order  $t^2$  points on  $\mathbb{S}^2$  achieves this optimal rate [15]. Assuming that the approximated spherical  $t$ -designs used in Figure 4.1(b) have optimal order number of points, the lower curve for the  $t_{0,1}$ -designs may indicate a better constant but not a better rate of convergence. There is perhaps insufficient data as one might need to go to a higher number of points to see the optimal rate of convergence. However, one should also take into account that for higher smoothness index  $s$  (as it is the case here) the numerics is much more delicate than for lower values of  $s$ .

## 5. POLYNOMIAL APPROXIMATION ON THE SPHERE USING SPHERICAL $t_\epsilon$ -DESIGNS

**5.1. Regularized weighted least squares approximation using spherical  $t_\epsilon$ -designs.** In this section we consider the restoration of a continuous function  $f \in C(\mathbb{S}^2)$  from its noisy values  $f^\delta$  given at  $N$  points  $X_N = \{\mathbf{x}_1, \dots, \mathbf{x}_N\} \subset \mathbb{S}^2$  by the  $l_2 - l_1$  regularized weighted discrete least squares form

$$(5.1) \quad \min_{\alpha_{\ell,k} \in \mathbb{R}} \frac{1}{2} \sum_{j=1}^N \mu_j \left( \sum_{\ell=0}^L \sum_{k=1}^{2k+1} \alpha_{\ell,k} Y_{\ell,k}(\mathbf{x}_j) - f^\delta(\mathbf{x}_j) \right)^2 + \lambda \sum_{\ell=0}^L \sum_{k=1}^{2\ell+1} |\beta_{\ell,k} \alpha_{\ell,k}|,$$

where  $\mu_j > 0, j = 1, \dots, N$  are the weights for each term of the least squares model,  $\lambda > 0$  is the regularization parameter, and  $\beta_{\ell,k} \geq 0, \ell = 0, \dots, L, k = 1, \dots, 2\ell + 1$  are usually chosen with the meaning of certain polynomial operators such as Laplace-Beltrami operator and filtered operator [3, 34]. In [28] both a priori choice based physical reason in satellite gravity gradiometry problem and a posteriori choice based on reproducing kernel theory are considered to choose  $\beta_{\ell,k}$ .

Note that  $\{Y_{\ell,k}, \ell = 0, \dots, L, k = 1, \dots, 2\ell + 1\}$  is a basis of  $\mathbb{P}_L$ . Problem (5.1) is to find a good approximation of  $f$  in  $\mathbb{P}_L$  in the form

$$p_{L,N}(\mathbf{x}) := \sum_{\ell=0}^L \sum_{k=1}^{2\ell+1} \alpha_{\ell,k} Y_{\ell,k}(\mathbf{x}).$$

Let the entries of matrix  $\mathbf{Y}_L \in \mathbb{R}^{N \times (L+1)^2}$  be

$$(\mathbf{Y}_L)_{i,\ell^2+k} = Y_{\ell,k}(\mathbf{x}_i), \quad i = 1, \dots, N, \ell = 0, \dots, L, k = 1, \dots, 2\ell + 1,$$

and  $\mathbf{f}^\delta = (f^\delta(\mathbf{x}_1), \dots, f^\delta(\mathbf{x}_N))^T$ . Problem (5.1) can be reformulated as

$$(5.2) \quad \min_{\alpha \in \mathbb{R}^{(L+1)^2}} \frac{1}{2} \|\mathbf{\Lambda}^{\frac{1}{2}}(\mathbf{Y}_L \alpha - \mathbf{f}^\delta)\|_2^2 + \lambda \|\mathbf{D} \alpha\|_1,$$

where

$$\mathbf{\Lambda} = \begin{bmatrix} \mu_1 & & \\ & \ddots & \\ & & \mu_N \end{bmatrix} \in \mathbb{R}^{N \times N},$$

and  $\mathbf{D}$  is a diagonal matrix satisfying  $\mathbf{D}_{\ell^2+k,\ell^2+k} = \beta_{\ell,k}$  with  $\beta_{\ell,k} \geq 0$ . For polynomial approximation on the sphere, an  $l_2$ -regularized weighted least squares model has also been considered [3, 28]:

$$(5.3) \quad \min_{\alpha \in \mathbb{R}^{(L+1)^2}} \frac{1}{2} \|\mathbf{\Lambda}^{\frac{1}{2}}(\mathbf{Y}_L \alpha - \mathbf{f}^\delta)\|_2^2 + \lambda \|\mathbf{D} \alpha\|_2^2.$$

The regularization of this model is of  $l_2$  norm, which can be seen as a measure of energy. It is known that the  $l_1$  regularization has desirable properties in approximation of nonsmooth continuous functions. An  $l_1$  regularization term is preferable to be considered here. By choosing a suitable penalization term, the  $l_2 - l_1$  regularized model is usually supposed to achieve a more sparse solution than the  $l_2 - l_2$  regularized one, which means that the target function is approximated by fewer basis spherical polynomials. Additionally, for functions which are globally continuous but locally nondifferentiable on the sphere, the  $l_2 - l_1$  regularization is better than the  $l_2 - l_2$  regularization.

**Theorem 5.1.** *Let  $X_N^\epsilon$  be a spherical  $t_\epsilon$ -design with  $t \geq 2L$  and let  $\mathbf{w}$  be the vector of weights satisfying (1.3) and (1.2) with respect to  $X_N^\epsilon$ . For model (5.2) set  $\mu_j = w_j$  for  $j = 1, \dots, N$ . Then*

$$(5.4) \quad \mathbf{H}_L := \mathbf{Y}_L^T \mathbf{\Lambda} \mathbf{Y}_L = \mathbf{I}_{(L+1)^2},$$

and (5.2) has the unique solution

$$(5.5) \quad \alpha_{\ell,k} = \max\{0, s_{\ell,k} - \lambda \beta_{\ell,k}\} + \min\{0, s_{\ell,k} + \lambda \beta_{\ell,k}\},$$

for  $\ell = 0, \dots, L, k = 1, \dots, 2k + 1$ , where  $s_{\ell,k} = \sum_{i=1}^N w_i Y_{\ell,k}(\mathbf{x}_i) f^\delta(\mathbf{x}_i)$ .

*Proof.* Note that when  $X_N^\epsilon$  is a spherical  $t_\epsilon$ -design,

$$\begin{aligned}
 (\mathbf{H}_L)_{\ell^2+k,(\ell')^2+k'} &= \sum_{i=1}^N w_i Y_{\ell,k}(\mathbf{x}_i) Y_{\ell',k'}(\mathbf{x}_i) \\
 (5.6) \qquad \qquad \qquad &= \int_{\mathbb{S}^2} Y_{\ell,k}(\mathbf{x}) Y_{\ell',k'}(\mathbf{x}) d\omega(\mathbf{x}) = \delta_{\ell\ell'} \delta_{kk'},
 \end{aligned}$$

where the third equality is established by the orthonormality of spherical harmonics. Problem (5.2) is strictly convex by the fact that  $\mathbf{H}_L$  is nonsingular and so it has a unique optimal solution. Since  $A(\alpha) = \frac{1}{2} \|\mathbf{A}^{\frac{1}{2}}(\mathbf{Y}_L \alpha - \mathbf{f}^\delta)\|_2^2$  is strictly differentiable, by deriving the first optimality condition of (5.2) and Corollary 1 in [19, Section 2.3], we obtain that its unique optimal solution satisfies

$$(5.7) \qquad \qquad \qquad 0 \in \mathbf{H}_L \alpha - \mathbf{Y}_L^T \mathbf{W} \mathbf{f}^\delta + \lambda \partial(\|\mathbf{D} \alpha\|_1),$$

where  $\partial(\cdot)$  denotes the subdifferential. By (5.6) which implies  $\mathbf{H}_L = \mathbf{I}_{(L+1)^2}$  and the fact that  $\mathbf{D}$  is diagonal, problem (5.7) is separable and thus  $\alpha$  is a solution of (5.7) if and only if it is a solution of

$$(5.8) \qquad 0 \in \alpha_{\ell,k} - s_{\ell,k} + \lambda \beta_{\ell,k} \partial|\alpha_{\ell,k}|, \quad \ell = 0, \dots, L, k = 1, \dots, 2\ell + 1.$$

Denote  $\tau_{\ell,k} = \partial|\alpha_{\ell,k}|$  and hence  $-1 \leq \tau_{\ell,k} \leq 1$ . Let  $\alpha_{\ell,k}^*$  be the optimal solution of (5.8) with corresponding  $\ell$  and  $k$  and hence

$$(5.9) \qquad \qquad \qquad \alpha_{\ell,k}^* = s_{\ell,k} - \lambda \beta_{\ell,k} \tau_{\ell,k} \quad \text{with } \tau_{\ell,k} \in [-1, 1].$$

When  $s_{\ell,k} > \lambda \beta_{\ell,k}$  we can set  $\tau_{\ell,k} = 1$  and obtain

$$\alpha_{\ell,k}^* = s_{\ell,k} - \lambda \beta_{\ell,k} > 0,$$

which together with  $\beta_{\ell,k} \geq 0$  satisfies (5.5) and (5.9). When  $s_{\ell,k} < -\lambda \beta_{\ell,k}$  similarly we set  $\tau_{\ell,k} = -1$  and get

$$\alpha_{\ell,k}^* = s_{\ell,k} + \lambda \beta_{\ell,k} < 0,$$

which also satisfies (5.5) and (5.9). Then when  $s_{\ell,k} \in [-\lambda \beta_{\ell,k}, \lambda \beta_{\ell,k}]$  we set  $\tau_{\ell,k} = \frac{s_{\ell,k}}{\lambda \beta_{\ell,k}} \in [-1, 1]$  and get that

$$\alpha_{\ell,k}^* = 0,$$

which also satisfies (5.5) and (5.9). Hence, the theorem is proved. □

Denote the approximation residual as  $A(\alpha) = \sum_{j=1}^N (p_{L,N}(\mathbf{x}_j) - f^\delta(\mathbf{x}_j))^2$ . Let  $\alpha^*(\lambda)$  be the optimal solution of (5.2) with different regularized parameters  $\lambda$ . The following proposition indicates that  $A(\alpha^*(\lambda))$  is monotonically increasing with respect to  $\lambda$ .

**Proposition 5.2.** *Let  $X_N^\epsilon$  be a spherical  $t_\epsilon$ -design with  $t \geq 2L$  and  $\mu_j = w_j$  for  $j = 1, \dots, N$ . Then  $A(\alpha^*(\lambda))$  is increasing in  $\lambda$ .*

*Proof.* Let  $\lambda, \tilde{\lambda}$  be given with  $0 < \lambda \leq \tilde{\lambda}$  and denote the optimal solution of problem (5.2) with  $\lambda, \tilde{\lambda}$  as  $\alpha^*, \tilde{\alpha}^*$ , respectively. Denote  $E(\lambda, \alpha) = \lambda \|\mathbf{D} \alpha\|_1$  and the minimization property of (5.2) for  $\lambda$  gives

$$(5.10) \qquad \qquad \qquad A(\alpha^*) + E(\lambda, \alpha^*) \leq A(\tilde{\alpha}^*) + E(\lambda, \tilde{\alpha}^*),$$

which implies that

$$(5.11) \qquad \qquad \qquad A(\alpha^*) - A(\tilde{\alpha}^*) \leq E(\lambda, \tilde{\alpha}^*) - E(\lambda, \alpha^*).$$

From (5.5) we have

$$(5.12) \quad \alpha_{\ell,k}^* = \begin{cases} s_{\ell,k} - \lambda\beta_{\ell,k}, & s_{\ell,k} > \lambda\beta_{\ell,k}, \\ s_{\ell,k} + \lambda\beta_{\ell,k}, & s_{\ell,k} < -\lambda\beta_{\ell,k}, \\ 0, & -\lambda\beta_{\ell,k} \leq s_{\ell,k} \leq \lambda\beta_{\ell,k}. \end{cases}$$

Since  $\lambda\beta_{\ell,k} \geq 0$ , we have  $|\alpha_{\ell,k}^*| = \max(0, |s_{\ell,k}| - \lambda\beta_{\ell,k})$ . Together with the fact that

$$E(\lambda, \alpha^*) = \lambda \sum_{\ell=0}^L \sum_{k=1}^{2\ell+1} \beta_{\ell,k} |\alpha_{\ell,k}^*|,$$

we have

$$|\tilde{\alpha}_{\ell,k}^*| = \max(0, |s_{\ell,k}| - \tilde{\lambda}\beta_{\ell,k}) \leq \max(0, |s_{\ell,k}| - \lambda\beta_{\ell,k}) = |\alpha_{\ell,k}^*|.$$

Hence, it is obtained that  $E(\lambda, \tilde{\alpha}^*) \leq E(\lambda, \alpha^*)$ . Together with (5.10) we complete the proof. □

**5.2. Numerical experiments.** In this subsection we report the numerical results to test the efficiency of the  $l_2 - l_1$  regularized model (5.2) using spherical  $t_\epsilon$ -designs.

**Example 5.1.** In the first numerical test, the target function is selected as spherical polynomials with degree no higher than  $L$ . Obviously using both models (5.2) and (5.3) the target function can be exactly restored when  $\lambda = 0$  and the data  $\mathbf{f}^\delta$  is noise free and the optimal values of the two models equal to 0 in such case. However, due to the noise in the data vector  $\mathbf{f}^\delta$ , it is necessary to use the regularization models.

In this experiment we will use the spherical  $t_{0.1}$ -designs which is calculated by solving a system of nonlinear equation (2.10) and the approximate spherical  $t$ -designs proposed in [33] as the point set for polynomial approximation. Both the uniform errors and  $\mathbb{L}_2$  errors are recorded to measure the approximation quality. We choose a large-scaled and well-distributed point set  $X_t \subset \mathbb{S}^2$  to be the test set and use it to estimate the errors. Then the uniform error and  $\mathbb{L}_2$  error of the approximation are estimated by


$$(5.13) \quad \|f - p_{L,N}\|_{C(\mathbb{S}^2)} \approx \max_{\mathbf{x}_i \in X_t} |f(\mathbf{x}_i) - p_{L,N}(\mathbf{x}_i)|$$

and


$$(5.14) \quad \|f - p_{L,N}\|_{\mathbb{L}_2} \approx \left( \frac{4\pi}{N_t} \sum_{i=1}^{N_t} (f(\mathbf{x}_i) - p_{L,N}(\mathbf{x}_i))^2 \right)^{\frac{1}{2}},$$

where  $N_t$  denotes the number of points  $\mathbf{x}_i$  in  $X_t$ . In this experiment, we choose  $X_t$  to be an equal area partitioning point set [31] with  $10^5$  points. The entries of the matrix  $\mathbf{D}$  in the experiment is always selected as  $\beta_{\ell,k} = \ell(\ell + 1)$  for  $\ell = 0, \dots, L, k = 2\ell + 1$ , inspired by the Laplace-Beltrami operator; see [3].


Figure 5.1 shows the approximation errors using both  $l_2 - l_1$  model (5.2) and  $l_2 - l_2$  model (5.3) with different  $\lambda$  and different noise scales  $\delta$ . The noise of the data  $\mathbf{f}^\delta$  obeys a uniform distribution in  $[-\delta, \delta]$ . In this numerical experiment a spherical  $37_{0.1}$ -design with only 514 points, which is much fewer than  $\lceil \frac{(t+1)^2}{2} \rceil$ , is applied to approximate a randomly generated spherical polynomial with degree  $\lfloor \frac{37}{2} \rfloor = 18$  (the polynomial is generated with all of its Fourier coefficients obeying the standard normal distribution). The regularization parameter  $\lambda$  is chosen from  $10^{-20}$  to  $10^{0.5}$ . Figure 5.1(a)(b) give the errors of the approximation with different  $\lambda$  for  $\delta = 0.1$  using the two models. From the two subfigures it can be seen that




(a) Uniform errors with different  $\lambda$  for  $\delta = 0.1$



(b)  $L_2$  errors with different  $\lambda$  for  $\delta = 0.1$



(c) Minimal uniform errors with different noise scales  $\delta$



(d) Minimal  $L_2$  errors with different noise scales  $\delta$

FIGURE 5.1. Errors for restoring 18-degree polynomials

model (5.2) can restore the 18-degree polynomial more accurately than model (5.3). The minimal error with respect to different  $\lambda$  can be achieved at about  $\lambda = 10^{-6}$ . Figure 5.1(c)(d) show the errors of the restoration results with different noise scales. It can be seen that the model (5.2) performs better in each noise scale than (5.3).

**Example 5.2.** In the second numerical experiment we test the numerical performance of model (5.2) using spherical  $t_\epsilon$ -designs and spherical  $t$ -designs. We select the Franke function [30],

$$\begin{aligned}
 f_1(\mathbf{x}) := f(x, y, z) = & 0.75 \exp(-(9x - 2)^2/4 - (9y - 2)^2/4 - (9z - 2)^2/4) \\
 & + 0.75 \exp(-(9x + 1)^2/49 - (9y + 1)/10 - (9z + 1)/10) \\
 & + 0.5 \exp(-(9x - 7)^2/4 - (9y - 3)^2/4 - (9z - 5)^2/4) \\
 & - 0.2 \exp(-(9x - 4)^2 - (9y - 7)^2 - (9z - 5)^2), \quad (x, y, z)^T \in \mathbb{S}^2,
 \end{aligned}
 \tag{5.15}$$

to be the target function which is not a spherical polynomial but continuously differentiable on the whole sphere. We set  $\epsilon = 0.1$  and also  $\delta = 0.1$  in this experiment and the scheme of choosing  $\lambda$  is the same as in Example 5.1. For spherical  $t_{0.1}$ -designs, we select those point sets constructed with possible least points. As is mentioned above, a spherical  $t_{0.1}$ -design may be constructed using less than  $\lceil \frac{(t+1)^2}{2} \rceil$  points. Approximate spherical  $t$ -designs proposed in [33] are also applied for comparison.

Note that the minimizer of model (5.2) has an explicit form (5.5) only when  $t \geq 2L$ , so for different  $t$  we choose  $L = \lfloor \frac{t}{2} \rfloor$ .




FIGURE 5.2. Errors for approximating Franke function with different scales of point sets

For Example 5.2, the approximation errors using both spherical  $t_{0,1}$ -designs and approximate spherical  $t$ -designs are shown in Figure 5.2. The X-axis represents the number of points in the data sets and the Y-axis represents the minimal uniform errors. From the figure we can see that approximation using spherical  $t_{0,1}$ -designs achieves smaller errors than using approximate spherical  $t$ -designs in most cases. Based on the numerical results in Figure 5.2, the approximation quality can be improved with the relaxation of weight vector using model (5.2).

**Example 5.3.** In the third experiment, a continuous but nondifferentiable function

$$(5.16) \quad f_2 = f_1(\mathbf{x}) + f_{\text{cap}}(\mathbf{x}),$$

is selected as the target function to approximate, with

$$(5.17) \quad f_{\text{cap}}(\mathbf{x}) = \begin{cases} \rho \cos\left(\frac{\pi \cos^{-1}(\mathbf{x}_c \cdot \mathbf{x})}{2r}\right), & \mathbf{x} \in \mathcal{C}(\mathbf{x}_c, r), \\ 0, & \text{otherwise,} \end{cases}$$

where  $\rho > 0$ ,  $0 < r < \pi$ . The function is nondifferentiable at the edge of the spherical cap  $\mathcal{C}(\mathbf{x}_c, r)$ . Since the basis functions applied for approximation are spherical harmonic polynomials which is globally differentiable on  $\mathbb{S}^2$ , restoration of the edge of  $\mathcal{C}(\mathbf{x}_c, r)$  turns to be a challenging problem when the data has noise.

A spherical  $37_{0,1}$ -design with 514 points is used as the data point set in this experiment. Other settings in this experiment are  $\delta = 0.5$ ,  $\lambda = 10^{-20}, 10^{-19.5}, \dots, 10^5$ ,  $\mathbf{x}_c = (-0.5, -0.5, \sqrt{0.5})^T$ ,  $r = 0.5$  and  $\rho = 1$ . The restorations of  $f_2$  using both models (5.2) and (5.3) are depicted in Figure 5.3. Similar with previous experiments, we choose the values of  $\lambda$  resulting in minimal uniform errors for each model and plot the shape of the restoration function on the sphere. From Figure 5.3(c)(d)(e)(f), restoration by model (5.2) is not as smooth as restoration by model (5.3) but has smaller errors. And more notably, as highlighted by the rectangle in Figure 5.3(c)(d), model (5.2) restores the nonsmooth edges of the spherical cap more accurately than model (5.3).




FIGURE 5.3. Restoration of  $f_2$  using spherical  $37_{0.1}$ -design with models (5.2) and (5.3)

ACKNOWLEDGMENTS

We would like to thank Professor Ian Sloan and the two referees for their valuable and helpful comments on spherical  $t_\epsilon$ -designs.

## REFERENCES

- [1] M. Abramowitz and I. A. Stegun (eds.), *Handbook of Mathematical Functions with Formulas, graphs, and mathematical tables*, Dover Publications, Inc., New York, 1992. Reprint of the 1972 edition. MR1225604
- [2] C. An, X. Chen, I. H. Sloan, and R. S. Womersley, *Well conditioned spherical designs for integration and interpolation on the two-sphere*, SIAM J. Numer. Anal. **48** (2010), no. 6, 2135–2157, DOI 10.1137/100795140. MR2763659
- [3] C. An, X. Chen, I. H. Sloan, and R. S. Womersley, *Regularized least squares approximations on the sphere using spherical designs*, SIAM J. Numer. Anal. **50** (2012), no. 3, 1513–1534, DOI 10.1137/110838601. MR2970753
- [4] K. Atkinson and W. Han, *Spherical harmonics and approximations on the unit sphere: an introduction*, Lecture Notes in Mathematics, vol. 2044, Springer, Heidelberg, 2012. MR2934227
- [5] B. Bajnok, *Construction of spherical  $t$ -designs*, Geom. Dedicata **43** (1992), no. 2, 167–179, DOI 10.1007/BF00147866. MR1180648
- [6] E. Bannai, *On tight spherical designs*, J. Combin. Theory Ser. A **26** (1979), no. 1, 38–47, DOI 10.1016/0097-3165(79)90052-9. MR525085
- [7] E. Bannai and E. Bannai, *A survey on spherical designs and algebraic combinatorics on spheres*, European J. Combin. **30** (2009), no. 6, 1392–1425, DOI 10.1016/j.ejc.2008.11.007. MR2535394
- [8] E. Bannai and E. Bannai, *Remarks on the concepts of  $t$ -designs*, J. Appl. Math. Comput. **40** (2012), no. 1-2, 195–207, DOI 10.1007/s12190-012-0544-1. MR2965326
- [9] B. J. C. Baxter and S. Hubbert, *Radial basis functions for the sphere*, Recent progress in multivariate approximation (Witten-Bommerholz, 2000), Internat. Ser. Numer. Math., vol. 137, Birkhäuser, Basel, 2001, pp. 33–47. MR1877496
- [10] A. Bondarenko, D. Radchenko, and M. Viazovska, *Optimal asymptotic bounds for spherical designs*, Ann. of Math. (2) **178** (2013), no. 2, 443–452, DOI 10.4007/annals.2013.178.2.2. MR3071504
- [11] P. Borwein and T. Erdélyi, *Polynomials and Polynomial Inequalities*, Graduate Texts in Mathematics, vol. 161, Springer-Verlag, New York, 1995. MR1367960
- [12] J. S. Brauchart and J. Dick, *A simple proof of Stolarsky’s invariance principle*, Proc. Amer. Math. Soc. **141** (2013), no. 6, 2085–2096, DOI 10.1090/S0002-9939-2013-11490-5. MR3034434
- [13] J. S. Brauchart and J. Dick, *A characterization of Sobolev spaces on the sphere and an extension of Stolarsky’s invariance principle to arbitrary smoothness*, Constr. Approx. **38** (2013), no. 3, 397–445, DOI 10.1007/s00365-013-9217-z. MR3122277
- [14] J. S. Brauchart and K. Hesse, *Numerical integration over spheres of arbitrary dimension*, Constr. Approx. **25** (2007), no. 1, 41–71, DOI 10.1007/s00365-006-0629-4. MR2263736
- [15] J. S. Brauchart, E. B. Saff, I. H. Sloan, and R. S. Womersley, *QMC designs: optimal order quasi Monte Carlo integration schemes on the sphere*, Math. Comp. **83** (2014), no. 290, 2821–2851, DOI 10.1090/S0025-5718-2014-02839-1. MR3246811
- [16] X. Chen, S. Du, and Y. Zhou, *A smoothing trust region filter algorithm for nonsmooth least squares problems*, Sci. China Math. **59** (2016), no. 5, 999–1014, DOI 10.1007/s11425-015-5116-z. MR3484496
- [17] X. Chen, A. Frommer, and B. Lang, *Computational existence proofs for spherical  $t$ -designs*, Numer. Math. **117** (2011), no. 2, 289–305, DOI 10.1007/s00211-010-0332-5. MR2754852
- [18] X. Chen and R. S. Womersley, *Existence of solutions to systems of underdetermined equations and spherical designs*, SIAM J. Numer. Anal. **44** (2006), no. 6, 2326–2341, DOI 10.1137/050626636. MR2272596
- [19] F. H. Clarke, *Optimization and Nonsmooth Analysis*, 2nd ed., Classics in Applied Mathematics, vol. 5, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1990. MR1058436
- [20] P. Delsarte, J. M. Goethals, and J. J. Seidel, *Spherical codes and designs*, Geometriae Dedicata **6** (1977), no. 3, 363–388. MR0485471
- [21] P. J. Grabner and R. F. Tichy, *Spherical designs, discrepancy and numerical integration*, Math. Comp. **60** (1993), no. 201, 327–336, DOI 10.2307/2153170. MR1155573



- [22] M. Gräf and D. Potts, *On the computation of spherical designs by a new optimization approach based on fast spherical Fourier transforms*, Numer. Math. **119** (2011), no. 4, 699–724, DOI 10.1007/s00211-011-0399-7. MR2854125
- [23] R. H. Hardin and N. J. A. Sloane, *McLaren's improved snub cube and other new spherical designs in three dimensions*, Discrete Comput. Geom. **15** (1996), no. 4, 429–441, DOI 10.1007/BF02711518. MR1384885
- [24] K. Hesse and I. H. Sloan, *Worst-case errors in a Sobolev space setting for cubature over the sphere  $S^2$* , Bull. Austral. Math. Soc. **71** (2005), no. 1, 81–105, DOI 10.1017/S0004972700038041. MR2127668
- [25] K. Hesse and I. H. Sloan, *Cubature over the sphere  $S^2$  in Sobolev spaces of arbitrary order*, J. Approx. Theory **141** (2006), no. 2, 118–133, DOI 10.1016/j.jat.2006.01.004. MR2252093
- [26] J. Korevaar and J. L. H. Meyers, *Spherical Faraday cage for the case of equal point charges and Chebyshev-type quadrature on the sphere*, Integral Transform. Spec. Funct. **1** (1993), no. 2, 105–117, DOI 10.1080/10652469308819013. MR1421438
- [27] C. Müller, *Spherical Harmonics*, Lecture Notes in Mathematics, vol. 17, Springer-Verlag, Berlin-New York, 1966. MR0199449
- [28] S. V. Pereverzyev, I. H. Sloan, and P. Tkachenko, *Parameter choice strategies for least-squares approximation of noisy smooth functions on the sphere*, SIAM J. Numer. Anal. **53** (2015), no. 2, 820–835, DOI 10.1137/140964990. MR3324977
- [29] M. Reimer, *Quadrature rules for the surface integral of the unit sphere based on extremal fundamental systems*, Math. Nachr. **169** (1994), 235–241, DOI 10.1002/mana.19941690117. MR1292809
- [30] R. J. Renka, *Multivariate interpolation of large sets of scattered data*, ACM Trans. Math. Software **14** (1988), no. 2, 139–148, DOI 10.1145/45054.45055. MR946761
- [31] E. B. Saff and A. B. J. Kuijlaars, *Distributing many points on a sphere*, Math. Intelligencer **19** (1997), no. 1, 5–11, DOI 10.1007/BF03024331. MR1439152
- [32] P. D. Seymour and T. Zaslavsky, *Averaging sets: a generalization of mean values and spherical designs*, Adv. in Math. **52** (1984), no. 3, 213–240, DOI 10.1016/0001-8708(84)90022-7. MR744857
- [33] I. H. Sloan and R. S. Womersley, *A variational characterisation of spherical designs*, J. Approx. Theory **159** (2009), no. 2, 308–318, DOI 10.1016/j.jat.2009.02.014. MR2562747
- [34] I. H. Sloan and R. S. Womersley, *Filtered hyperinterpolation: a constructive polynomial approximation on the sphere*, GEM Int. J. Geomath. **3** (2012), no. 1, 95–117, DOI 10.1007/s13137-011-0029-7. MR2915950
- [35] G. W. Stewart and J. G. Sun, *Matrix Perturbation Theory*, Computer Science and Scientific Computing, Academic Press, Inc., Boston, MA, 1990. MR1061154
- [36] D. S. Watkins, *Fundamentals of Matrix Computations*, 3rd ed., Pure and Applied Mathematics (Hoboken), John Wiley & Sons, Inc., Hoboken, NJ, 2010. MR2778339
- [37] H. Wendland, *Scattered Data Approximation*, Cambridge Monographs on Applied and Computational Mathematics, vol. 17, Cambridge University Press, Cambridge, 2005. MR2131724

SCHOOL OF MATHEMATICS AND STATISTICS, SHANDONG NORMAL UNIVERSITY, JINAN, SHANGDONG, CHINA 250000

*Email address:* andres.zhou@connect.polyu.hk

DEPARTMENT OF APPLIED MATHEMATICS, THE HONG KONG POLYTECHNIC UNIVERSITY, HUNG HOM, HONG KONG

*Email address:* maxjchen@polyu.edu.hk