

1019-13-207

**C-Y. Jean Chan\*** ([chan@math.purdue.edu](mailto:chan@math.purdue.edu)) and **Claudia Miller**. *Local cohomology and the constant term of a Hilbert-Samuel polynomial*. Preliminary report.

Let  $(A, \mathfrak{m})$  be a regular local ring. Let  $P_M(t)$  be the Hilbert-Samuel polynomial of a finitely generated  $A$ -module  $M$  such that  $P_M(n) = \ell(M/\mathfrak{m}^n M)$  for  $n \gg 0$ . The *Rees module*  $R_{\mathfrak{m}}(M) = \bigoplus_{n \geq 0} \mathfrak{m}^n M$  of  $M$  is a graded module over the Rees algebra of  $A$ .

We will prove that the difference between the Hilbert function and the Hilbert-Samuel polynomial of  $M$ , at any non-negative integer, is the alternating sum of the length of the graded pieces of the local cohomology of  $R_{\mathfrak{m}}(M)$ . This proof is an application of a well-known formula due to Serre and is modified from the work of Johnston and Verma whose results are mainly on  $\mathfrak{m}$ -primary ideals.

By the Grothendieck-Riemann-Roch formula for the projective spaces, we relate the Hilbert-Samuel polynomial  $P_M(t)$  to the Chern characters of the graded Rees module  $R_{\mathfrak{m}}(M)$  up to a constant term. The constant term of  $P_M(t)$  can be recovered using the graded pieces of the local cohomology of  $R_{\mathfrak{m}}(M)$ . (Received August 15, 2006)