

1041-55-98

**Ethan S. Devinatz\*** (devinatz@math.washington.edu), Department of Mathematics, Box 354350, University of Washington, Seattle, WA. *Finiteness properties of certain homotopy fixed point spectra.*

Let  $G$  be a closed subgroup of the  $n$ th Morava stabilizer group  $S_n$ , and let  $E_n^{hG}$  denote the continuous homotopy fixed point spectrum of Devinatz and Hopkins. We conjecture that, for "most"  $G$ ,  $\pi_*(E_n^{hG} \wedge X)$  is of essentially finite rank—this will be defined—whenever  $X$  is a  $K(n-2)_*$ -acyclic finite spectrum annihilated by  $p$ . This implies in particular that  $\pi_*(L_{K(n)}X)$  is of essentially finite rank. We show that this conjecture is true when  $G$  is a topologically cyclic group whose generator is non-torsion in the quotient of the  $p$ -Sylow subgroup of  $S_n$  by its center. We also show that any appropriate  $G$  in  $S_2$  contains an open subnormal subgroup  $U$  such that  $\pi_*(E_2^{hU} \wedge M)$  is of essentially finite rank, where  $M$  is the mod  $(p)$  Moore spectrum. Finally, we indicate a strategy for establishing the conjecture in general. (Received August 11, 2008)