

1050-20-59

Paul-Hermann Zieschang* (zieschang@utb.edu), Department of Mathematics, University of Texas at Brownsville, Brownsville, TX 78520. *On association schemes with thin thin residue.*

Let S be an association scheme. The smallest closed subset of S with thin quotient scheme is called the thin residue of S and is denoted by $O^\theta(S)$; cf. [2]. We shall look at schemes the thin residue of which is thin. If $O^\theta(S)$ is thin, $O^\theta(S)$ can be viewed as a group. In [1] it was shown that S is schurian if the set of all normal subgroups of $O^\theta(S)$ is linearly ordered. As a consequence, one obtains that S is schurian if $O^\theta(S)$ is a finite simple group. In [3], it is shown that S is schurian if $O^\theta(S)$ the direct product of two simple groups of different order. In my talk, I will discuss the main idea of the proof of this latter result. The emphasis will be on ideas how to generalize this result.

[1] Hirasaka, M. and Zieschang, P.-H.: Sufficient conditions for a scheme to originate from a group, *J. Combin. Theory Ser. A* **104**, 17-27 (2003)

[2] Zieschang, P.-H.: *Theory of Association Schemes*. Springer Monographs in Mathematics, Berlin Heidelberg New York (2005)

[3] Zieschang, P.-H.: On association schemes with thin thin residue, *J. Algebra*, to appear (Received February 22, 2009)