

Groupoids: Unifying Internal and External Symmetry

A Tour through Some Examples

Alan Weinstein

Introduction

Mathematicians tend to think of the notion of symmetry as being virtually synonymous with the theory of *groups* and their actions, perhaps largely because of the well-known Erlanger program of F. Klein and the related work of S. Lie, which virtually defined geometric structures by their groups of automorphisms. (See, for example, Yaglom's account in [27].) In fact, though groups are indeed sufficient to characterize homogeneous structures, there are plenty of objects which exhibit what we clearly recognize as symmetry, but which admit few or no nontrivial automorphisms. It turns out that the symmetry, and hence much of the structure, of such objects can be characterized algebraically if we use *groupoids* and not just groups.

The aim of this paper is to explain, mostly through examples, what groupoids are and how they describe symmetry. We will begin with elementary examples, with discrete symmetry, and end with examples in the differentiable setting which involve *Lie groupoids* and their corresponding infinitesimal objects, *Lie algebroids*. These objects play a role in the study of general differentiable manifolds and partial differential equations, which is, in a sense, an extension of the role which Lie groups play in the geometry and analysis of highly symmetric manifolds.

Some History

The following historical remarks are not intended to be complete but merely to indicate the breadth of areas where groupoids have been used. An extensive survey of groupoids as of 1986 can be found in R. Brown's article [2].

Alan Weinstein is professor of mathematics at the University of California at Berkeley. His e-mail address is a1anw@math.berkeley.edu.

Research partially supported by NSF Grant DMS-93-09653. This article is based on a talk given first at UCB (the University of California at Berkeley) in April 1995 during Mathematics Awareness Week, the theme of which was Mathematics and Symmetry. I would like to thank the Department of Mathematics of another UCB (the University of Colorado at Boulder) for an invitation to Boulder in November 1995, where a similar talk was most recently given and where the first draft of this manuscript was largely written in the downtown cafés. Thanks should also be given to a third UCB (Université Claude-Bernard, Lyon), where I started my intensive work on groupoids in the summer of 1986. George Bergman, Ronnie Brown, Pierre Dazord, Jack Feldman, Kirill Mackenzie, George Mackey, Saunders Mac Lane, Jean Pradines, Arlan Ramsay, Sasha Voronov, and the referees made helpful comments on the original manuscript. Finally, I would like to thank Paul Brown for preparing the figures.—A. Weinstein

Groupoids were introduced (and named) by H. Brandt [1] in a 1926 paper on the composition of quadratic forms in four variables. C. Ehresmann [6] added further structures (topological and differentiable as well as algebraic) to groupoids, thereby introducing them as a tool in differentiable topology and geometry.

In algebraic geometry, A. Grothendieck used groupoids extensively and, in particular, showed how they could be used to tame the unruly equivalence relations which arise in the construction of moduli spaces. (See [8], with further details in [7].) As the role of moduli spaces expands in physics as well as mathematics, groupoids continue to play an essential role.

In analysis, G. Mackey [16] used groupoids under the name of *virtual groups* to allow the treatment of ergodic actions of groups, “as if” they were transitive, while it was observed that the convolution operation, extended from groups to groupoids, made possible the construction of a multitude of interesting noncommutative algebras. (See [21] for a survey.) In this context, the use of groupoid convolution algebras as a substitute for the algebras of functions on badly behaved quotient spaces is a central theme in the noncommutative geometry of A. Connes [4]. The book just cited also shows the extent to which groupoids provide a framework for a unified study of operator algebras, foliations, and index theory.

In algebraic topology, the *fundamental groupoid* of a topological space has been exploited by P. Higgins, R. Brown, and others (see Brown’s textbook [3], or Higgins’ book [9], which is a good general introduction to groupoids as well) in situations where the use of a fixed base point as imposed by the usual fundamental group would be too restrictive. Groupoid methods are thus well adapted to disconnected spaces (which arise frequently when connected spaces are cut into pieces, as in the van Kampen theorem) and to spaces with fixed-point-free group actions.

The extension of Lie theory from differentiable groups to groupoids was carried out for the most part by J. Pradines, as described in a series of notes ending with [20]. An exposition of this work, together with a detailed study of the role of groupoids and Lie algebroids in differential geometry, can be found in the book of K. Mackenzie [13].¹ There is also a close relation between Lie algebroids and Lie groupoids and the




Figure 1. Tiling of the plane by 2×1 rectangles.

better-known pseudogroups of differentiable transformations. (See the brief remark at the end of the following section or [8] for more details.)

Finally, the author’s own interest in groupoids arose from the discovery [24, 25], made independently at around the same time by Karasev [10] and Zakrzewski [28], that a groupoid with compatible *symplectic* structure is the appropriate geometric model for a family of noncommutative algebras obtained by deformation from the algebra of functions on a manifold with a Poisson bracket structure.

Global and Local Symmetry Groupoids

We begin our exposition with a simple example which will lead to the definition of a groupoid. Consider a tiling of the euclidean plane \mathbb{R}^2 by identical 2×1 rectangles, specified by the set X (idealized as 1-dimensional) where the grout between the tiles lies: $X = H \cup V$, where $H = \mathbb{R} \times \mathbb{Z}$ and $V = 2\mathbb{Z} \times \mathbb{R}$. (See Figure 1.) We will call each connected component of $\mathbb{R}^2 \setminus X$ a *tile*.

The symmetry of this tiling is traditionally described by the group Γ consisting of those rigid motions of \mathbb{R}^2 which leave X invariant. It consists of the normal subgroup of translations by elements of the lattice $\Lambda = H \cap V = 2\mathbb{Z} \times \mathbb{Z}$ (corresponding to corner points of the tiles), together with reflections through each of the points in $\frac{1}{2}\Lambda = \mathbb{Z} \times \frac{1}{2}\mathbb{Z}$ and across the horizontal and vertical lines through those points.

Now consider what is lost when we pass from X to Γ .

- The same symmetry group would arise if we had replaced X by the lattice Λ of corner points, even though Λ looks quite different

¹We note that, before the appearance of [5], Lie groupoids were called differentiable groupoids, the term “Lie” being reserved for those satisfying a certain transitivity property.




Figure 2. Multiplication in a groupoid.

from X . (Of course, there are reasons why this loss of detail in passing from X to Γ is welcome, as is the case for many mathematical abstractions, but read on.)

- The group Γ retains no information about the local structure of the tiled plane, such as the fact that neighborhoods of all the points inside the tiles look identical if the tiles are uniform, while they may look different if the tiles are painted with a design (which could still be invariant under Γ).
- If, as is the case on real bathroom floors, the tiling is restricted to a finite part of the plane, such as $B = [0, 2m] \times [0, n]$, the symmetry group shrinks drastically. The subgroup of Γ leaving $X \cap B$ invariant contains just 4 elements, even though a repetitive pattern on the bathroom floor is clearly visible.

Our first groupoid will enable us to describe the symmetry of the finite tiled rectangle. We first define the *transformation groupoid* of the action of Γ on \mathbb{R}^2 to be the set²

$$G(\Gamma, \mathbb{R}^2) = \{(x, y, \gamma) \mid x \in \mathbb{R}^2, y \in \mathbb{R}^2, \gamma \in \Gamma, \text{ and } x = \gamma y\}$$

with the partially defined binary operation

$$(x, y, \gamma)(y, v, \delta) = (x, y\delta, \gamma\delta).$$

This operation on $G = G(\Gamma, \mathbb{R}^2)$ has the following properties.

1. It is defined only for certain pairs of elements: gh is defined only when $\beta(g) = \alpha(h)$ for certain maps α and β from G onto \mathbb{R}^2 ; here $\alpha : (x, y, \gamma) \mapsto x$ and $\beta : (x, y, \gamma) \mapsto y$.
2. It is associative: if either of the products $(gh)k$ or $g(hk)$ is defined, then so is the other, and they are equal.
3. For each g in G , there are left and right identity elements λ_g and ρ_g such that $\lambda_g g = g = g \rho_g$.
4. Each g in G has an inverse g^{-1} for which $gg^{-1} = \lambda_g$ and $g^{-1}g = \rho_g$.

More generally, a *groupoid* with base B is a set G with mappings α and β from G onto B and a partially defined binary operation

²As a set, $G(\Gamma, \mathbb{R}^2)$ is isomorphic to $\Gamma \times \mathbb{R}^2$, but we prefer the following more symmetric description, which makes the composition law more transparent.

$(g, h) \mapsto gh$ satisfying the conditions 1–4 above. We may think of each element g of G as an arrow pointing from $\beta(g)$ to $\alpha(g)$ in B ; arrows are multiplied by placing them head to tail, as in Figure 2.

Many properties of groupoids suggested by this picture are easily proven. For instance, $\alpha(g^{-1}) = \beta(g)$ (since gg^{-1} is defined), and $\alpha(gh) = \alpha(g)$ (since $(\lambda_g g)h = \lambda_g(gh)$ implies that $\lambda_g(gh)$ is defined). One also shows easily that $\alpha(g)$ and λ_g (or $\beta(g)$ and ρ_g) determine one another, thus producing a bijective mapping ι from the base B to the subset $G^{(0)} \subseteq G$ consisting of the identity elements. The reader familiar with categories will probably have realized by now that a *groupoid is just a category in which every morphism has an inverse*.³

Now we return to our transformation groupoid $G(\Gamma, \mathbb{R}^2)$ and form its *restriction* to $B = [0, 2m] \times [0, n]$ (or any other subset B of \mathbb{R}^2) by defining

$$G(\Gamma, \mathbb{R}^2)|_B = \{g \in G(\Gamma, \mathbb{R}^2) \mid \alpha(g) \text{ and } \beta(g) \text{ belong to } B\}.$$

The following concepts from groupoid theory, applied to this example, now show that $G(\Gamma, \mathbb{R}^2)|_B$ indeed captures the symmetry which we see in the tiled floor.

- An *orbit* of the groupoid G over B is an equivalence class for the relation

$$x \sim_G y \text{ if and only if there is a groupoid element } g \text{ with } \alpha(g) = x \text{ and } \beta(g) = y.$$

In the example, two points are in the same orbit if they are similarly placed within their tiles or within the grout pattern. (We are allowed to turn the tiles over, as well as to translate them or rotate them by 180° .)

- The *isotropy group* of $x \in B$ consists of those g in G with $\alpha(g) = x = \beta(g)$. In the example, the isotropy group is trivial for every point except those in $\frac{1}{2}\Lambda \cap B$, for which it is $\mathbb{Z}_2 \times \mathbb{Z}_2$.

We can describe local symmetries of our tiling by introducing another groupoid. Consider the plane \mathbb{R}^2 as the disjoint union of $P_1 = B \cap X$ (the grout), $P_2 = B \setminus P_1$ (the tiles), and $P_3 = \mathbb{R}^2 \setminus B$ (the exterior of the tiled room). Let E be the group of all euclidean motions of the plane, and define the local symmetry groupoid G_{loc} as the set of triples (x, γ, y) in $B \times E \times B$ for which $x = \gamma y$, and for which y has a neighborhood U

³Actually, not quite. The elements of a category can constitute a class rather than a set, so a groupoid is a small category with every morphism invertible. This is just a question of terminology, though; many authors allow their groupoids to be classes.

in \mathbb{R}^2 such that $\gamma(U \cap P_i) \subseteq P_i$ for $i = 1, 2, 3$. The composition is given by the same formula as for $G(\Gamma, \mathbb{R}^2)$.

For this groupoid, there are only a finite number of orbits (see Figure 3).

- \mathcal{O}_1 = interior points of tiles
- \mathcal{O}_2 = interior edge points
- \mathcal{O}_3 = interior crossing points
- \mathcal{O}_4 = boundary edge points
- \mathcal{O}_5 = boundary “T” points
- \mathcal{O}_6 = boundary corner points

The isotropy group of a point in \mathcal{O}_1 is now isomorphic to the entire rotation group $O(2)$; it is $\mathbb{Z}_2 \times \mathbb{Z}_2$ for \mathcal{O}_2 , the 8-element dihedral group D_4 for \mathcal{O}_3 , and \mathbb{Z}_2 for \mathcal{O}_4 , \mathcal{O}_5 , and \mathcal{O}_6 .

$G_{loc}|_L$ now has just three orbits, with isotropy groups isomorphic to D_4 , \mathbb{Z}_2 , and \mathbb{Z}_2 as before. (See Figure 4.) At this point, we have something like the smile which remains after the Cheshire Cat has disappeared. The set L has no structure at all except for that provided by the groupoid $G_{loc}|_L$, which is a sort of relic of the symmetry structure which was given by the original decomposition into grout, tiles, and exterior. We could say that the points of L have “internal symmetry” specified by the isotropy groups of $G_{loc}|_L$ and that the “types” of these points are classified by the orbit structure of the groupoid. This internal symmetry and type classification can be viewed either as a relic of the original tiling or simply as a “geometric structure” on L .

The situation here is similar to that in gauge theory, where the points of space-time are equipped with internal symmetry groups whose representations correspond to physical particles. But gauge groups and gauge transformations are applicable only because all the points of space-time look alike. To describe nonhomogeneous structures like the tiled floor requires groupoids rather than groups.⁴

To close this section, we mention that another approach to local symmetry is through the theory of **pseudogroups**, whose elements typically consist of selected homeomorphisms between open subsets of a space of interest, with the composition of two transformations defined on the largest possible domain. The “germs” of elements of a pseudogroup form a groupoid, while in the other direction certain subsets of a groupoid admit a multiplication which makes them the abstract counterpart of a pseudogroup (in the same sense that a group is the abstract counterpart of a set of invertible mappings closed under composition) called an **inverse semigroup**. We refer to [21] for an exposition of inverse semigroups and their relation to groupoids.




Figure 3. The point labelled by j belongs to the orbit \mathcal{O}_j of the groupoid $G(\Gamma, \mathbb{R}^2)|_B$.




Figure 4. Orbit decomposition of the finite groupoid $G_{loc}|_L$.

Groupoids and Equivalence Relations

If B is any set, the product $B \times B$ is a groupoid over B with $\alpha(x, y) = x$, $\beta(x, y) = y$, and $(x, y)(y, z) = (x, z)$. The identities are the (x, x) , and $(x, y)^{-1} = (y, x)$. We call this the *pair groupoid* of B .⁵ Note that a subgroupoid of $B \times B$, i.e., a subset closed under product and inversion and containing all the identity elements,⁶ is nothing but an equivalence relation on B . The orbits are the equivalence classes, and the isotropy subgroups are all trivial.

If G is any groupoid over B , the map $(\alpha, \beta) : G \rightarrow B \times B$ is a morphism from G to the pair groupoid of B . (A groupoid morphism from G over B to G' over B' is a pair of maps $G \rightarrow G'$ and $B \rightarrow B'$ compatible with the multiplication, “head,” and “tail” maps of the two groupoids.) The image of (α, β) is the orbit equivalence relation \sim_G , and the kernel is the union of the isotropy groups.

The morphism (α, β) suggests another way of looking at groupoids. Until now, we have been describing them as generalized groups, but now

⁵Other authors use the terms coarse groupoid or banal groupoid for this object.


⁶One often omits this last condition from the definition of a subgroupoid. To emphasize it, we can refer to a wide subgroupoid.

⁴Actually, groupoids are useful even in the realm of gauge theory. (See [14].)

we want to think of groupoids as generalized equivalence relations as well. From this point of view, a groupoid over B tells us not only *which* elements of B are equivalent to one another (or “isomorphic”), but it also parametrizes the *different ways* (“isomorphisms”) in which two elements can be equivalent. This leads us to the following guiding principle of Grothendieck, Mackey, Connes, Deligne,...

Almost every interesting equivalence relation on a space B arises in a natural way as the orbit equivalence relation of some groupoid G over B . Instead of dealing directly with the orbit space B/G as an object in the category S_{map} of sets and mappings, one should consider instead the groupoid G itself as an object in the category \mathcal{G}_{htp} of groupoids and homotopy classes of morphisms.

Here is the definition of homotopy. If



are morphisms of groupoids for $i = 1, 2$, a *homotopy* (or *natural transformation* in the language of categories) from f^1 to f^2 is a map $h : B \rightarrow G'$ with the following properties.

- For each $x \in B$, $\alpha'(h(x)) = f_B^1(x)$ and $\beta'(h(x)) = f_B^2(x)$.
- For each $g \in G$, $h(\alpha(g))f_G^2(g) = f_G^1(g)h(\beta(g))$.

An isomorphism in \mathcal{G}_{htp} between the groupoids G and G' (sometimes called an *equivalence of groupoids*) gives a bijection between the orbit spaces B/G and B/G' , as well as group isomorphisms between corresponding isotropy subgroups in G and G' . In the absence of further structure on the groupoids, an equivalence is essentially no more than this, but when the groupoids (and their bases) have measurable, topological, differentiable, algebraic, or symplectic (and Poisson) structures, the quotient spaces can be rather nasty objects, outside the category originally under consideration, and it is essential to focus attention on the groupoids themselves. One tool which makes this focus effective is an algebra associated to the groupoid G over B which is a useful substitute for the space of functions on B/G when the quotient is a “bad” space. We begin to describe this algebra in the next section.

Groupoid Convolution and Matrices

The *convolution* of two complex valued functions on a group G is defined as the sum

$$(a * b)(g) = \sum_{k \in G} a(k)b(k^{-1}g),$$

at least when the support of each function (i.e., the set where it is not zero) is finite. When the supports are infinite, the sum may still be defined if the functions are absolutely summable. One obtains in this way, for instance, the rules for multiplying Fourier series as functions on the additive group \mathbb{Z} .

To extend the definition of convolution from groups to groupoids, one replaces the range of summation G by the α -*fibre* $G_g = \alpha^{-1}(\alpha(g))$. Alternatively, one may use the more symmetric formula for convolution:

$$(a * b)(g) = \sum_{\{(k,\ell) | k\ell=g\}} a(k)b(\ell).$$

For instance, if $G = B \times B$ is the pair groupoid over $B = \{1, 2, \dots, n\}$, the convolution operation (defined on all functions) is

$$(a * b)(k, j) = \sum_{(i,k)(k,j)=(i,j)} a(i, k)b(i, j);$$

and if we write a_{ij} for $a(i, j)$, we get

$$(a * b)_{ij} = \sum_{k=1}^n a_{ik}b_{kj},$$

which exhibits $\mathcal{F}(\{1, 2, \dots, n\} \times \{1, 2, \dots, n\})$ as the algebra of $n \times n$ matrices.

This may look like a peculiar way of viewing the algebra of matrices, but Connes ([4], pp. 33–39, and elsewhere) has made the point that it was precisely as a groupoid algebra that Heisenberg constructed his original formulation of quantum mechanics. (The elements of the groupoid were transitions between excited states of an atom. Only later were the elements of the groupoid algebra identified as matrices.) The noncommutativity of the algebra of observables in quantum mechanics is then seen to be a direct consequence of the noncommutativity of the pair groupoid $B \times B$, as compared with the commutativity of the group \mathbb{Z} underlying the Fourier series representation of oscillatory motion in classical mechanics.

Topological Groupoids and Groupoid Algebras

For many interesting groupoids, such as the pair groupoid of the real numbers, the functions with finite or even countable support are rather unnatural objects, and it is more common to consider continuous functions. To define the con-

volution of such functions, it is necessary to replace the sum in the definition by an integral. An appropriate general setting for this extended notion of convolution is that of topological groupoids.

A groupoid G is a *topological groupoid* over B if G and B are topological spaces and α, β , and multiplication are continuous maps. It turns out that this notion is useful for many purposes beyond the definition of convolution. A criticism sometimes applied to the theory of groupoids is that their classification up to isomorphism is nothing other than the classification of equivalence relations (via the orbit equivalence relation) and groups (via the isotropy groups). The imposition of a compatible topological structure produces a nontrivial interaction between the two structures. (The theory of principal bundles is just a special case of this.⁷)

For instance, the groupoid $G_{\text{loc}}|_L$ of local symmetries introduced in the section entitled “Global and Local Symmetry Groupoids” can be entirely decomposed into its restrictions to the orbits $\mathcal{O}_3, \mathcal{O}_5$, and \mathcal{O}_6 . In other words, the groupoid contains no information about the relation of the orbits to one another. This relation can be encoded, however, in a suitable topology on the groupoid.

To return to our discussion of convolution on groupoids, we consider an example. If $G = \mathbb{R}^n \times \mathbb{R}^n$, we can define a convolution operation on the space $C_c(G)$ of continuous functions with *proper* support (i.e., for each compact subset K of \mathbb{R}^n , the intersections of $K \times \mathbb{R}^n$ and $\mathbb{R}^n \times K$ with $\{(x, y) | a(x, y) \neq 0\}$ have compact closure). Multiplication in $C_c(G)$ is given by the integral formula

$$(1) \quad (a * b)(x, y) = \int_{\mathbb{R}^n} a(x, z)b(z, y) dz,$$

where dz is Lebesgue measure.⁸

This “continuous matrix multiplication” is precisely the composition law which we obtain by thinking of each $a \in C_c(G)$ as the “kernel” (here the term is used in the sense of Schwartz [23]) of an operator \tilde{a} on $L^2(\mathbb{R}^n)$:

$$(\tilde{a}\psi)(x) = \int_{\mathbb{R}^n} a(x, y)\psi(y)dy.$$

Not every operator (not even the identity) is realized by a continuous kernel, but all operators

⁷Of course, a completely different retort could be made to the criticism above. By the same criterion, finite-dimensional vector spaces should be completely uninteresting, but the study of their morphisms is a very rich subject.

⁸Dirac’s notation $\langle x|a|y \rangle$ for $a(x, y)$ emphasizes the idea that the values of a are “transition amplitudes” between “states” and so fits nicely with the groupoid interpretation of (1).

(including unbounded ones) satisfying reasonable continuity properties can be realized by kernels which are *distributions*, or generalized functions on $\mathbb{R}^n \times \mathbb{R}^n$. (See [23].)

To define convolution operations on functions on a general locally compact topological groupoid, we need to start with a family $\{\mu_x\}$ of measures along the α -fibres, which will fill the role played by the measure dz in (1) above. Convolution is then defined⁹ by

$$(a * b)(g) = \int_{G_g} a(k)b(k^{-1}g)d\mu_g(k).$$

This operation is associative if the measures μ_g satisfy a suitable left-invariance property.

From continuous functions on G , one can pass by various processes of completion and localization to larger *groupoid algebras*, which play the role of continuous functions, measurable functions, or even distributions on the orbit space of G . We refer to the books [4, 19], and [21] for more details.

A more intrinsic construction of groupoid algebras, free of the choice of measures, can also be given. It includes as special cases the convolution of measures on a group G and the multiplication of kernels of operators on Mackey’s “intrinsic Hilbert spaces” [15]. The definition can be found on p. 101 of [4] for the case of Lie groupoids (see next section “Lie Groupoids and Lie Algebroids” below). A construction modeled on the definition of the intrinsic Hilbert spaces should extend this idea to a general locally compact groupoid with a prescribed left-invariant family of measure *classes* along the α -fibres.

Lie Groupoids and Lie Algebroids

A groupoid G over B is a *Lie groupoid* if G and B are differentiable manifolds, and α, β , and multiplication are differentiable maps.¹⁰ For example, the pair groupoid $M \times M$ over any differentiable manifold M is a Lie groupoid, and any Lie group is a Lie groupoid over a single point.

The infinitesimal object associated with a Lie groupoid is a *Lie algebroid*. A Lie algebroid over a manifold B is defined to be a vector bundle A over B with a Lie algebra structure $[\cdot, \cdot]$ on its space of smooth sections, together with a bundle map ρ (called the *anchor* of the Lie algebroid) from A to the tangent bundle TB satisfying the conditions:

1. $[\rho(X), \rho(Y)] = \rho([X, Y]);$
2. $[X, \varphi Y] = \varphi[X, Y] + (\rho(X) \cdot \varphi)Y.$

⁹This definition seems to be due to J. Westman [26].

¹⁰A further technical condition is included in the definition to insure that the domain of multiplication is a smooth submanifold of $G \times G$; the derivatives of α and β are required to have maximal rank everywhere.




Figure 5. The groupoid G with an element of \mathcal{A} and its image under ρ .

Here, X and Y are smooth sections of A , φ is a smooth function on B , and the bracket on the left-hand side of the first condition is the usual Jacobi-Lie bracket of vector fields. For example, the tangent bundle TB with the usual bracket, and the identity map as anchor, is a Lie algebroid over B , while any Lie algebra is a Lie algebroid over a single point.

The Lie algebroid of a Lie groupoid is defined via left-invariant vector fields, just as it is in the case of Lie groups. Since the left translation by a groupoid element g maps only from $G_{\beta(g)}$ to $G_{\alpha(g)}$, we must first restrict the values of our vector fields to lie in the subbundle $T^\alpha G = \ker(T\alpha)$ of TG consisting of vectors tangent to the α -fibres. (Here, $T\alpha$ is the derivative of α .) For every x in B and every g in the β -fibre $\beta^{-1}(x)$, the fibre $T_g^\alpha G$ can be identified with $T_{i(x)}^\alpha G$ via left translation by g . Thus, the space of left-invariant sections of $T^\alpha G$ along $\beta^{-1}(x)$ can be identified with the single space $\mathcal{A}_x G = T_{i(x)}^\alpha G$, and $T\beta$ gives a well-defined map $\rho_x : \mathcal{A}_x G \rightarrow T_x B$. The (disjoint) union of these objects for all x in B forms a vector bundle $\mathcal{A}G$ over B with a bundle map $\rho : \mathcal{A}G \rightarrow TB$. (See Figure 5.) Identifying the sections of $\mathcal{A}G$ with left-invariant sections of $T^\alpha G \subseteq TG$ and using the usual bracket on sections of TG defines a bracket of sections on $\mathcal{A}G$ making it into a Lie algebroid with anchor ρ .

For instance, the Lie algebroid of $B \times B$ is TB , and by extension of a well-known idea from Lie group theory, one may think of the Jacobi-Lie bracket on sections of TB as being the infinitesimal remnant of the noncommutativity of the composition law $(x, y)(y, z) = (x, z)$.

The local versions of Lie's three fundamental theorems all extend from groups to groupoids, but there are some interesting differences at the global level. Although every subalgebroid A' of $\mathcal{A}G$ is indeed $\mathcal{A}G'$ for a Lie groupoid G' ,

there may be no such G' for which the induced morphism $G' \rightarrow G$ is globally one-to-one. More seriously (and contrary to an incorrect assertion in [20]), there exist Lie algebroids which are "nonintegrable" in the sense that they are not the Lie algebroid of any global Lie groupoid at all. (There is a *local* Lie groupoid corresponding to any Lie algebroid, though.) We refer to Chapter V of [13] for a discussion of all this, together with further references.

Boundary Lie Algebroids

In this section we will explain how Lie algebroids provide a natural setting for understanding some recent work by R. Melrose [17, 18] on the analysis of (pseudo)differential operators on manifolds with boundary. (Some prior knowledge about such operators on manifolds *without* boundary will be assumed here.)

The space $\mathcal{X}(M, \partial M)$ of vector fields which are tangent to the boundary ∂M of a manifold M forms a Lie algebra over \mathbb{R} and a module over $C^\infty(M)$, and the condition $[X, \varphi Y] = \varphi[X, Y] + (X \cdot \varphi)Y$ is satisfied (since it is satisfied for all vector fields and functions). Remarkably, there is a vector bundle ${}^b TM$ over M whose sections are in one-to-one correspondence with the elements of $\mathcal{X}(M, \partial M)$ via a bundle map $\rho : {}^b TM \rightarrow TM$ which is an isomorphism over the interior of M and whose image over a boundary point is the tangent space to ∂M at that point. These structures make ${}^b TM$ into a Lie algebroid. We refer to [17] for a precise definition of ${}^b TM$, mentioning here only that, when (y_1, \dots, y_{n-1}, x) are local coordinates for M defined on an open subset of the half space $x \geq 0$, a local basis over $C^\infty(M)$ for the sections of ${}^b TM$ is given by $(\frac{\partial}{\partial y_1}, \dots, \frac{\partial}{\partial y_{n-1}}, x \frac{\partial}{\partial x})$.

Melrose develops analysis on M by regarding the b -tangent bundle ${}^b TM$ as the "correct" tangent bundle for this manifold with boundary. Thus, the algebra of b -differential operators is the algebra of operators on $C^\infty(M)$ generated by the sections of ${}^b TM$ (acting on functions via ρ) and $C^\infty(M)$ (acting on itself by multiplication).¹¹ The principal symbols of these operators are homogeneous functions on the dual, or b -cotangent bundle ${}^b T^*M$. Inverting (modulo smoothing operators) elliptic b -differential operators leads to the notion of b -pseudodifferential op-

¹¹Such a "universal enveloping algebra" was already defined for general Lie algebroids by Rinehart [22].

erators, whose symbols are again functions on the b -cotangent bundle.

Since the b -pseudodifferential operators act on $C^\infty(M)$, their Schwartz kernels are distributions on $M \times M$, which is a manifold with corners. Analysis of these kernels is facilitated, though, by lifting them to a certain manifold with boundary, $b(M \times M)$, obtained from $M \times M$ by a blowing-up operation along the corners. The case where M is a half-open interval is illustrated in Figure 6. The general case is locally the product of this one with a manifold without boundary.

It turns out that $b(M \times M)$ is a groupoid whose Lie algebroid is bTM and that the composition law for b -kernels ([17], Equation (5.66)) is precisely the convolution operation in the groupoid algebra of $b(M \times M)$. Thus, we can think of the groupoid $b(M \times M)$ and its Lie algebroid bTM as providing a new “internal symmetry structure” appropriate to our manifold with boundary.

It is therefore interesting to understand the structure of the groupoid $b(M \times M)$. In fact, it decomposes algebraically (though not topologically) as the disjoint union of the pair groupoid over the interior of M and the groupoid G over ∂M for which $(\alpha, \beta)^{-1}(x, y)$ consists of the orientation-preserving linear isomorphisms between the normal lines $T_y M / T_y(\partial M)$ and $T_x M / T_x(\partial M)$. Thus, with the “geometric structure” on M provided by the enlarged pair groupoid $b(M \times M)$, each point of the boundary ∂M which has the internal symmetries are those of an oriented 1-dimensional vector space. We should think of the vectors in this space as being points “infinitesimally far” from the boundary, so that the boundary of M becomes an “infinitesimal boundary layer”. The section $x \frac{\partial}{\partial x}$ of bTM can now be seen as pointing into this layer in a nontrivial way along ∂M ; that is why it is not zero there.

At this point, our use of Lie algebroids and Lie groupoids adds only a geometric picture to Melrose’s analysis. It does, however, suggest the possibility of extending the theory of pseudodifferential operators to algebras of operators associated with other Lie algebroids. In fact, the theory of longitudinal operators on foliated manifolds (see Section 1.5. γ of [4]) is an example of such an extension, but the class of Lie algebroids which can be treated in this way remains to be determined. An interesting test case should be the Lie algebroids associated with the Bruhat-Poisson structures on flag manifolds [12], since the corresponding orbit decomposition on a flag manifold is the Bruhat decomposition, which plays a central role in representation theory.




Figure 6. The groupoids $G = M \times M$ and $bG = b(M \times M)$, where $M = (0, 1)$.

Conclusion

This article has presented a small sample of the applications of groupoids in analysis, algebra, and topology. R. Brown has reported in [2] the suggestion of F. W. Lawvere that groupoids should perhaps be renamed “groups” and those special groupoids with just one base point be given a new name to reflect their singular nature. Even if this is too far to go, I hope to have convinced the reader that groupoids are worth knowing about and worth looking out for.¹²

References

- [1] W. BRANDT, *Über eine Verallgemeinerung des Gruppenbegriffes*, Math. Ann. **96** (1926), 360–366.
- [2] R. BROWN, *From groups to groupoids: A brief survey*, Bull. London Math. Soc. **19** (1987), 113–134.
- [3] ———, *Topology: A geometric account of general topology, homotopy types, and the fundamental groupoid*, Halsted Press, New York, 1988.
- [4] A. CONNES, *Noncommutative geometry*, Academic Press, San Diego, 1994.
- [5] A. COSTE, P. DAZORD, and A. WEINSTEIN, *Groupoïdes symplectiques*, Publications du Département de Mathématiques, Université Claude Bernard-Lyon I **2A** (1987), 1–67.
- [6] C. EHRESMANN, *Oeuvres complètes et commentées*, (A. C. Ehresmann, ed.) Suppl. Cahiers Top. Géom. Diff., Amiens, 1980–1984.
- [7] P. GABRIEL, *Construction de préschémas quotient*, Schémas en Groupes, Sémin. Géométrie Algébrique, Inst. Hautes Études Scientifiques, 1963/64, Fasc. 2a, Exposé 5.
- [8] A. GROTHENDIECK, *Techniques de construction et théorèmes d’existence en géométrie algébrique III: préschémas quotients*, Séminaire Bourbaki 13e année 1960/61, no. 212 (1961).
- [9] P. J. HIGGINS, *Notes on categories and groupoids*, Van Nostrand Reinhold, London, 1971.
- [10] M. V. KARASEV, *Analogues of objects of Lie group theory for nonlinear Poisson brackets*, Math. USSR Izvestia **28** (1978) 497–527.

¹²Any reader stimulated to look further may wish to consult the Groupoid Home Page, <http://amath-www.colorado.edu:80/math/department/groupoids/groupoids.shtml> (under construction in May 1996).

- [11] A. KUMPERA and D. SPENCER, *Lie equations*, Princeton University Press, Princeton, NJ, 1972.
- [12] J.-H. LU and A. WEINSTEIN, *Poisson Lie groups, dressing transformations, and the Bruhat decomposition*, J. Differential Geom. **31** (1990), 501–526.
- [13] K. MACKENZIE, *Lie groupoids and Lie algebroids in differential geometry*, LMS Lecture Note Ser., vol. 124, Cambridge Univ. Press, 1987.
- [14] ———, *Classification of principal bundles and Lie groupoids with prescribed gauge group bundle*, J. Pure Appl. Algebra **58** (1989), 181–208.
- [15] G. W. MACKEY, *The mathematical foundations of quantum mechanics*, W. A. Benjamin, New York, 1963.
- [16] ———, *Ergodic theory and virtual groups*, Math. Ann. **166** (1966), 187–207.
- [17] R. B. MELROSE, *The Atiyah-Patodi-Singer index theorem*, A. K. Peters, Wellesley, 1993.
- [18] ———, *Geometric scattering theory*, Cambridge Univ. Press, Cambridge, 1995.
- [19] C. C. MOORE and C. SCHOCHET, *Global analysis on foliated spaces*, MSRI Publications, vol. 9, Springer-Verlag, New York, 1988.
- [20] J. PRADINES, *Troisième théorème de Lie sur les groupoïdes différentiables*, C. R. Acad. Sci. Sér. I Math. Paris **267** (1968), 21–23.
- [21] J. RENAULT, *A groupoid approach to C^* algebras*, Lecture Notes in Math. **793** (1980).
- [22] G. S. RINEHART, *Differential forms on general commutative algebras*, Trans. Amer. Math. Soc. **108** (1963), 195–222.
- [23] L. SCHWARTZ, *Théorie des distributions*, Nouvelle édition, Hermann, Paris, 1966.
- [24] A. WEINSTEIN, *Symplectic groupoids and Poisson manifolds*, Bull. Amer. Math. Soc. (N.S.) **16** (1987), 101–104.
- [25] ———, *Noncommutative geometry and geometric quantization*, Symplectic Geometry and Mathematical Physics: Actes du Colloque en l'Honneur de Jean-Marie Souriau (P. Donato et al. eds.), Birkhäuser, Boston, 1991, pp. 446–461.
- [26] J. WESTMAN, *Harmonic analysis on groupoids*, Pacific J. Math. **27** (1968), 621–632.
- [27] I. M. YAGLOM, *Felix Klein and Sophus Lie: Evolution of the idea of symmetry in the nineteenth century*, Birkhäuser, Boston, 1988.
- [28] S. ZAKRZEWSKI, *Quantum and classical pseudogroups, I and II*, Commun. Math. Phys. **134** (1990), 347–370, 371–395.