

Differential Topology Forty-six Years Later

John Milnor

In the 1965 Hedrick Lectures,¹ I described the state of differential topology, a field that was then young but growing very rapidly. During the intervening years, many problems in differential and geometric topology that had seemed totally impossible have been solved, often using drastically new tools. The following is a brief survey, describing some of the highlights of these many developments.

Major Developments

The first big breakthrough, by Kirby and Siebenmann [1969, 1969a, 1977], was an obstruction theory for the problem of triangulating a given topological manifold as a PL (= piecewise-linear) manifold. (This was a sharpening of earlier work by Casson and Sullivan and by Lashof and Rothenberg. See [Ranicki, 1996].) If B_{Top} and B_{PL} are the stable classifying spaces (as described in the lectures), they showed that the relative homotopy group $\pi_j(B_{\text{Top}}, B_{\text{PL}})$ is cyclic of order two for $j = 4$, and zero otherwise. Given an n -dimensional topological manifold M^n , it follows that there is an obstruction $\mathfrak{o} \in H^4(M^n; \mathbb{Z}/2)$ to triangulating M^n as a PL-manifold. In dimensions $n \geq 5$ this is the only obstruction. Given such a triangulation, there is a similar obstruction in $H^3(M^n; \mathbb{Z}/2)$ to

John Milnor is professor of mathematics at Stony Brook University. His email address is jack@math.sunysb.edu.

The author would like to thank Mike Hopkins, Rob Kirby, Mark Mahowald, John Morgan, Larry Siebenmann, and Jonathan Sondow for their help. He also wants to thank the NSF for its support under grant DMS0757856.

¹*These lectures have recently been digitized by MSRI and should be available soon. Thanks to Dusa McDuff for unearthing the original tapes. (With regard to Wilder's introduction, compare [Milnor, 1999].)*

its uniqueness up to a PL-isomorphism that is topologically isotopic to the identity. In particular, they proved the following.

Theorem 1. *If a topological manifold M^n without boundary satisfies*

$$H^3(M^n; \mathbb{Z}/2) = H^4(M^n; \mathbb{Z}/2) = 0 \quad \text{with } n \geq 5,$$
then it possesses a PL-manifold structure that is unique up to PL-isomorphism.

(For manifolds with boundary one needs $n > 5$.) The corresponding theorem for all manifolds of dimension $n \leq 3$ had been proved much earlier by Moise [1952]. However, we will see that the corresponding statement in dimension 4 is false.

An analogous obstruction theory for the problem of passing from a PL-structure to a smooth structure had previously been introduced by Munkres [1960, 1964a, 1964b] and Hirsch [1963]. (See also [Hirsch-Mazur, 1974].) Furthermore, Cerf had filled in a crucial step, proving by a difficult geometric argument that the space of orientation-preserving diffeomorphisms of the three-sphere is connected. (See the Cartan Seminar Lectures of 1962/63, as well as [Cerf, 1968].) Combined with other known results, this led to the following.

Theorem 2. *Every PL-manifold of dimension $n \leq 7$ possesses a compatible differentiable structure, and this structure is unique up to diffeomorphism whenever $n < 7$.*

For more information see **Further Details** at the end of this article.

The next big breakthrough was the classification of simply connected closed topological 4-manifolds by Freedman [1982]. He proved, using wildly nondifferentiable methods, that such a manifold is uniquely determined by

- (1) the isomorphism class of the symmetric bilinear form $H^2 \otimes H^2 \rightarrow H^4 \cong \mathbb{Z}$, where $H^k = H^k(M^4; \mathbb{Z})$, together with
- (2) the Kirby-Siebenmann invariant $\circ \in H^4(M^4; \mathbb{Z}/2) \cong \mathbb{Z}/2$.

These can be prescribed arbitrarily, except for two restrictions: The bilinear form must have determinant ± 1 ; and in the “even case” where $x \cup x \equiv 0 \pmod{2H^4}$ for every $x \in H^2$, the Kirby-Siebenmann class must be congruent to $(1/8)$ th of the signature. As an example, the Poincaré hypothesis for 4-dimensional topological manifolds is an immediate consequence. For if M^4 is a homotopy sphere, then both H^2 and the obstruction class must be zero.

One year later, Donaldson [1983] used gauge theoretic methods to show that many of these topological manifolds cannot possess any smooth structure (and hence by Theorem 2 cannot be triangulated as PL-manifolds). More explicitly, if M^4 is smooth and simply connected with positive definite bilinear form, he showed that this form must be diagonalizable; hence M^4 must be homeomorphic to a connected sum of copies of the complex projective plane. There are many positive definite bilinear forms with determinant 1, and with signature divisible by 16 in the even case, which are not diagonalizable. (See, for example, [Milnor-Husemoller, 1973].) Each of these corresponds to a topological manifold M^4 with no smooth structure but such that $M^4 \times \mathbb{R}$ does have a smooth structure that is unique up to diffeomorphism.

The combination of Freedman’s topological results and Donaldson’s analytic results quickly led to rather amazing consequences. For example, it followed that there are uncountably many nonisomorphic smooth or PL structures on \mathbb{R}^4 . (Compare [Gompf, 1993].) All other dimensions are better behaved: For $n > 4$, Stallings [1962] showed that the topological space \mathbb{R}^n has a unique PL-structure up to PL-isomorphism. Using the Moise result for $n < 4$ together with the Munkres-Hirsch-Mazur obstruction theory, it follows that the differentiable structure of \mathbb{R}^n is unique up to diffeomorphism for all $n \neq 4$.

A satisfactory theory of three-dimensional manifolds took longer. The first milestone was the geometrization conjecture by Thurston [1982, 1986], which set the goal for what a theory of three-manifolds should look like. This conjecture was finally verified by Perelman [2002, 2003a, 2003b], using a difficult argument based on the “Ricci flow” partial differential equation. (Compare the expositions of Morgan-Tian [2007] and Kleiner-Lott [2008].) The three-dimensional Poincaré hypothesis followed as a special case.

The Poincaré Hypothesis: Three Versions

First consider the purely topological version.

Theorem 3. *The topological Poincaré hypothesis is true in all dimensions.*

That is, every closed topological manifold with the homotopy type of an n -sphere is actually homeomorphic to the n -sphere. For $n > 4$ this was proved by Newman [1966] and by Connell [1967], both making use of the “engulfing method” of Stallings [1960]. For $n = 4$ it is of course due to Freedman. For $n = 3$ it is due to Perelman, using Moise [1952] to pass from the topological to the PL case, and then using the Munkres-Hirsch-Mazur obstruction theory to pass from PL to smooth. \square

Theorem 4. *The piecewise-linear Poincaré hypothesis is true for n -dimensional manifolds except possibly when $n = 4$.*

That is, any closed PL manifold of dimension $n \neq 4$ with the homotopy type of an n -sphere is PL-homeomorphic to the n -sphere. For $n > 4$ this was proved by Smale [1962]; while for $n = 3$ it follows from Perelman’s work, together with the Munkres-Hirsch-Mazur obstruction theory. \square

The differentiable Poincaré hypothesis is more complicated, being true in some dimensions and false in others, while remaining totally mysterious in dimension 4. We can formulate the question more precisely by noting that the set of all oriented diffeomorphism classes of closed smooth homotopy n -spheres (= topological n -spheres) forms a commutative monoid S_n under the connected sum operation. In fact this monoid is actually a finite abelian group except possibly when $n = 4$. Much of the following outline is based on [Kervaire-Milnor, 1963], which showed in principle how to compute these groups² in terms of the stable homotopy groups of spheres for $n > 4$. Unfortunately, many proofs were put off to part 2 of this paper, which was never completed. However, the missing arguments have been supplied elsewhere; see especially [Levine, 1985].

Using Perelman’s result for $n = 3$, the group S_n can be described as follows for small n (Table 1). (Here, for example, $2 \cdot 8$ stands for the group $\mathbb{Z}/2 \oplus \mathbb{Z}/8$, and 1 stands for the trivial group.)

Thus the differentiable Poincaré hypothesis is true in dimensions 1, 2, 3, 5, 6, and 12, but unknown in dimension 4. I had conjectured that it would be false in all higher dimensions. However, Mahowald has pointed out that there is at least

²The Kervaire-Milnor paper worked rather with the group Θ_n of homotopy spheres up to h -cobordism. This makes a difference only for $n = 4$, since it is known, using the h -cobordism theorem of Smale [1962], that $S_n \xrightarrow{\cong} \Theta_n$ for $n \neq 4$. However, the difference is important in the four-dimensional case, since Θ_4 is trivial, while the structure of S_4 is the great unsolved problem.

Table 1

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
S_n	1	1	1	?	1	1	28	2	2·2·2	6	992	1	3	2	2·8128	2	2·8	2·8

one more exceptional case: *The group S_{61} is also trivial.* (Compare **Further Details** below.)

Problem. Is the group S_n nontrivial for all $n > 6, n \neq 12, 61$?

(Any precise computation for large n is impossible at the present time, since the stable homotopy groups of spheres have been computed completely only up to dimension 64. However, it seems possible that enough is known to decide this question one way or another.)

Denote the stable homotopy groups of spheres by

$$\Pi_n = \pi_{n+q}(S^q) \quad \text{for } q > n + 1,$$

and let $J_n \subset \Pi_n$ be the image of the stable Whitehead homomorphism $J : \pi_n(\mathbf{SO}) \rightarrow \Pi_n$. (See [Whitehead, 1942].) This subgroup J_n is cyclic of order³

$$|J_n| = \begin{cases} \text{denominator} \left(\frac{B_k}{4k} \right) & \text{for } n = 4k - 1, \\ 2 & \text{for } n \equiv 0, 1 \pmod{8}, \text{ and} \\ 1 & \text{for } n \equiv 2, 4, 5, 6 \pmod{8}, \end{cases}$$

where the B_k are Bernoulli numbers, for example,

$$B_1 = \frac{1}{6}, B_2 = \frac{1}{30}, B_3 = \frac{1}{42}, \\ B_4 = \frac{1}{30}, B_5 = \frac{5}{66}, B_6 = \frac{691}{2730},$$

and where the fraction $\frac{B_k}{4k}$ must be reduced to lowest terms. (Compare [Milnor-Stasheff, 1974, Appendix B].)

According to Pontrjagin and Thom, the stable n -stem Π_n can also be described as the group of all framed cobordism classes of framed manifolds. (Here one considers manifolds smoothly embedded in a high-dimensional Euclidean space, and a *framing* means a choice of trivialization for the normal bundle.) Every homotopy sphere is stably parallelizable, and hence possesses such a framing. If we change the framing, then the corresponding class in Π_n will be changed by an element of the subgroup J_n . Thus there is an exact sequence

$$(1) \quad 0 \rightarrow S_n^{\text{bp}} \rightarrow S_n \rightarrow \Pi_n/J_n,$$


³This computation of $|J_{4k-1}|$ is a special case of the **Adams conjecture** [Adams, 1963, 1965]. The proof was completed by Mahowald [1970], and the full Adams conjecture was proved by Quillen [1971], Sullivan [1974], and by Becker-Gottlieb [1975]. Adams also showed that J_n is always a direct summand of Π_n .

where $S_n^{\text{bp}} \subset S_n$ stands for the subgroup represented by homotopy spheres that bound parallelizable manifolds. This subgroup is the part of S_n that is best understood. It can be partially described as follows.

Theorem 5. For $n \neq 4$ the group S_n^{bp} is finite cyclic with an explicitly known generator. In fact this group is:

- trivial when n is even,
- either trivial or cyclic of order two when $n = 4k - 3$, and
- cyclic of order $2^{2k-2}(2^{2k-1} - 1)$ numerator $\left(\frac{4B_k}{k} \right)$ when $n = 4k - 1 > 3$.

(This last number depends on the computation of $|J_{4k-1}|$ as described above.) In the odd cases, setting $n = 2q - 1$, an explicit generator for the S_{2q-1}^{bp} can be constructed using one basic building block, namely the tangent disk-bundle of the q -sphere, together with one of the following two diagrams.



Here each circle represents one of our $2q$ -dimensional building blocks, which is a $2q$ -dimensional parallelizable manifold with boundary, and each dot represents a plumbing construction in which two of these manifolds are pasted across each other so that their central q -spheres intersect transversally with intersection number $+1$. The result will be a smooth parallelizable manifold with corners. After smoothing these corners we obtain a smooth manifold X^{2q} with smooth boundary.

For q odd, use the left diagram, and for q even use the right diagram. In either case, if $q \neq 2$, the resulting smooth boundary ∂X^{2q} will be a homotopy sphere representing the required generator of S_{2q-1}^{bp} . (The case $q = 2$ is exceptional since ∂X^4 has only the homology of the 3-sphere. In all other cases where S_{2q-1}^{bp} is trivial, the boundary will be diffeomorphic to the standard $(2q - 1)$ -sphere.)

The exact sequence (1) can be complemented by the following information.

Theorem 6. For $n \not\equiv 2 \pmod{4}$, every element of Π_n can be represented by a topological sphere.

Hence the exact sequence (1) takes the more precise form

$$(2) \quad 0 \rightarrow S_n^{\text{bp}} \rightarrow S_n \rightarrow \Pi_n/J_n \rightarrow 0.$$

However, for $n = 4k - 2$, it rather extends to an exact sequence

$$(3) \quad 0 = S_{4k-2}^{\text{bp}} \rightarrow S_{4k-2} \rightarrow \Pi_{4k-2}/J_{4k-2} \xrightarrow{\Phi_k} \mathbb{Z}/2 \rightarrow S_{4k-3}^{\text{bp}} \rightarrow 0.$$

Brumfiel [1968, 1969, 1970] sharpened this result by showing that the exact sequence (2) is split exact, except possibly in the case where n has the form $2^k - 3$. (In fact it could fail to split only in the cases $n = 2^k - 3 \geq 125$. See the discussion below.)

The **Kervaire homomorphism** Φ_k in (3) was introduced in [Kervaire, 1960]. (The image $\Phi_k(\theta) \in \mathbb{Z}/2$ is called the **Kervaire invariant** of the homotopy class θ .) Thus there are two possibilities:

- If $\Phi_k = 0$, then $S_{4k-3}^{\text{bp}} \cong \mathbb{Z}/2$, generated by the manifold ∂X^{4k-2} described above, and every element of Π_{4k-2} can be represented by a homotopy sphere.

- If $\Phi_k \neq 0$, then $S_{4k-3}^{\text{bp}} = 0$. This means that the boundary of X_{4k-2} is diffeomorphic to the standard S^{4k-3} . We can glue a $4k - 2$ ball onto this boundary to obtain a framed $(4k - 2)$ -manifold that is not framed cobordant to any homotopy sphere. In this case the kernel of Φ_k forms a subgroup of index two in Π_{4k-2}/J_{4k-2} consisting of those framed cobordism classes that can be represented by homotopy spheres.

The question as to just when $\Phi_k = 0$ was the last major unsolved problem in understanding the group of homotopy spheres. It has recently been solved in all but one case by Hill, Hopkins, and Ravenel:

Theorem 7. *The Kervaire homomorphism Φ_k is nonzero for $k = 1, 2, 4, 8, 16$, and possibly for $k = 32$, but is zero in all other cases.*

In fact Browder [1969] showed that Φ_k can be nonzero only if n is a power of two, and Barratt, Jones, and Mahowald [1984] completed the verification that Φ_k is indeed nonzero for $k = 1, 2, 4, 8, 16$. Finally, Hill, Hopkins, and Ravenel [2010] have shown that $\Phi_k = 0$ whenever $k > 32$. (Their basic tool is a carefully constructed generalized cohomology theory of period 256.)

Thus only the case $k = 32$, with $4k - 2 = 126$, remains unsettled. In particular, for $n \neq 4, 125, 126$, if the order $|\Pi_n|$ is known, then we can compute the number $|S_n|$ of exotic n -spheres precisely. In fact, if we exclude 4, 126, and numbers of the form $2^k - 3 \geq 125$, then the group S_n can be described completely whenever the structure of Π_n is known.⁴

⁴One also needs the fact that the kernel of Φ_k is always a direct summand of Π_{4k-2} (at least for $4k - 2 \neq 126$).

Further Details

Here is a brief outline of the current knowledge of Π_n . Since the direct summand J_n is known precisely, we need only look at the quotient Π_n/J_n . The most difficult part is the 2-primary component, which has been computed by Kochman [1990], with corrections by Kochman and Mahowald [1995], in all cases with $n \leq 64$. The 3 and 5 primary components have been computed in a much larger range by Ravenel [1986]. The primary components for $p \geq 7$ are trivial for $n < 82$. (In fact, for any p , the p -primary component of Π_n/J_n is trivial whenever $n < 2p(p - 1) - 2$ and is cyclic of order p when $p = 2p(p - 1) - 2$.)

Thus the stable stem Π_n is precisely known for $n \leq 64$, and hence the group S_n is precisely known for $n \leq 64$, $n \neq 4$. In Table 2 the notation b_k stands for the order of the subgroup $S_{4k-1}^{\text{bp}} \subset S_{4k-1}$, whereas a notation such as $2^3 \cdot 4$ stands for the direct sum of three copies of $\mathbb{Z}/2$ and one copy of $\mathbb{Z}/4$. The trivial group is indicated by a heavy dot. All entries corresponding to the subgroups S_n^{bp} have been underlined. (Note that S_{4k-3}^{bp} is cyclic of order two, indicated by a 2 in the 1 or 5 column, except in the cases $k = 1, 2, 4, 8, 16$.) Within this range, the group S_n is trivial only in the cases

$$n = 1, 2, 3, 5, 6, 12, 61 \quad (\text{and possibly } 4).$$

The corresponding values of b_k are not difficult to compute but grow very rapidly. See Table 3 (with approximate values for $k > 5$). Note that those b_k for which k has many divisors tend to be somewhat larger.

In conclusion, here is an argument that was postponed above.

Outline Proof of Theorem 2. It is not difficult to check that the group $\pi_0(\text{Diff}^+(\mathbb{S}^n))$ consisting of all smooth isotopy classes of orientation preserving diffeomorphisms of the unit n -sphere is abelian. Define Γ_n to be the quotient of $\pi_0(\text{Diff}^+(\mathbb{S}^{n-1}))$ by the subgroup consisting of those isotopy classes that extend over the closed unit n -disk. There is a natural embedding $\Gamma_n \subset S_n$ that sends each $(f) \in \Gamma_n$ to the “twisted n -sphere” obtained by gluing the boundaries of two n -disks together under f . It followed from [Smale, 1962] that $\Gamma_n = S_n$ for $n \geq 5$, and from [Smale, 1959] that $\Gamma_3 = 0$. Since it is easy to check that $\Gamma_1 = 0$ and $\Gamma_2 = 0$, we have

$$\Gamma_n = S_n \quad \text{for every } n \neq 4.$$

Mahowald tells me that this is true in dimension 62, and the remaining four cases are straightforward.

Table 2

n	0	1	2	3	4	5	6	7
0+	—	•	•	•	?	•	•	\underline{b}_2
8+	2	$\underline{2} \cdot 2^2$	6	\underline{b}_3	•	3	2	$\underline{b}_4 \cdot 2$
16+	2	$\underline{2} \cdot 2^3$	$2 \cdot 8$	$\underline{b}_5 \cdot 2$	24	$\underline{2} \cdot 2^2$	2^2	$\underline{b}_6 \cdot 2 \cdot 24$
24+	2	$\underline{2} \cdot 2$	$2 \cdot 6$	\underline{b}_7	2	3	3	$\underline{b}_8 \cdot 2^2$
32+	2^3	$\underline{2} \cdot 2^4$	$2^3 \cdot 4$	$\underline{b}_9 \cdot 2^2$	6	$\underline{2} \cdot 2 \cdot 6$	$2 \cdot 60$	$\underline{b}_{10} \cdot 2^4 \cdot 6$
40+	$2^4 \cdot 12$	$\underline{2} \cdot 2^4$	$2^2 \cdot 24$	\underline{b}_{11}	8	$\underline{2} \cdot 2^3 \cdot 720$	$2^3 \cdot 6$	$\underline{b}_{12} \cdot 2^3 \cdot 12$
48+	$2^3 \cdot 4$	$\underline{2} \cdot 6$	$2^2 \cdot 6$	$\underline{b}_{13} \cdot 2^2 \cdot 4$	$2^2 \cdot 6$	$\underline{2} \cdot 2^4$	$2 \cdot 4$	$\underline{b}_{14} \cdot 3$
56+	2	$\underline{2} \cdot 2^3$	2^2	$\underline{b}_{15} \cdot 2^2$	4	•	$2 \cdot 12$	$\underline{b}_{16} \cdot 2^3$

Table 3

k	2	3	4	5	6	7	8	9
b_k	28	992	8128	261632	1.45×10^9	6.71×10^7	1.94×10^{12}	7.54×10^{14}

k	10	11	12	13	14	15	16
b_k	2.4×10^{16}	3.4×10^{17}	8.3×10^{21}	7.4×10^{20}	3.1×10^{25}	$5. \times 10^{29}$	1.8×10^{31}

On the other hand, Cerf proved⁵ that $\pi_0(\text{Diff}^+(\mathbb{S}^3)) = 0$ and hence that $\Gamma_4 = 0$ (although S_4 is completely unknown). Using results about S_n as described above, it follows that $\Gamma_n = 0$ for $n < 7$, and that Γ_n is finite abelian for all n .

The Munkres-Hirsch-Mazur obstructions to the existence of a smooth structure on a given PL-manifold M^n lie in the groups $H^k(M^n; \Gamma_{k-1})$, whereas obstructions to its uniqueness lie in $H^k(M^n; \Gamma_k)$. (Unlike most of the constructions discussed above, this works even in dimension 4.) Evidently Theorem 2 follows. \square

For further historical discussion see Milnor [1999, 2007, 2009].

References

[Adams, 1963] J. F. ADAMS, On the groups $J(X)$ I, *Topology* 2 (1963), 181-195.
 [Adams, 1966] ———, On the groups $J(X)$ II, *Topology* 3 (1965), 137-171.
 [Antonelli, Burghlea and Kahn, 1972] P. L. ANTONELLI, D. BURGHELEA, and P. J. KAHN, The nonfinite homotopy type of some diffeomorphism groups, *Topology* 11 (1972), 1-49.
 [Barratt-Jones-Mahowald, 1984] M. G. BARRATT, J. D. S. JONES, and M. E. MAHOWALD, Relations amongst Toda brackets and the Kervaire invariant in dimension 62, *J. London Math. Soc.* 30 (1984), 533-550.

[Becker-Gottlieb, 1975] J. C. BECKER and D. H. GOTTLIEB, The transfer map and fiber bundles, *Topology* 14 (1975), 1-12.
 [Browder, 1969] W. BROWDER, The Kervaire invariant of framed manifolds and its generalization, *Ann. of Math.* 90 (1969), 157-186.
 [Brumfiel, 1968] G. BRUMFIEL, On the homotopy groups of BPL and PL/O, *Ann. of Math.* 88 (1968), 291-311.
 [Brumfiel, 1969] ———, On the homotopy groups of BPL and PL/O. II, *Topology* 8 (1969), 305-311.
 [Brumfiel, 1970] ———, On the homotopy groups of BPL and PL/O. III, *Michigan Math. J.* 17 (1970), 217-224.
 [Cerf, 1968] J. CERF, *Sur les difféomorphismes de la sphère de dimension trois* ($\Gamma_4 = 0$), Lecture Notes in Mathematics, 53, Springer-Verlag, Berlin-New York, 1968, xii+133 pp.
 [Connell, 1967] E. H. CONNELL, A topological H -cobordism theorem for $n \geq 5$, *Illinois J. Math.* 11 (1967), 300-309.
 [Donaldson, 1983] S. K. DONALDSON, An application of gauge theory to four-dimensional topology, *J. Differential Geom.* 18 (1983), 279-315.
 [Freedman, 1982] M. H. FREEDMAN, The topology of four-dimensional manifolds, *J. Differential Geom.* 17 (1982), 357-453.
 [Gompf, 1993] R. GOMPf, An exotic menagerie, *J. Differential Geom.* 37 (1993), 199-223.
 [Hatcher, 1983] A. E. HATCHER, A proof of the Smale conjecture, $\text{Diff}(\mathbb{S}^3) \simeq O(4)$, *Ann. of Math.* 117 (1983), 553-607.
 [Hill-Hopkins-Ravenel, 2010] M. A. HILL, M. J. HOPKINS, and D. C. RAVENEL, On the nonexistence of elements of Kervaire invariant one, arXiv 0908.3724v2 [math.AT] (November, 2010).
 [Hirsch, 1963] M. W. HIRSCH, Obstruction theories for smoothing manifolds and maps, *Bull. Amer. Math. Soc.* 69 (1963), 352-356.
 [Hirsch-Mazur, 1974] M. W. HIRSCH and B. MAZUR, *Smoothings of Piecewise Linear Manifolds*, Annals of Mathematics Studies, No. 80. Princeton University Press,

⁵Hatcher [1983] later proved the sharper result that the inclusion $SO(4) \rightarrow \text{Diff}^+(\mathbb{S}^3)$ is a homotopy equivalence. On the other hand, for $n \geq 7$, Antonelli, Burghlea, and Kahn [1972] showed that $\text{Diff}^+(\mathbb{S}^n)$ does not have the homotopy type of any finite complex. (For earlier results, see [Novikov, 1963].) For $n = 6$ the group $\text{Diff}^+(\mathbb{S}^n)$ is not connected since $\Gamma_7 \neq 0$; but I am not aware of any results for $n = 4, 5$.

- Princeton, N.J.; University of Tokyo Press, Tokyo, 1974. ix+134 pp.
- [Kervaire, 1960] M. A. KERVAIRE, A manifold which does not admit any differentiable structure, *Comment. Math. Helv.* **34** (1960), 257-270.
- [Kervaire-Milnor, 1963] M. A. KERVAIRE and J. MILNOR, Groups of homotopy spheres, I, *Ann. of Math.* **77** (1963), 504-537.
- [Kirby-Siebenmann, 1969] R. C. KIRBY and L. C. SIEBENMANN, On the triangulation of manifolds and the Hauptvermutung, *Bull. Amer. Math. Soc.* **75** (1969), 742-749.
- [Kirby-Siebenmann, 1969a] ———, For manifolds the Hauptvermutung and the triangulation conjecture are false, *Notices Amer. Math. Soc.* **16** (1969), 695.
- [Kirby-Siebenmann, 1977] ———, *Foundational Essays on Topological Manifolds, Smoothings, and Triangulations. With Notes by John Milnor and Michael Atiyah*, Annals of Mathematics Studies, 88, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, vii+355 pp.
- [Kleiner-Lott, 2008] B. KLEINER and J. LOTT, Notes on Perelman's papers, *Geom. Topol.* **12** (2008), 2587-2855.
- [Kochman, 1990] S. O. KOCHMAN, *Stable Homotopy Groups of Spheres. A Computer-assisted Approach*, Lecture Notes in Mathematics, 1423, Springer-Verlag 1990, viii+330 pp.
- [Kochman and Mahowald, 1995] S. O. KOCHMAN and M. E. MAHOWALD, On the computation of stable stems, *Contemporary Mathematics* **181** (1995), 299-316.
- [Levine, 1985] J. P. LEVINE, Lectures on groups of homotopy spheres. In: *Algebraic and Geometric Topology*, 62-95, Lecture Notes in Math., 1126, Springer, 1985.
- [Mahowald, 1970] M. MAHOWALD, The order of the image of the J -homomorphism, *Proc. Advanced Study Inst. on Algebraic Topology* (Aarhus, 1970), **II**, 376-384, Mat. Inst., Aarhus Univ., 1970.
- [Milnor, 1999] J. MILNOR, Growing up in the old Fine Hall. In: *Prospects in Mathematics*, 1-11, Amer. Math. Soc., Providence, RI, 1999.
- [Milnor, 2007] ———, *Collected Papers*. III, Differential Topology, Amer. Math. Soc., Providence, RI, 2007.
- [Milnor, 2009] ———, *Collected Papers*. IV, Homotopy, Homology, and Manifolds, Amer. Math. Soc., Providence, RI, 2009.
- [Milnor-Husemoller, 1973] J. MILNOR and D. HUSEMOLLER, *Symmetric Bilinear Forms*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 73, Springer-Verlag, 1973, viii+147 pp.
- [Milnor-Stasheff, 1974] J. MILNOR and J. STASHEFF, *Characteristic Classes*, Annals of Math. Studies, 76, Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974, vii+331 pp.
- [Moise, 1952] E. E. MOISE, Affine structures in 3-manifolds. V. The triangulation theorem and Hauptvermutung, *Ann. of Math.* **56** (1952), 96-114.
- [Morgan-Tian 2007] J. MORGAN and G. TIAN, *Ricci Flow and the Poincaré Conjecture*, Clay Mathematics Monographs, 3, Amer. Math. Soc., Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2007, xii+521 pp.
- [Munkres 1960] J. MUNKRES, Obstructions to the smoothing of piecewise-differentiable homeomorphisms, *Ann. of Math.* **72** (1960), 521-554.
- [Munkres 1964a] ———, Obstructions to imposing differentiable structures, *Illinois J. Math.* **8** (1964), 361-376.
- [Munkres 1964b] ———, Obstructions to extending diffeomorphisms, *Proc. Amer. Math. Soc.* **15** (1964), 297-299.
- [Newman, 1966] M. H. A. NEWMAN, The engulfing theorem for topological manifolds, *Ann. of Math.* **84** (1966), 555-571.
- [Novikov, 1963] S. P. NOVIKOV, Homotopy properties of the group of diffeomorphisms of the sphere (Russian), *Dokl. Akad. Nauk SSSR* **148** (1963), 32-35.
- [Perelman, 2002] G. PERELMAN, *The entropy formula for the Ricci flow and its geometric applications*, 2002, arXiv: math.DG/0211159.
- [Perelman, 2003a] ———, *Ricci flow with surgery on three-manifolds*, 2003, arXiv:math.DG/0303109.
- [Perelman, 2003b] ———, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, 2003, arXiv:math.DG/0307245.
- [Quillen, 1971] D. QUILLEN, The Adams conjecture, *Topology* **10** (1971), 67-80.
- [Ravenel, 1986] D. C. RAVENEL, *Complex Cobordism and Stable Homotopy Groups of Spheres*, Pure and Applied Mathematics, 121, Academic Press, Orlando, FL, 1986, xx+413 pp.
- [Ranicki, 1996] A. Ranicki (editor) *The Hauptvermutung Book*, Kluwer Academic Publ., Dordrecht, 1996.
- [Smale, 1959] S. SMALE, Diffeomorphisms of the 2-sphere, *Proc. Amer. Math. Soc.* **10** (1959), 621-626.
- [Smale, 1962] ———, On the structure of manifolds, *Amer. J. Math.* **84** (1962), 387-399.
- [Stallings, 1960] J. STALLINGS, Polyhedral homotopy-spheres, *Bull. Amer. Math. Soc.* **66** (1960), 485-488.
- [Stallings, 1962] ———, The piecewise-linear structure of Euclidean space, *Proc. Cambridge Philos. Soc.* **58** (1962), 481-488.
- [Sullivan, 1974] D. SULLIVAN, Genetics of homotopy theory and the Adams conjecture, *Annals of Math.* **100**, 1-79.
- [Thurston, 1982] W. THURSTON, Hyperbolic geometry and 3-manifolds, In: Low-dimensional topology, (Bangor, 1979), 9-25, *London Math. Soc. Lecture Note Ser.*, 48, Cambridge Univ. Press, Cambridge-New York, 1982.
- [Thurston, 1986] ———, Hyperbolic structures on 3-manifolds. I. Deformation of acylindrical manifolds. *Ann. of Math.* (2) **124** (1986), 203-246.
- [Whitehead, 1942] G. W. WHITEHEAD, On the homotopy groups of spheres and rotation groups, *Ann. of Math.* **43** (1942), 634-640.