

# ANOTHER SUBDIVISION WHICH CAN NOT BE SHELLED

CHANDLER DAVIS<sup>1</sup>

**1. Introduction.** Let  $B$  be a 3-cell, and consider a subdivision of  $B$  into 3-cells  $B_1, \dots, B_k$ . Let  $K_i$  denote the union of the  $B_j$  other than  $B_i$ . I will say that the subdivision *can not be shelled* in case, for each  $i$ ,  $K_i$  is not a 3-cell. Several examples of such subdivisions have been constructed, see [3]; the most remarkable is M. E. Rudin's [1], in which  $B$  and all the  $B_i$  are tetrahedra ( $k$  is 41). The example given here is quite simple, and the value of  $k$  is obviously minimal! However, it is not a *simplicial* decomposition.

**THEOREM 1.** *There exists a subdivision of the 3-cell into three pieces which can not be shelled.*

**2. The example.** Introduce Cartesian coordinates  $x_1, x_2, x_3$ . Let  $B$  be  $x_1^2 + x_2^2 + x_3^2 \leq M^2$ , where  $M$  is a sufficiently large positive number. Let  $T_i$  be the line  $x_{i+1} = 1, x_{i+2} = -1$ ; these subscripts are modulo 3. Let  $B_i$  consist of those points  $p$  of  $B$  such that

$$d(p, T_i) = \min_{j=1,2,3} d(p, T_j).$$

(Here  $d(\cdot, \cdot)$  is Euclidean distance between closed sets.)

$B_1$  is a 3-cell by the following argument. It is permissible to ignore the small caps where  $x_1^2 \geq M^2 - 1$ . Consider any other cross-section of  $B_1$  by a plane  $x_1 = \text{constant}$ . It is star-shaped about the point where  $T_1$  cuts it; for going from  $p = (x_1, x_2, x_3) \in B_1$  to  $p_\lambda = (x_1, 1 + \lambda(x_2 - 1), -1 + \lambda(x_3 + 1))$  with  $0 \leq \lambda < 1$  reduces the distance to  $T_1$  by  $d(p, p_\lambda)$  and can not reduce the distance to  $T_2$  or  $T_3$  by more. It contains a centered disk, namely  $(x_2 - 1)^2 + (x_3 + 1)^2 \leq 1$ . It is closed. This gives an obvious method for deforming  $B_1$  onto a cylinder.

$K_1 = B_2 \cup B_3$  is not a cell, for its fundamental group does not vanish. Namely, consider  $x_1 = 0, x_2^2 + x_3^2 = M^2$ . If  $p$  is on this circle,  $d(p, T_1) \geq M - 2^{1/2}$ , while at least one of  $d(p, T_2)$  and  $d(p, T_3)$  is  $\leq (M^2/2 + 1)^{1/2}$ . Since  $M$  is large,  $p \in K_1$ . This circle is not contractible in  $K_1$ , for  $K_1$  has no points in  $(x_2 - 1)^2 + (x_3 + 1)^2 < 1$ .

Because of the evident symmetry between the  $B_i$ , this proves Theorem 1.

### 3. Generalization.

**THEOREM 2.** *For  $k \geq 3$  there exists a subdivision of the 3-cell into  $k$  pieces which can not be shelled.*

Received by the editors March 25, 1958.

<sup>1</sup> National Science Foundation Fellow.

This result, which was proposed by O. G. Harrold, is clearly a consequence of the 3-dimensional case of the following theorem. (If the conclusion of Theorem 3 was simply that the subdivision could not be shelled, it could conversely be deduced from Theorem 2.)

**THEOREM 3.** *For  $k \geq n \geq 3$  there exists a subdivision of the  $n$ -cell into  $k$  pieces such that the union of any  $k-1$  of the pieces has nontrivial  $n-2$ th homotopy group.*

The idea of the construction is the same as for Theorem 1. Choose  $k$  vectors, the first  $n$  of which are the coordinate unit vectors, and no  $n$  of which are linearly dependent. Choose any nonintersecting lines  $T_1, \dots, T_k$  in these respective directions, such that each  $d(T_i, T_j)$  is at least 2. There exists  $D > 0$  such that each  $T_i$  passes within  $D$  of the origin. Imitate the definitions above of  $M, B$  and the  $B_i$ . The proof that  $B_i$  is a cell is not much affected.

The conclusion of the theorem concerns  $\pi_{n-2}(K_i)$ . We may choose Cartesian coordinates  $y_1, \dots, y_n$  such that  $T_i$  is the  $y_1$ -axis, and such that the  $n-2$ -sphere

$$y_1 = 0, y_2^2 + \dots + y_n^2 = (M - D)^2$$

lies in  $B$ . Then this sphere lies in  $K_i$ . (This is proved separately for  $i \leq n$  and  $i > n$ .) It is not contractible in  $K_i$ , for  $K_i$  has no points in  $y_2^2 + \dots + y_n^2 < 1$ .

**4. Another version of the example.** Let  $C_{1i}$  (for  $i = 1, 2, 3$ ) be the set of points with cylindrical coordinates

$$0 \leq z \leq 1, \quad 0 \leq r \leq 1,$$

$$\frac{2\pi}{3}(i - 1) \leq \theta - \frac{\pi}{2}r \leq \frac{2\pi}{3}i.$$

Similarly define  $C_{2i}$  by

$$1 \leq z \leq 2, \quad 0 \leq r \leq 1,$$

$$\frac{2\pi}{3}(i - 1) \leq \theta + \frac{\pi}{2}r \leq \frac{2\pi}{3}i.$$

Now if  $B_i = C_{1i} \cup C_{2i}$  and  $B = B_1 \cup B_2 \cup B_3$ , a figure is obtained which is homeomorphic to the one so labeled in §2. The two versions yield Theorem 1 equally easily, but this one leads to a different generalization.

**THEOREM 4.** *For any  $\nu=1, 2, 3, \dots$ , there exists a subdivision of the 3-cell into three pieces such that omitting any of the pieces leaves a set whose fundamental group is the free group on  $\nu$  generators.*

In fact, define  $C_{2\mu+1,i}$  as the result of translating  $C_{1,i}$  along the  $z$ -axis a distance  $2\mu$ ;  $C_{2\mu+2,i}$  similarly in terms of  $C_{2,i}$ . If  $B_i = C_{1,i} \cup \dots \cup C_{\nu+1,i}$  and  $B = B_1 \cup B_2 \cup B_3$ , the resulting figure may be shown to provide the example for Theorem 4.

I thank several colleagues, and especially R. H. Bing, for valuable conversations.

#### BIBLIOGRAPHY

1. M. E. Rudin, *An unshellable triangulation of a tetrahedron*, Bull. Amer. Math. Soc. vol. 64 (1958) pp. 90-91.
2. M. E. Rudin, to appear.
3. D. E. Sanderson, *Isotopy in 3-manifolds, I, Isotopic deformations of 2-cells and 3-cells*, Proc. Amer. Math. Soc. vol. 8 (1957) pp. 912-922.

INSTITUTE FOR ADVANCED STUDY