

SHORTER NOTES

The purpose of this department is to publish very short papers of an unusually elegant and polished character, for which there is normally no other outlet.

A NOTE ON CONTINUED FRACTIONS

S. DROBOT

It is well known that the convergents p_n/q_n of a continued fraction of a real number α are its best approximations, i.e. that for every rational $a/b \neq p_n/q_n$ with $1 \leq b \leq q_n$ and $n \geq 1$ there is

$$(1) \quad |q_n \alpha - p_n| < |b \alpha - a|.$$

The usually produced proofs of this fact use rather intricate arguments.¹ Here is a proof of (1) based on the two following elementary properties of the continued fraction

$$(i) \quad 1/q_{n+1} < |q_{n-1} \alpha - p_{n-1}| < 1/q_n,$$

$$(ii) \quad q_n |q_{n-1} \alpha - p_{n-1}| + q_{n-1} |q_n \alpha - p_n| = 1.$$

If $a/b = p_{n-1}/q_{n-1}$ inequality (1) holds by (i):

$$|q_n \alpha - p_n| < 1/q_{n+1} < |q_{n-1} \alpha - p_{n-1}|.$$

If $|aq_{n-1} - bp_{n-1}| \geq 1$ then

$$|a/b - \alpha| + |\alpha - p_{n-1}/q_{n-1}| \geq |a/b - p_{n-1}/q_{n-1}| \geq 1/(bq_{n-1})$$

i.e.

$$b |q_{n-1} \alpha - p_{n-1}| + q_{n-1} |b \alpha - a| \geq 1,$$

while the assumption $1 \leq b \leq q_n$ implies by (ii)

$$1 \geq b |q_{n-1} \alpha - p_{n-1}| + q_{n-1} |q_n \alpha - p_n|$$

whence

$$(1') \quad |q_n \alpha - p_n| \leq |b \alpha - a|.$$

Received by the editors February 19, 1962.

¹ See e.g. O. Perron, *Die Lehre von den Kettenbrüchen*, 1929, pp. 52-53; G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, 3rd ed., 1956, pp. 151-152; I. Niven, *Irrational numbers*, 1956, pp. 62-63; J. W. S. Cassels, *An introduction to Diophantine approximations*, 1957, pp. 2-4; A. Ya. Khintchine, *Continued fractions* (in Russian), 2nd ed., 1949, pp. 38-40.

Equality in (1') is for irrational α impossible in view of $a/b \neq p_n/q_n$ assumed. For rational α excluding equality in (1') requires, strangely enough, additional argument which may run as follows. Substitute into (1') with equality presumed

$$\alpha = \frac{P}{Q} = \frac{p_{n-1}r_n + p_{n-2}}{q_{n-1}r_n + q_{n-2}}$$

with rational r_n , and $Q \geq q_n$, to get

$$|bP - aQ| = |r_n - a_n|,$$

where a_n is the n th partial quotient in the continued fraction expansion of α . The last equality shows that r_n is an integer, thus, by Euclid's algorithm, $r_n = a_n$ whence $a/b = P/Q = p_n/q_n$ contrary to assumption.

UNIVERSITY OF NOTRE DAME