HOMOGENEOUS ALMOST COMPLEX SPACES OF POSITIVE CHARACTERISTIC

SAMUEL PASIENCIER

In this paper we give a necessary and sufficient condition for a homogeneous space of positive Euler characteristic to admit an invariant almost complex structure (i.a.c.s.), Theorem 4.1. A classification of these spaces has already been accomplished by R. Hermann [3], but no characterization of them has been given, to date. This work is a portion of the author's dissertation directed by H. C. Wang.

1. Let G be a compact, connected Lie group, and L a closed subgroup of G , $\mathfrak{g}$ and $\mathfrak{l}$ their Lie algebras. Then as is well known, L is reductive in G and we may write $\mathfrak{g} = \mathfrak{l} + \mathfrak{m}$, where $\text{Ad } L: \mathfrak{m} \rightarrow \mathfrak{m}$. Moreover by homogeneity, a necessary and sufficient condition that there exist an almost complex structure on G/L , is that there exist a linear transformation $J: \mathfrak{m} \rightarrow \mathfrak{m}$, such that $J^2 = -I$ (I , the identity), and such that $\text{Ad } lJ = J \text{ Ad } l$ for all $l \in L$. See for example, A. Frölicher [2].

2. We give first a sufficient condition in the general case.

THEOREM 2.1. *Let Z be a closed, abelian subgroup of a compact Lie group G , and L the identity component of the centralizer of Z in G . If each 1-dimensional subspace of $\mathfrak{g}$ invariant by $\text{Ad } Z$ is pointwise fixed, then G/L has an i.a.c.s.*

PROOF. Z being compact and abelian, $\text{Ad } Z$ is completely reducible; with notation as in 1, there cannot exist any 1-dimensional subspaces of $\mathfrak{m}$ reducing $\text{Ad } Z$, for if $X \neq 0$ belonged to such a subspace, then by hypothesis we would have $\text{Ad } z(X) = X$ for all $z \in Z$, and thus $X \in \mathfrak{l}$, which is impossible.

If $C\mathfrak{m}$ denotes the complexification of $\mathfrak{m}$, and $\text{Ad } Z$ and $\text{Ad } L$ are extended by linearity to $C\mathfrak{m}$, then we know by Schur's Lemma that $\text{Ad } Z$ acting on $C\mathfrak{m}$ has a weight vector v_i . That is, $\text{Ad } z(v_i) = \psi_i(z)v_i$ for all $z \in Z$, ψ_i being a character on Z . Moreover, since $\text{Ad } Z$ is real, $\text{Ad } z(\bar{v}_i) = \bar{\psi}_i(z)\bar{v}_i$, thus the weight vectors and roots appear in conjugate pairs.

Let $v_1, \dots, v_n, \bar{v}_1, \dots, \bar{v}_n$ be the weight vectors with characters $\psi_1, \dots, \psi_n, \bar{\psi}_1, \dots, \bar{\psi}_n$, where some of the ψ_i 's may be identical. If

Received by the editors September 4, 1962 and, in revised forms, December 3, 1962 and April 9, 1963.

we let $\mathfrak{U}$ be the subspace of $C\mathfrak{M}$ spanned by the v_i 's, then each $x \in \mathfrak{M}$ has a unique expression of the form $x = v + \bar{v}$, where $v \in \mathfrak{U}$ and $\bar{v} \in \bar{\mathfrak{U}}$. For putting $x_j = v_j + \bar{v}_j$, and $x'_j = iv_j - i\bar{v}_j$, the x_j 's and x'_j 's span the irreducible subspaces of $\mathfrak{M}$. And thus any $x \in \mathfrak{M}$ has a unique expression

$$x = \sum a_j x_j + \sum b_j x'_j,$$

so

$$x = \sum \lambda_j v_j + \sum \bar{\lambda}_j \bar{v}_j.$$

Furthermore, $\mathfrak{U}$ is invariant by $\text{Ad } L$, for if $z \in Z$, $l \in L$, then $\text{Ad } z \text{ Ad } l(v_i) = \text{Ad } l \text{ Ad } z(v_i) = \psi_i(z) \text{ Ad } l(v_i)$, that is, $\text{Ad } l(v_i)$ belongs to the character ψ_i , and by hypothesis there can be no real characters, thus $\psi_i \neq \bar{\psi}_j$. Now defining $J: \mathfrak{M} \rightarrow \mathfrak{M}$ by $J(x) = J(v + \bar{v}) = iv - i\bar{v}$, defines an i.a.c.s. on G/L .

COROLLARY 2.1. *If the number of connected components in $\text{Ad } Z$ is odd, then G/L has an i.a.c.s., G, L, Z as above.*

PROOF. Suppose there exists a 1-dimensional subspace of $\mathfrak{M}$, with basis X , reducing $\text{Ad } Z$, then for each $z \in Z$, $\text{Ad } z(X) = \psi(z)X$, where $\psi(z)$ is a homomorphism of $Z \rightarrow Z_2 = \{+1, -1\}$. By continuity ψ is constant on components of $\text{Ad } z$, and takes the value $+1$ on the identity component $(\text{Ad } z)^0$. It thus induces a homomorphism $\psi: \text{Ad } z/(\text{Ad } z)^0 \rightarrow Z_2$, but $\text{Ad } z/(\text{Ad } z)^0$ is a finite group of odd order, therefore its image is identically 1, so that $\psi \equiv 1$, and thus $\text{Ad } z$ is pointwise fixed on each 1-dimensional subspace reducing it.

3. Next we consider the case that L is semi-simple.

LEMMA 3.1. *Let L, H be two closed subgroups of G , $L \subset H \subset G$. And suppose that G/L has an i.a.c.s. defined by $J: \mathfrak{M} \rightarrow \mathfrak{M}$, where $\mathfrak{g} = \mathfrak{L} + \mathfrak{M}$. If further $\mathfrak{K} = \mathfrak{L} + \mathfrak{M}_1$, $\mathfrak{M}_1 \subset \mathfrak{M}$, and $J(\mathfrak{M}_1) = \mathfrak{M}_1$, then H/L has an i.a.c.s.*

PROOF. Since H is a subgroup of G , $\text{Ad } L: \mathfrak{M}_1 \rightarrow \mathfrak{M}_1$. Thus applying 1, we see that H/L has an i.a.c.s.

LEMMA 3.2. *Let L be a closed semi-simple subgroup of G , of maximal rank, such that G/L has an i.a.c.s.; then L is not a symmetric subgroup.*

PROOF. Let $\mathfrak{g} = \mathfrak{L} + \mathfrak{M}$, and $J: \mathfrak{M} \rightarrow \mathfrak{M}$ define the i.a.c.s. as in §1. Since the torsion tensor of J is given by

$$\mathfrak{F}(X, Y) = [JX, JY] - J[JX, Y] - J[X, JY] + [X, Y] \bmod \mathfrak{L}$$

for $X, Y \in \mathfrak{M}$ [2], if we assume G/L is symmetric $\mathfrak{F}(X, Y) \equiv 0$, for

symmetric implies $[\mathfrak{M}, \mathfrak{M}] \subset L$. Thus J defines a complex structure on G_0/L_0 (where subscripts denote identity components), but then a classical result of E. Cartan says that L_0 (and thus L) cannot be semi-simple.

LEMMA 3.3. *If G/L has an i.a.c.s. and L is connected semi-simple of maximal rank, Z the center of L ; then $\text{Ad}(Z)$ acting on $\mathfrak{M}$ has no 1-dimensional invariant subspaces.*

PROOF. Suppose by contradiction that $\mathfrak{R} \subset \mathfrak{M}$ is 1-dimensional and reduces $\text{Ad}(Z)$. Let $x \in \mathfrak{R}$, $x \neq 0$, then

$$\text{Ad } z(x) = \psi(z)x, \quad z \in Z, \text{ and } \psi \text{ is a real character on } Z.$$

Since Z is compact, either $\psi(Z) \equiv 1$, or $\psi(Z) = \{+1, -1\}$. From Borel-deSiebenthal [1], we know that L is the connected component of the centralizer of its own center Z , and thus $\mathfrak{L} = \{u: u \in \mathfrak{g}, \text{Ad } Zu = u\}$. If $\psi(Z) \equiv 1$, then $\mathfrak{R} \subset \mathfrak{L}$, which is impossible. Thus $\psi(Z) = \{1, -1\}$.

Let $\mathfrak{Q} = \{y: y \in \mathfrak{M}, \text{Ad } z(y) = \psi(z)y, z \in Z\}$ be the weight space corresponding to the character ψ . Then $J(\mathfrak{Q}) = \mathfrak{Q}$, for if $y \in \mathfrak{Q}$,

$$\text{Ad } z J(y) = J \cdot \text{Ad } z(y) = J\psi(z)y = \psi(z)Jy.$$

Furthermore, $\text{Ad } L(\mathfrak{Q}) = \mathfrak{Q}$, for if $y \in \mathfrak{Q}$, $l \in L$, then

$$\text{Ad } z \text{Ad } ly = (\text{Ad } l)\psi(z)y = \psi(z)\text{Ad } ly.$$

But then

$$[\mathfrak{L}, \mathfrak{Q}] \subset \mathfrak{Q}.$$

Finally if $y, y' \in \mathfrak{Q}$, $\text{Ad } z[y, y'] = \psi(z)^2[y, y'] = [y, y']$, that is, $[y, y']$ is fixed under all $\text{Ad } z$, so that

$$[y, y'] \in \mathfrak{L}, \text{ that is } [\mathfrak{Q}, \mathfrak{Q}] \subset \mathfrak{L}.$$

Therefore, $\mathfrak{H} = \mathfrak{L} + \mathfrak{Q}$ is a subalgebra and H/L is a symmetric space, $H = \exp \mathfrak{H}$. Since H is of maximal rank, H is closed by the theorem in §8 of [1]. By Lemma 3.1, H/L has an invariant almost complex structure, and H/L is nontrivial since $\mathfrak{Q} \neq 0$, for $\mathfrak{R} \subset \mathfrak{Q}$. This contradicts Lemma 3.2, thus Lemma 3.3 is proven.

LEMMA 3.4. *Under the hypothesis of Lemma 3.3, $\text{Ad}(Z)$ cannot be a group of order 2^k , $k > 0$.*

PROOF. (a) If $k = 1$, G/L is symmetric and this is impossible by Lemma 3.2.

(b) Suppose the lemma is true for all $k \leq n$, and consider the case that $\text{Ad } Z$ is of order 2^{n+1} . Since $\text{Ad } Z$ contains a subgroup of order 2,

there exists an element $z_0 \in Z$ such that $\text{Ad } z_0 \neq 1$ and $(\text{Ad } z_0)^2 = 1$. Let G^* be the 1-component of the centralizer of z_0 in G . Then $L \subset G^*$, and we have

$$\mathfrak{G} = \mathfrak{L} + \mathfrak{M}, \quad \mathfrak{M} = \mathfrak{Q}_1 + \mathfrak{Q}_2, \quad \mathfrak{G}^* = \mathfrak{L} + \mathfrak{Q}_1,$$

where $\text{Ad } z_0|_{\mathfrak{Q}_1} = 1$ and $\text{Ad } z_0|_{\mathfrak{Q}_2} = -1$, $\mathfrak{G}^*$ being the Lie algebra of G^* . Moreover $J(\mathfrak{Q}_1) = \mathfrak{Q}_1$ for if $m \in \mathfrak{Q}_1$, $(\text{Ad } z)J(m) = J(\text{Ad } z)m = J(m)$. Similarly we show that $\text{Ad } L(\mathfrak{Q}_1) = \mathfrak{Q}_1$.

G^*/L has i.a.c.s. by Lemma 3.1. But since $L \subset G^*$, Z is the center of L in G^* . Since $\text{Ad } z_0|_{\mathfrak{G}^*} = 1$, $\text{Ad } Z|_{\mathfrak{G}^*}$ is of order 2^m , $m \leq n$. Thus by the inductive hypothesis $m = 0$ and $G^* = L$. But then G/L is a symmetric space, contradicting Lemma 3.2. Thus G/L cannot have invariant almost complex structure, and the lemma is proved.

THEOREM 3.5. *Let G be a compact connected Lie group and L a connected semi-simple subgroup of G of maximal rank; then G/L has invariant almost complex structure if and only if L is the identity component of the centralizer in G of a finite subgroup F , of its center, such that $\text{Ad}(F)$ is of odd order.*

PROOF. The sufficiency of this condition was proved in Theorem 2.1. To prove the necessity, let Z be the center of L ; then by the fundamental theorem of Abelian groups we may write $\text{Ad } Z = \text{Ad } F_1 \times \text{Ad } F_2$, where $\text{Ad } F_1$ is of odd order and $\text{Ad } F_2$ is of order 2^k .

Let G_1 be the identity component of the centralizer of F_1 in G . We shall show that $G_1 = L$. Suppose on the contrary that $L \subsetneq G_1$. Then G_1/L has i.a.c.s. by Lemma 3.1. If $\text{Ad}(F_2): \mathfrak{G}_1 \rightarrow \mathfrak{G}_1$ is identically 1, then we are done because $\mathfrak{G}_1$ centralizes both F_1 and F_2 and by the result of Borel and de Siebenthal [1], $G_1 = L$. But $\text{Ad}(F_1)$ is identically 1 on $\mathfrak{G}_1$, and $\text{Ad}(F_2)$, if nontrivial on $\mathfrak{G}_1$, must be of order 2^n , for some $n > 1$. But then $\text{Ad } Z$ acting on $\mathfrak{G}_1$ would be of order 2^n , which is impossible. Therefore $\text{Ad}(F_2) \equiv 1$ on $\mathfrak{G}_1$.

4. THEOREM 4.1. *Let G be a compact connected Lie group, L a connected subgroup of maximal rank; then G/L possesses an invariant almost complex structure if and only if L is the connected component of the centralizer in G of a finite subgroup F , of its center, such that $\text{Ad}(F)$ is a finite group of odd order.*

The sufficiency of this condition follows from Corollary 2.1. In this general case when L is not necessarily semi-simple, we proceed by proving the following two lemmas.

LEMMA 4.2. *Let G be compact, connected, L a connected subgroup of maximal rank such that G/L has an i.a.c.s.; if $L = T^a \times L_1$, local direct*

product, where T^a is toral and L_1 is connected semi-simple, then L is the centralizer in G of $T^a \cup F_1$, where F_1 is a finite subgroup of L , such that $\text{Ad}(F_1)$ is of odd order.

PROOF. $L = T^a \times L_1$, $\mathfrak{L} = R^a \oplus \mathfrak{L}_1$, let $G_2 = \text{centralizer of } T^a \text{ in } G$, then G_2 is a connected subgroup of maximal rank in G , and

$$G_2 = T^a \times G_3, \quad \mathfrak{G}_2 = R^a \oplus \mathfrak{G}_3, \quad \text{where } G_3 \text{ is closed in } G_2.$$

By hypothesis there exists J , commuting with $\text{Ad}(L)$ such that

$$J: \mathfrak{G}_2/\mathfrak{L} \rightarrow \mathfrak{G}_2/\mathfrak{L},$$

but $\mathfrak{L}_1 \subset \mathfrak{L}$, and there exists an isomorphism

$$\phi: \mathfrak{G}_2/\mathfrak{L} \rightarrow \mathfrak{G}_3/\mathfrak{L}_1.$$

Define J' on $\mathfrak{G}_3/\mathfrak{L}_1$ by $J' = \phi J \phi^{-1}$, then J' defines an almost complex structure on G_3/L_1 . Then, by 3.5, L_1 is the connected component of the centralizer in G_3 of a finite subgroup F_1 of its center, such that $\text{Ad}(F_1)$ is of odd order.

Now, if $x \in L$, then $x = tl_1$, where $t \in T^a$ and $l_1 \in L_1$. If $f_1 \in F_1$, then $(t \cdot l_1)f_1 = tf_1l_1 = f_1(tl_1)$ since t is central, and $f_1l_1 = l_1f_1$ since $l_1 \in L_1$. Thus x commutes with F_1 , and x commutes with T^a .

Conversely, if x belongs to the connected component of the centralizer of $T^a \cup F_1$, then $x \in G_2$, so that $x = tg_3$. Moreover, since x commutes with F_1 , we have $xf_1 = f_1x$, so that $tg_3f_1 = f_1tg_3 = tf_1g_3$, so that $g_3f_1 = f_1g_3$, thus $g_3 \in L_1$, and $x \in T^a \times L_1$, that is, $x \in L$.

LEMMA 4.3. *Let G be compact, connected Lie group and L a subgroup of G , such that L is the centralizer in G of a toral subgroup T^a . Then there exists an $x \in T^a$ such that (1) L is the connected component of the centralizer of x in G , and (2) $\text{Ad}(x)$ is of odd order.*

PROOF OF (1). $T^a \subset T^b$, where T^b is a maximal torus of G . If θ_i , $i = 1, \dots, m$, are the root forms, or angular parameters of T^b , and R^b is the covering space of T^b , then R^a , the covering space of T^a , is characterized by $R^a = \{x: \theta_i(x) = 0, i = 1, \dots, m', m' \leq m\}$, where θ_i , $i = 1, \dots, m'$, are angular parameters of T^a . If $x \in T^a$, and $Z(x)$ denotes the connected component of the centralizer of x in G , then $L \subset Z(x)$. Moreover $Z(x)$ is minimal if $\theta_i(x) \neq 0$, $i = m' + 1, \dots, m$. But each plane $\theta_i(x) = 0$, $i = m' + 1, \dots, m$, determines a hyperplane of R^a , and the union of these finitely many hyperplanes cannot be all of R^a , thus there exist $x \in R^a$, $\theta_i(x) \neq 0$, $i = m' + 1, \dots, m$.

PROOF OF (2). Given any Euclidean space R , the set of elements with coordinates whose denominators are all odd is dense in R , because the set of points with rational coordinates is dense in R . And

if $x = (r_1, \dots, r_n)$ where $r_i = p_i/q_i$, p_i, q_i relatively prime, is a point with rational coordinates and

$$x_m = \left(\frac{2mp_1}{2mq_1 + 1}, \frac{2mp_2}{2mq_2 + 1}, \dots, \frac{2mp_n}{2mq_n + 1} \right),$$

then $\lim_{m \rightarrow \infty} x_m = x$.

Now let $\bar{\theta}_i, i = 1, \dots, b$, be a basis for R^b , consisting of root vectors. Thus if $x \in R^b$, then $\langle \bar{\theta}_i, x \rangle = \theta_i(x)$ defines an inner product on R^b , and if $x = \sum x_i \bar{\theta}_i$, $\langle \bar{\theta}_i, x \rangle = x_i$ so that all $\theta_i(x) \equiv 0 \pmod{1}$, if x has integral coordinates in this basis, that is, $\text{Ad } x = 1$.

Now if $x \in T^a$, such that $Z(x) = \mathcal{L}$, then it does not lie on any of the planes $\theta_i(x) = 0, i = m' + 1, \dots$, which is a closed set, and there exists a point z with rational coordinates, with odd denominators, arbitrarily close to x , so that $Z(x) = \mathcal{L}$. But $\text{Ad}(z^k) = 1$, where k = the least common multiple of the denominators of the coordinates of z . And, $\text{Ad}(z^k) = \text{Ad}^k(z)$.

Combining the last two lemmas, we have that since L is the centralizer of $x \in T^a$, such that $\text{Ad}(x)$ is of odd order and L is the connected component of the centralizer of F_1 , then L is the connected component of the centralizer, in G , of the group generated by x and F_1 , whose adjoint is of odd order.

BIBLIOGRAPHY

1. A. Borel and J. deSiebenthal, *Les sous-groupes fermés de rang maximum des groupes de Lie clos*, Comment. Math. Helv. **23** (1949), 200–221.
2. A. Frölicher, *Zur Differentialgeometrie der komplexen Strukturen*, Math. Ann. **129** (1955), 50–95.
3. R. Hermann, *Compact homogeneous almost complex spaces of positive characteristic*, Trans. Amer. Math. Soc. **83** (1956), 471–481.
4. H. C. Wang, *Closed manifolds with homogeneous complex structure*, Amer. J. Math. **76** (1954), 1–32.

LAKE FOREST COLLEGE
AND NEW YORK UNIVERSITY