

## MODULAR FORMS ON HECKE'S MODULAR GROUPS

RONALD J. EVANS

**ABSTRACT.** Let  $H = \{\tau = x + iy : y > 0\}$ . Let  $\lambda > 0$ ,  $k > 0$ ,  $\gamma = \pm 1$ . Let  $M(\lambda, k, \gamma)$  denote the set of functions  $f$  for which  $f(\tau) = \sum_{n=0}^{\infty} a_n e^{2\pi i n \tau / \lambda}$  and  $f(-1/\tau) = \gamma(\tau/i)^k f(\tau)$ , for all  $\tau \in H$ . Let  $M_0(\lambda, k, \gamma)$  denote the set of  $f \in M(\lambda, k, \gamma)$  for which  $f(\tau) = O(y^c)$  uniformly for all  $x$  as  $y \rightarrow 0^+$ , for some real  $c$ . We give a new proof that if  $\lambda = 2 \cos(\pi/q)$  for an integer  $q \geq 3$ , then  $M(\lambda, k, \gamma) = M_0(\lambda, k, \gamma)$ .

Petersson [5, p. 176] and Ogg [4] filled a gap in Hecke's work [2, p. 21] by establishing analytically the theorem below. We present here a short, elementary proof which uses no non-Euclidean geometry.

**THEOREM.** Let  $\lambda = 2 \cos(\pi/q)$  for an integer  $q \geq 3$ . Then  $M(\lambda, k, \gamma) = M_0(\lambda, k, \gamma)$ .

**PROOF.** Let  $f \in M(\lambda, k, \gamma)$ . Let  $H_1 = \{\tau \in H : |x| \leq \lambda/2, y \leq 1\}$ . Since  $f(\tau) = f(\tau + \lambda)$ , it suffices to show that  $|y^k f(\tau)|$  is uniformly bounded for all  $\tau \in H_1$ .

Let  $B(\lambda) = \{\tau \in H : |x| < \lambda/2, |\tau| > 1\}$  and let  $\text{Cl}(B(\lambda))$  denote the closure of  $B(\lambda)$ . For large  $y$ ,  $|f(\tau)| < |a_0| + 1$ , and since  $f$  is bounded on compact subsets of  $H$ , there is a constant  $A$  such that  $|f(\tau)| \leq A$  for all  $\tau \in \text{Cl}(B(\lambda))$ .

Hecke's modular group  $G(\lambda)$  is defined to be the group of linear fractional transformations generated by  $S_\lambda : \tau \rightarrow \tau + \lambda$  and  $T : \tau \rightarrow -1/\tau$ . We shall identify the transformation  $\tau \rightarrow (a\tau + b)/(c\tau + d)$  with the matrix  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ . Hecke proved [2, pp. 11–20] that  $B(\lambda)$  is a fundamental region (as defined in [3, p. 20]) for  $G(\lambda)$ . (For an elementary proof, see [1].) Thus for each  $\tau \in H$ , there exists

$$V_\tau = \begin{pmatrix} a_\tau & b_\tau \\ c_\tau & d_\tau \end{pmatrix} \in G(\lambda)$$

such that  $V_\tau \tau \in \text{Cl}(B(\lambda))$ .

It can be easily shown that for all  $\tau \in H$  and for all  $V = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G(\lambda)$ ,

$$|f(\tau)| = |f(V\tau)| \cdot |c\tau + d|^{-k}.$$

---

Presented to the Society March 31, 1972; received by the editors May 11, 1972.

AMS (MOS) subject classifications (1970). Primary 10D05, 10D15; Secondary 30A20, 30A58.

*Key words and phrases.* Modular form, Hecke modular groups, fundamental region, equivalent points.

© American Mathematical Society 1973

Thus for all  $\tau \in H$ ,

$$\begin{aligned} |y^k f(\tau)| &= y^k |f(V_\tau \tau)| \cdot |c_\tau \tau + d_\tau|^{-k} \leq y^k A \cdot |c_\tau \tau + d_\tau|^{-k} \\ &= A |ic_\tau + (c_\tau x + d_\tau)/y|^{-k}. \end{aligned}$$

We shall now show that for all  $\tau \in H_1$  and for all  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G(\lambda)$ ,

$$|ic + (cx + d)/y|^2 \geq 1 - \lambda/2.$$

This will show that  $|y^k f(\tau)| \leq A(1 - \lambda/2)^{-k/2}$  for all  $\tau \in H_1$ , which proves our theorem. Fix  $\tau \in H_1$  and  $V = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G(\lambda)$ . Then

$$\begin{aligned} (cx + d)^2/y^2 + c^2 &\geq (cx + d)^2 + c^2 = c^2(x^2 + 1) + d^2 + 2cdx \\ &\geq c^2 + d^2 - \lambda |cd| \geq c^2 + d^2 - (\lambda/2)(c^2 + d^2) \\ &= (1 - \lambda/2)(c^2 + d^2). \end{aligned}$$

It remains to show that  $c^2 + d^2 \geq 1$ . Suppose that  $c^2 + d^2 < 1$ . Then  $\text{Im}(Vi) = 1/(c^2 + d^2) > 1$ , so  $i$  is  $G(\lambda)$ -equivalent to a point  $\tau_1 \in \text{Cl}(B(\lambda))$  such that  $\text{Im}(\tau_1) > 1$ . Thus, by continuity, some point in  $B(\lambda)$  close to  $i$  is  $G(\lambda)$ -equivalent to another point in  $B(\lambda)$  close to  $\tau_1$ , contradicting the fact that  $B(\lambda)$  is a fundamental region.  $\square$

#### REFERENCES

1. R. J. Evans, *A fundamental region for Hecke's modular group*, J. Number Theory (to appear).
2. E. Hecke, *Dirichlet series*, Planographed Lecture Notes, Princeton Institute for Advanced Study, Edwards Brothers, Ann Arbor, Mich., 1938.
3. J. Lehner, *Discontinuous groups and automorphic functions*, Math. Surveys, no. 8, Amer. Math. Soc., Providence, R.I., 1964. MR 29 #1332.
4. A. Ogg, *On modular forms with associated Dirichlet series*, Ann. of Math. (2) 89 (1969), 184–186. MR 38 #3232.
5. H. Petersson, *Über die Berechnung der Skalarprodukte ganzer Modulformen*, Comment. Math. Helv. 22 (1949), 168–199. MR 10, 445.

DEPARTMENT OF MATHEMATICS, JACKSON STATE COLLEGE, JACKSON, MISSISSIPPI 39217

Current address: Department of Mathematics, University of Illinois, Urbana, Illinois 61801