

## A REMARK ON $C_\sigma$ SPACES

SIMEON REICH

**ABSTRACT.** We give a simple new proof of the following result, conjectured by Effros and proved by Fakhoury: Let  $E$  be a  $C_\sigma$  space and  $Z$  the set of extreme points of the unit ball of  $E^*$ . Then  $Z \cup \{0\} = \{p \in E^*: \langle fgh, p \rangle = \langle f, p \rangle \langle g, p \rangle \langle h, p \rangle \text{ for all } f, g, h \text{ in } E\}$ .

Let  $C(X)$  be the Banach space of all continuous real valued functions on a compact Hausdorff space  $X$ , equipped with the supremum norm. If  $\sigma: X \rightarrow X$  is an involutory homeomorphism, then  $C_\sigma(X)$  is the subspace of  $C(X)$  consisting of those  $f$  in  $C(X)$  which satisfy  $f(\sigma x) = -f(x)$  for all  $x$  in  $X$ .  $e_x$  will stand for the point functional corresponding to a point  $x$  in  $X$ . If  $E$  is a Banach space, then we shall denote its unit ball by  $B(E)$  and its conjugate space by  $E^*$ . The set of extreme points of a subset  $Q$  of  $E$  will be denoted by  $\text{ext } Q$ .

The following result was conjectured by Effros [2, Remark 8.4] and proved by Fakhoury [4, Theorem 15].

**THEOREM.** *If  $E$  is a  $C_\sigma$  space, then  $\text{ext } B(E^*) \cup \{0\} = \{p \in E^*: \langle fgh, p \rangle = \langle f, p \rangle \langle g, p \rangle \langle h, p \rangle \text{ for all } f, g, h \text{ in } E\}$ .*

Fakhoury's proof is measure-theoretic in nature. Since this theorem appears to be useful (see, for instance, [4, Theorem 25] and [3, Theorem 11]), a simple different proof might perhaps be of some interest. The purpose of this note is to present such a proof. We shall need a few auxiliary propositions.

**LEMMA 1** [1, p. 89]. *If  $E = C_\sigma(X)$ , then  $\text{ext } B(E^*) = \{e_x: x \in X \text{ and } x \neq \sigma x\}$ .*

**LEMMA 2** [5, PROPOSITION 3.5]. *Let  $E$  be a Banach space and  $J^*$  the adjoint of the canonical map  $J: E \rightarrow E^{**}$ . If  $y \in \text{ext } B(E^{***})$ , then  $J^*y$  belongs to the weak star closure of  $\text{ext } B(E^*)$ .*

Since the dual of a  $C_\sigma$  space is isometric to an  $L$  space, its second dual is isometric to  $C(Y)$  for some (extremally disconnected) compact Hausdorff space  $Y$ .

---

Received by the editors November 9, 1972.

AMS (MOS) subject classifications (1970). Primary 46E15.

Key words and phrases.  $C_\sigma$  space, extreme point.

© American Mathematical Society 1973

LEMMA 3. Let  $E = C_\sigma(X)$  and  $E^{**} = C(Y)$ . Then  $J(fgh) = J(f)J(g)J(h)$  for all  $f, g, h$  in  $E$ .

PROOF. Let  $y$  belong to  $Y$ . The previous lemmas imply that  $J^*e_y$  belongs to  $\{e_x: x \in X \text{ and } x \neq \sigma x\} \cup \{0\}$ . Therefore  $J(fgh)(y) = \langle J(fgh), e_y \rangle = \langle fgh, J^*e_y \rangle = \langle f, J^*e_y \rangle \langle g, J^*e_y \rangle \langle h, J^*e_y \rangle = J(f)(y)J(g)(y)J(h)(y)$ .

LEMMA 4. If  $E = C(X)$ , then the Theorem is true.

PROOF. If  $p \neq 0$ , then the equality  $\langle f, p \rangle = \langle f, p \rangle \langle 1, p \rangle^2$ , valid for each  $f$  in  $E$ , implies that either  $\langle 1, p \rangle = 1$ , or  $\langle 1, p \rangle = -1$ . Clearly we may assume that the former possibility holds. Then  $p$  is multiplicative on  $C(X)$ , hence an extreme point of the positive face of  $B(E^*)$ .

PROOF OF THE THEOREM. By Lemma 3,  $\langle p, J(f)J(g)J(h) \rangle = \langle p, J(fgh) \rangle = \langle fgh, p \rangle = \langle f, p \rangle \langle g, p \rangle \langle h, p \rangle = \langle p, J(f) \rangle \langle p, J(g) \rangle \langle p, J(h) \rangle$  for all  $f, g, h$  in  $E$ . Let  $K: E^* \rightarrow E^{***}$  be the canonical map. For each  $r$  in  $E^{**} = C(Y)$ , the operator defined on  $C(Y)$  by  $s \rightarrow rs$  is weak star continuous. Therefore  $\langle p, rst \rangle = \langle p, r \rangle \langle p, s \rangle \langle p, t \rangle$  for all  $r, s, t$  in  $C(Y)$ . By Lemma 4,  $Kp$  belongs to  $\text{ext } B(C(Y)^*) \cup \{0\}$ . Since  $p = J^*Kp$ , an appeal to Lemmas 1 and 2 concludes the proof.

ACKNOWLEDGEMENT. I wish to thank my friend, Micha Sharir, for his valuable help.

#### REFERENCES

1. M. M. Day, *Normed linear spaces*, 2nd rev. ed., Academic Press, New York; Springer-Verlag, Berlin, 1962. MR 26 #2847.
2. E. G. Effros, *On a class of real Banach spaces*, Israel J. Math. 9 (1971), 430–458.
3. H. Fakhoury, *Préduaux de L-espaces et éléments extrémaux*, C. R. Acad. Sci. Paris Sér. A-B 272 (1971), A1703–A1706. MR 43 #6700.
4. ———, *Préduaux de L-espaces: notion de centre*, J. Functional Analysis 9 (1972), 189–207.
5. M. Sharir, *Extremal structure in operator spaces* (to appear).

DEPARTMENT OF MATHEMATICS, THE TECHNION-ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA, ISRAEL

DEPARTMENT OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, TEL AVIV, ISRAEL