

SHORTER NOTES

The purpose of this department is to publish very short papers of an unusually elegant and polished character, for which there is no other outlet.

ON A CLASS OF ANALYTIC FUNCTIONS

NOBUYUKI SUITA

ABSTRACT. We show that the class \mathfrak{E}_0 of analytic functions f in a plane region $\Omega \notin O_{AB}$ vanishing at $z_0 \in \Omega$ and such that $1/f$ omits a set of values of area $\geq \pi$ is not compact. Here O_{AB} denotes the class of Riemann surfaces which have no nonconstant bounded analytic functions. We remark that the extremal functions maximizing $|f'(z_0)|$ in \mathfrak{E}_0 coincide with linear transformations $w/(1-cw)$ of those for the same problem in the class \mathfrak{B}_0 consisting of functions f such that $f(z_0)=0$ and $|f(z)| \leq 1$, i.e. so-called Ahlfors functions. Here $1/c$ is an omitted value of the Ahlfors function.

Under the notations in the above abstract Ahlfors and Beurling [1] stated that the classes \mathfrak{B}_0 and \mathfrak{E}_0 are both compact and proved that the maxima of $|f'(z_0)|$ in \mathfrak{B}_0 and \mathfrak{E}_0 are equal. However, we can show that the alleged compactness of \mathfrak{E}_0 is not true by constructing a counterexample: For the annulus $\frac{1}{2} < |z| < 2$, the functions $f_n = (\frac{3}{2})^n(z^n - (\frac{3}{2})^n)/z^n$, $n=1, 2, \dots$, belong to \mathfrak{E}_0 with $z_0 = \frac{3}{2}$. Then $\{f_n\}$ tends to zero for $\frac{9}{8} + \delta < |z| < 2$ and to infinity for $\frac{1}{2} < |z| < \frac{9}{8} - \delta$, $\delta > 0$ as $n \rightarrow \infty$.

If $\Omega \notin O_{AB}$, there exist the extremal functions $A(z)$ which maximize $|f'(z_0)|$ in \mathfrak{B}_0 . Those functions are called the *Ahlfors functions* which are unique save for rotations [3]. If $1/c$ is an omitted value of $A(z)$, $A(z)/(1-cA(z))$ belongs to \mathfrak{E}_0 . By the result cited above it is extremal for the problem in \mathfrak{E}_0 .

For any extremal $g \in \mathfrak{E}_0$, let E be the set of all omitted values of g . From the extremality of g the area of E is equal to π . They used a transformation

$$\Phi\left(\frac{1}{g}\right) = \frac{1}{\pi} \iint_E \frac{du \, dv}{\frac{1}{g} - w}, \quad w = u + iv,$$

Received by the editors April 17, 1973 and, in revised form, June 4, 1973.

AMS (MOS) subject classifications (1970). Primary 30A40, 30A42.

Key words and phrases. Extremal problem, compact family, bounded functions.

and proved $\Phi(1/g) \in \mathfrak{B}_0$ with $\Phi'(1/g(z_0)) = g'(z_0)$ [1]. Hence $\Phi(1/g)$ is an Ahlfors function and there exists a point $\omega \in E$ such that $|\Phi(\omega)| = 1$. Note that $\Phi(w)$ is continuous in the whole plane [2]. Without loss of generality (by a rotation) we set $\Phi(\omega) = 1$ and $E_+ = E \cap \{\operatorname{Re}(w - \omega) \geq 0\}$. Then from the equality statement for Schwarz's inequality in their proof we infer that E_+ coincides with the disc $r \leq 2 \cos \theta$, $|\theta| \leq \pi/2$ ($w - \omega = re^{i\theta}$), except for a set of area zero. We can deduce, from $\Phi(\omega) = 1$, that the area of $E - E_+$ vanishes. Denoting by c the center of the above disc, by a direct calculation we see that $\Phi(w)$ reduces to a linear transformation $1/(w - c)$ for $|w - c| \geq 1$. Hence we have $g = \Phi(1/g)/(1 + c\Phi(1/g))$. Clearly $-1/c$ is an omitted value of the Ahlfors function $\Phi(1/g)$ and therefore g is of the form stated in the abstract.

REFERENCES

1. L. V. Ahlfors and A. Beurling, *Conformal invariants and function-theoretic null-sets*, Acta Math. **83** (1950), 101–129. MR **12**, 171.
2. A. Denjoy, *Sur fonctions analytiques uniformes qui restent continues sur un ensemble parfait discontinu de singularités*, C.R. Acad. Sci. Paris **149** (1909), 1154–1156.
3. S. Ja. Havinson, *Analytic capacity of sets, joint nontriviality of various classes of analytic functions and the Schwarz lemma in arbitrary domains*, Mat. Sb. **54** (96) (1961), 3–50; English transl., Amer. Math. Soc. Transl. (2) **43** (1964), 215–266. MR **25** #182.

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, OH-OKAYAMA, MEGURO-KU, TOKYO, JAPAN