

## $K_1$ OF THE COMPACT OPERATORS IS ZERO<sup>1</sup>

L. G. BROWN AND CLAUDE SCHOCHET

**ABSTRACT.** We prove that  $K_1$  of the compact operators is zero. This theorem has the following operator-theoretic formulation: *any invertible operator of the form (identity) + (compact) is the product of (at most eight) multiplicative commutators  $(A_j B_j A_j^{-1} B_j^{-1})^{\pm 1}$ , where each  $B_j$  is of the form (identity) + (compact).* The proof uses results of L. G. Brown, R. G. Douglas, and P. A. Fillmore on essentially normal operators and a theorem of A. Brown and C. Pearcy on multiplicative commutators.

**1. Statement of results.** Let  $\mathcal{L}$  be the bounded operators on a separable, infinite dimensional Hilbert space,  $\mathcal{K}$  the closed two-sided ideal of compact operators, and  $\mathfrak{A} = \mathcal{L}/\mathcal{K}$  the Calkin algebra.

**THEOREM.**  $K_1(\mathcal{K}) = 0$ .

That is, the “algebraic  $K_1$ ” of  $\mathcal{K}$ , regarded as an ideal in  $\mathcal{L}$ , is zero. The result may be interpreted as follows. Let  $G$  be the set of invertible operators in  $\mathcal{L}$  of the form  $I + K$ , where  $K \in \mathcal{K}$ . Let  $H$  denote the subgroup of  $G$  generated by all multiplicative commutators  $(u, g) = ugu^{-1}g^{-1}$  where  $u \in \mathcal{L}$  is invertible and  $g \in G$ . Then  $G = H$ . (This uses the definition of  $K_1$  [8, p. 36] and the fact that the matrix rings  $M_n(\mathcal{K})$  and  $M_n(\mathcal{L})$  are isomorphic to  $\mathcal{K}$  and  $\mathcal{L}$  respectively.) So the theorem is equivalent to the following operator-theoretic proposition.

**PROPOSITION.** *Let  $I + K$  be invertible with  $K \in \mathcal{K}$ . Then there exist invertible operators  $A_j$  and  $B_j = I + K_j$  ( $j = 1, \dots, n$ ) with  $K_j \in \mathcal{K}$  such that*

$$I + K = \prod_{j=1}^n (A_j, B_j)^{\pm 1}.$$

*In fact  $n \leq 8$ . This proposition is proved in §2.*

Combining the theorem with known information yields the first six terms of the Milnor long exact sequence in algebraic  $K$ -theory associated to  $\mathcal{K} \hookrightarrow \mathcal{L} \rightarrow \mathfrak{A}$ . It reads:

$$\begin{array}{ccccccccc} K_1(\mathcal{K}) & \longrightarrow & K_1(\mathcal{L}) & \longrightarrow & K_1(\mathfrak{A}) & \longrightarrow & K_0(\mathcal{K}) & \longrightarrow & K_0(\mathcal{L}) & \longrightarrow & K_0(\mathfrak{A}) & \longrightarrow & 0 \\ \parallel & & \\ 0 & & 0 & & \mathbf{Z} & & \mathbf{Z} & & 0 & & 0 & & \end{array}$$

Received by the editors April 2, 1975 and, in revised form, September 22, 1975.

AMS (MOS) subject classifications (1970). Primary 47B47, 18F25.

Key words and phrases. Compact operator, essentially normal operator, algebraic  $K$ -theory, extensions of  $C^*$ -algebras.

<sup>1</sup> Research partially supported by NSF GP 29006.

**2. Proof of the proposition.** Let  $I + K$  be invertible with  $K \in \mathfrak{K}$ . Write  $I + K = UP$  in polar decomposition. Then each of the operators  $U$  and  $P$  is of the form  $I + N$  where  $N$  is compact normal. We may assume that  $N$  has an infinite dimensional null-space  $\mathfrak{K}_0$ . So it suffices to show that any invertible operator of the form  $I + N$  with  $N$  compact normal with infinite dimensional null-space  $\mathfrak{K}_0$  is a product of (two) multiplicative commutators of the correct sort.

A. Brown and C. Pearcy show [1, Theorem 3] that  $I + N = (G_1, G_2)$  where  $G_1$  and  $G_2$  are invertible and  $G_1$  is a bilateral shift. An examination of their proof shows that  $G_2$  is also normal. Then the  $G_j$  live on  $\mathfrak{K}_0^\perp$ . Let  $H_1$  and  $H_2$  be commuting normals living on  $\mathfrak{K}_0$  such that  $\sigma(G_j) \subset \sigma_e(H_j) =$  an annulus (where  $\sigma_e$  denotes essential spectrum), and  $\sigma_e(H_1) \times \sigma_e(H_2) = \text{joint } \sigma_e\{H_1, H_2\} \equiv X$ . Then  $I + N = (G_1 \oplus H_1, G_2 \oplus H_2)$ . The operators  $G_1 \oplus H_1$  and  $G_2 \oplus H_2$  thus essentially commute.

Let  $\tau \in \text{Ext}(X)$  be the extension

$$0 \rightarrow \mathfrak{K} \rightarrow C^*\{I, \mathfrak{K}, G_1 \oplus H_1, G_2 \oplus H_2\} \rightarrow C(X) \rightarrow 0,$$

where  $C^*\{T_j\}$  denotes the  $C^*$ -algebra generated by  $\{T_j\}$  [4], [5]. We claim that  $\tau = 0$ ; the extension splits. The proof is as follows. The space  $X$  is homotopy equivalent to a torus, hence  $\text{Ext}(X) \cong \mathbf{Z} \oplus \mathbf{Z}$  via the index map [5]. A direct check shows that the index of  $\pi(G_j \oplus H_j)$  is zero for  $j = 1, 2$ , hence  $\tau = 0$ . (A more economical choice of  $H_j$  using the fact that  $G_1$  is unitary would yield  $X \subset R^3$  and allow avoidance of a homotopy argument.)

By the basic Brown-Douglas-Fillmore theorem [5], there exist commuting normals  $N_1, N_2$  and compact operators  $C_j$  such that

$$G_j \oplus H_j = N_j(I + C_j), \quad j = 1, 2.$$

Then

$$\begin{aligned} I + N &= (G_1 \oplus H_1, G_2 \oplus H_2) = (N_1(I + C_1), N_2(I + C_2)) \\ &= (B_1, A_1)(A_2, B_2) = (A_1, B_1)^{-1}(A_2, B_2) \end{aligned}$$

by direct computation, where

$$\begin{aligned} A_1 &= N_1 N_2 (I + C_2) N_1^{-1}, & A_2 &= N_1, \\ B_1 &= N_1 (I + C_1) N_1^{-1} = I + N_1 C_1 N_1^{-1} \in I + \mathfrak{K}, \\ B_2 &= N_2 (I + C_2) N_2^{-1} = I + N_2 C_2 N_2^{-1} \in I + \mathfrak{K}. \end{aligned}$$

This completes the proof.

**3. Remarks.**

**REMARK 1.** Our interest in  $K_1(\mathfrak{K})$  arose from the following considerations (inspired by Helton and Howe [6]; see also [2], [3]). Let  $\mathcal{A}$  be a  $*$ -subalgebra of  $\mathcal{L}$  containing the trace class  $\mathfrak{T}$  and suppose that  $\mathcal{A}/\mathfrak{T}$  is commutative. Let  $\mathfrak{K}_0 = \mathfrak{K} \cap \mathcal{A}$ . An invertible operator  $T$  in  $I + \mathfrak{K}_0$  represents zero in  $K_1(\mathfrak{K}_0)$  if  $T$  can be represented as a product of commutators  $(A_j, B_j)^{\pm 1}$  as in the

proposition, but with  $A_j, B_j \in \mathcal{Q}$ . If this is so, then  $\det T = 1$ . The same holds true if  $\mathcal{Q}$  is replaced by  $M_n(\mathcal{Q})$  provided  $T$  is also in the determinant class  $I + \mathfrak{T}$ .

Which hypotheses on  $\mathcal{Q}$  are really necessary for these conclusions to hold? Our proof shows that if  $K$  is compact normal then  $I + K$  is a product of commutators  $(A_j, B_j)^{\pm 1}$  where the  $A_j$  and  $B_j$  lie in a  $*$ -algebra which is commutative mod  $\mathfrak{K}$ . Thus the assumption that  $\mathcal{Q}$  be commutative mod  $\mathfrak{T}$  is necessary. A more interesting and difficult question is whether the hypothesis that  $\mathcal{Q}$  be closed under  $*$  can be eliminated. The special case where  $T$  is a single commutator is equivalent to the corresponding question for traces of additive commutators. In the case  $T \in \mathcal{Q}$ , rather than  $T \in M_n(\mathcal{Q})$ , this special case is equivalent to the general case.

REMARK 2. The inequality  $n \leq 8$  of the proposition can be improved to  $n \leq 6$  by means of a trick used by Radjavi:

$$\begin{pmatrix} S & 0 \\ 0 & T \end{pmatrix} = \begin{pmatrix} S & 0 \\ 0 & S^{-1} \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 \\ 0 & ST \end{pmatrix}.$$

Also

$$\begin{pmatrix} S & 0 \\ 0 & S^{-1} \end{pmatrix} = (A, B), \quad \text{where } A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} S^{-1} & 0 \\ 0 & 1 \end{pmatrix}.$$

REMARK 3. A more constructive proof of the proposition (which yields  $n \leq 8$ ) can be given by using an idea from [9]. At the cost of increasing  $n$  to 24, one can also require that both  $A_j$  and  $B_j$  be in  $I + \mathfrak{K}$ .

REMARK 4. The fact that  $K_0(\mathfrak{K}) = \mathbf{Z}$  has a number of generalizations. If  $\mathfrak{b}$  is any proper two sided ideal of  $\mathfrak{L}$ , then  $K_0(\mathfrak{b}) = \mathbf{Z}$ . Similarly,  $K_0(\mathfrak{c}) = \mathbf{Z}$  for a large class of dense  $*$ -subalgebras of  $\mathfrak{K}$ —in particular for  $\mathfrak{c} = \mathcal{Q} \cap \mathfrak{K}$  where  $\mathcal{Q}/\mathfrak{T}$  is commutative as before and  $\mathcal{Q}$  is maximal in a certain sense. The situation for  $K_1$  is quite different. Our result that  $K_1(\mathfrak{K}) = 0$  contrasts with the fact that  $K_1(\mathfrak{T}) \neq 0$  (by a determinant argument [2]). However, the methods alluded to in Remark 3 do apply to the Schatten classes  $\mathcal{C}_p$  for some  $p$ . The fact that  $K_1(\mathfrak{b})$  depends upon the ring in which  $\mathfrak{b}$  is an ideal complicates such considerations.

REFERENCES

1. A. Brown and C. Pearcy, *Multiplicative commutators of operators*, *Canad. J. Math.* **18** (1966), 737–749. MR **34** #608.
2. L. G. Brown, *The determinant invariant for operators with trace class self-commutators*, *Proc. Conf. on Operator Theory, Lecture Notes in Math.*, vol. 345, Springer-Verlag, New York, 1973.
3. ———, *Operator algebras and algebraic K-theory*, *Bull. Amer. Math. Soc.* **81** (1975), 1119–1121.
4. L. G. Brown, R. G. Douglas and P. A. Fillmore, *Unitary equivalence modulo the compact operators and extensions of  $C^*$ -algebras*, *Proc. Conf. on Operator Theory, Lecture Notes in Math.*, vol. 345, Springer-Verlag, New York, 1973.
5. ———, *Extensions of  $C^*$ -algebras, operators with compact self-commutators, and K-homology*, *Bull. Amer. Math. Soc.* **79** (1973), 973–978.
6. J. W. Helton and R. E. Howe, *Integral operators: commutators, traces, index and homology*, *Proc. Conf. on Operator Theory, Lecture Notes in Math.*, vol. 345, Springer-Verlag, New York, 1973.

7. J. Kaminker and C. Schochet, *Steenrod homology and operator algebras*, Bull. Amer. Math. Soc. **81** (1975), 431–434.

8. J. Milnor, *Introduction to algebraic K-theory*, Ann. of Math. Studies, no. 72, Princeton Univ. Press, Princeton, N. J., 1971.

9. C. Percy and D. Topping, *On commutators in ideals of compact operators*, Michigan Math. J. **18** (1971), 247–252. MR **44** #2077.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA 47907 (Current address of L. Brown)

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47401

*Current address:* (C. Schochet): Department of Mathematics, Wayne State University, Detroit, Michigan 48202